

Distinct patterns of cognitive outcome in young children with autism spectrum disorder receiving the Early Start Denver Model

Godel Michel*¹ • Robain François*¹, Kojovic Nada¹, Franchini Martina¹, Wood de Wilde Hilary¹, Schaer Marie¹

*Equal contribution

¹ Department of Psychiatry, University of Geneva School of Medicine, Geneva, Switzerland

Abstract

Evidence-based, early intervention significantly improves developmental outcome in young children with autism. Nonetheless, there is high interindividual heterogeneity in developmental trajectories during the therapy. It is currently not possible to predict a child's developmental outcome after receiving an early intervention based on his skills and symptoms at baseline. In this study, we analyze developmental trajectories of preschoolers with autism who received 2 years of intervention using the Early Start Denver Model (ESDM) approach in Geneva, Switzerland (n = 47). Our aim was to identify early predictors of response to treatment. We applied a cluster analysis to distinguish between 3 subgroups based on their cognitive level at intake, and rates of cognitive change over the course of treatment. The first subgroup of children only had a mild cognitive delay at intake, and nearly no cognitive delay by the end of treatment (Higher Cognitive at baseline: HC). The children in the two other subgroups all presented with severe cognitive delay at baseline, however, they had two very different patterns of response to treatment. The majority significantly improved developmental scores within their first year of intervention (Optimal Responders: OptR). Whereas a minority of children showed little to no improvement over the course of treatment (Minimal Responders: MinR). Our findings suggest that children who end up having an optimal two-year treatment outcome show rapid developmental improvement. Inversely, less significant progress by the sixth month of intervention may predict a less optimal response to treatment.

Keywords: Autism Spectrum Disorders, Early intervention, Predictors, Response to treatment, Heterogeneity, Minimal responder

1 Introduction

2

3 Autism spectrum disorder (ASD) is characterized by impairments in communication and
4 social interactions, along with restricted and repetitive behaviors (1). Over the last three
5 decades, several comprehensive, evidence-based early intensive behavioral intervention
6 (EIBI) approaches have been developed for young children with ASD, with the aim to
7 improve their social communication, cognitive functioning, and adaptive skills (2–7). As the
8 name suggests, the principles of EIBI comprise a significant number of hours (usually more
9 than 15 hours per week) as well as an early age of onset (usually younger than 4 or 5 years
10 old) (8). Systematic reviews and meta-analyses showed positive effects of EIBI on cognition,
11 adaptive skills and communication at the group level (9,10). Nevertheless, many EIBI studies
12 have reported a relatively heterogeneous response to these interventions, where most
13 children show significant improvements, while others make smaller gains (11,12). Despite
14 important efforts to better understand variables affecting treatment response, it is currently
15 not possible to predict to what extent a child will respond to intervention based on his or her
16 behavioral characteristics at intake (13). In the current therapeutic context and in the
17 absence of additional knowledge about individual predictors of outcome, many authors
18 suggest that intensive early intervention should be an intervention of choice for young
19 children diagnosed with ASD (14,15) regardless of their specific behavioral or symptom
20 profile. Yet, in the global framework of precision medicine (16), there is an urge to develop
21 more individualized guidelines for intervention in ASD. Given the importance of providing
22 effective programming for children with ASD as early as possible, and because of the costs
23 and parental investment associated with early intervention, it is crucial that we move away
24 from a “one size fits all” service provision model, and find ways to tailor a child’s
25 intervention to their specific needs, choosing therapy approaches based on the child’s
26 individual profile at diagnosis (17–19).

27

28 During the last decade, Early Start Denver Model (ESDM) has emerged as a promising EIBI
29 approach in the framework of the Naturalistic Developmental Behavioral Interventions
30 (NDBI, 7), notably because of its overall effectiveness, its emphasis on natural environment
31 teaching, comprehensive learning objectives and parental involvement. ESDM intervention
32 has originally been implemented in an individualized setting (one therapist for one child),
33 but other applications have been developed such as G-ESDM where one therapist works
34 with a little group of children and P-ESDM where parents/caregivers actually provide the
35 intervention under supervision. In their 2010 landmark randomized controlled trial (RCT, 2),
36 Rogers and Dawson reported a mean increase of 18 IQ points in a sample of 24 toddlers with
37 ASD receiving the ESDM intervention over two years. Subsequent studies have replicated
38 these results (for a review see 18) and demonstrated the cost-effectiveness of ESDM
39 intervention (18,19). Overall, the ESDM approach has been shown to significantly increase
40 cognitive, communication and adaptive skills at the group level (20). However, the inter-
41 individual variability in child response to treatment (RTT) is high, as with all types of EIBI (21).

42 To date, research about RTT in ESDM remains sparse and most studies focusing on
43 homogeneous and individualized therapy settings comprised limited sample size. To better
44 understand heterogeneity in treatment outcome, Vivanti et al. (22) attempted to identify
45 predictors of RTT in children receiving G-ESDM intervention. Their study showed that
46 developmental gains after one year of treatment were best predicted by higher imitation
47 skills, goal understanding and more advanced skills in the functional use of objects at
48 baseline. This study offered insight into how children with certain baseline competencies
49 might progress faster in a G-ESDM setting. However, outcomes were assessed after only one
50 year of intervention and baseline measures used in this study were based on original tasks
51 (i.e., specially developed for this study and not available in the common practice), making its
52 results poorly reproducible. In addition, its group setting makes its conclusion hardly
53 generalizable to the canonical individualized setting of ESDM. To the best of our knowledge
54 there are no identified child characteristics or symptom profiles based on standardized
55 clinical assessments that are known to predict child response to the individualized ESDM
56 intervention. It is therefore currently not possible to know whether a child will respond or
57 not to the ESDM intervention when advising it. Again, given the considerable investment
58 that intensive early intervention represents for therapists, families, and healthcare systems
59 (19,23), the ability to predict whether it will work for a child would drastically change how
60 early autism treatment programs are prescribed.

61

62 In the present study, we aim to identify early predictors of outcome after receiving a specific
63 and homogeneous EIBI. To do so we explored the developmental trajectories of 47
64 preschoolers with ASD who completed two years of individualized and intensive (20 hours
65 per week) ESDM intervention available in Geneva, Switzerland. We used a longitudinal single
66 group design without a control population, similar to previous studies in the field (21,22,24).
67 Indeed, because of ethical as well as logistic considerations, a random referencing to either
68 the ESDM intervention program or any other community treatment was not achievable. We
69 first investigated if our sample's outcome data, in terms of cognition, symptom severity and
70 adaptive functioning, reflected findings described in the ESDM literature. We then parsed
71 the heterogeneity in our sample's outcome by using cluster analysis (CA) and cognitive
72 scores as the main outcome measure. CA highlighted three different groups based on
73 cognitive outcome. We further explored baseline differences as well as early rates of change
74 between the three groups to identify potential predictors of two-year treatment outcome.

75

76 **Method**

77

78 **Participants**

79

80 Our original sample included 52 participants who completed two years of ESDM intervention
81 in Geneva, Switzerland. Four participants were not included in the analyses because of
82 missing data regarding their developmental assessment at baseline and one participant
83 because of missing data at the end of the intervention. Our analyses were thus based on the

84 data collected from 47 participants (see Table 1). No children were affected by either Fragile
85 X, Rett, Phelan-McDermid syndromes or neurofibromatosis. No participant suffered from
86 any major somatic disorders or epilepsy. All children were referred to the intervention
87 program after receiving a clinical diagnosis of ASD according to Diagnostic and Statistical
88 Manual of mental disorders, 5th edition (1) criteria and Autism Diagnosis Observation
89 Schedule-Generic (ADOS-G, 29) or 2nd edition (ADOS-2, 30) diagnosis cut-offs. Inclusion
90 criteria to the intervention program and research study included a confirmed diagnosis of
91 ASD, being able to participate for two full preschool years prior to age of school entry (which
92 in Switzerland is at 4 years old by July 31st), and parental consent. Enrollment also depended
93 on place availability at time of referral. The parents of all participants gave their written
94 informed consent to the research protocol that was approved by the institutional review
95 board of the University of Geneva. All participants were assessed in the context of the
96 ongoing longitudinal Geneva Autism Cohort study. Twenty-two children from this same
97 sample were already included in a previous study measuring outcome after one year of
98 ESDM intervention (27). Baseline evaluations were completed at the start of the
99 intervention and comprised behavioral measures that are detailed below. Parents also filled
100 out questionnaires regarding medical history, as well as demographic information detailed
101 below. Children were then assessed at 3 other time points at 6, 12 and 24 months of
102 therapy, for a total of 4 assessments. Post-intervention data about subsequent school
103 placement and support needs were collected. Children went onto either regular educational
104 classrooms with varied levels of in-class paraprofessional support or special education
105 classrooms.

106
107
108

107 **Intervention**

109 The 47 participants were enrolled in one of the 4 units of the *Centre d'Intervention Précoce*
110 *en Autisme* (CIPA) in Geneva, Switzerland (Fondation Pôle Autisme ([http://www.pole-](http://www.pole-autisme.ch)
111 [autisme.ch](http://www.pole-autisme.ch)) & Office Médico-Pédagogique), where they received 20 hours a week of daily,
112 individual intervention sessions using the Early Start Denver Model (ESDM). The ESDM is a
113 comprehensive, evidence-based early intervention approach that promotes child learning
114 through naturalistic developmental, and behavioral techniques (7,28). Parents of the
115 participants were provided with 12 hours of once-a-week parent coaching sessions in the
116 use of the ESDM model at the start of their child's program, and continued parent support
117 sessions as needed throughout the two-year period. The children were evaluated every 3
118 months using the Early Start Denver Model Curriculum Checklist for Young Children with
119 Autism (ESDM-CC) to establish targeted and measurable learning objectives. The
120 intervention services were provided by graduate-level therapists, who were trained within
121 the CIPA program in the use of the ESDM approach, meeting ESDM fidelity on the ESDM
122 Fidelity Rating System (28). Today, the team consists of 20 credentialed ESDM therapists,
123 and the program is overseen by an ESDM certified trainer.

124
125

125 **Measures**

126

127 The ADOS (which refers to the ADOS-G and its later version, the ADOS-2), is a standardized
128 assessment which comprises a series of semi-structured social presses aimed to elicit and
129 measure ASD symptoms (25,26). The schedule comprises 5 different modules, adapted to
130 the person's age and level of language. The calibrated severity score (CSS) was used to
131 compare the total severity score as well as the restricted and repetitive behaviors (RRB) and
132 social affect (SA) symptoms severity scores (29,30). The ADOS were administered by a
133 trained examiner and filmed. The members of the team who rated the video recordings
134 were not implicated in the delivery of the ESDM intervention.

135

136 The Mullen Scales of Early Learning (MSEL) is a standardized assessment for children aged
137 from birth to 68 months (31). It measures the child's development in five developmental
138 domains: expressive language (EL), receptive language (RL), visual reception (VR), fine motor
139 (FM), as well as gross motor skills (GM).

140

141 The Psychoeducational Profile – third edition (PEP-3) is a standardized assessment tool that
142 evaluates cognitive, motor, and adaptive domains in children 2 to 7 years of age (32). These
143 domains include EL, RL, FM and cognitive verbal and preverbal (CVP). The PEP-3 as well as
144 the MSEL were administered by psychologists following the standard instructions of both
145 evaluations.

146

147 Developmental quotient scores (DQ) were computed for each subdomain of the MSEL by
148 dividing the individual developmental age by the chronological age and multiplying by 100 as
149 described in 2006 by Lord and colleagues (33). The composite DQ was computed by
150 calculating the average of all four subdomains' developmental ages, then dividing by the
151 chronological age and multiplying by 100. Similarly, DQ scores were computed for the
152 subdomains of the PEP-3 that assess domains equivalent to those of the MSEL, namely EL,
153 RL, CVP and FM. The PEP-3 composite DQ was derived using the same formula as described
154 for the MSEL, and has already been used for the PEP-3 subdomains in previous studies (27).
155 For our analyses of cognitive skills, we used the MSEL Early Learning Composite DQ. Since
156 the MSEL was not administered for some participants (n = 7 at baseline, n = 7 after 6 months
157 of therapy, n = 3 after 12 months of therapy, n = 2 after 24 months of therapy) we replaced
158 the missing DQ scores by their equivalent DQ scores derived from the PEP-3. It is important
159 to keep in mind that DQ is normalized for the age at the time of evaluation. Hence, a DQ that
160 remains stable over time does not reflect stagnation but rather continued developmental
161 progress. Also, a loss of DQ over time does not necessarily imply regression (a loss of skills)
162 but rather slower skill acquisition, leading to a widening of the gap between the child's
163 current abilities and what would be expected in typical development.

164

165 The Vineland Adaptive Behavior Scales – 2nd edition (VABS-II) is a semi-structured interview
166 administered by a trained clinician that assesses a person's adaptive behavior (34). The

167 domains assessed comprise communication, socialization, daily living skills (DLS) and motor
168 skills. An overall adaptive behavior composite score (ABC) of all these 4 domains is
169 computed.

170

171 The ADOS, VABS-II, PEP-3 and MSEL were administered at baseline, after 12 months and
172 after 24 months of therapy. Assessment at six months only comprised the VABS-II and the
173 MSEL.

174

175 We measured participant socio-economic using the total household yearly income and the
176 highest level of education achieved by parents at baseline. The household income was
177 divided into three subgroups that are detailed in Table 1. Parental educational level was first
178 coded using the seven categories of the four-factor index of social status developed by
179 Hollingshead (35). We then divided these categories into two groups: 1) elementary school
180 or high school completed, and 2) college and/or graduate degree completed.

181

182 **Rate of change**

183

184 For all behavioral measures acquired longitudinally (ADOS, VABS-II and DQ), we computed
185 an individual rate of change using the following Symmetrized Percent Change (SPC) formula:

186

$$SPC [\%/year] = 100 \times \frac{(B_y - B_x) / [(B_x + B_y) / 2]}{(age_y - age_x)}$$

187

188 Where B_x and B_y represent the behavioral measure acquired when the participant was aged
189 of age_x and age_y respectively. In other words, SPC is the behavioral difference between
190 two timepoints relatively to the mean of the scores across these two timepoints, then
191 divided by the time interval (in years). This results in a yearly rate of change that can be
192 expressed as a percentage when multiplied by 100. The main advantages of using
193 symmetrized measures of change over absolute differences (such as $B_y - B_x$) or non-
194 symmetrized percentages (such as $(B_y - B_x) / B_x$) comprise increased statistical
195 robustness, higher reliability in small samples, limited sensitivity to outliers and equivalent
196 consideration of both B_x and B_y measures (36).

197

198 **Statistical Analyses**

199

200 IBM®SPSS® Statistics v26.0.0.0 for macOS (Armonk, NY: IBM Corp.) was used for all analyses.
201 Statistical significance threshold was set at $\alpha = 0.05$. Graphs were obtained with Prism®
202 v8.3.0 (GraphPad Software, La Jolla California USA, www.graphpad.com) and Matlab R2018b
203 for macOS (MathWorks).

204

205 To test for an effect of time a repeated measure ANOVA was performed on the whole
206 sample for each longitudinal behavioral measure using the scores collected at baseline, 12

207 months, and 24 months after the start of the intervention services. Greenhouse-Geisser
208 correction was applied whenever the assumption of sphericity was violated according to
209 Mauchly test.

210

211 Until now, methodological strategies to identify intervention-specific predictors of EIBI
212 outcome include whole-sample correlations between baseline and outcome measures
213 (22,37), comparison between subgroups defined based on an arbitrary cut-off such as rapid
214 vs slow learners (38) or best vs non-best outcome (39,40). A promising alternative relies on
215 the identification of distinct phenotypic subgroups within ASD (41). Defining more
216 homogeneous subgroups based on behavioral characteristics in a data-driven manner can be
217 achieved by applying cluster analyses (CA), a strategy that has already been used in ASD
218 preschool studies (for a review see 27). To date, CA has only been applied once on children
219 with ASD participating in an EIBI program (Applied Behavioral Analysis: ABA) with a special
220 focus on language development (43). We here performed a cluster analysis (CA) using
221 cognition (assessed with the composite DQ measure) as our main outcome measure. There
222 are several reasons why we chose DQ over other parameters. First, it is generally the main
223 outcome measure reported in early intervention studies, and it displays the most variability
224 (2,44,45). Second, cognition has been shown to be the domain that improves the most after
225 early intervention (46). Third, studies investigating possible ASD subtypes within ASD have
226 shown that the most salient group differences emerge when categorized by cognitive skills
227 (47). We used a *k*-means clustering approach to identify subgroups in terms of DQ
228 trajectories with a maximal number of iterations set to 10 (48). We chose two variables that
229 capture individual DQ trajectories: the composite DQ at baseline and the composite DQ SPC
230 over the two-year intervention period. To objectively determine the number of clusters *k* we
231 used a two-step clustering approach as suggested by Kodinariya (49). We used the two-step
232 clustering algorithm developed by Chiu et al. (50) as it is implemented in IBM®SPSS®
233 Statistics. Briefly, this method firstly divides the sample into a set of sub-clusters through a
234 sequential approach and secondly merges the sub-clusters through a hierarchical technique
235 based on the log-likelihood distance between them. Finally, the Akaike's information
236 criterion is used to objectively determine the optimal number of clusters.

237

238 The cluster analysis (CA) yielded 3 optimal clusters based on the baseline composite DQ and
239 the composite DQ SPC over 2 years (Fig 1). The ANOVA revealed that one of these clusters
240 exhibited significantly greater composite DQ at baseline compared to the others and was
241 therefore named "higher cognitive at baseline" (HC, *n* = 18). Its average DQ at baseline was
242 78.8 ± 10.9 with a range between 64 and 108 with a SPC of 10.2 ± 5.8 %/yr. This corresponds
243 to an average 19.2 gain for a final DQ of 98.0 ± 14.7 with a range between 64 and 124. The
244 second "optimal responders" cluster (OptR, *n* = 20) was characterized by high rates of
245 progress within the two-year program. DQ at baseline was 50.0 ± 10.8 with a range between
246 22 and 67, and its average SPC was 23.8 ± 8.4 %/year. This corresponds to an average 33.9
247 gain for a final DQ of 83.9 ± 21.8 with a range between 33 and 130. The third "minimal

248 responders” cluster (MinR, $n = 9$) was characterized by decreased rates of progress
249 compared to the two other clusters with an average SPC of -10.3 ± 11.3 %/yr. Its composite
250 DQ at baseline was 46.0 ± 8.3 with a range between 32 and 60. The average loss was 8.5 for
251 a final DQ of 37.5 ± 8.9 with a range between 27 and 58. The OptR and MinR subgroups did
252 not differ in composite DQ at baseline, with an average of 50.0 and 46.0 respectively.
253 Together, they form a group of children with lower cognitive scores (LC) at baseline. Cluster
254 differences over composite DQ at baseline and composite DQ SPC are illustrated on Fig 1.
255 Detailed analyses are reported in supplementary Table 1 and 2.

256

257 **Figure 1: A.** Composite DQ trajectory of the total sample over the two years of intervention.
258 Significant results of repeated measure ANOVA are displayed. **B.** Composite DQ trajectories
259 of the three subgroups parsed by the cluster analysis. **C.** Individual values of the two
260 measures used in the clustering analysis algorithm (composite DQ at baseline and DQ SPC
261 over the two years of interventions). Color code represents the cluster membership of each
262 participant after the application of the cluster analysis. **D.** Differences between the three
263 subgroups on the two measures that were used to parse them. *** $p < .001$. DQ:
264 Developmental Quotient. HC: Higher Cognitive. MinR: Minimal Responders. OptR: Optimal
265 Responders. SPC: Symmetrized Percent Change.

266

267 Demographic, socio-economic measures and behavioral measures at baseline were
268 compared between clusters using one-way analysis of variance (ANOVA) or chi-square test.
269 We used a Bonferroni correction for multiple testing on the subdomains of the same clinical
270 evaluation (e.g., the subdomains of the VABS-II), setting the statistical significance at
271 $0.05/\text{number of subdomains}$. When an ANOVA reached statistical significance, post-hoc
272 comparisons between clusters were performed using multiple T-tests with Bonferroni
273 correction and statistical significance set at $0.05/\text{number of clusters}$.

274

275 We then applied the same strategy to compare the SPC between clusters. We performed
276 analyses on the following SPC: from baseline to 6 months, from baseline to 12 months, from
277 baseline to 24 months of therapy.

278

279 Finally, we focused on the two LC clusters which showed no differences in their composite
280 DQ at baseline to explore whether any other behavioral measure could help classifying
281 them. To do so, we used binary logistic regression models. More specifically, we selected all
282 behavioral measures that differed between OptR and MinR on post-hoc T-tests at baseline.
283 Then, we performed a multivariate logistic regression using the selected measures.
284 Whenever a composite score and one or more subdomain scores of the same test were
285 selected, we preferred the composite measure to minimize potential collinearity between
286 variables in the model. Then, we used the same strategy for the SPC measures during the 6
287 first months of intervention, and ultimately with those of the 12 first months.

288

289 **Results**

290

291 **Whole sample trajectories**

292

293 Descriptive measures collected at each visit are reported in Table 1 for the total sample. The
294 children were aged from 15.3 to 42.0 months at the beginning of the intervention (average:
295 28.9 ± 5.3 months). The average composite DQ of the entire group at baseline was $60.3 \pm$
296 18.0 (range: 22 - 108). As a group, all 47 children receiving ESDM showed a significant
297 decrease in their total level of symptom severity (ADOS CSS) (see Table 1). This improvement
298 was driven by a decrease in the Social Affect (SA) domain. On the contrary, the RRB
299 symptom severity increased over time. We found that these changes occurred mainly during
300 the first year of intervention and that CSS (both RRB and SA) were stable during the second
301 year of intervention. In parallel, participants' developmental scores improved. This
302 improvement was significant in all subdomains (i.e., VR, RL and EL) except for fine motor
303 (FM). As for the measures of symptom severity, all changes in cognition were significant
304 during the first year of therapy but not the second one. Finally, there was no significant
305 change regarding the adaptive behavior composite (ABC) scores over time. Nevertheless,
306 participants made significant gains in the communication subdomain which occurred both
307 during the first and the second year of intervention. Participants also increased their delay in
308 the motor skills subdomain. This effect on motor skills was only significant during the first
309 year of intervention and its effect size was the smallest (see Table 1) of all the mentioned
310 significant behavioral changes. All statistically significant results are detailed in Table 1.
311 Concerning the type of schooling after the intervention, 29 participants (61.7%) joined a
312 public regular education classroom with individual educational support. One participant
313 (2.1%) joined a regular education classroom without any support. Four children (8.5%)
314 entered a private school that provided in-class support in a regular education classroom.
315 Finally, 13 participants (27.7%) entered special education program within the public-school
316 system.

317

318

319 **Table 1** Sample characteristics over the two years of ESDM intervention. Scores at 6 months
320 (in italic) are indicative and were not used in the repeated measure (R.M.) ANOVA. Scores
321 with a significant difference are highlighted in bold; * $p < .05$ ** $p < .01$ *** $p < .001$. G:
322 *Greenhouse-Geisser correction applied, ESDM = Early Start Denver Model*

MEASURE	At Baseline	+ 6 months	+ 12 months	+24 months	Pval (R. M. ANOVA)	Partial Eta Squared	0 - 24mo	0 - 12mo	12m 24m
Clinical Description									
ADOS CSS total [Mean (SD)]	8.0 (1.9)		6.5 (2.0)	7.0 (1.9)	<0.001***	.241	.011*	<0.001***	.05
ADOS CSS SA	7.7 (2.0)		5.32 (1.8)	5.8 (1.9)	<0.001*** (G)	.365	<0.001***	<0.001***	.26
ADOS CSS RRB	7.9 (2.1)		9.0 (1.4)	9.0 (1.3)	<0.001*** (G)	.199	.003**	.001**	1.0
ADI-R subdomains [Mean (SD)] (n = 43)									
ADI-R Social interactions	13.7 (4.9)								
ADI-R RRB	4.0 (2.3)								
VABS-II Adaptative Behavior Composite [Mean (SD)]	79.8 (9.1)	<i>81.0 (12.2)</i>	80.1 (12.6)	82.6 (15.5)	.102 (G)				
VABS-II socialization	81.1 (9.2)	<i>81.0 (10.5)</i>	78.6 (10.5)	79.5 (12.9)	.201 (G)				
VABS-II communication	75.2 (12.1)	<i>80.2 (15.4)</i>	83.7 (18.5)	87.7 (18.5)	<0.001*** (G)	.359	<0.001***	<0.001***	.06
VABS-II daily living skills	84.0 (10.8)	<i>84.9 (11.4)</i>	82.7 (12.4)	83.1 (16.5)	.569				
VABS-II motor skills	90.5 (9.6)	<i>89.4 (11.1)</i>	85.1 (10.3)	88.3 (14.7)	.031* (G)	.092	1.0	.002**	
Composite DQ [Mean (SD)]	60.3 (18.0)	<i>72.6 (21.0)</i>	77.8 (25.5)	80.4 (27.9)	<0.001*** (G)	.480	<0.001***	<0.001***	
Fine Motricity DQ	75.0 (16.9)	<i>76.4 (16.8)</i>	76.7 (19.1)	81.9 (25.3)	.058 (G)				
Visual Reception DQ	75.3 (24.1)	<i>86.7 (23.8)</i>	89.3 (31.9)	91.2 (31.8)	<0.001***	.235	<0.001***	<0.001***	
Expressive Language DQ	43.6 (19.6)	<i>57.9 (26.3)</i>	66.9 (29.6)	68.8 (27.5)	<0.001*** (G)	.449	<0.001***	<0.001***	
Receptive Language DQ	47.0 (25.8)	<i>69.3 (28.7)</i>	78.1 (31.4)	79.6 (33.4)	<0.001*** (G)	0.591	<0.001***	<0.001***	
Demographics									
Chronological age [months Mean (SD)]	28.9 (5.3)								
Gender [females Number (percentage)]	6 (12.8%)								
Parental Education [Number (percentage)] (n = 46)									
Elementary School or High School	18 (39.1%)								
University or PhD	26 (60.9%)								
Household Income [Number (percentage)] (n = 45)									
<60k	12 (26.7%)								
60-140k	17 (37.8%)								
>140k	16 (35.6%)								

medRxiv preprint doi: <https://doi.org/10.1101/2021.04.05.21254908>; this version posted October 4, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. All rights reserved. No reuse allowed without permission.

324 Parsing the heterogeneity in treatment response

325

326 Difference between the three subgroups at baseline

327

328 We found no differences between clusters for parental educational attainment or household
329 income (see Supplementary Table 1). There were also no differences regarding age at
330 baseline. When looking at DQ at baseline, we found that HC showed higher scores in all DQ
331 subdomains compared to both other clusters. Considering adaptive behavior, HC exhibited
332 higher scores in ABC as well as in the communication subdomain compared to both other
333 clusters. HC also showed a higher score in socialization compared to MinR. All statistically
334 significant results of analyses on the DQ and the VABS-II across the three subgroups at
335 baseline are illustrated on Fig 2. There was no difference in the total ADOS CSS. In the ADOS
336 subdomains, we found that HC exhibited lower RRB compared to MinR. Besides, there was
337 no difference in any clinical measure between MinR and OptR.

338

339 **Figure 2: A.** Statistically significant differences in DQ subdomains between subgroups at
340 baseline. **B.** Statistically significant differences in VABS-II ABC and VABS-II subdomains
341 between subgroups at baseline. * $p < .05$. ** $p < .01$. *** $p < .001$. ABC: Adaptive Behavior
342 Composite. DQ: Developmental Quotient. EL: Expressive Language. FM: Fine Motor. HC:
343 Higher Cognitive. MinR: Minimal Responders. OptR: Optimal Responders. RL: Receptive
344 Language. VABS-II: Vineland Adaptive Behavior Scale. VR: Visual Reception.

345

346 Differences between subgroups in rates of change over 6, 12 and 24 months of intervention

347

348 We found that over the two years of therapy, OptR exhibited higher rates of change
349 compared to the other two subgroups in cognition (composite DQ as well as VR, FM and RL
350 subdomains) as well as in adaptive behavior compared to MinR (VABS-II ABC, in socialization,
351 communication and DLS subdomains) (see Supplementary table 2). Differences in rates of
352 change for all these domains except for socialization, were already present within the 12
353 first months of therapy. Also, we found that MinR exhibited slower rates of change during
354 the total time of intervention compared to both other subgroups in cognition (composite
355 DQ, VR, FM and EL) as well as in adaptive behavior (VABS-II ABC, socialization,
356 communication and DLS). These differences in the rates of change were already present
357 during the first 12 months of therapy, except for socialization, VR and EL.

358

359 Finally, we found that OptR already exhibited faster rates of change in composite DQ, and
360 adaptive communication compared to MinR (Fig 3) after only 6 months of intervention.
361 During the same early period, HC exhibited higher SPC in receptive language (RL) DQ
362 compared to OptR.

363

364 We did not find any difference in the rates of change of symptom severity (total ADOS, SA
365 and RRB) between the three subgroups during the time of intervention.

366

367 **Figure 3: A.** Statistically significant differences between subgroups in the rates of change of
368 behavioral measures (DQ and VABS-II) within the first 6 months of intervention. **B.**
369 Statistically significant differences between subgroups in the rates of change of behavioral
370 measures (DQ and VABS-II) within the first 12 months of intervention. * $p < .05$. ** $p < .01$.
371 *** $p < .001$. ABC: Adaptive Behavior Composite. DLS: Daily Living Skills. DQ: Developmental
372 Quotient. EL: Expressive Language. FM: Fine Motor. HC: Higher Cognitive. MinR: Minimal
373 Responders. OptR: Optimal Responders. RL: Receptive Language. SPC: Symmetrized Percent
374 Change. VABS-II: Vineland Adaptive Behavior Scale. VR: Visual Reception.

375

376 Subgroup classification based on early rates of change

377

378 Minimal responders (MinR) and optimal responders (OptR) showed no difference on the
379 composite DQ at baseline and were both considered to have lower cognitive scores at
380 baseline (LC). They were thus selected for our classification analyses to address the potential
381 of clinical measures at baseline as well as their early rates of progress to classify them. Since
382 we found no differences between the two subgroups in any of the clinical measures at
383 baseline, no logistic regression was performed on this time point. Within the first 6 months
384 of therapy, MinR showed slower SPC in the VABS-II communication subdomain and in the
385 composite DQ. Logistic regression based on these two variables allowed a partition of MinR
386 and OptR with a 95.5 % overall correct classification rate (see Fig 4A). All the 17 OptR
387 included in the model and 4 out of the 5 MinR included were classified correctly. The logistic
388 regression model was statistically significant ($\chi^2 = 10.501$, $p = .005$) and explained 38% of the
389 variance (Nagelke R^2). The prediction equation was the following: $O = -.046 * DQ\ SPC - .065$
390 $VABS-II\ communication\ SPC + .561$. Within the first 12 months of therapy, OptR exhibited
391 higher SPC in both the VABS-II ABC and the composite DQ. Logistic regression performed
392 with both measures reached a 96.4 % rate of overall correct classification between OptR and
393 MinR (see Fig 4B). All the 19 OptR included in the model and 8 out of the 9 MinR included
394 were classified correctly. The logistic regression model was statistically significant ($\chi^2 =$
395 27.274 , $p < .001$) and explained 87% of the variance (Nagelke R^2). The prediction equation
396 was the following: $O = -.198 * DQ\ SPC - .222\ VABS-II\ ABC\ SPC + 1.838$. In other words, it
397 would have already been possible for a clinician to classify a child as being an OptR or MinR
398 with more than 95% of accuracy after six months of intervention based on the child's rates
399 of change in the VABS-II communication and composite DQ, and after 12 months of
400 intervention based on their rates of change in VABS-II ABC and composite DQ.

401

402 **Figure 4: A.** Logistic regression between MinR and OptR using their rates of change within
403 the 6 first months of intervention in Composite DQ and VABS-II communication. Color code
404 corresponds to subgroup membership (red: MinR, green: OptR). Decision boundary is
405 represented in a dotted line. **B.** Logistic regression between MinR and OptR using their rates
406 of change within the 12 first months of intervention in Composite DQ and VABS ABC. ABC:
407 Adaptive Behavior Composite. DQ: Developmental Quotient. SPC: Symmetrized Percent
408 Change. VABS-II: Vineland Adaptive Behavior Scale.

409

410 **Discussion**

411

412 In the present study, we analyzed data from one of the largest samples of children who
413 underwent two years of intensive (20h per week) and individualized ESDM intervention to
414 identify predictors of their developmental outcome. Overall, we observed that, preschoolers
415 in our sample made significant cognitive progress and adaptive skill gains over the two years
416 of intervention (see Fig 1A). Improvements in the current sample allowed 72.3% of the
417 children to enter a regular education classroom post-intervention, which in Geneva requires
418 the child to have near peer-level functioning. These results are consistent with those
419 reported in other studies on ESDM-based intervention (2,51). More specifically, our sample
420 exhibited an average change in DQ (+20 points) that is very close to the one described in the
421 randomized controlled trial (RCT) study by Dawson et al. (2), which reported 18 points of
422 cognitive gain, an average significantly greater than that of their control group. In parallel, a
423 naturalistic study that explored developmental trajectories in preschoolers with ASD who
424 were not enrolled in any specific therapeutic program reported an average DQ gain of only
425 6.3 points between 24 and 48 months of age (52). The average initial DQ in the cited study
426 (63.6 ± 11.5) was similar to ours (60.3 ± 18.0). Considering similarities in the outcome
427 between our results and previous ESDM studies as well as differences with naturalistic
428 studies, one can infer that ESDM intervention in our study had a causal effect on the
429 improvements observed at the whole group level. These results therefore highlight the
430 possibility of implementing ESDM in Europe as effectively as in the US, despite differences in
431 culture and health care system. They also support the cost-effectiveness of ESDM
432 intervention, with improvements in cognition and adaptive behavior known to reduce
433 subsequent school-based support needs, offsetting costs associated with early intensive
434 intervention (17–19).

435

436 This study also aimed to determine whether preschoolers with ASD who participated in a
437 two-year EIBI intervention program (here ESDM in an individual setting) could be separated
438 into distinct subgroups based on their cognitive trajectories over time. To achieve this, we
439 used a *k*-means cluster analysis (CA) approach with cognitive abilities at baseline and
440 cognitive rates of change over time as variables. CA yielded three groups: ~38% ($n = 18$) of
441 children with a mild cognitive delay at baseline that displayed a globally good outcome
442 (Higher cognitive at baseline: HC), and two groups of children constituting the lower
443 cognitive scores at baseline group (LC) that had very different outcomes. The first group of
444 LC, which represented ~43% ($n = 20$) of the entire sample, underwent significant cognitive
445 and adaptive skill improvements (Optimal responders: OptR) while the second group of LC,
446 which represented ~19% ($n = 9$) of our sample, showed slower overall progress, and saw a
447 widening of the developmental gap over time in cognition and adaptive behavior compared
448 to same aged peers (Minimal responders: MinR). The clear distinction between toddlers with
449 mild cognitive delay (HC) and those with the more severe cognitive delay at baseline (LC)
450 observed in our sample is also reported in previous studies that applied CA to preschoolers
451 with ASD (53–55). These studies identified at least two subgroups categorized as “high” and

452 “low-functioning” based on early cross-sectional cognitive measures. One of the main
453 differences in the present study is that we included a longitudinal variable in our CA (i.e., the
454 rate of cognitive change) and were therefore able to further define our subgroups of LC
455 children based on individual cognitive trajectories over time that a cross-sectional CA would
456 have failed to capture. In our second analysis, we aimed to uncover potential predictors of
457 outcome by evaluating how we can predict a child’s cluster membership. Amongst the LC
458 subgroups (MinR and OptR), we found neither behavioral nor demographic differences at
459 baseline (see Supplementary Table 1). However, we noted a significant difference between
460 their rates of change in cognition and communication skills within the first 6 months of
461 intervention. Using a logistic regression model, we showed that these early rates of change
462 predicted at 95% attrition to either the MinR or OptR subgroups. This means that despite the
463 absence of differences at baseline that could predict outcome in LC children, early, rapid
464 developmental progress by 6 months of intervention allowed an accurate classification of
465 subsequent developmental pattern.

466
467 Our analyses of the HC subgroup suggest that a mild cognitive delay (78.8 ± 10.9 of
468 composite DQ) at the start of an ESDM intervention is associated with an alleviation of the
469 delay in cognitive skills (+10% DQ per year) and adaptive behavior (+4.1% ABC per year) over
470 the course of treatment. In addition, children in the HC group exhibited higher levels of
471 adaptive skills compared to other subgroups (84.9 ± 9.2 of ABC) at baseline, especially in the
472 VABS-II domain of communication (85.7 ± 11.0). All HC children except for one were able to
473 continue into a regular education classroom following the intervention. With a DQ of 64 at
474 both the beginning and the end of the intervention, this child was the only one in the HC
475 group with a DQ value lower than 80 at the end of the intervention. Overall, our HC
476 subgroup results suggest a positive outcome in preschoolers with a mild delay in cognition
477 and communication at baseline after receiving an individualized and intensive ESDM
478 intervention. A recent review concluded that a higher cognitive level at baseline is a good
479 predictor of positive outcome after various types of EIBI (56). Also, previous studies focusing
480 on another type of intervention (Applied Behavioral Analysis: ABA) reported that higher
481 abilities in adaptive behavior (38,57) as well as in language (58) constitute predictors of good
482 outcome. This might suggest that mild delays in cognition and communication could
483 represent a common predictor of good outcome among various EIBI approaches. These
484 findings will need to be further explored with future RCT that assess the specific causality of
485 ESDM intervention within these results. A practical implication of our findings concerning
486 the HC subgroup is that clinicians who refer a toddler with a mild developmental delay at
487 baseline to an ESDM program can be relatively confident that there will be a good outcome
488 in cognition and adaptive behavior by the end of the intervention.

489
490 Apart from the HC subgroup, the rest of the sample included children presenting a severe
491 cognitive delay at baseline (Lower Cognitive, or LC). These children presented drastically
492 different cognitive trajectories of change over time and were attributed to two distinct

493 subgroups: OptR and MinR. Despite their severe cognitive delay at baseline (average DQ of
494 50.0 ± 10.8), the 15 children that composed the OptR subgroup greatly improved their
495 cognitive and adaptive skills over time and 75% of them were able to join a regular
496 education classroom with in-class support. On the other hand, the 9 children in the MinR
497 subgroup had a similar level of cognitive impairment at baseline (average DQ of 46.0 ± 8.3),
498 however their cognitive and adaptive functioning scores did not improve over time.
499 Furthermore, the developmental gap between continued to widen, despite receiving
500 intensive early intervention. Only 2 out of 9 (22.2%) MinR children joined a regular
501 education classroom following the intervention. Here, we did not identify any differences
502 (i.e., in cognition, adaptive behavior or autism severity) that could accurately discriminate
503 between the OptR and MinR subgroups at baseline. There is currently no clear clinical
504 marker within LC toddlers that could help clinicians predict how the child will respond to
505 ESDM intervention. One clinical implication of our analyses of OptR and MinR at baseline is
506 that the OptR constituted most of the LC children (~69%) thus supporting the *a priori* that
507 most toddlers who present with lower cognitive scores at intake display a positive outcome
508 after receiving an individual intensive ESDM intervention. Nonetheless, a better
509 understanding of the factors (behavioral, biological, and environmental) that are associated
510 with MinR remains necessary to develop more targeted clinical recommendations. For
511 instance, future studies including more participants with comorbid conditions such as
512 epilepsy should investigate whether they represent moderator of outcome. Furthermore,
513 they could determine whether the additive effect of various genetic mutations may
514 moderate the outcome (59). The observation of two distinct trajectories of change in
515 children with larger cognitive impairments at baseline could shed new light on the
516 inconsistencies that exist between various studies that measured cognitive response to EIBI
517 within LC preschoolers with ASD. For instance, one previous study concluded that children
518 with this kind of profile only improve in fine motor skills and receptive language but not in
519 adaptive behavior after receiving an early and intensive ABA intervention (60). Other studies
520 focusing in ESDM reported an association between lower cognitive level at baseline and high
521 cognitive gains (61,62). One can hypothesize that the inter-individual heterogeneity of
522 outcome reported by previous studies (60–62), as well as the differences in their results
523 were potentially due to the existence of two latent subgroups (MinR and OptR) that may
524 have driven results in opposite directions. Our results thus advocate for a more systematic
525 subgroup phenotyping, including longitudinal variables, in future studies focusing on the
526 clinical outcome of EIBI to better describe the phenotypic heterogeneity within LC
527 preschoolers with ASD.

528

529 Finally, our results suggest that despite the lack of significant differences at baseline
530 between OptR and MinR children, the outcome after two years of intervention can be
531 predicted by the end of the first year of intervention with high accuracy. The OptR group's
532 rates of change appear to be significantly higher than those observed in the MinR group
533 within the first year for cognitive and adaptive skills (especially in communication and daily

534 living skills). Our results are in line with those of Sallows et al. (38), who reported cognitive
535 gain during the first year of an early and intensive ABA intervention as one of the best
536 predictor of outcome at the end of the intervention. Furthermore, we found that based on
537 the rates of change in cognition and communication within the first six months of
538 intervention, we could infer the outcome after two years of intervention. Together, these
539 conclusions might shed light on the timing of RTT, and when children can be considered as
540 “non responders” as raised by Vivanti et al. (13). Indeed, our analyses suggest that the first 6
541 months to a year of intervention offers critical information about how a child will respond to
542 ESDM intervention over time, and leads us to question whether an early, clear response to
543 intervention can predict an optimal response overall. The emphasis on this early response to
544 intervention as a predictor of long-term outcome has several clinical implications. One of
545 them would be the importance of implementing regular, standardized follow-ups to
546 measure children’s cognition and adaptive behavior in the first 6 and 12 months, in addition
547 to the systematic ESDM *Curriculum Checklist* that is currently used in the model. An
548 alternative could lie in the development of a standardized way to use the ESDM-CC to track
549 the rate of developmental change and ultimately the post-intervention outcome. This type
550 of early standardized follow-up could potentially alert the clinician of difficulties a MinR child
551 might face and give the child’s team more time to consider alternative supports or enhanced
552 service provision long before the end of the intervention. More research is needed to
553 understand what supports or program enhancements would allow a child with a slower
554 response to intervention to have a more optimal outcome.

555
556 In summary, our results show that despite the lack of reliable outcome predictors for
557 children with ASD who present severe cognitive delays at baseline, the consideration of their
558 early dynamic behavioral parameters may help predict their overall response to
559 intervention. Further RCTs that explore the trajectories of subgroups similar to ours are
560 needed to determine the precise effect of the ESDM on children with MinR and OptR
561 profiles. More specifically, we need to understand whether ESDM helps OptR improve their
562 outcome or if it prevents MinR from falling even further behind developmentally, or both.
563 Another hypothesis to be addressed is whether ESDM has an influence in the relative
564 number of participants that are affected to each subgroup – i.e., whether some OptR
565 participants would have been MinR if they had not undergone an ESDM based therapy.
566 Future research on the specific effects of ESDM on each subgroup could result in improved
567 therapeutic guidelines that are more tailored to each child’s individual developmental
568 trajectory. Our study provides relevant variables that should be explored by future research
569 at the beginning and during the very first months of an ESDM intervention.

570

571 **Limitations**

572

573 Despite being one of the largest samples of preschoolers who benefited from a two-year
574 intensive and individualized ESDM program, the sample size of the present study limits the
575 number as well as the size of subgroups that can be detected by a cluster analysis.

576 Nevertheless, we took care to respect the commonly accepted prerequisites of cluster
577 analyses, including the minimum sample size in each group or the number of factors in the
578 analyses given the overall sample size (63,64). It is possible that studies performed on larger
579 samples could achieve more fine-grained subgrouping on a similar population based on the
580 same measures and could lead to bigger subgroups, in turn increasing the statistical power
581 to detect differences at baseline between lower cognitive clusters that we could not
582 highlight.

583

584 Another limitation that is a direct consequence of the previous one lies in the choice of the
585 main outcome. We chose parameters related to cognitive skills as the main clustering
586 factors. However, it would have been possible to use other measures such as level of ASD
587 symptoms, adaptive skills or even a combination of these two. The inclusion of more
588 variables in the model could help in defining a larger number of clusters and therefore
589 increase our understanding of the heterogeneity of ASD in a refined manner. However, this
590 was not possible in the present study, because of the limited sample size. The addition of
591 more variables in the model and the multiplication of clusters would have violated the
592 cluster analysis assumptions, making its interpretation invalid. Studies with larger samples
593 should include more clinical parameters and could also use outcome variables suggested by
594 parents (65).

595

596 A last limitation here lies in a lack of a non-ASD lower cognitive (LC) control group making
597 difficult to evaluate the causality of ESDM intervention in the observed outcome of this
598 specific population. Nevertheless, Hedval et al. 2014 reported that 87.7% of the
599 preschoolers with ASD and LC at baseline (<70 of DQ) still had a DQ lower than 70 when
600 assessed after two years without receiving any EIBI (66). Moreover, their delay in adaptive
601 functioning worsened in all the VABS-II subdomains except for communication at the group
602 level. In contrast, in the present study LC children with similar developmental pattern (MinR)
603 only constituted 31% of our LC group, while the other LC participants (OptR) exhibited large
604 improvements in DQ as well as in adaptive behaviors. Considering these results, one can
605 infer a causal effect of ESDM in the progress made by children with important cognitive
606 delay at start. The specific effects of ESDM compared to other types of EIBI still needs to be
607 addressed with future RCT.

608

609 **Conclusion**

610

611 In this study, we applied a cluster analysis to the largest European sample of preschoolers
612 with ASD who participated in an ESDM program for 20 hours a week over a two-year period.
613 Overall, we found that ASD symptom severity decreased, and cognitive delay improved over
614 the intervention period. Furthermore, the cluster analysis suggested three main patterns of
615 cognitive trajectories over time. First, children who displayed mild cognitive and adaptive
616 behavioral delays at baseline tended to have a good developmental prognosis, finishing their
617 two years of early intervention with cognitive and adaptive behavior scores within the

618 normal range. Second, children who presented with severe cognitive delays at the start of
619 their early intervention exhibited two dramatically different patterns of developmental
620 trajectories. About a third of these children continued to fall behind developmentally,
621 despite intensive therapy services. The two remaining thirds of the children, who presented
622 with lower cognitive and adaptive behavior scores at the beginning of treatment, exhibited
623 early and important gains in cognition and adaptive behavior which continued for the
624 duration of the 2 years of intervention. We did not find any significant demographic or
625 behavioral differences between these two lower cognitive subgroups at baseline, suggesting
626 that it is not yet possible to predict how a child with a more severe presentation of ASD will
627 respond to treatment based solely on developmental assessment measures commonly used
628 at intake in clinical practice. Nevertheless, our results suggest that it may be possible to
629 predict, after only 6 months of early intervention, and with very high levels of accuracy,
630 whether a child will have an overall minimal or optimal response to treatment, based on
631 their early gains in cognition and communicative adaptive behavior. These results advocate
632 for close monitoring using standardized cognitive and adaptive behavioral testing during the
633 first six months of intervention, especially for children that exhibit a clinically significant
634 cognitive delay at baseline. Having an understanding early-on of how a child is responding to
635 early intervention could alert clinicians and parents to the need to adapt and enhance the
636 child's treatment plan Future studies are needed to replicate these findings, and to evaluate
637 the kinds of treatment adaptations that would optimize child outcome for each ASD
638 subgroup. Also, there is a need for longitudinal studies that provide a long-term follow-up in
639 the years following the end of early intervention, to be able to assess whether the patterns
640 of cognitive profiles and response to treatment observed remain stable over time. Overall,
641 our results advocate for a more systematic use of subgroup phenotyping that includes
642 longitudinal parameters when assessing the efficacy of an early intensive intervention, to
643 better decipher the great heterogeneity of behavioral dynamics in treatment response.

644

645 **Supplementary Material**

646 Supplementary table 1

647 Supplementary table 2

648

649 **Declaration**

650

651 **Author's contribution**

652

653 M.S. conceived and designed the study. M.F, M.G., N.K., and F.R. participated in the data
654 acquisition. M.G. and F.R. prepared and analyzed the data under the supervision of M.S. All
655 authors participated in interpretation of results. M.G. and F.R. wrote the manuscript with
656 the inputs from all other authors. All authors read and approved the final manuscript.

657

658 **Funding**

659

660 This research was supported by the Swiss National Foundation Synapsy Grant No. (51NF40 –
661 185897) and the Swiss National Foundation for Scientific Research Grant (No. 323630-
662 191227 to M.G., and #163859 & #190084 to M.S.). and by the “Fondation Pôle Autisme”
663 (<https://www.pole-autisme.ch>). The funders were not involved in this study and had no role
664 other than to provide financial support.

665

666 **Availability of data and materials**

667 The datasets generated in the current study are available from the corresponding author on
668 reasonable request.

669

670 **Ethics approval and consent to participate**

671 Informed consent was obtained from the parents of all participants included in the study.
672 Swissethics - Commission d'éthique Suisse relative à la recherche sur l'être humain approved
673 this study (Protocole 12-163/Psy 12-014), referred under the number PB_2016-01880 and
674 accepted on september the 25th, 2012.

675 **Competing interests**

676 The authors have no competing interests to report.

677

678 **Acknowledgments**

679





680 The authors would like to thank all the families who kindly participated in the study, all
681 therapists at the Centre d'Intervention Précoce en Autisme in Geneva, as well as Alexandra
682 Bastos, Stéphanie Baudoux, Lylia Ben Hadid, Aurélie Bochet, Léa Chambaz, Flore Couty,
683 Sophie Diakonoff, Lisa Esposito, Constance Ferrat, Marie-Agnès Graf, Oriane Grosvernier,
684 Kenza Latrèche, Sara Maglio, Matthieu Mansion, Eva Micol, Irène Pittet, Sonia Richetin,
685 Laura Sallin, Stefania Solazzo, Myriam Speller, Chiara Usuelli and Ornella Vico Begara for
686 their help with data collection.



- 687 1. American Psychiatric Association, editor. Diagnostic and statistical manual of mental
688 disorders: DSM-5. 5th ed. Washington, D.C: American Psychiatric Association; 2013. 947 p.
- 689 2. Dawson G, Rogers SJ, Munson J, Smith M, Winter J, Greenson J, et al. Randomized,
690 Controlled Trial of an Intervention for Toddlers With Autism: The Early Start Denver Model.
691 *Pediatrics*. 2010 Jan 1;125(1):e17–23.
- 692 3. Lovaas OI, Schreibman L, Koegel RL. A behavior modification approach to the
693 treatment of autistic children. *J Autism Dev Disord*. 1974 Jun 1;4(2):111–29.
- 694 4. Freitag CM, Feineis-Matthews S, Valerian J, Teufel K, Wilker C. The Frankfurt early
695 intervention program FFIP for preschool aged children with autism spectrum disorder: a pilot
696 study. *J Neural Transm*. 2012 Sep 1;119(9):1011–21.
- 697 5. Koegel LK, Koegel RL, Harrower JK, Carter CM. Pivotal Response Intervention I:
698 Overview of Approach. *Journal of the Association for Persons with Severe Handicaps*. 1999
699 Sep 1;24(3):174–85.
- 700 6. Prizant BM, Wetherby AM, Rubin E, Laurent AC, Rydell P. *The Scerts Model: A*
701 *Comprehensive Educational Approach for Children With Autism Spectrum Disorders*.
702 Baltimore, Md: Brookes Publishing Co; 2006. 774 p.
- 703 7. Schreibman L, Dawson G, Stahmer AC, Landa R, Rogers SJ, McGee GG, et al.
704 *Naturalistic Developmental Behavioral Interventions: Empirically Validated Treatments for*
705 *Autism Spectrum Disorder*. *J Autism Dev Disord*. 2015;45(8):2411–28.
- 706 8. Corsello CM. *Early Intervention in Autism: Infants & Young Children*. 2005
707 Apr;18(2):74–85.
- 708 9. Eldevik S, Hastings RP, Hughes JC, Jahr E, Eikeseth S, Cross S. Meta-Analysis of
709 Early Intensive Behavioral Intervention for Children With Autism. *Journal of Clinical Child*
710 *& Adolescent Psychology*. 2009 May 19;38(3):439–50.
- 711 10. Fuller EA, Kaiser AP. The Effects of Early Intervention on Social Communication
712 Outcomes for Children with Autism Spectrum Disorder: A Meta-analysis. *J Autism Dev*
713 *Disord*. 2020 May;50(5):1683–700.
- 714 11. Howlin P, Magiati I, Charman T. Systematic Review of Early Intensive Behavioral
715 Interventions for Children With Autism. MacLean Jr William E, editor. *American Journal on*
716 *Intellectual and Developmental Disabilities*. 2009 Jan 1;114(1):23–41.
- 717 12. Warren Z, McPheeters ML, Sathe N, Foss-Feig JH, Glasser A, Veenstra-VanderWeele
718 J. A Systematic Review of Early Intensive Intervention for Autism Spectrum Disorders.
719 *Pediatrics*. 2011 May 1;127(5):e1303–11.
- 720 13. Vivanti G, Prior M, Williams K, Dissanayake C. Predictors of Outcomes in Autism
721 Early Intervention: Why Don't We Know More? *Front Pediatr* [Internet]. 2014 [cited 2017
722 Nov 27];2. Available from: <https://www.frontiersin.org/articles/10.3389/fped.2014.00058/full>
- 723 14. Zwaigenbaum L, Bauman ML, Choueiri R, Kasari C, Carter A, Granpeesheh D, et al.
724 *Early Intervention for Children With Autism Spectrum Disorder Under 3 Years of Age:*
725 *Recommendations for Practice and Research*. *Pediatrics*. 2015 Oct 1;136(Supplement 1):S60–
726 81.
- 727 15. Fuentes J, Hervás A, Howlin P, (ESCAP ASD Working Party). ESCAP practice
728 guidance for autism: a summary of evidence-based recommendations for diagnosis and
729 treatment. *Eur Child Adolesc Psychiatry*. 2021 Jun;30(6):961–84.
- 730 16. König IR, Fuchs O, Hansen G, Mutius E von, Kopp MV. What is precision medicine?
731 *European Respiratory Journal* [Internet]. 2017 Oct 1 [cited 2021 Mar 25];50(4). Available
732 from: <https://erj.ersjournals.com/content/50/4/1700391>
- 733 17. Peters-Scheffer N, Didden R, Korzilius H, Matson J. Cost comparison of early
734 intensive behavioral intervention and treatment as usual for children with autism spectrum
735 disorder in the Netherlands. *Research in Developmental Disabilities*. 2012 Nov 1;33(6):1763–
736 72.

- 737 18. Penner M, Rayar M, Bashir N, Roberts SW, Hancock-Howard RL, Coyte PC. Cost-
738 Effectiveness Analysis Comparing Pre-diagnosis Autism Spectrum Disorder (ASD)-Targeted
739 Intervention with Ontario's Autism Intervention Program. *J Autism Dev Disord*. 2015
740 Sep;45(9):2833–47.
- 741 19. Cidav Z, Munson J, Estes A, Dawson G, Rogers SJ, Mandell D. Cost Offset
742 Associated With Early Start Denver Model for Children With Autism. *J Am Acad Child*
743 *Adolesc Psychiatry*. 2017 Sep;56(9):777–83.
- 744 20. Fuller EA, Oliver K, Vejnaska SF, Rogers SJ. The Effects of the Early Start Denver
745 Model for Children with Autism Spectrum Disorder: A Meta-Analysis. *Brain Sci [Internet]*.
746 2020 Jun 12 [cited 2021 Mar 24];10(6). Available from:
747 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7349854/>
- 748 21. Contaldo A, Colombi C, Pierotti C, Masoni P, Muratori F. Outcomes and moderators
749 of Early Start Denver Model intervention in young children with autism spectrum disorder
750 delivered in a mixed individual and group setting. *Autism*. 2020 Apr;24(3):718–29.
- 751 22. Vivanti G, Dissanayake C, Zierhut C, Rogers SJ, Team VA. Brief Report: Predictors
752 of Outcomes in the Early Start Denver Model Delivered in a Group Setting. *J Autism Dev*
753 *Disord*. 2013 Jul 1;43(7):1717–24.
- 754 23. Estes A, Swain DM, MacDuffie KE. The effects of early autism intervention on
755 parents and family adaptive functioning. *Pediatr Med*. 2019 Jun;2.
- 756 24. Lombardo MV, Busuoli EM, Schreibman L, Stahmer AC, Pramparo T, Landi I, et al.
757 Pre-treatment clinical and gene expression patterns predict developmental change in early
758 intervention in autism. *Mol Psychiatry [Internet]*. 2021 Aug 2 [cited 2021 Oct 3]; Available
759 from: <https://www.nature.com/articles/s41380-021-01239-2>
- 760 25. Lord C, Risi S, Lambrecht L, Cook EH, Leventhal BL, DiLavore PC, et al. The autism
761 diagnostic observation schedule-generic: a standard measure of social and communication
762 deficits associated with the spectrum of autism. *J Autism Dev Disord*. 2000 Jun;30(3):205–
763 23.
- 764 26. Lord C, Rutter M, DiLavore PC, Risi S, Gotham K, Bishop SL. *Autism Diagnostic*
765 *Observation Schedule, 2nd edition (ADOS-2)*. Los Angeles, CA: Western Psychological
766 Services; 2012.
- 767 27. Franchini M, Zöllner D, Gentaz E, Glaser B, Wood de Wilde H, Kojovic N, et al. Early
768 Adaptive Functioning Trajectories in Preschoolers With Autism Spectrum Disorders. *J*
769 *Pediatr Psychol*. 2018 Apr 25;
- 770 28. Rogers SJ, Dawson G. *Early Start Denver Model for young children with autism:*
771 *promoting language, learning, and engagement*. New York: Guilford Press; 2010. 297 p.
- 772 29. Gotham K, Pickles A, Lord C. Standardizing ADOS scores for a measure of severity
773 in autism spectrum disorders. *J Autism Dev Disord*. 2009 May;39(5):693–705.
- 774 30. Hus V, Gotham K, Lord C. Standardizing ADOS domain scores: separating severity
775 of social affect and restricted and repetitive behaviors. *J Autism Dev Disord*. 2014
776 Oct;44(10):2400–12.
- 777 31. Mullen EM. *Mullen Scales of Early Learning*. AGS ed. MN: American Guidance
778 Service Inc.; 1995. (Circle Pines).
- 779 32. Schopler E, Lansing MD, Reichler RJ, Marcus LM. *Psychoeducational Profile:*
780 *TEACCH individualized assessment for children with autism spectrum disorders (3rd ed.)*.
781 Austin, TX: Pro-Ed, Inc.; 2005.
- 782 33. Lord C, Risi S, DiLavore PS, Shulman C, Thurm A, Pickles A. Autism from 2 to 9
783 years of age. *Arch Gen Psychiatry*. 2006 Jun;63(6):694–701.
- 784 34. Sparrow SS, Balla DA, Cicchetti DV. *Vineland II: Vineland adaptive behavior scales*.
785 American Guidance Service. 2005.
- 786 35. Hollingshead AA. *Four-factor index of social status*. Yale University.



- 787 1975;Unpublished manuscript.
- 788 36. Berry DA, Ayers GD. Symmetrized Percent Change for Treatment Comparisons. *The*
789 *American Statistician*. 2006;60(1):27–31.
- 790 37. Klintwall L, Eikeseth S. Number and controllability of reinforcers as predictors of
791 individual outcome for children with autism receiving early and intensive behavioral
792 intervention: A preliminary study. *Research in Autism Spectrum Disorders*. 2012
793 Jan;6(1):493–9.
- 794 38. Sallows GO, Graupner TD. Intensive Behavioral Treatment for Children With Autism:
795 Four-Year Outcome and Predictors. MacLean Jr. WE, editor. *American Journal on Mental*
796 *Retardation*. 2005 Nov 1;110(6):417–38.
- 797 39. Sallows GO, Graupner TD. Replicating Lovaas ' Treatment and Findings: Preliminary
798 Results 1. 1999.
- 799 40. Perry A, Cummings A, Geier J, Freeman N, Hughes S, LaRose L, et al. Effectiveness
800 of Intensive Behavioral Intervention in a large, community-based program. *Research in*
801 *Autism Spectrum Disorders*. 2008 Oct 1;2:621–42.
- 802 41. Ousley O, Cermak T. Autism Spectrum Disorder: Defining Dimensions and
803 Subgroups. *Curr Dev Disord Rep*. 2014 Mar 1;1(1):20–8.
- 804 42. Syriopoulou-Delli CK, Papaefstathiou E. Review of cluster analysis of phenotypic
805 data in Autism Spectrum Disorders: distinct subtypes or a severity gradient model?
806 *International Journal of Developmental Disabilities*. 2020 Jan 1;66(1):13–21.
- 807 43. Frazier TW, Klingemier EW, Anderson CJ, Gengoux GW, Youngstrom EA, Hardan
808 AY. A Longitudinal Study of Language Trajectories and Treatment Outcomes of Early
809 Intensive Behavioral Intervention for Autism. *J Autism Dev Disord*. 2021 Feb 8;1–17.
- 810 44. Landa RJ. Efficacy of early interventions for infants and young children with, and at
811 risk for, autism spectrum disorders. *International Review of Psychiatry*. 2018 Jan 2;30(1):25–
812 39.
- 813 45. Smith T, Groen AD, Wynn JW. Randomized trial of intensive early intervention for
814 children with pervasive developmental disorder. *Am J Ment Retard*. 2000 Jul;105(4):269–85.
- 815 46. Rodgers M, Simmonds M, Marshall D, Hodgson R, Stewart LA, Rai D, et al.
816 Intensive behavioural interventions based on applied behaviour analysis for young children
817 with autism: An international collaborative individual participant data meta-analysis. *Autism*.
818 2021 May 1;25(4):1137–53.
- 819 47. Witwer AN, Lecavalier L. Examining the validity of autism spectrum disorder
820 subtypes. *J Autism Dev Disord*. 2008 Oct;38(9):1611–24.
- 821 48. Hartigan JA, Wong MA. Algorithm AS 136: A K-Means Clustering Algorithm.
822 *Applied Statistics*. 1979;28(1):100.
- 823 49. Kodinariya T, Makwana PR. Review on determining number of Cluster in K-Means
824 Clustering. undefined [Internet]. 2013 [cited 2021 May 26]; Available from: /paper/Review-
825 on-determining-number-of-Cluster-in-K-Means-Kodinariya-
826 Makwana/1a34936bffe558a380168b790dc37956813514ba
- 827 50. Chiu T, Fang D, Chen J, Wang Y, Jeris C. A robust and scalable clustering algorithm
828 for mixed type attributes in large database environment. *KDD '01* [Internet]. 2001 [cited 2021
829 May 26];Proceedings of the seventh QCM SIGKDD internqtionql conference on Knowledge
830 discovery and data mining. Available from: /paper/A-robust-and-scalable-clustering-
831 algorithm-for-type-Chiu-Fang/a1ddee7a0c4885b46891d5d8f873e9388d42a4c1
- 832 51. Rogers SJ, Estes A, Lord C, Vismara L, Winter J, Fitzpatrick A, et al. Effects of a
833 Brief Early Start Denver Model (ESDM)–Based Parent Intervention on Toddlers at Risk for
834 Autism Spectrum Disorders: A Randomized Controlled Trial. *Journal of the American*
835 *Academy of Child & Adolescent Psychiatry*. 2012 Oct 1;51(10):1052–65.
- 836 52. Clark MLE, Barbaro J, Dissanayake C. Continuity and Change in Cognition and

- 837 Autism Severity from Toddlerhood to School Age. *J Autism Dev Disord*. 2017
838 Feb;47(2):328–39.
- 839 53. Fein D, Stevens M, Dunn M, Waterhouse L, Allen D, Rapin I, et al. Subtypes of
840 Pervasive Developmental Disorder: Clinical Characteristics. *Child Neuropsychology -*
841 *CHILD NEUROPSYCHOL*. 1999 Mar 1;5:1–23.
- 842 54. Stevens MC, Fein DA, Dunn M, Allen D, Waterhouse LH, Feinstein C, et al.
843 Subgroups of children with autism by cluster analysis: a longitudinal examination. *J Am Acad*
844 *Child Adolesc Psychiatry*. 2000 Mar;39(3):346–52.
- 845 55. Zheng S, Hume KA, Able H, Bishop SL, Boyd BA. Exploring Developmental and
846 Behavioral Heterogeneity among Preschoolers with ASD: A Cluster Analysis on Principal
847 Components. *Autism Research*. 2020;13(5):796–809.
- 848 56. Zachor DA, Ben-Itzhak E. Variables Affecting Outcome of Early Intervention in
849 Autism Spectrum Disorder. *Journal of Pediatric Neurology*. 2017 Jun;15(03):129–33.
- 850 57. Eldevik S, Hastings RP, Hughes JC, Jahr E, Eikeseth S, Cross S. Using participant
851 data to extend the evidence base for intensive behavioral intervention for children with
852 autism. *Am J Intellect Dev Disabil*. 2010 Sep;115(5):381–405.
- 853 58. Magiati I, Moss J, Charman T, Howlin P. Patterns of change in children with Autism
854 Spectrum Disorders who received community based comprehensive interventions in their pre-
855 school years: A seven year follow-up study. *Research in Autism Spectrum Disorders*. 2011
856 Jul 1;5(3):1016–27.
- 857 59. Warriar V, Zhang X, Reed P, Havdahl A, Moore TM, Cliquet F, et al. Genetic
858 correlates of phenotypic heterogeneity in autism [Internet]. *Genetic and Genomic Medicine*;
859 2020 Jul [cited 2021 Oct 3]. Available from:
860 <http://medrxiv.org/lookup/doi/10.1101/2020.07.21.20159228>
- 861 60. Ben-Itzhak E, Watson LR, Zachor DA. Cognitive Ability is Associated with
862 Different Outcome Trajectories in Autism Spectrum Disorders. *J Autism Dev Disord*. 2014
863 Sep 1;44(9):2221–9.
- 864 61. Devescovi R, Monasta L, Mancini A, Bin M, Vellante V, Carrozzi M, et al. Early
865 diagnosis and Early Start Denver Model intervention in autism spectrum disorders delivered
866 in an Italian Public Health System service. *Neuropsychiatr Dis Treat*. 2016;12:1379–84.
- 867 62. Robain F, Franchini M, Kojovic N, Wood de Wilde H, Schaer M. Predictors of
868 Treatment Outcome in Preschoolers with Autism Spectrum Disorder: An Observational Study
869 in the Greater Geneva Area, Switzerland. *J Autism Dev Disord*. 2020 Mar 12;
- 870 63. Formann AK. *Die Latent-Class-Analyse: Einführung in die Theorie und Anwendung*
871 [Latent class analysis: Introduction to theory and application]. Weinheim. Beltz; 1984.
- 872 64. Qiu W, Joe H. Generation of Random Clusters with Specified Degree of Separation.
873 *Journal of Classification*. 2006 Feb 1;23:315–34.
- 874 65. McConachie H, Livingstone N, Morris C, Beresford B, Couteur AL, Gringras P, et al.
875 Parents Suggest Which Indicators of Progress and Outcomes Should be Measured in Young
876 Children with Autism Spectrum Disorder. *J Autism Dev Disord*. 2018 Apr 1;48(4):1041–51.
- 877 66. Hedvall Å, Westerlund J, Fernell E, Holm A, Gillberg C, Billstedt E. Autism and
878 developmental profiles in preschoolers: stability and change over time. *Acta Paediatr*. 2014
879 Feb;103(2):174–81.
- 880

A**B****C****D**

A**B**

A



B

