Single Prime hAd5 Spike (S) + Nucleocapsid (N) Dual Antigen Vaccination of Healthy Volunteers Induces a Ten-Fold Increase in Mean S- and N- T-Cell Responses Equivalent to T-Cell Responses from Patients Previously Infected with SARS-CoV-2

Peter Sieling[1], Thomas King[1], Raymond Wong[1], Andy Nguyen[1], Kamil Wnuk[1], Elizabeth Gabitzsch[1], Adrian Rice[1], Helty Adisetiyo[1], Melanie Hermreck[1], Mohit Verma[1], Lise Zakin[1], Annie Shin[1], Brett Morimoto[1], Wendy Higashide[1], Kyle Dinkins[1], Joseph Balint[1], Victor Peykov[1], Justin Taft[2,3], Roosheel Patel[2,3], Sofija Buta[2,3,4], Marta Martin-Fernandez[2,3,4,5,6], Dusan Bogunovic[2,3,4,5,6], Patricia Spilman[1], Lennie Sender[7], Sandeep Reddy[7], Philip Robinson[8], Shahrooz Rabizadeh[1], Kayvan Niazi[1], and Patrick Soon-Shiong[1]*

Affiliations

[1]ImmunityBio, LLC, 9920 Jefferson Blvd., Culver City, CA 90232, USA

[2]Center for Inborn Errors of Immunity, [3]Department of Pediatrics, [4]Precision Immunology Institute, [5]Mindich Child Health and Development Institute, [6]Department of Microbiology, Icahn School of Medicine at Mount Sinai, 1 Gustave Lane, Levy Place, New York, NY 10029-5674, USA SPA

[7]NantKwest, Inc., 9920 Jefferson Blvd., Culver City, CA 90232, USA

[8]Hoag Hospital, Newport Beach, CA 92663, USA

*Corresponding author: Patrick@Nantworks.com

ABSTRACT

In response to the need for a safe, efficacious vaccine that provides broad immune protection against SARS-CoV-2 infection, we have developed a dual-antigen COVID-19 vaccine. The vaccine delivers both the viral spike (S) protein modified to increase cell-surface expression (S-Fusion) and the viral nucleocapsid (N) protein with an Enhanced T-cell Stimulation Domain (N-ETSD) to enhance MHC class I and II presentation and T-cell responses. The vaccine antigens are delivered using a human adenovirus serotype 5 (hAd5) platform with E1, E2b, and E3 regions deleted that has been shown in previous cancer vaccine studies to be effective in the presence of pre-existing hAd5 immunity. Here, we demonstrate the hAd5 S-Fusion + N-ETSD (hAd5 S + N) vaccine antigens when expressed by dendritic cells (DCs) of previously SARS-CoV-2-infected patients elicit Th1 dominant activation of autologous patient T cells, indicating the vaccine antigens have the potential for generating immune responses in patients previously infected or vaccinated. We further demonstrate that participants in our open-label Phase 1b study of the dual-antigen hAd5 S + N vaccine generate Th1 dominant S- and N- specific T cells after a single prime subcutaneous injection and that the magnitude of these responses were comparable to those seen for T cells from previously infected patients. We further present our *in silico* prediction of T-cell epitope HLA binding for both the first-wave SARS-CoV-2 'A' strain and the K417N, E484K, and N501Y S as well as the T201I N variants that suggests T-cell responses to the hAd5 S + N vaccine will retain efficacy against these variants. These findings that the dual-antigen hAd5 S + N vaccine elicits SARS-CoV-2-relevant T-cell responses and that such cell-mediated protection is likely to be sustained against emerging variants supports the testing of this vaccine as a universal booster that would enhance and broaden existing immune protection conferred by currently approved S-based vaccines.

INTRODUCTION

To address the need for an efficacious COVID-19 vaccine that elicits broad immunity against SARS-CoV-2, is suitable for global distribution and use, and has a high likelihood of maintaining efficacy against emerging SARS-CoV-2 variants, we have developed a dual-antigen COVID-19 'T cell' vaccine. This dual-antigen vaccine expresses both the virus spike (S) and nucleocapsid (N) proteins using a next-generation human adenovirus serotype 5 (Ad5) platform. The S antigen is full-length S including SD1 receptor binding domain, S1 and S2 domains modified to enhance surface expression (S-Fusion); [1] and full-length N protein modified with an Enhanced T cell Stimulation Domain (ETSD) to direct N to the endo-/lysosomal compartment for increased MHC class I and II expression. The hAd5 platform used has deletions in the E1, E2b, and E3 gene regions, thereby minimizing host anti-vector immune responses [2,3] and enabling efficient antigen cargo expression and cognate T cell activation even in the presence of existing anti-adenovirus immunity, as demonstrated previously in clinical studies targeting tumor-associated antigens in cancer patients. [4,5]

The emergence of SARS-CoV-2 variants that have the potential to evade immune responses generated in response to currently available vaccines has spurred renewed interest in the potential role of T cells in conferring long-term protection against COVID-19. [6-9] SARS-CoV-2 vaccine development has focused largely on eliciting neutralizing antibodies against the S protein to inhibit infection and reduce disease severity. [10-14] S-specific T cell responses have been reported for two available vaccines, mRNA1273 and BNT162b1, [15,16] however, their role in conferring protective immunity against COVID-19 is unclear.

SARS-CoV-2 strain analysis in participants in Phase 3 studies in South Africa has revealed that some vaccines show diminished protection against the B.1.351 variant strain, [17] including ChAdOx, [18] NVX-CoV2373, [19,20] and Ad26.COV2-S. [21] In addition, *in vitro* studies show that wild type S-specific antibodies elicited by mRNA-1273 and BNT162b1 show reduced binding to the

B.1.351 variant S protein. [22-24] Taken together, these findings raise the concern that monovalent

vaccines targeting only the S protein may not be the most optimal strategy for conferring protection

against continually emerging variants.

SARS-CoV-2 expresses a number of other immunogenic proteins that may induce protective

antibody and/or T cell responses. Among these, the N protein is the most abundantly expressed, is

highly conserved among coronaviruses, and has been studied previously as an antigen for

therapeutics and vaccines for SARS-CoV.[25-29] Inclusion of N-derived antigens into vaccine

designs may be a rational strategy to elicit broader protective immunity against SARS-CoV-2 with

a high probability of being sustained against variants. The N protein associates with viral RNA

within the SARS-CoV-2 virion and is necessary for viral RNA replication, virion assembly, and

release from host cells.[28,29] Virtually all SARS-CoV-2 convalescents develop N-specific antibody

and $CD8^+$ and/or $CD4^+$ T cell responses.[30] Remarkably, SARS-CoV convalescents from 2003

maintain robust cognate N-specific $CD8^+$ and/or $CD4^+$ T cell responses that were detectable up to

17 years later.[31] These long-lived memory T cells also cross-react with N and other proteins of

SARS-CoV-2.[31] To a lesser extent, individuals with no history of exposure to SARS-CoV, SARS-

CoV-2, or Middle Eastern Respiratory Syndrome (MERS) also show cross-reactive SARS-CoV-

2 N-specific $CD8^+$ and/or $CD4^+$ T cell responses, likely due to prior exposure to related endemic

coronaviruses including OC43, HKU1, 229E, and NL63.[30] These data suggest that induction of

N-specific memory T cell responses via vaccination may at least partly recapitulate the natural

course of SARS-CoV-2 infection and recovery from illness, and could be beneficial for achieving

more optimal disease-limiting immunity against the virus.

Our addition of N to our dual antigen hAd5 S-Fusion + N-ETSD vaccine leverages the ability

to N to generate protective, durable T-cell responses. Recent characterization of COVID-19 patient

immune responses to SARS-CoV-2 indicates T cell activation is critical for clearance of infection

and production of long-term immunity to coronavirus infections.[31-34] Both CD4+ and CD8+ T cells

underpin durable humoral responses because CD4+ T cells, while not effector cells like CD8+ T

cells, are critical to the generation of robust and long-lasting immunity afforded by antibody-

secreting plasma cells and the elimination of infected cells by memory cytotoxic CD8+ T cells.[35,36]

In the studies reported here, we first confirm that the addition of the ETSD to N, intended to

increase MHC class I and II responses and T-cell activation, directs N to the cell endo-/lysosomal

compartment. We then demonstrate that the hAd5-S-Fusion + N-ETSD COVID-19 vaccine

antigens, when expressed by monocyte-derived dendritic cells (MoDCs) from individuals

previously infected with SARS-CoV-2, can activate and induce recall of S- and N-specific

autologous CD8$^+$ and CD4$^+$ T cells. We specifically show that overall, N-ETSD elicits greater T-

cell responses than N without the ETSD. This study demonstrates the vaccine-expressed antigens

are 'recalled' by T cells and thus suggests they have the potential to induce relevant, protective

immune responses in previously SARS-CoV-2 infected or vaccinated individuals who receive the

hAd5 S-Fusion + N-ETSD vaccine.

We next show that T cells isolated 14 and 23 days after prime vaccination alone from

participants in our ongoing Phase 1b clinical trials of the hAd5 S-Fusion + N-ETSD vaccine are

activated by both S and N peptides pools and that these T-cell responses are comparable in

magnitude to those of T cells from previously SARS-CoV-2 infected patients.

To support our proposition that a 'T cell' vaccine that provides both S and N antigens will

sustain efficacy against variants, we also present our *in silico* prediction of T cell epitope HLA

binding based on analyses of all possible 9-mer epitopes from both first-wave 'A' strain SARS-

CoV-2 and the variants that are found in the B.1.351 strain discovered in South Africa – spike

K417N, E484K, and N501Y and nucleocapsid T205I. This analysis shows that memory T-cell-

mediated immunity has a very low probability of being significantly affected by the B.1.351

variants and that the hAd5 S + N COVID-19 T-cell vaccine has the potential of being protective

against these variants.

RESULTS

N-ETSD is directed to endo-/lysosomal and autophagosomal compartments

The hAd5 dual antigen vaccine construct includes sequences designed to target N to MHC class II antigen loading compartments. To confirm this occurs, we evaluated the localization of N-ETSD in human monocyte-derived dendritic cells (MoDCs) compared to its untargeted cytoplasmic counterpart (N). MoDCs from healthy subjects were infected with hAd5 N-ETSD or hAd5 N and localization was determined by immunocytochemistry. N-ETSD showed localization to discrete vesicles, some coincident with CD71, a marker of recycling endosomes (Fig. 1A-C), and LAMP-1, a marker for late endosome/lysosomes (Fig. 1G-I), whereas N was expressed diffusely and uniformly throughout the cytoplasm (Fig. 1D-F; J-L).

Studies have shown that lysosomes fuse with autophagosomes to enhance peptide processing and MHC class II presentation.[37,38] Thus we examined whether N-ETSD localized in autophagosomes in MoDCs by co-labeling with the autophagosome marker LC3a/b [39] to identify another potential site of localization relevant to MHC class II antigen presentation.[40] We found that N-ETSD also displayed some co-localization with the autophagosome marker (Fig. 1M-O).

Protein processing in autophagosomes plays a key role in MHC-mediated antigen presentation in DCs,[41-43] providing a potential mechanism of enhanced CD4+ T cells induced by N-ETSD in the vaccine construct. Evidence of this T cell interaction with a MoDC infected with N-ETSD translocated to autophagosomes (and, it is assumed, also endosomes and lysosomes) is seen in this phase-contrast microscopy of the N-ETSD and LC3a/b co-labeled cells, which reveals the elongated DC morphology in contrast to the spherical morphology of what are posited to be undifferentiated lymphocytes.

Fig. 1 *N-ETSD localizes to endosomes, lysosomes, and autophagosomes.* MoDCs were infected
with Ad5 N-ETSD or N without ETSD and were co-labeled with anti-flag (N, N-ETSD here have
a flag tag) and anti-CD71 (endosomal marker), anti-Lamp 1 (lysosomal marker), or anti-LC3a/b
antibodies. (A) N-ETSD, (B) CD71, and (C) overlay. (D) N, (E) CD71, and (F) overlay. (G) N-
ETSD, (H) Lamp-1, and (I) overlay. (J) N, (K) Lamp-1, and (L) overlay. (M) N-ETSD, (N)
LC3a/b, and (O) overlay. N/N-ETSD is red, other markers green, co-localization indicated by
yellow arrows, and white arrows indicate suspected undifferentiated lymphocytes.

The Immune Recall Study with Previously SARS-CoV-2 Infected Patients

Blood samples from previously SARS-CoV-2 infected patients confirmed to show humoral and

T-cell responses and enhanced binding to hAd5 S-Fusion + N-ETSD expressing cells

For the 'recall' studies described below, plasma samples were collected from four individuals

convalescing from SARS-CoV-2 infection as confirmed by antibody assays and patient history.

The presence of anti-Spike IgG, and neutralizing antibodies by both a surrogate SARS-CoV-1

neutralization [44] and live virus assays, were confirmed in all patient samples (Suppl. Fig. S1).

Samples from four virus-naïve individuals were used as controls.

In additional studies to validate immune responses to SARS-CoV-2 antigens, the binding of

previously SARS-CoV-2 infected patient and virus-naïve control individual plasma to human

embryonic kidney (HEK) 293T cells transfected with either hAd5 S-Fusion alone or hAd5 S-

Fusion + N-ETSD was assessed (Suppl. Fig. S2). This binding reflects the presence of antibodies

in plasma that recognize antigens expressed by the hAd5 vectored vaccines. Quantification of

histograms showed little or no binding of virus-naïve plasma antibodies to cells expressing

constructs, with the highest previously SARS-CoV-2 infected patient plasma binding to cells

expressing the dual antigen S-Fusion + N-ETSD construct (Suppl. Fig. S2R).

To confirm the T cells from previously SARS-CoV-2-infected patients were reactive to

SARS-CoV-2 peptides, isolated T cells from the patients were incubated with S1 (containing the

S receptor binding domain), S2 and N peptides pools. T cells from all four patients showed

reactivity (Suppl. Fig. S3). Selected CD4+ and CD8+ T cells from 2 patients were activated by S1,

S2, and N peptides, revealing CD8+ T cells were more reactive to N (Suppl. Fig. S3D and E).

Previously SARS-CoV-2 infected patient Th1 dominant CD3+ and CD4+/CD8+ memory T-cells

recall N-ETSD antigens expressed by autologous hAd5-infected MoDCs

To evaluate the immune significance of endo/lysosome-localized N-ETSD versus cytoplasmic

N, MoDCs were infected with hAd5 constructs (Null, N-ETSD or N) then incubated with

autologous CD3+ and CD4+- or CD8+-selected T cells (Fig. 2A). CD3+ T cells from previously

infected SARS-CoV-2 patients showed significantly greater IFN-γ secretion in response to N-

ETSD than both Null and cytoplasmic N in the two patients where N-ETSD and N were compared

(Fig. 2C and D). There were relatively few interleukïn-4 (IL-4) secreting CD3+ T cells for all

patients (Fig. 2E-G); the high IFN-γ and low IL-4 responses show Th1 dominance for both N-

ETSD and N. Both CD4+ and CD8+ selected T-cell populations showed significantly greater IFN-

γ responses to N-ETSD than Null (Fig. 2H-M) and in two of three patients, CD4+ and CD8+ T

cells showed greater recognition of N-ETSD compared to N.

189
Fig. 2 *Previously SARS-CoV-2 infected patient T-cell IFN-γ secretion is greater in response to*
MoDC-expressed N-ETSD than N and is Th1 dominant. (A) Experimental design. Secretion of
IFN-γ (B-D) or interleukin-4 (E-G) by autologous CD3+ T cells in response to hAd5 N-ETSD-/N-
/Null- expressing MoDCs is shown (same scale for IFN-γ and IL-4 for each Pt). IFN-γ secretion
by CD4+ (H-J) and CD8+ (K-M) T cells in response to hAd5-N-ETSD/N/Null are shown.
Statistical analysis performed using One-way ANOVA and Tukey's post-hoc multiple comparison
analysis, where *p<=0.05; **p<0.01, ***p<0.001 and ****p<0.0001. Comparison to Null shown
above bars, comparison between N-ETSD and N indicated with a line. Data graphed as mean and
SEM; n = 3-4.

CD4+ and CD8+ memory T-cells from previously SARS-CoV-2 infected patients recall

nucleocapsid and spike antigens expressed by autologous hAd5-infected MoDCs

To compare CD3+, CD4+, CD8+ memory T-cell responses to the components of the hAd5 S-

Fusion + N-ETSD vaccine, MoDCs were infected with hAd5 S-Fusion, hAd5 N-ETSD, or the dual

antigen vaccine (Fig. 3A).

For unselected CD3+ T cells, IFN-γ responses were similar to S-Fusion + N-ETSD and N-

ETSD with responses to S-Fusion being relatively low (Fig. 3A-C). The number of IL-4 secreting

T cells in response to any infected MoDC was very low (Fig. 3D-F). Based on the increased

expression of S in the dual antigen vaccine compared to monovalent S-Fusion, the increased T-

cell responses could be explained by either T cells recognizing increased S or the presence of N.

Importantly, these T-cell responses were characterized by a predominance of IFN-γ (Th1) relative

to IL-4 (Th2).

CD4+ T cells from all three patients showed significantly greater recognition of all three

constructs compared to Null (Fig. 3G-I). While there were greater responses to specific constructs

in some individuals, overall the responses to S-Fusion, S-Fusion + N-ETSD and N-ETSD were

similar. CD8+ T cells from all three patients recognized the dual antigen and N-ETSD vaccines at

a significantly higher level than Null; in only two of three patients did CD8+T cells recognize S-

Fusion to a significant degree above Null (Fig. 3J-L). These data indicate that T cells from

previously infected SARS-CoV-2 patients have reactivity and immune memory recall to both of

the vaccine antigens (S and N) in our vaccine vector.

219
Fig. 3 *Previously SARS-CoV-2 infected patient T-cell responses to the dual antigen vaccine and its individual components reveal distinct antigen specificity of T-cell populations.* (A-C) CD3+ T cell IFN-γ responses for three patients. (D-F) CD3+ T cell IL-4 responses. (G-I) CD4+ IFN-γ responses. (J-L) CD8+ IFN-γ responses. Statistical analysis performed using One-way ANOVA and Tukey's post-hoc multiple comparison analysis to compared each antigen-containing construct to the Null construct, where *p<0.05 and ****p<0.00001. Comparison to Null only above bars; comparison between antigen-expressing vaccines above lines. Data graphed as mean and SEM; n = 3-4.

T-Cell Responses after a Single Prime Vaccination in a Phase 1 Clinical Trial

For the ELISpot assay results shown here, T cells were isolated from patients in cohort 2 of the Phase 1b study of the ImmunityBio hAd5 COVID-19 'T cell' vaccine (NCT04591717). These participants were vaccinated subcutaneously with 1×10^{11} hAd5 S-Fusion + N-ETSD viral particles (VP).

The studies presented here are experimental and all assays and analyses were performed in-house at ImmunityBio, Inc. Formal analyses of the primary and secondary end points are ongoing pending completion of the trial.

T cells from vaccinated participants show Th1 dominant responses to S and N by Day 14 after a single prime injection, with a more than 10-fold increase in mean responses to N

T-cell responses to S, N, and SARS-CoV-2 membrane (M) peptides were determined by ELISpot for clinical study participants pre-vaccination (Day 1) and at two time points following the prime injection: Days 14-16 and 21-23.

By Day 14-16, T-cell secretion of IFN-γ in response to N peptides increased more than 10-fold compared to pre-vaccination (Day 1); these responses were sustained on Day 21-23 and were significant (Fig. 4A). The changes in T-cell responses to N for individual participants from pre- to post-vaccination are shown in Fig. 4B.

T-cell secretion of IFN-γ in response to S peptides also increased post-prime only vaccination, increasing more than 3-fold by Day 14-16 and more than 10-fold by Day 21-23, when differences were significant (Fig. 4C). T-cell responses to S for individual participants are shown in Fig. 4D.

When responses to S, N and SARS-CoV-2 membrane (M) peptides are compared, the greatest responses are seen to N peptides (Fig. 4F and G). As expected, there were very low/no responses to M peptide, not delivered in the vaccine. Importantly, IL-4 secretion in response to peptide stimulation was very low, indicating all responses to vaccine antigens were highly Th1 dominant.

Fig. 4 *Vaccinated participant T-cell responses to S, N, and M peptides.* (A) T-cell interferon-γ
(IFN-γ) secretion in response to nucleocapsid (N) peptide is shown for pre-vaccination (Day 1)
and post-prime only Day 14-16 and 21-23. (B) Individual participant T-cell responses to N. (C)T-
cell IFN-γ secretion in response to spike (S) peptide is shown for pre-vaccination (Day 1) and post-
prime only Day 14-16 and 21-23. (D) Individual participant T-cell responses to S. T-cell IFN-γ
secretion in response to no stimulation (Ctl), S, N and membrane (M) (E) pre-vaccination and on
(F) Day 14-16 and (G) 21-23 post-prime only. (H-J) Interleukin-4 (IL-4) secretion. Statistical
analysis performed using One-way ANOVA and Sidak's post-hoc analysis to compare pre-to post-
vaccination (A, C) or peptide stimulation to Ctl (E-J), where *p , 0.05, **p ≤ 0.01, and ***p ≤
0.001. Data graphed as mean and SD; baseline n = 9, Day 14-16 n = 7, Day 21-23 n = 8.

T-cell responses to S and N antigens generated from hAd5 prime vaccination alone were

equivalent to those from previously SARS-CoV-2 infected patients

As shown in Figure 5, T-cell responses to both N and S for previously SARS-CoV-2 infected

patients (Pt) and hAd5 S-Fusion + N-ETSD vaccinated participants (IB Vac) on Day 21-23 were

comparable. For responses to N, both Pt and IB Vac were ~40-fold greater than Unex; and for

responses to S, both Pt and IB Vac were >15-fold higher than Unex. As expected, T cells from

previously SARS-CoV-2 infected patients, but not vaccinated participants, reacted to M due to its

presence in the live virus. The M antigen is not present in the vaccine.

273
Fig. 5 *T-cells from vaccinated and previously SARS-CoV-2 infected individuals show similar*
responses to S and N. T-cell secretion of interferon-γ (IFN-γ) in response to (A) N, (B) S, and (C)
M as determined by ELISpot shown for SARS-CoV-2 unexposed (Unex) and previously infected
(Pt) individuals as compared to participants receiving a prime vaccination alone (IB Vac) from
Day 21-23. Secretion of interleukin-4 (IL-4) in response to (D) N, (E) S, and (F) M. Statistical
analysis performed using One-Way ANOVA with Dunnett's post-hoc analysis comparing Pt and
IB Vac to Unex; for IFN-γ secretion in response to M, Pt compared to Unex, p = 0.0303. Data
graphed as mean and SEM; Unex n = 4, Pt n = 5, IB vac n = 8.
282
283

T-Cell Epitope HLA Binding Prediction

In silico identification of 9-mer T-cell epitopes and HLA binding prediction suggests this binding is not significantly affected by B.1.351 strain variants

To understand and predict the possible effects emerging SARS-CoV-2 variants on vaccine-elicited T-cell protection, we utilized our in-house T-cell epitope identification and binding analysis pipeline to compare the first wave 'A' strain epitopes to emerging variant strains.

In silico epitope analysis of the reference strain assembled from 75,000 individual sequences deposited in the GISAID database [45,52] revealed 2479 MHC class I and 20963 MHC class II potential binders. Altogether, the SARS-CoV-2 variants (K417N, E484K and N501Y mutations) generate 10970 MHC class I and 18017 MHC class II unique epitopes.

We compared these unique epitopes to the reference strain and counted the different epitopes that are bound by HLAs, finding 1659 unique MHC class I and 2887 unique MHC class II epitopes that only exist in the variants and not the original strain. These epitopes are from the S, envelope (E), N and membrane (M) proteins, but the most unique epitopes were found in variant S due to the higher numbers of coding mutations.

For HLA binding prediction, sequences and predicted binding are compared and aligned and show that, for example, as compared to the 'A' strain, an B.1.351 sequence has a 3 amino acid deletion that results in a unique sequence RFQTLHRSYL predicted to bind HLA-C*14:02. By similar determination of the epitopes and HLAs from the reference strain and the variant strains, were identified differences in binding affinity or HLAs. Similarly at the N501Y site, the reference strain generates an epitope (TNGVGYQPY) predicted to bind HLA-A*30:02, while the B.1.351 strain would generate a different yet similar epitope (TYGVGYQPY) also predicted to bind HLA-A*30:02.

The key questions asked here were: how does mutation change HLA availability and which mutations favor immune evasion? Not all mutations lead to evasion; while a mutation may lead to

an amino acid substitution that has potential functional consequences, that same mutation may still

generate an epitope that retains HLA binding affinity and does not contribute to escape. The unique

binding HLAs for each S variants when compared to the A strain are shown in Figure 1. For class

I, there are only 2 K417N (B.1.351) HLAs and for the A strain there are 6 (Fig. 6A); for E484K

there are 8 and for the A strain 7 (Fig. 6B); and for N501Y there are 8 and only 1 for the A strain

(Fig. 6C). There are no unique class II HLAs for these same comparisons (Fig. 6D-F).

In the B.1.351 strain, there is only one N mutation – T205I. Similarly to the S variants, there

are few unique binding HLAs. For class I, the T201I mutation generates 2 and the A strain 3 unique

binding HLAs (Fig. 6G). For class II, the T205I generates 4 and the A strain just 1 unique binding

HLA(s) (Fig. 6H).

Fig. 6 *A high degree of overlap is seen for HLA binding of predicted T-cell epitopes from variants*
and the first-wave A strain SARS-CoV-2. Comparison of 'A' and B.1.351 strain predicted class I
T-cell epitopes reveals significant overlap for the spike (A) K417N (50 of 58 epitopes), (B) E484K
(31 of 46 epitopes), and (C) N501Y (52 of 61 epitopes) variants. (D-F) Complete overlap is seen
for class II epitopes for those variants, respectively. (G) For T205I class I, 30 of 35 epitopes overlap
and (H) for class II, 44 of 49. Unique HLA class I binding is shown in the tables.
328

DISCUSSION

The emergence and rapid spread of SARS-CoV-2 variants is raising concern that first

generation monovalent wild-type S-targeting vaccines may eventually lose effectiveness at

protecting against COVID-19. In addition to updating S sequences to include variants in current

vaccines, alternate or complimentary vaccine strategies with enhanced focus on eliciting protective

T cell responses and possibly inclusion of other immunogenic structural proteins, including M

and/or N. Evidence indicates that virtually all COVID-19 convalescents develop IFN-γ-secreting

CD4[+] and/or CD8[+] T cell responses not only against S, but also against the M and N proteins of

SARS-CoV-2. [30] Significant T cell reactivity in COVID-19 convalescents has also been observed

against a wide breadth of other SARS-CoV-2 proteins, particularly nsp6 and ORF3a.[27,30,46]

Collectively, the total magnitude of the T cell response against non-S proteins such as N appears

at least equal to or possibly greater than the T cell response against S in COVID-19

convalescents.[27,30,46] Further, in a recent report by Redd *et al.* [7] the authors assessed reactivity of

CD8+ T cells from thirty COVID-19 convalescent individuals to forty-five different mutations and

concluded that virtually all CD8+ T cell responses to SARS-CoV-2 antigens should recognize the

variants. Taken together, these data suggest that monovalent S-targeting vaccines designed to elicit

humoral responses show a relatively narrow T cell response compared to the natural T cell

responses to N and other antigens observed in COVID-19 convalescents.

Here we show that our strategy of including N-ETSD, confirmed to be directed to the endo-

/lysosomal compartment with S optimized for cell-surface display [1,47] results in recognition of the

N-ETSD and S-Fusion + N-ETSD antigen(s) by previously SARS-CoV-2 patient T cells and in

generation of S- and N-reactive T cells in participants receiving the hAd5 S-Fusion + N-ETSD

vaccine. Not only does the enhancement of T-cell responses by the addition of N in this vaccine

confer a greatly likelihood of sustained protection against variants as our T-cell epitope HLA

binding prediction analysis would suggest, it presents the possibility that this vaccine can be

effective as a 'universal' booster. Not only would boosting with this vaccine likely increase cell-

mediated protection for individuals who have already received a first-generation S-only vaccine,

because the N protein is highly conserved among coronaviruses,[48] the hAd5-S-Fusion + N-ETSD

vaccine might provide protection against viruses similar to SARS-CoV-2 that may arise.

The hAd5 S-Fusion + N-ETSD COVID-19 T cell vaccine is currently undergoing clinical

testing with delivery by a variety of routes, including sublingual, SC, intranasal, and a thermally-

stable oral tablet as a prime/boost to provide protection against SARS-CoV-2. We anticipate our

testing its use as a universal boost for current first generation vaccines in the near future.

ACKNOWLEDGEMENTS

We thank the volunteers participating in our Phase I COVID-19 vaccine clinical trial and

Deborah Fridman, Director of Clinical Research at Hoag Hospital (Irvine/Newport Beach, CA).

We also thank Phil Yang of ImmunityBio for his ongoing coordination of project updates for the

studies described herein.

MATERIALS and METHODS

Production hAd5-S-Fusion+N-ETSD and related SARS-CoV-2 antigen expressing vectors.

Production of all hAd5 [E1-, E2b-, E3-] (Fig. 7A) constructs and virus particles were carried

out as previously described.[49] In brief, high titer adenoviral stocks were generated by serial

propagation in the E1- and E2b-expressing E.C7 packaging cell line, followed by $CsCl_2$

purification, and dialysis into storage buffer (2.5% glycerol, 20 mM Tris pH 8, 25 mM NaCl) by

ViraQuest Inc. (North Liberty, IA). Viral particle counts were determined by sodium dodecyl

sulfate disruption and spectrophotometry at 260 and 280 nm. Viral titers were determined using

the Adeno-X™ Rapid Titer Kit (Takara Bio).

The constructs created included:

B. S-Fusion: S optimized to enhance surface expression and display of RBD.

C. N (N without ETSD): Nucleocapsid (wild type) sequences with tags for immune detection, but

without ETSD modification, and predominantly cytoplasmic localization.

D. N with the Enhanced T cell Stimulation Domain (N-ETSD): Nucleocapsid (wild type) with

ETSD to direct lysosomal/endosomal localization and tags for immune detection.

E. The dual-antigen hAd5 S-Fusion + N-ETSD vaccine.

385
386 **Fig. 7** *Human adenovirus serotype 5 (hAd5) platform and constructs*. (A) hAd5 with E1, E2b and
387 E3 regions deleted (hAd5[E1-, E2b-, E3-]). The site of antigen sequence insertion is shown with a
388 red arrow. Constructs tested included (B) spike (S)-Fusion, (C) nucleocapsid (N) without the
389 Enhanced T-cell Stimulation Domain (ETSD), (D) N-ETSD, and (E) the hAd5 S-Fusion + N-
390 ETSD vaccine.

391
The T-cell recall studies of unexposed and previously SARS-CoV-2 infected individuals

Collection of plasma and PBMCs for differentiation of MoDCs from patients with confirmed

previous SARS-CoV-2 infection and from virus-naïve volunteers

Blood was collected with informed consent via venipuncture from volunteer patients (Pt) who

had either not been exposed (UNEX) to SARS-CoV-2 as confirmed by ELISA and multiple

negative SARS-CoV-2 tests or who had recovered from COVID-19 as indicated by recent medical

history and a positive SARS-CoV-2 antibody test. The presence of anti-S IgG and of neutralizing

antibodies in plasma from previously SARS-CoV-2 infected patients was confirmed by ELISA, a

surrogate neutralization assay,[44] and a live virus assay as shown in Supplementary Fig. S1. A third

source of whole blood was apheresis of healthy subjects from a commercial source (HemaCare).

Peripheral blood mononuclear cells (PBMCs) were isolated from whole blood by density

gradient centrifugation and plasma was collected after density gradient centrifugation.

Monocyte-derived dendritic cells (MoDC) were differentiated from PBMC using GM-CSF

(200U/ml) and IL-4 (100U/ml) as previously described [50]. Briefly, monocytes were enriched by

adherence on plastic, while the non-adherent cells were saved and frozen as a source of

lymphocytes, specifically T cells. Adherent cells were differentiated into dendritic cells (3-5 d in

RPMI containing 10% FBS), then frozen in liquid nitrogen for later use. T cells were enriched

from the non-adherent fraction of PBMC using MojoSort (BioLegend CD3 enrichment). CD4+

and CD8+ T cells were enriched using analogous kits from the same manufacturer. Efficiency of

the cell separations was evaluated by flow cytometry.

Transduction of MoDCs with hAd5 N-WT or N-ETSD and labeling with anti-N, anti-CD71, anti-

LAMP-1, and Anti-LC3a/b antibodies

Freshly thawed MoDCs were plated on 4-well Lab-Tek II CC2 Chamber Slides, using 3 x10^4

cells per well and transduction performed at MOI 5000 one hour after plating using hAd5 N-ETSD

or hAd5 N. Slides were incubated o/n at 37°C, fixed in 4% paraformaldehyde for 15 minutes, then

permeabilized with 1% Triton X100, in PBS) for 15 min. at room temperature. To label N, cells

were then incubated with an anti-flag monoclonal (Anti-Flag M2 produced in mouse) antibody at

1:1000 in phosphate buffered saline (PBS) with 3% BSA, 0.5% Triton X100 and 0.01% saponin

overnight at 4°C, followed by three washes in PBS and a 1 hour incubation with a goat anti-Mouse

IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 555 (Life

Technologies) at 1:500. For co-localization studies, cells were also incubated overnight at 4ºC with

a rabbit anti-CD71 (transferrin receptor, recycling/sorting endosomal marker) antibody

(ThermoFisher) at 1:200; sheep anti-Lamp1 Alexa Fluor 488-conjugated (lysosomal marker)

antibody (R&D systems) at 1:10; or a rabbit monoclonal anti human LC3a/b (Light Chain 3,

autophagy marker) antibody (Cell Signaling Tech #12741S) used at 1:100. After removal of the

primary antibody, two washes in PBS and three washes in PBS with 3% BSA, cells were incubated

with fluor-conjugated secondary antibodies when applicable at 1:500 (Goat anti-Rabbit IgG (H+L)

secondary antibody, Alexa Fluor 488; 1:500 dilution) for 1 hour at room temperature. After brief

washing, cells were mounted with Vectashield Antifade mounting medium with DAPI (Fisher

Scientific) and immediately imaged using a Keyence all-in-one Fluorescence microscope camera

and Keyence software.

T cell and selected CD4+ and CD8+ T cell secretion of IFN-γ in response to MoDCs pulsed with

SARS-CoV-2 peptide antigen pools

The ability of T cells from previously infected patients used in these studies to recognize

SARS-CoV-2 antigens *in vitro* was validated and then similar analyses were performed for

selected CD4+ and CD8+ T cells. Briefly, MoDC (2×10^4) were pulsed with SARS-CoV-2 peptide

antigens (1μg/ml, PepMix S comprising the S1 and S2 pools PM-WCPV-S-1; and N PM-WCPV-

NCAP-1, both JPT Peptide Technologies) then autologous T cells (1×10^5), enriched from the

non-adherent fraction of PBMC using MojoSort (BioLegend CD3 enrichment) were added in

enriched RPMI (10% human AB serum). Cells were cultured in a microtiter plate (Millipore)

containing an immobilized primary antibody to target IFN-γ, overnight (37°C), then IFN-γ spot

forming cells enumerated by ELISpot. For ELISpot detection, after aspiration and washing to

remove cells and media, IFN-γ was detected by a secondary antibody to cytokine conjugated to

biotin. A streptavidin/horseradish peroxidase conjugate was used detect the biotin-conjugated

secondary antibody. The number of spots per well (1 x 10^5 cells), was counted using an ELISpot

plate reader. IL-4 was measured by ELISpot using a kit (MabTech) with wells precoated with anti-

IL-4 antibody and following the manufacturer's instructions. Remaining steps for IL-4 detection

were identical to those for IFN-γ, but with alkaline phosphatase detection rather than peroxidase.

Determination of previously SARS-CoV-2 infected patient-derived T-cell reactivity in response to

autologous hAd5 vaccine-transduced MoDCs

MoDCs were transduced with hAd5 S-Fusion, S-Fusion + N-ETSD, N-ETSD, N or GFP/Null

constructs and incubated overnight at 37°C. The transduced MoDCs were cultured with CD3+,

CD4+, or CD8+ T-cells from the same individuals overnight. Antigen specific T-cell responses

were enumerated using ELISpot as described above.

Phase 1b Clinical Trail and Analyses

Vaccination of healthy adult subjects with hAd5-S-Fusion+N-ETSD.

The cGMP vaccine used in the clinical trial was prepared using the generated high titer

adenoviral stocks described above in *Production hAd5-S-Fusion+N-ETSD and related SARS-*

CoV-2 antigen expressing vectors.

In a phase 1b, open-label study (QUILT 4.001 in adults conducted at Hoag Hospital in

Orange County, California, USA, hAd5-S-Fusion+N-ETSD was administered to cohort 2, the

participants evaluated in the studies here, subcutaneously at a dosage of $1x10^{11}$ VP. An initial

priming dose was administered on Day 1.

Study participants comprised healthy adult volunteers between the ages of 18 and 55. Subjects

were randomized to cohorts described above. All the participants provided informed, written

consent before enrollment.

Trial Oversight

The trial protocol was approved by the Western IRB for the Chan Soon-Shiong Institute for

Medicine (CSSIFM) and the Hoag Hospital Providence IRB at Hoag Hospital Newport Beach.

ImmunityBio, Inc. was the regulatory sponsor of the trial and holder of the Investigational New

Drug (IND) application. The trial was funded by ImmunityBio, Inc.. The vaccine was designed by

ImmunityBio, Inc. and manufacture by ImmunityBio. *The studies presented here are experimental*

and all assays and analyses were performed in-house at ImmunityBio, Inc. Formal analyses of the

primary and secondary end points are not reported in this manuscript.

T-cell analyses

T-cell responses in PBMCs were measured at baseline (Day 1) and after prime vaccination

alone on Day 14-16 and 21-23 by ELISpot for secretion of interferon γ (IFNγ) and interleukin 4

(IFN-4). T helper cell type 1 dominance was determined based on the IFNγ/IL-4 ratio.

IFN-γ ELISpot for enumeration of S- and N-specific T cells in PBMC

ELISpot assays were used to enumerate S- and N-specific IFN-γ-secreting T cells in fresh

PBMCs isolated from the blood of subjects vaccinated with hAd5-S-Fusion+N-ETSD. Whole

blood was collected by venipuncture pre- and post-vaccination. PBMCs were isolated from whole

blood by standard density gradient centrifugation and frozen in liquid nitrogen until use. On the

day of assay, PBMCs were thawed and re-suspended in RPMI 10% human AB serum, then

incubated with 2 μg/ml of SARS-CoV-2 S or N peptide pools (JPT Peptide Technologies catalogue

PM-WCPV-S and PM-WCPV-NCAP-1, respectively). Internal positive controls were incubated

with 2 μg/ml of CEFT peptide pool consisting of epitopes derived from human Cytomegalovirus,

Epstein-Barr virus, Influenza A, and Clostridium tetani (JPT Peptide Technologies catalogue #

PM-CEFT-4). Positive assay controls were incubated with a CD3/CD28/CD2 agonist (StemCell

Technologies catalogue # 10970). Negative controls were incubated with media alone or 2 μg/ml

of SARS-CoV-2 membrane peptide pool (JPT Peptide Technologies catalogue # PM-WCPV-

VME). The number of spot-forming cells (SFCs) per well (2.5-4 x 10^5 PBMCs) were enumerated

using an ELISpot plate reader (Cellular Technologies Limited model # S6UNIV-01-7115). All

IFN-γ ELISpot reagents were obtained from a commercial kit and used according to manufacturer

protocol (Cellular Technologies Limited catalogue # Hu IFN-g/IL-4).

In silico T-Cell Epitope HLA Binding Prediction

SARS-CoV-2 reference sequences

The reference strain and COVID19 variants used for epitope analysis was assembled from

75,000 individual sequences deposited in the GISAID database. Binding analysis was run on the

consensus sequence generated using our in-house binding algorithm.[51] Briefly, machine learning

classifiers were trained on experimental binding data deposited in the IEDB database.[52]

The variants assessed included (name/mutation(s)): B.1.177/A222V mutation; B.1.1.7

/N501Y mutation; B.1.351 and B.1.1.28/K417N (or K417T for B.1.1.28), E484K, and the N501Y

mutations; and P.2/E484K. Binding efficiencies as well as HLA-epitope pairs were compared

between reference sequences and mutations to determine potential immune evasion due to the

reported variants.

REFERENCES

1   Gabitzsch, E. *et al.* Complete Protection of Nasal and Lung Airways Against SARS-
CoV-2 Challenge by Antibody Plus Th1 Dominant N- and S-Specific T-Cell Responses
to Subcutaneous Prime and Thermally-Stable Oral Boost Bivalent hAd5 Vaccination in
an NHP Study. *bioRxiv*, 2020.2012.2008.416297, doi:10.1101/2020.12.08.416297
(2021).
2   Amalfitano, A. Next-generation adenoviral vectors: new and improved. *Gene Therapy* **6**,
1643-1645, doi:10.1038/sj.gt.3301027 (1999).
3   Gabitzsch, E. S. & Jones, F. R. New recombinant SAd5 vector overcomes Ad5 immunity
allowing for multiple safe, homologous, immunizations. *J Clin Cell Immunol* **S4**, 001,
doi:doi:10.4172/2155-9899 (2011).

4   Gabitzsch, E. S. *et al.* Anti-tumor immunotherapy despite immunity to adenovirus using a
novel adenoviral vector Ad5 [E1-, E2b-]-CEA. *Cancer Immunology, Immunotherapy* **59**,
1131-1135, doi:10.1007/s00262-010-0847-8 (2010).

5   Gatti-Mays, M. E. *et al.* A Phase I Trial Using a Multitargeted Recombinant Adenovirus
5 (CEA/MUC1/Brachyury)-Based Immunotherapy Vaccine Regimen in Patients with
Advanced Cancer. *Oncologist* **25**, 479, doi:10.1634/theoncologist.2019-0608 (2019).

6   Hellerstein, M. What are the roles of antibodies versus a durable, high quality T-cell
response in protective immunity against SARS-CoV-2? *Vaccine X* **6**, 100076,
doi:10.1016/j.jvacx.2020.100076 (2020).

7   Redd, A. D. *et al.* CD8+ T cell responses in COVID-19 convalescent individuals target
conserved epitopes from multiple prominent SARS-CoV-2 circulating variants. *Open
Forum Infect Dis*, doi:10.1093/ofid/ofab143 (2021).

8   NIH. T cells recognize recent SARS-CoV-2 variants. *News Release*
https://www.nih.gov/news-events/news-releases/t-cells-recognize-recent-sars-cov-2-
variants (2021).

9   Tarke, A. *et al.* Negligible impact of SARS-CoV-2 variants on CD4+ and CD8+ T cell
reactivity in COVID-19 exposed donors and vaccinees. *bioRxiv*, 2021.2002.2027.433180,
doi:10.1101/2021.02.27.433180 (2021).

10   Baden, L. R. *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N
Engl J Med* **384**, 403-416, doi:10.1056/NEJMoa2035389 (2021).

11   Polack, F. P. *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N
Engl J Med* **383**, 2603-2615, doi:10.1056/NEJMoa2034577 (2020).

12   Sadoff, J. *et al.* Interim Results of a Phase 1-2a Trial of Ad26.COV2.S Covid-19
Vaccine. *N Engl J Med*, doi:10.1056/NEJMoa2034201 (2021).

13   Ewer, K. J. *et al.* T cell and antibody responses induced by a single dose of ChAdOx1
nCoV-19 (AZD1222) vaccine in a phase 1/2 clinical trial. *Nat Med* **27**, 270-278,
doi:10.1038/s41591-020-01194-5 (2021).

14   Keech, C. *et al.* Phase 1-2 Trial of a SARS-CoV-2 Recombinant Spike Protein
Nanoparticle Vaccine. *N Engl J Med* **383**, 2320-2332, doi:10.1056/NEJMoa2026920
(2020).

15   Anderson, E. J. *et al.* Safety and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine
in Older Adults. *N Engl J Med* **383**, 2427-2438, doi:10.1056/NEJMoa2028436 (2020).

16   Sahin, U. *et al.* COVID-19 vaccine BNT162b1 elicits human antibody and T(H)1 T cell
responses. *Nature* **586**, 594-599, doi:10.1038/s41586-020-2814-7 (2020).

17   Karim, S. S. A. Vaccines and SARS-CoV-2 variants: the urgent need for a correlate of
protection. *The Lancet*, doi:10.1016/S0140-6736(21)00468-2.

18   Madhi, S. A. *et al.* Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine against the
B.1.351 Variant. *New England Journal of Medicine*, doi:10.1056/NEJMoa2102214
(2021).

19   Shinde, V. *et al.* Preliminary Efficacy of the NVX-CoV2373 Covid-19 Vaccine Against
the B.1.351 Variant. *medRxiv*, 2021.2002.2025.21252477,
doi:10.1101/2021.02.25.21252477 (2021).

20   Novavax. Novavax COVID-19 Vaccine Demonstrates 89.3% Efficacy in UK Phase 3
Trial. (2021).

21   Johnson, J. Johnson & Johnson Announces Single-Shot Janssen COVID-19 Vaccine
Candidate Met Primary Endpoints in Interim Analysis of its Phase 3 ENSEMBLE Trial.
(2021).

22 Edara, V. V. *et al.* Reduced binding and neutralization of infection- and vaccine-induced
antibodies to the B.1.351 (South African) SARS-CoV-2 variant. *bioRxiv*,
doi:10.1101/2021.02.20.432046 (2021).
23 Wang, Z. *et al.* mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating
variants. *Nature* **Preview**, doi:10.1038/s41586-021-03324-6 (2021).
24 Wu, K. *et al.* mRNA-1273 vaccine induces neutralizing antibodies against spike mutants
from global SARS-CoV-2 variants. *bioRxiv*, doi:10.1101/2021.01.25.427948 (2021).
25 Azizi, A. *et al.* A combined nucleocapsid vaccine induces vigorous SARS-CD8+ T-cell
immune responses. *Genet Vaccines Ther* **3**, 7, doi:10.1186/1479-0556-3-7 (2005).
26 Shang, B. *et al.* Characterization and application of monoclonal antibodies against N
protein of SARS-coronavirus. *Biochemical and Biophysical Research Communications*
336, 110-117, doi:https://doi.org/10.1016/j.bbrc.2005.08.032 (2005).
27 Peng, H. *et al.* Long-lived memory T lymphocyte responses against SARS coronavirus
nucleocapsid protein in SARS-recovered patients. *Virology* **351**, 466-475,
doi:https://doi.org/10.1016/j.virol.2006.03.036 (2006).
28 Narayanan, K., Chen, C.-J., Maeda, J. & Makino, S. Nucleocapsid-independent specific
viral RNA packaging via viral envelope protein and viral RNA signal. *Journal of
virology* **77**, 2922-2927, doi:10.1128/jvi.77.5.2922-2927.2003 (2003).
29 Zeng, W. *et al.* Biochemical characterization of SARS-CoV-2 nucleocapsid protein.
Biochemical and biophysical research communications, S0006-0291X(0020)30876-
30877, doi:10.1016/j.bbrc.2020.04.136 (2020).
30 Grifoni, A. *et al.* Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans
with COVID-19 Disease and Unexposed Individuals. *Cell* **181**, 1489,
doi:10.1016/j.cell.2020.05.015 (2020).
31 Le Bert, N. *et al.* SARS-CoV-2-specific T cell immunity in cases of COVID-19 and
SARS, and uninfected controls. *Nature* **584**, 457, doi:10.1038/s41586-020-2550-z (2020).
32 Sekine, T. *et al.* Robust T cell immunity in convalescent individuals with asymptomatic
or mild COVID-19. *Cell* **183**, 158, doi:10.1101/2020.06.29.174888 (2020).
33 Altmann, D. M. & Boyton, R. J. SARS-CoV-2 T cell immunity: Specificity, function,
durability, and role in protection. *Sci Immunol* **5**, eabd6160,
doi:10.1126/sciimmunol.abd6160 (2020).
34 Chen, Z. & John Wherry, E. T cell responses in patients with COVID-19. *Nat Rev
Immunol*, 1-8, doi:10.1038/s41577-020-0402-6 (2020).
35 Matloubian, M., Concepcion, R. J. & Ahmed, R. CD4+ T cells are required to sustain
CD8+ cytotoxic T-cell responses during chronic viral infection. *Journal of virology* **68**,
8056-8063, doi:10.1128/jvi.68.12.8056-8063.1994 (1994).
36 Laidlaw, B. J., Craft, J. E. & Kaech, S. M. The multifaceted role of CD4(+) T cells in
CD8(+) T cell memory. *Nat Rev Immunol* **16**, 102-111, doi:10.1038/nri.2015.10 (2016).
37 Nakamura, S. & Yoshimori, T. New insights into autophagosome-lysosome fusion. *J Cell
Sci* **130**, 1209-1216, doi:10.1242/jcs.196352 (2017).
38 Lőrincz, P. & Juhász, G. Autophagosome-Lysosome Fusion. *J Mol Biol* **432**, 2462-2482,
doi:10.1016/j.jmb.2019.10.028 (2020).
39 Kabeya, Y. *et al.* LC3, a mammalian homologue of yeast Apg8p, is localized in
autophagosome membranes after processing. *Embo j* **19**, 5720-5728,
doi:10.1093/emboj/19.21.5720 (2000).
40 Schmid, D., Pypaert, M. & Münz, C. Antigen-loading compartments for major
histocompatibility complex class II molecules continuously receive input from
autophagosomes. *Immunity* **26**, 79-92, doi:10.1016/j.immuni.2006.10.018 (2007).

41   Patterson, N. L. & Mintern, J. D. Intersection of autophagy with pathways of antigen
presentation. *Protein & Cell* **3**, 911-920, doi:10.1007/s13238-012-2097-3 (2012).
42   Crotzer, V. L. & Blum, J. S. Autophagy and Its Role in MHC-Mediated Antigen
Presentation. *The Journal of Immunology* **182**, 3335, doi:10.4049/jimmunol.0803458
(2009).
43   You, L. *et al.* The crosstalk between autophagic and endo-/exosomal pathways in antigen
processing for MHC presentation in anticancer T cell immune responses. *Journal of
Hematology & Oncology* **10**, 165, doi:10.1186/s13045-017-0534-8 (2017).
44   Tan, C. W. *et al.* A SARS-CoV-2 surrogate virus neutralization test based on antibody-
mediated blockage of ACE2-spike protein-protein interaction. *Nat Biotechnol* **38**, 1073-
1078, doi:10.1038/s41587-020-0631-z (2020).
45   GISAID. SARS-CoV-2 Lineages. (2020-2021).
46   Peng, Y. *et al.* Broad and strong memory CD4(+) and CD8(+) T cells induced by SARS-
CoV-2 in UK convalescent individuals following COVID-19. *Nat Immunol* **21**, 1336-
1345, doi:10.1038/s41590-020-0782-6 (2020).
47   Rice, A. *et al.* The Dual-Antigen Ad5 COVID-19 Vaccine Delivered as an Intranasal Plus
Subcutaneous Prime Elicits Th1 Dominant T-Cell and Humoral Responses in CD-1 Mice.
bioRxiv, 2021.2003.2022.436476, doi:10.1101/2021.03.22.436476 (2021).
48   Tilocca, B. *et al.* Comparative computational analysis of SARS-CoV-2 nucleocapsid
protein epitopes in taxonomically related coronaviruses. *Microbes Infect*, S1286-
4579(1220)30053-30058, doi:10.1016/j.micinf.2020.04.002 (2020).
49   Amalfitano, A. *et al.* Production and Characterization of Improved Adenovirus Vectors
with the E1, E2b, and E3 Genes Deleted. *Journal of virology* **72**, 926,
doi:10.1128/JVI.72.2.926-933.1998 (1998).
50   Kasinrerk, W., Baumruker, T., Majdic, O., Knapp, W. & Stockinger, H. CD1 molecule
expression on human monocytes induced by granulocyte-macrophage colony-stimulating
factor. *The Journal of Immunology* **150**, 579-584 (1993).
51   Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative
contribution to global health. *Glob Chall* **1**, 33-46, doi:10.1002/gch2.1018 (2017).
52   Vita, R. *et al.* The Immune Epitope Database (IEDB): 2018 update. *Nucleic Acids Res* **47**,
D339-d343, doi:10.1093/nar/gky1006 (2019).
53   Tan, C. W. *et al.* A SARS-CoV-2 surrogate virus neutralization test based on antibody-
mediated blockage of ACE2-spike protein-protein interaction. *Nat Biotechnol* **38**, 1973,
doi:10.1038/s41587-020-0631-z (2020).
653

SUPPLEMENTARY MATERIALS for

Single Prime hAd5 Spike (S) + Nucleocapsid (N) Dual Antigen Vaccination of
Healthy Volunteers Induces a Ten-Fold Increase in Mean S- and N- T-Cell Responses
Equivalent to T-Cell Responses from Patients Previously Infected with SARS-CoV-2

Supplementary Materials

Methods

Quantification of anti-SARS-CoV-2 spike antibodies in plasma

The presence of anti-SARS-CoV-2 spike antigen antibodies in previously infected patient plasma used in studies here was validated. IgG against SARS-CoV-2 spike was detected in plasma using ELISA. Briefly, EIA/RIA plates were coated with a solution of purified recombinant SARS-CoV-2-derived Spike protein (1μg/ml, S-Fusion. ImmunityBio, Inc.) suspended in coating buffer (0.05 M carbonate-bicarbonate, pH 9.6) and incubated overnight at 4°C. Plates were washed three times with TPBS solution (PBS + 0.05% Tween 20). Blocking solution (2% non-fat milk in TPBS) was added and incubated at room temperature (RT, 1h). Serial dilutions of plasma were prepared in 1% non-fat milk in TPBS. Plates were washed as described above and serial dilutions of plasma were added to the plate and incubated at RT (1h). Plates were washed three times. Goat anti-Human IgG (H+L) HRP-conjugated secondary antibody (1:6000 dilution) prepared in 1% non-fat milk/TPBS was added and incubated at RT (1h). Plates were washed three times. Substrate (3,3',5,5'-tetramethylbenzidine (TMB)) was added to each well and incubated at RT (10min). The reaction was stopped by addition of sulfuric acid (1N H_2SO_4). The optical density (450 nm) was measured by a Synergy 2 (BioTek Instruments, Inc.) plate reader. Data were analyzed using Prism 8 (GraphPad Software, LLC).

cPass^{TM} neutralizing antibody detection

The GenScript cPass^{TM} (https://www.genscript.com/cpass-sars-cov-2-neutralization-antibody-detection-Kit.html) kit for detection of neutralizing antibodies was used according to the manufacturer's instructions.[44] The kit detects circulating neutralizing antibodies against SARS-CoV-2 that block the interaction between the S RBD with the ACE2 cell surface receptor. It is suitable for all antibody isotypes.

To evaluate the levels of anti-SARS-CoV-2 neutralizing antibodies in plasma, dilutions of plasma were incubated with horseradish peroxidase-conjugated spike RBD (37°C, 30min). The RBD-plasma mixture was added to a microtiter plate coated with ACE2 and incubated at 37°C (15 min). Plates were washed and substrate (TMB) added at room temperature (15 min). The reaction was stopped and plates read on a plate reader at 450 nm.

Live virus assay of plasma neutralization of infection

The ability of previously SARS-CoV-2 infected patient plasma used in studies here to neutralize SARS-CoV-2 infection *in vitro* was also validated. All aspects of the assay utilizing virus were performed in a BSL3 containment facility according to the ISMMS Conventional Biocontainment Facility SOPs for SARS-CoV-2 cell culture studies. Vero E6 kidney epithelial cells from *Cercopithecus aethiops* (ATCC CRL-1586) were plated (20,000 cells/well) in a 96-well

format and 24 hours later, cells were incubated with antibodies or heat inactivated plasma previously serially diluted in 3-fold steps in DMEM containing 2% FBS, 1% NEAAs, and 1% Pen-Strep; the diluted samples were mixed 1:1 with SARS-CoV-2 in DMEM containing 2% FBS, 1% NEAAs, and 1% Pen-Strep at 10,000 TCID 50/mL for 1 hr at 37°C, 5% CO_2. This incubation did not include cells to allow for neutralizing activity to occur prior to infection. For detection of neutralization, the virus/sample mixture (120 µL) was transferred to the Vero E6 cells and incubated (48 hours) before fixation with 4% PFA. Each well received virus (60 µl) or an infectious dose of 600 TCID50. Control wells, including six wells on each plate for no virus and virus-only controls, were used. The percent neutralization was calculated as 100-((sample of interest-[average of "no virus"])/[average of "virus only"])*100) with a stain for CoV-2 Np imaged on a Celigo Imaging Cytometer (Nexcelom Bioscience).

Transfection of HEK 293T cells and detection of plasma antibody binding

To assess binding of plasma from a previously infected patient to antigens expressed by hAd5 S-Fusion and hAd5 S-Fusion + N-ETSD vaccines, plasma was incubated with construct-transfected HEK 293T cells and binding determined by flow cytometry. HEK 293T cells (2.5 x 10^5 cells/well in 24 well plates) were grown in DMEM (Gibco) with 10% FBS and 1X PSA (100 units/mL penicillin, 100 µg/mL streptomycin, 0.25 µg/mL Amphotericin B) at 37°C. Cells were either left untransfected or were transfected with 0.5 µg of S-Fusion or S-Fusion + N-ETSD hAd5 plasmid DNA using a JetPrime transfection reagent (Polyplus) according to the manufacturer's instructions. Twenty-four hours later, cells were incubated for 30 min. with previously infected patient or healthy (unexposed, UnEx) plasma that had been serially diluted 10-fold for 30 min. Plasma IgG was labeled using a goat anti-human IgG-phycoerythrin conjugated and labeled cells were acquired using the Thermo-Fisher Attune NxT flow cytometer and analyzed using FlowJo Software to determine Mean Fluorescence Intensity (MFI) values of both the untransfected and transfected cells. Results were graphed as the difference in MFI between untransfected and transfected cells for the S-Fusion and S-Fusion + N-ETSD constructs over the plasma dilutions.

Results

Previously SARS-CoV-2-infected patient plasma confirmed to have neutralizing anti-S antibodies

For the studies described below, plasma samples were collected from four individuals convalescing from SARS-CoV-2 infection as confirmed by antibody assays and patient history as described Methods. The presence of anti-Spike IgG, and of neutralizing antibodies by both a surrogate SARS-CoV-1 neutralization assay [53] and live virus assays, were confirmed in all patient samples (Suppl. Fig. S1**).** Samples were also collected from four virus-naïve individuals and were used as controls.

750   **Fig. S1** *Plasma samples from patients (Pt) previously infected with SARS-CoV-2 contain*
751   *neutralizing antibodies.* (A) The levels of anti-spike IgG as determined by OD at 450 nm are
752   shown. (B) All four Pt plasma samples contained antibodies that blocked RBD-ACE2 binding and
753   (C) contained SARS-CoV-2 neutralizing antibodies.
754   *Previously SARS-CoV-2-infected patient plasma binds to vaccine-transfected HEK-293T cells*
755
In additional studies to validate immune responses to SARS-CoV-2 antigens, the binding of
previously SARS-CoV-2 infected patient and virus-naïve control individual plasma to human
embryonic kidney (HEK) 293T cells transfected with either hAd5 S-Fusion alone or hAd5 S-
Fusion + N-ETSD was assessed (Suppl. Fig. S2). This binding reflects the presence of antibodies
in plasma that recognize antigens expressed by the hAd5 vectored vaccines. Quantification of
histograms showed little or no binding of virus-naïve plasma antibodies to cells expressing either
construct, and the highest binding of plasma antibodies from a previously SARS-CoV-2 infected
patient to cells expressing the dual antigen S-Fusion + N-ETSD construct (Suppl. Fig. S2R**).**

764
Fig. S2 *Plasma from previously SARS-CoV-2 infected patients shows greater binding to S-Fusion*
+ N-ETSD surface antigens compared to S-Fusion alone after transfection of HEK 293T cells.
HEK 293T cells that were either uninfected (pink) or infected (blue) with S-Fusion alone or
bivalent S-Fusion + N-ETSD were exposed to plasma from either unexposed (UnEx) controls or
previously infected SARS-CoV-2 patients (Pt). Flow histograms for (A-D) UnEx with S-Fusion;
(E-H) UnEx with S-Fusion + N-ETSD; (I-L) Pt with S-Fusion; and (M-P) Pt with S-Fusion + N-
ETSD, all n = 4, are shown. When the ΔMFI (difference in binding to transfected versus
untransfected cells) is graphed, it is apparent that compared to (Q) S-Fusion, (R) binding of Pt
plasma to S-Fusion + N-ETSD transfected cells is much higher and that plasma from UnEx
individuals shows little binding.
775
776   *Previously SARS-CoV-2-infected patient T cells are activated by S and N peptide pools*
777
778   The reactivities of T cells from the previously SARS-CoV-2-infected patients was confirmed
779   by incubation with S1 (containing the spike receptor binding domain), S2 and N peptides pools

780   (Fig. S3A). The specific reactivities of CD4+ (Fig. S3B and C) and CD8+ T cells was then
781   determined, showing CD8+ T cells were more selectively activated by the N peptide pool.
782

Fig. S3. *T-cell responses to MoDCs pulsed with SARS-CoV-2 peptides.* Unselected T cells from
all four previously SARS-CoV-2 infected patients (Pt) show significant IFN-γ responses to S1, S2,
and N peptide pool-pulsed MoDCs as compared to 'none'. T cells from virus-naïve (unexposed,
UnEx) control individuals showed far lower responses. Statistical analysis performed using One-
way ANOVA and Tukey's post-hoc analysis for samples from each patient compared only to
'none' where * $p \leq 0.05$, **$p<0.01$, ***$p<0.001$, and ****$p<0.0001$. Data graphed as the mean
and SEM; n = 3-4. Patient MoDCs pulsed with SARS-CoV-2 peptide mixes overnight were
incubated with autologous CD4+ (B, Pt 4; C, Pt 3) or CD8+ (D, Pt 4; E, Pt 3) T cells. IFN-γ levels
were determined by ELISpot. Statistical analysis performed using One-way ANOVA and
Dunnett's post-hoc multiple comparison analysis to compare each peptide pool to Veh (shown
above the bar) or between peptide pools (above line) where *$p<0.05$, **$p<0.01$ and
****$p<0.00001$. Data graphed as mean and SEM; n = 3.