

1 Article type: Original research reports

2

3 Article title: **Statistical Modeling of deaths due to COVID-19 influenced by social isolation**
4 **in Latin American countries**

5

6 Rafael André da Silva^{1*}, Luiz Philipe de Souza Ferreira², Jean Michel Rocha Sampaio Leite³,
7 Fernanda Assunção Tiraboschi², Thiago Maciel Valente², Vinicius Moraes de Paiva Roda¹,
8 Jeniffer Johana Duarte Sanchez⁴.

9

10 Affiliation

11

12 1. Institute of Biomedical Sciences, University of São Paulo (ICB/USP), São Paulo, SP, Brazil.

13 2. University of Fortaleza (UNIFOR), Fortaleza, CE, Brazil

14 3. Faculty of Public Health, University of Sao Paulo (FSP/USP), São Paulo, SP, Brazil.

15 4. Federal University of Ceará (UFC), Fortaleza, CE, Brazil.

16

17 *Corresponding author:

18

19 Rafael André da Silva, Institute of Biomedical Sciences, University of São Paulo (ICB/USP),
20 São Paulo, SP, Brazil. E-mail: rafael@usp.br .

21

22 Number of figures: 2

23 Number of tables: 1

24 Number of words: 1.786

25

26 **Abstract**

27

28 Social isolation is extremely important to minimize the effects of a pandemic. Latin American
29 (LA) countries have similar socioeconomic characteristics and health system infrastructures.
30 These countries face difficulties to deal with the COVID-19 pandemic and some of them had
31 very high death rates. Government stringency index (GSI) of twelve LA countries was gathered
32 from the Oxford COVID-19 Government Response Tracker (OxCGRT) project. GSI was
33 calculated considering nine metrics such as school and work closures, stay-at-home
34 requirements, among others types of social distancing and isolation measures. Population data
35 from the United Nations Population Fund and number of deaths data was collected from the
36 dashboard of the World Health Organization (WHO). We performed an analysis of the period
37 March-December using a mixed linear model approach. Peru, Brazil, Chile, Bolivia, Colombia,
38 Argentina and Ecuador had the highest death rates with an increasing trend over time, while
39 Suriname, Venezuela, Uruguay, Paraguay and Guyana had the lowest ones, which remained
40 steady. GSI in most countries followed the same pattern during the analyzed months. i.e., high
41 indices at the beginning of the pandemic and lower ones in the last evaluated months, while the
42 number of deaths increased over the whole period. Almost no country kept its GSI high for
43 much time, especially from October to December. Time and GSI as well as their interaction
44 were highly significant. As their interaction increases, death rate decreases. In conclusion, our
45 statistical model explains and substantiates the need for maintaining social distancing and
46 isolation measures over time during the pandemic.

47

48 **Keywords:** COVID-19, SARS-CoV-2, Coronavirus, Latin American countries, Social isolation,
49 Government stringency index

50

51 **Introduction**

52

53 The COVID-19 pandemic has affected healthcare systems and caused collapses across the
54 globe. In Latin America (LA), the first case of SARS-CoV-2 infection was recorded on

55 February 25th in the City of São Paulo. In less than a month after the first case, all LA countries
56 had confirmed cases of COVID-19^{1,2}.

57 The LA region has several obstacles that make it difficult for countries to take action
58 against the spread of the virus. Precarious conditions, such as poverty, lack of hospital
59 infrastructure, low sanitary conditions, high prevalence of chronic diseases and government's
60 tardy responses are factors that make it difficult to prevent contamination by the virus, so that
61 they facilitate transmission and directly impact the hospital system³⁻⁵. Through predictive
62 models' studies, it has been suggested that the virus could spread aggressively through LA^{6,7}.
63 Moreover, analyses of the initial cases of the COVID-19 pandemic in LA estimated an
64 unfavorable scenario for the countries, and also evidenced aggressive dynamics of the disease
65 outbreak in Brazil and Ecuador compared to Italy and Spain⁷. Above all, among the LA
66 countries, Brazil was considered a major epicenter of the disease⁸.

67 Although there are measures aimed at reducing the spread of the new coronavirus such
68 as social distancing, school closures, cancellation of public events and, sometimes, severe
69 methods such as lockdown, these measures have been relaxed, in addition to noncompliance by
70 the population and poor governmental management⁹. Considering that social distancing and
71 isolation are important protective measures for the containment of SARS-CoV-2 infection, and
72 that there is lack of studies demonstrating the relationship between the social isolation and death
73 rate due to COVID-19, based on the Government Stringency Index (GSI) from The Coronavirus
74 Government Response Tracker (OxCGRT)¹⁰, the objective of this work was to analyze the
75 relationship between GSI and time, and the death rate from COVID-19 in 12 LA countries,
76 using a mixed linear model approach. Through this model, the need for maintaining social
77 distancing and isolation measures over time during the pandemic is explained and substantiated.

78

79 **Material and Methods**

80 **Data Sources**

81 Government stringency index of twelve LA countries was developed by the Oxford University
82 and gathered from the platform Policy Responses to the Coronavirus Pandemic on the website
83 (<https://ourworldindata.org/policy-responses-covid>). This index represents the strictness of
84 government policies and was calculated considering nine metrics of social distancing and
85 isolation, such as school and work closures, stay-at-home requirements, transport restrictions,
86 restrictions on public gatherings, cancellation of public events, public information campaigns;
87 restrictions on internal movements; and international travel controls. The index on any given
88 day is calculated as the mean score of nine policy measures, each taking a value between 0 and
89 100. The detailed methodology for calculating indices is described elsewhere
90 ([https://github.com/OxCGRT/covid-policy-
91 tracker/blob/master/documentation/index_methodology.md](https://github.com/OxCGRT/covid-policy-tracker/blob/master/documentation/index_methodology.md)). To use that data, we first
92 calculated the mean of the GSI for each month. The number of cumulative monthly deaths data
93 was collected from the dashboard of the World Health Organization (WHO). Population data
94 were obtained from the United Nations Population Fund available on
95 (<https://www.unfpa.org/about-us>) and used to calculate the proportion of the number of deaths
96 for each country. This data was collected from the first day of March until the last day of
97 December of 2020.

98

99 **Statistical analysis**

100

101 We evaluated the relationship between GSI and time, and the death rates from COVID-19.
102 Since we analyzed repeated measures over a period of time (time was uniformly measured
103 across all countries), this is a longitudinal analysis requiring a mixed linear model approach,

104

$$y_{ij} = \mu + X_{ij}^T \beta + Z\gamma + \varepsilon_{ij},$$

105

where

$$y_{ij} = \frac{N^{\circ} \text{ of deaths related to COVID - 19}}{N^{\circ} \text{ of inhabitants in the country}} \times 1000000$$

106

107 μ = mean of the death ratio adjusted by total population for each million inhabitants,
108 X_{ij} = vector of covariates, β = vector of the regression parameters for the covariates. Z = matrix
109 of covariates, γ = vector of random effects and ε_{ij} = vector of random errors. γ and ε are
110 uncorrelated.

111 In the model, the variables time (month) and GSI were considered both as fixed. In
112 addition, the country was included as a random effect. All statistical analysis was performed
113 under the most commonly used significance levels (1%, 5% and 10%) using the RStudio
114 statistical software v.3.6.

115

116 Results

117

118 We analyzed data of deaths related to COVID-19 in twelve LA countries in order to evaluate the
119 relationship between death rates, government stringency index and time progression. In this
120 context, time and isolation index are useful to explain the dispersion of the data.

121 Figure 1a displays death rates from March to December. Peru, Brazil, Chile, Bolivia,
122 Colombia, Argentina and Ecuador have high death rates, with an increasing trend over time,
123 while Suriname, Venezuela, Uruguay, Paraguay and Guyana presented low death rates, which
124 remained stable. Figure 1b displays GSI from March to December. It is noticeable that there
125 was much fluctuation in the GSI for most countries, but with a large decrease from October to
126 December. The only country whose GSI was maintained high for the whole period was
127 Venezuela.

128 Also, in Figure 1a, it is possible to observe the asymmetry of the data, so a skew- t
129 distribution was adopted for the model's error. Since the model presents a variable dispersion,
130 we used a linear regression model for the dispersion.

131

132

$$y_{ij}|time_i, country_j, GSI_{ij} \sim ST(\mu_{ij}, \sigma_{ij}, \nu, \tau)$$
$$\mu_{ij}^{-1} = \beta_0 + \beta_1 time_i + \beta_2 GSI_{ij} + \beta_3 time_i \times GSI_{ij} + \gamma_0 + \gamma_1 country_j$$
$$\log(\sigma_{ij}) = \alpha_0 + \alpha_1 time_i + \alpha_2 GSI_{ij}.$$

133

134 where $i = 1, \dots, 10$ represents each of the months, starting from March, and $j =$
135 $1, \dots, 12$ corresponds to each of the countries. $time_i$ represents the i -th month, $country_j$
136 represents the j -th country, GSI_{ij} represents the GSI in the i -th month in the j -th country. σ_{ij}
137 corresponds to the standard deviation in the i -th month in the j -th country, with its
138 corresponding parameters α .

139

140 Table 1 displays the coefficient estimates of the mixed linear model and the
141 dispersion model. Time and GSI, as well as their interaction, were highly significant to explain
142 the rate death under all assumed significance levels. As time progresses, the death rate per
143 million inhabitants increases. In addition, as the interaction of time and GSI increases, the death
144 rate decreases.

145 In Figure 2a, the QQ-plot envelope shows there is no evidence that the skew- t
146 distribution is inappropriate to explain the death rate for each million inhabitants. Other aspects
147 of the model were analyzed by the quantile residuals (Figure 2b), such as the correct
148 specification of the model's dispersion and distribution. We can conclude from these graphs that
149 the model satisfies the assumptions so that the model specification is appropriate.

150

151 Discussion

152

153 In the COVID-19 pandemic, prolonged periods of social isolation were adopted across
154 the globe, as recommended by the WHO. Social isolation can have a dual impact. It has been
155 observed that, in this period, there is an increase in the rate of suicides, mental disorders, and
156 depression, which are explained by human hyper sociability. In addition, it can trigger physical
157 effects that impact children, young people and the elderly^{11,12}. In contrast, we know that social

158 isolation is extremely important to decrease the spread of the SARS-CoV-2 virus. It is important
159 to note that, during the pandemic caused by Influenza A (H1N1), social isolation was also
160 adopted¹³. Mortality from influenza and pneumonia during the 1918-1919 pandemic was lower
161 in civilians in rural areas when compared to those in urban areas. These observations have led to
162 the planning of strategies for pandemics, suggesting that social distancing interventions have a
163 potential effect on mortality by reducing the number of deaths¹⁴. In addition to social distancing
164 and isolation, large-scale testing is fundamental to fight against the pandemic. However,
165 addressing the influence of this factor on death rates remains a big challenge because countries
166 publish their testing data at different time points: some provide daily updates, while others
167 provide only on a weekly-basis, and some only publish figures on an ad-hoc basis at longer
168 intervals.

169 Based on GSI data extracted from the OxCGRT project¹⁰, it is possible to propose
170 statistical models to evaluate how closely these variables are related to time. Herein, the model
171 shows that the relationship between time and GSI is highly significant. When analyzing time
172 and GSI together, it was observed that, as the interaction of these two variables increases, a drop
173 in the death rate is detected. For instance, according to this model, with a GSI set to 0 in March
174 and a GSI set to 80 in April. i.e. a 80% increase in the strictness of government policies, we
175 could have observed a reduction of approximately 32 deaths per million inhabitants. In Brazil,
176 whose population is near 212 million, that would represent 6,784 lives that could have been
177 saved at the beginning of the pandemic. This figure would be even higher in other months
178 within the analyzed period. Furthermore, once these two variables were analyzed separately, as
179 time increases, the rate of deaths per million inhabitants increases as well. Surprisingly, the
180 same happens with the GSI reported by the countries. In light of these observations, we can
181 make two hypotheses: 1) GSI alone may not entirely represent the reality regarding social
182 isolation and the death rate from COVID-19, since this condition depends on other factors, such
183 as the infrastructure of the countries' public hospitals, government management, and the
184 population's compliance with the rules. 2) Restriction policies as measured by GSI do not have
185 immediate effects and must be maintained over longer periods in order to decrease death rates
186 by COVID-19. Hence, the problem is complex and deserves to be studied in detail taking into
187 account other aspects that may be influencing the death rates.

188 LA countries present problematic issues, such as social inequality and less access to
189 healthcare. However, complying with social isolation is difficult for individuals when work is
190 the only source of income¹⁵. In our analyses, countries such as Peru, Brazil, Chile, Bolivia,
191 Colombia, Argentina and Ecuador have the highest death rates. Peru presents low conditions to
192 face a pandemic and even in lockdown at the beginning of the pandemic², it presented high
193 death rates. In a prediction study with data from the first 10 days of the pandemic, it was
194 estimated that Peru had the lowest effective reproductive number (R_e), parameter used to keep
195 track of epidemics⁷, therefore, the country had these numbers accentuated during the pandemic
196 period.

197 Brazil was the first LA country to report cases of COVID-19¹⁶⁻¹⁸. Since it presents
198 favorable conditions to face a pandemic when compared to other LA countries, it was expected
199 to have lower rates. However, in the present study, Brazil and Chile presented higher death
200 rates, followed by Peru. It is important to consider that, although Peru's president has played a
201 germane role helping to control the number of deaths from COVID-19, there has not been
202 neither a national strategic plan to guide communication and educational health policies nor a
203 large-scale awareness campaign to stimulate people to protect their health and abide to the
204 protective measures. This lack of policies is also a current problem in Brazil². For instance,
205 through GSI e and COVID-19 Community Mobility Reports from Google, daily new cases and
206 real time R_e were calculated, showing that Brazil is not doing very well regarding its response to
207 COVID-19 pandemic¹⁹. Although Brazil presents a robust public health system and reasonable
208 GSI, the high death rates may be deeply connected to inadequate policy management that has
209 received several criticisms^{5,15}. In comparison to Brazil, Suriname had a similar GSI but a low
210 death rate, remaining stable over time. Except for Venezuela, no other country kept a high GSI
211 for longer periods. Particularly, GSI decreased in the end of the year (October-December) in
212 most countries, while death rates increased. On the other hand, isolation in Venezuela was

213 maintained even in December (an atypical month because of Christmas and holiday season), and
214 its death rates were low and remained unchanged over time.

215 According to the present analysis, Uruguay followed a relatively lower GSI than other
216 LA countries but showed low death rates. Uruguay was a country that acted quickly, closing its
217 borders and schools, with insertion of screening tests, reducing SARS-CoV-2 infections and
218 controlling the outbreak very efficiently²⁰. In contrast, Ecuador started with high social
219 isolation, but a decrease in the isolation rate was observed later. On the other hand, Ecuador had
220 a high mortality rate, which is accentuated over time even with the adoption of lockdown. In
221 addition, it should be noted that this country had poor conditions of public health infrastructure
222 at the beginning of the pandemic². At the beginning of the COVID-19 pandemic, it had been
223 already suggested that closing public transportation, work places and schools is particularly
224 effective in reducing COVID-19 transmission²¹.

225 The rapidly evolving pandemic in LA countries is worthy of especial attention,
226 considering their often weak and low stringency responses to the current sanitary crisis. In this
227 study, GSI varies considerably in all LA countries over time. This variation can partially explain
228 why these countries have been differently impacted by COVID-19. In spite of not specifically
229 addressing and discussing the government policies adopted by each country, in this
230 investigation, we successfully show that social distancing and isolation measured by GSI
231 influences death rates from COVID-19 over time. For instance, the interaction between GSI and
232 time can decrease the number of deaths, which demonstrates the importance of maintaining
233 social distancing and isolation measures for longer periods, as opposed to what most LA
234 countries did. Almost no country kept its GSI high for much time, especially from October to
235 December.

236 Our results have significant implications; however, some limiting aspects must be
237 considered. 1) The GSI was extracted from the OxCGRT project. The curators of this database
238 emphasized how challenging the collection of information on the exact data was due to the
239 nature and extent of the policies of the different governments. This complex data set can
240 obscure the qualitative differences in each of the nine metrics GSI measures across countries. In
241 addition, many local and cultural factors can affect the implementation of interventions. 2) Our
242 data provide a general interpretation of the influence of time and GSI on death rates in LA.
243 Therefore, future studies can deepen the search for more specific interpretations for each
244 country, taking into account local aspects and other metrics not covered here. 3) The numbers of
245 deaths from COVID-19 can be easily underreported²²⁻²³, this is due to limited testing, problems
246 in determining the cause of death and the way in which COVID-19 deaths are recorded. Hence,
247 we cannot define the real impact of the GSI on death rates with perfect precision. 4) We know
248 that the differences in population size between countries are often large, and the COVID-19
249 death count in more populous countries tends to be greater. Thus, in order to perform a more
250 truthful comparison, we used the cumulative death data and calculated the death rate adjusted by
251 the population of each country.

252

253 **Conclusions**

254

255 We conclude that, in combination, time and GSI have beneficial effects on the decrease of death
256 rates from COVID-19 in LA countries. Higher strictness of social distancing and isolation, as
257 measured by the GSI, over time could have flattened mortality curves from COVID-19 from
258 March to December 2020. Our statistical model explains and substantiates the need for
259 maintaining social distancing and isolation measures over time during the pandemic.

260

261 **Acknowledgments**

262 Publication charges for this article were waived due to the ongoing COVID-19 pandemic.

263 **Authors' addresses**

264 Rafael André da Silva and Vinicius Moraes de Paiva Roda, Institute of Biomedical Sciences,
265 University of São Paulo (ICB/USP), São Paulo, SP, Brazil, E-mails: rafaels@usp.br,

266 vinicius.moraes95@gmail.com. Luiz Philipe de Souza Ferreira, Thiago Maciel Valente and
267 Fernanda Assunção Tiraboschi, University of Fortaleza (UNIFOR), Fortaleza, CE, Brazil, E-
268 mails: philipedesouza@edu.unifor.br, maciel.thiago@edu.unifor.br, tiraboschi@edu.unifor.br.
269 Jean Michel Rocha Sampaio Leite, Faculty of Public Health, University of Sao Paulo
270 (FSP/USP), São Paulo, SP, Brazil, E-mail: jmichelsampaio@hotmail.com. Jeniffer Johana
271 Duarte Sanchez, Federal University of Ceará (UFC), Fortaleza, CE, Brazil, E-mail:
272 jjduartes@dema.ufc.br.

273

274 **Authors' contributions**

275 Rafael André da Silva: Conceptualization, Methodology, Data curation, interpretation, Writing -
276 original draft, Supervision, critical review, Project administration. Luiz Philipe de Souza
277 Ferreira: Writing - original draft, Design of the study, Manuscript drafting, review & editing.
278 Jean Michel Rocha Sampaio Leite: Methodology, Data curation, Critical review, Interpretation,
279 Writing - review & editing. Fernanda Assunção Tiraboschi: Writing - review & editing. Thiago
280 Maciel Valente: Data curation, Writing - original draft. Vinicius Moraes de Paiva Roda: Writing
281 - original draft, critical review. Jeniffer Johana Duarte Sanchez: Conceptualization,
282 Methodology, Formal analysis, Data curation, interpretation, Validation, Supervision.

283


284 **References**

285

- 286 1. Rodriguez-Morales AJ, Gallego V, Escalera-Antezana JP, et al. COVID-19 in Latin
287 America: The implications of the first confirmed case in Brazil. *Travel Medicine and*
288 *Infectious Disease* 2020; 35. doi:10.1016/j.tmaid.2020.101613
- 289 2. Garcia PJ, Alarcón A, Bayer A, et al. COVID-19 Response in Latin America. *American*
290 *Journal of Tropical Medicine and Hygiene* 2020;103(5):1765-1772.
291 doi:10.4269/ajtmh.20-0765
- 292 3. Miller MJ, Loaiza JR, Takyar A, et al. Covid-19 in latin america: Novel transmission
293 dynamics for a global pandemic? *PLoS Neglected Tropical Diseases* 2020;14(5):3-7.
294 doi:10.1371/JOURNAL.PNTD.0008265
- 295 4. Pablos-Méndez A, Vega J, Aranguren FP, et al. Covid-19 in Latin America. *BMJ*
296 2020;369-370. doi:10.1136/bmj.m2939
- 297 5. Burky T. COVID-19 in Latin America Several. *The Lancet Infectious Diseases* 2020; 20
298 (5):547-548. doi: 10.1016/S1473-3099(20)30303-0
- 299 6. Pacheco-Barrios K, Cardenas-Rojas A, Giannoni-Luza S, et al. COVID-19 pandemic
300 and Farr's law: A global comparison and prediction of outbreak acceleration and
301 deceleration rates. *PLoS ONE* 2020;15(9 September):1-25.
302 doi:10.1371/journal.pone.0239175
- 303 7. Caicedo-ochoa Y, Rebellón-sánchez DE, Peñaloza-rallón M. Effective Reproductive
304 Number estimation for initial stage of COVID-19 pandemic in Latin American
305 Countries. *International Journal of Infectious Diseases* 2020; 95(January):316-318.
306 <https://doi.org/10.1016/j.ijid.2020.04.069>
- 307 8. Bermudi PMM, Lorenz C, Aguiar BS de, et al. Spatiotemporal ecological study of
308 COVID-19 mortality in the city of São Paulo, Brazil: Shifting of the high mortality risk
309 from areas with the best to those with the worst socio-economic conditions. *Travel*
310 *Medicine and Infectious Disease* 2021;39(October 2020):101945.
311 doi:10.1016/j.tmaid.2020.101945
- 312 9. Chen YT, Yen YF, Yu SH, et al. An examination on the transmission of covid-19 and
313 the effect of response strategies: A comparative analysis. *International Journal of*
314 *Environmental Research and Public Health* 2020;17(16):1-14.
315 doi:10.3390/ijerph17165687
- 316 10. Hale T, Angrist N, Goldszmidt R, et al. A global panel database of pandemic policies
317 (Oxford COVID-19 Government Response Tracker). *Nature Human Behaviour* 2021.
318 doi:10.1038/s41562-021-01079-8
- 319 11. Bzdok D, Dunbar RIM. The Neurobiology of Social Distance. *Trends in Cognitive*

- 320 *Sciences* 2020;24(9):717-733. doi:10.1016/j.tics.2020.05.016
- 321 12. Sepúlveda-Loyola W, Rodríguez-Sánchez I, Pérez-Rodríguez P, et al. Impact of Social
322 Isolation Due to COVID-19 on Health in Older People: Mental and Physical Effects and
323 Recommendations. *Journal of Nutrition, Health and Aging* 2020;24(9):938-947.
324 doi:10.1007/s12603-020-1469-2
- 325 13. Ahmed F, Zviedrite N, Uzicanin A. Effectiveness of workplace social distancing
326 measures in reducing influenza transmission: A systematic review. *BMC Public Health*
327 2018;18(1):1-13. doi:10.1186/s12889-018-5446-1
- 328 14. Paynter S, Ware RS, Shanks GD. Host and environmental factors reducing mortality
329 during the 1918-1919 influenza pandemic. *Epidemiology and Infection*
330 2011;139(9):1425-1430. doi:10.1017/S0950268811000367
- 331 15. The Lancet. COVID-19 in Latin America: a humanitarian crisis. *The Lancet*
332 2020;396(10261):1463. doi:10.1016/S0140-6736(20)32328-X
- 333 16. Ferreira LPS, Valente TM, Tiraboschi FA, et al. Description of Covid-19 Cases in Brazil
334 and Italy. *SN Comprehensive Clinical Medicine* 2020: 497–500. doi: 10.1007/s42399-
335 020-00307-y
- 336 17. Valente TM, Ferreira LPS, Silva RAD, et al. Brazil Covid-19: Change of
337 hospitalizations and deaths due to burn injury? *Burns* 2020:9-11.
338 doi:10.1016/j.burns.2020.10.009
- 339 18. de Souza WM, Buss LF, Candido D da S, et al. Epidemiological and clinical
340 characteristics of the COVID-19 epidemic in Brazil. *Nature Human Behaviour*
341 2020;4(8):856-865. doi:10.1038/s41562-020-0928-4
- 342 19. Zhu D, Mishra SR, Han X, et al. Social distancing in Latin America during the COVID-
343 19 pandemic: an analysis using the Stringency Index and Google Community Mobility
344 Reports. *International Society of Travel Medicine* 2020:1-25. doi: 10.1093/jtm/taaa125
- 345 20. Elizondo V, Harkins GW, Mabvakure B, et al. SARS-CoV-2 genomic characterization
346 and clinical manifestation of the COVID-19 outbreak in Uruguay. *Emerging Microbes*
347 *and Infections* 2021;10(1):51-65. doi:10.1080/22221751.2020.1863747
- 348 21. Vannoni M, McKee M, Semenza JC, et al. Using volunteered geographic information to
349 assess mobility in the early phases of the COVID-19 pandemic: A cross-city time series
350 analysis of 41 cities in 22 countries from March 2nd to 26th 2020. *Globalization and*
351 *Health* 2020;16(1):1-9. doi:10.1186/s12992-020-00598-9
- 352 22. Alves THE, de Souza TA, Silva SA, Ramos NA, de Oliveira SV. Underreporting of
353 Death by COVID-19 in Brazil Second Most Populous State. *Frontiers in Public Health.*
354 2020;8:578645. doi:10.3389/fpubh.2020.578645
- 355 23. Karlinsky A, Kobak D. The World Mortality Dataset: Tracking excess mortality across
356 countries during the COVID-19 pandemic. Preprint *medRxiv*
357 2021;2021.01.27.21250604. doi:10.1101/2021.01.27.21250604
- 358


Figure 1. Death rates from COVID-19 and Government Stringency index from March to December 2020 in Latin American countries.



Legend Figure 1a: Death rates adjusted by population size for each country per million inhabitants. Figure 1b: Mean Government Stringency index reported as percentage.

Figure 2. Analysis of the model fit; a) Q-Q plot envelope. b) Residual analysis.

a)



b)




Table 1. Estimates of the dispersion model and mixed linear model for death rates from COVID-19 in 2020 in Latin American countries.

	Estimate	Std Error	<i>t</i> Value	<i>p</i> -value
μ				
Intercept (β_0)	4,91,633.0	69,250.3	7.099	<0.005***
Time (β_1)	-70,118.5	12,794.0	-5,.81	<0.005***
GSI(β_2)	-6,721.4	984.6	-6.826	<0.005***
Time×GSI (β_3)	953.6	189.2	5.041	<0.005***
σ				
Intercept (α_0)	-389	0.59	-6.63	<0.005***
Time(α_1)	0.72	0.04	16.67	<0.005***
GSI (α_2)	0.05	0.01	8.00	<0.005***

Legend: GSI - Government stringency index. Significance levels: “***” 0.001, “**” 0.01, “*” 0.05, “.” 0.1. The relationship between the predictors and the original response variable is inversely proportional .i.e. a negative sign indicates an increase of death rates while the positive one indicates a decrease.