

1 **Title: Examining Recent Trends in the Racial Disparity Gap in Tap Water Consumption:**
2 **NHANES 2011–2018**

Authors: Asher Y. Rosinger^{1,2,*}, Anisha I. Patel³, and Francesca Weaks⁴

Affiliations:

¹Department of Biobehavioral Health, Pennsylvania State University, University Park, PA, 16802

²Department of Anthropology, Pennsylvania State University, State College PA, 16802

³Department of Pediatrics, Stanford School of Medicine, Stanford, CA 94305

⁴Health Department, National Association for the Advancement of Colored People, Baltimore, MD 21215

*Corresponding Author: Asher Y. Rosinger

The Pennsylvania State University

219 Biobehavioral Health Building

University Park, PA 16802

Email: arosinger@psu.edu

Phone: 814-865-5220

Under consideration at Public Health Nutrition

Short title: Trends in tap water consumption disparities

The authors declare they have no conflicts of interest.

Funding statement: This work was supported by the Ann Atherton Hertzler Early Career Professorship funds, and Penn State's Population Research Institute (NICHD P2CHD041025). The funders had no role in the research or interpretation of results.

3 **Abstract:**

4 **Objective.** As tap water distrust has grown in the US with greater levels among Black and
5 Hispanic households, we aimed to examine recent trends in not drinking tap water including the
6 period covering the US Flint Water Crisis and racial/ethnic disparities in these trends.

7 **Design, Setting, and Participants.** We analyzed data from the cross-sectional National Health
8 and Nutrition Examination Survey data (2011–2018) for 9,439 children aged 2-19 and 17,268
9 adults. Log-binomial regressions and marginal predicted probabilities examined US nationally-
10 representative trends in tap and bottled water consumption overall and by race/ethnicity.

11 **Results.** Among US children and adults, respectively, in 2017-2018 there was a 63% (adjusted
12 prevalence ratio [PR]:1.63, 95%CI: 1.25-2.12, p<0.001) and 40% (PR:1.40, 95%CI: 1.16-1.69,
13 p=0.001) higher prevalence of not drinking tap water compared to 2013-2014 (pre-Flint Water

14 Crisis). For Black children and adults, the probability of not drinking tap water increased
15 significantly from 18.1% (95%CI: 13.4-22.8) and 24.6% (95%CI: 20.7-28.4) in 2013–14 to
16 29.3% (95%CI: 23.5-35.1) and 34.5% (95%CI: 29.4-39.6) in 2017–2018. Among Hispanic
17 children and adults, not drinking tap water increased significantly from 24.5% (95%CI: 19.4-
18 29.6) and 27.1% (95%CI: 23.0-31.2) in 2013-14 to 39.7% (95%CI: 32.7-46.8) and 38.1%
19 (95%CI: 33.0-43.1) in 2017-2018. No significant increases were observed among Asian or white
20 persons between 2013-14 and 2017-18. Similar trends were found in bottled water consumption.

21 **Conclusions.** This study found persistent disparities in the tap water consumption gap from
22 2011–2018. Black and Hispanics’ probability of not drinking tap water increased following the
23 Flint Water Crisis.

24

25 **Keywords:** Tap water; health disparities; water insecurity; bottled water

26

27 **Introduction**

28 Approximately two million Americans lack basic access to drinking water¹. However,
29 this figure does not capture individuals who have access but do not drink their tap water. Not
30 drinking tap water occurs for many reasons, including aesthetic considerations or distrust due to
31 water quality violations in homes² or schools³. The Flint, Michigan Water Crisis⁴, in which the
32 Flint community was exposed to high concentrations of lead in their water from 2015 through
33 2016 raised national awareness⁵ of tap water safety concerns in US communities^{2,6}. Then,
34 Newark, New Jersey reported a lead crisis in 2016⁷ further stoking fears. Both Flint and Newark
35 are comprised predominantly of non-white residents.

36 Other households in the US lack consistent access to tap water at home because of water
37 shutoffs due to affordability issues⁸, which disproportionately affect residents in low-income
38 housing, and Black and Hispanic households^{9,10}. The differential treatment of individuals in these
39 categories increases exposure to environmental injustice and racism¹¹ which may affect both
40 access to and use of tap water. Consequently, these groups are also more likely to drink bottled
41 water¹² and sugar-sweetened beverages^{13,14} that are more costly¹⁵ and lack the protective benefits
42 of fluoride¹⁶ in tap water.

43 While previous work documented an increase in US children not drinking tap water in
44 2015-2016¹² during the Flint Water Crisis compared to prior years, it is critical to surveil this
45 nutritional behavior as a way to monitor trust and usage of tap water. Therefore, we examined
46 whether not drinking tap water in the US continued to increase in 2017-2018 by analyzing
47 nationally-representative trends among children/adolescents and adults, and whether disparities
48 in this behavior persisted by race/ethnicity.

49

50 **Methods**

51 Data come from the National Health and Nutrition Examination Survey (NHANES)¹⁷, a
52 cross-sectional survey of the civilian, non-institutionalized household population, which uses a
53 complex, multistage probability design. Detailed sampling procedures and methodology are
54 described elsewhere¹⁸. NHANES is conducted by the National Center for Health Statistics
55 (NCHS) and approved by their research ethics board. Children aged 7-17 years provided assent
56 and parents provided consent for children under 18; adults consented for themselves.

57 This analysis uses the four most recent two-year cycles (2011–2018) because
58 oversampling of non-Hispanic (NH) Asians began in 2011-12 allowing for an examination of
59 this population’s trends in tap water consumption, and because prior work already examined
60 trends in Black, Hispanic and white tap water consumption from 2007-2016¹². Response rates
61 ranged from 77.0% in 2011–12 to 54.6% in 2017–18 for the examination component for youth
62 aged 1-19 years, and from 64.5% in 2011–12 to 45.3% in 2017–18 for adults aged 20+.

63 To assess the prevalence of not drinking tap water, we analyzed responses to the dietary
64 recall question: “When you drink tap water, what is the main source of the tap water?” If
65 respondents reported not drinking tap water, they were coded as “did not drink tap.” All others
66 were coded as “drank tap water”^{16,19}. A proxy responded to interview questions for children aged
67 ≤ 5 years; interviews for children aged 6–11 years were conducted with proxy assistance while
68 children aged ≥ 12 years responded themselves. If they responded that they did not know their
69 primary source, they were excluded from analyses.

70 For robustness analyses, we analyzed day-one 24-hour dietary recall data and
71 dichotomized tap water consumption and bottled water consumption as >0 ml or 0 ml following
72 prior work¹³. This allowed for a comparison of consumption on a given day to self-reports of not
73 drinking water from their main source of tap and if not drinking tap water shifted consumption to
74 bottled water.

75 *Statistical analyses*

76 A 2-sided *P*-value of .05 was used to assess statistical significance. Multiple log-binomial
77 regression models were estimated since prevalence of not drinking tap was $>10\%$ ¹². As there
78 were no significant race/ethnicity by survey cycle interactions (all $p>0.05$), we estimated the
79 regressions and covariate-adjusted time-trends without interaction terms. We used the 2013-14
80 survey cycle as the reference category as it was the last cycle prior to the Flint Water Crisis. We
81 plotted trends from the log-binomial regressions using marginal standardization²⁰ to generate
82 predicted probabilities by self-reported race/ethnicity (white, Black, Asian, Hispanic, or other).
83 We adjusted for the range²¹ of the following covariates: age (2-5, 6-11, 12-19, 20-39, 40-59,
84 60+), sex, federal income to poverty ratio (FIPR; ratio of family income to the federal poverty
85 guideline²²: $\leq 130\%$, 131-350%, $>350\%$), household reference educational attainment (for
86 children/adolescents; high school graduate/GED equivalent and some college were released

87 combined) and educational attainment (for adults), and whether respondents were born in the US
88 or not^{12,19}.

89 As robustness analyses, we re-estimated log-binomial regressions models as described
90 above with the dichotomized outcome of not drinking any tap water (0 ml) on a given day using
91 the 24-hour dietary recall. We repeated this analysis with the dichotomized outcome of any (>0
92 ml) bottled water intake.

93 All analyses accounted for the complex survey design, including day-one dietary sample
94 weights, which adjusted for oversampling, non-response, non-coverage, and day of week^{18,23} and
95 incorporated guidelines from NCHS²⁴. Statistical analyses were conducted using Stata V15.1
96 (Statacorp, College Station, TX).

97

98 **Results**

99 Data for the primary analyses were analyzed for 9,439 children and 17,268 adults with
100 complete covariate information (**Table 1**). Overall, the probability of not drinking tap increased
101 significantly from 12.3% (95% confidence interval [CI]: 9.5-15.0) and 13.9% (95% CI: 11.8-
102 16.0) among children and adults, respectively in 2013-2014 to 19.9% (95% CI: 16.0-23.8) and
103 19.5% (95% CI: 16.5-22.5) in 2017-2018. Among children/adolescents and adults, respectively,
104 in the post-Flint period (2017-2018) there was a 63% (adjusted prevalence ratio [PR]: 1.63, 95%
105 CI: 1.25-2.12, p<0.001) and 40% (PR: 1.40, 95% CI: 1.16-1.69, p=0.001) higher prevalence of
106 not drinking tap water compared to pre-Flint (2013-2014), respectively (**Figure 1a-b**;
107 **Supplemental Table 1**: Models 1-2).

108 Hispanic, Black, and Asian children/adolescents and adults had significantly higher
109 prevalence of not drinking tap than white persons (all p<0.001, **Figure 1a-1b**; **Supplemental**
110 **Table 1**: Models 1-2). Lower income, less educational attainment, and being born outside the US
111 were associated with higher prevalence of not drinking tap water (**Figure 1a-1b**).

112 Covariate-adjusted trends by race/ethnicity of not drinking tap water are presented in
113 **Figure 2a-2b**. Among Black children/adolescents and adults, the probability of not drinking tap
114 water increased significantly from 18.1% (95% CI: 13.4-22.8) and 24.6% (95% CI: 20.7-28.4) in
115 2013–14 to 29.3% (95% CI: 23.5-35.1) and 34.5% (95% CI :29.4-39.6) in 2017-18, respectively.
116 Among Hispanic children/adolescents and adults, not drinking tap increased from 24.5% (95%
117 CI: 19.4-29.6) and 27.1% (95% CI: 23.0-31.2) in 2013-14 to 39.7% (95% CI: 32.7-46.8) and

118 38.1% (95% CI: 33.0-43.1) in 2017–2018, respectively (**Figure 2a-2b**). The probability of not
119 drinking tap water did not significantly increase during between 2013-14 and 2017-18 among
120 Asian or non-Hispanic white children/adolescents and adults, respectively.

121

122 *Robustness analysis*

123 In robustness analyses, we used data from 10,254 children/adolescents and 17,829 adults
124 with valid dietary recall data, slightly more than the primary analyses due to fewer missing
125 responses. Results of the log-binomial regressions examining not drinking tap water on a given
126 day were slightly attenuated but largely consistent with main analyses (**Table 3**). Overall for
127 children/adolescents and adults, the prevalence of not drinking tap water was 14.0% (PR=1.14;
128 95% CI: 0.98-1.33) and 12.6% (PR=1.13; 95% CI: 1.01-1.25) higher in 2017-2018 than 2013-
129 2014. As in primary analyses, Black and Hispanic children/adolescents and adults had
130 significantly higher prevalence of not drinking any tap water on a given day than white persons
131 (**Table 3**, Models 1-2).

132 In the second analysis, prevalence of any bottled water consumption on a given day was
133 significantly higher overall for both children (PR: 1.26; 95% CI: 1.12-1.41) and adults (PR: 1.25;
134 95% CI: 1.12-1.40) in 2017-18 than in 2013-14. Further, Black, Asian, and Hispanic persons all
135 had higher prevalence of drinking bottled water than white children and adults, respectively
136 (**Table 4**, Models 1-2).

137

138 **Discussion**

139 This study used nationally-representative data to examine trends and racial/ethnic
140 disparities in tap water consumption among US children and adults which included the pre- and
141 post-Flint period as national awareness about water safety concerns increased. There was little
142 change between 2011–14, but probability of not drinking tap water increased successively from
143 2013-14 to 2017-2018, with Hispanic, Black, and Asian individuals being more likely to not
144 drink tap water than whites. Self-reported race/ethnicity continued to be the largest predictor of
145 not drinking tap water of all covariates examined as the differential treatment of individuals in
146 these categories increases exposure to environmental injustice and racism¹¹.

147 Our findings demonstrate persistent racial/ethnic disparities in the tap water consumption
148 gap and that Hispanic and Black households' probability of not drinking tap has further increased

149 in recent years. Higher inequities in water access⁹ and distrust²⁵ has been pointed to in prior
150 research as reasons that Black and Hispanic populations are less likely to drink tap water^{12,19,26}. It
151 is critical for public health nutrition to examine how environmental racism and policies may
152 contribute to health disparities, including disparities in water access and consumption. Tap water
153 consumption has health benefits through fluoride delivery that lowers risk of dental caries¹⁶.
154 Moreover, when individuals do not drink water they consume double the calories from sugar-
155 sweetened beverages¹³.

156 Environmental injustice and racism contribute to these disparities⁹ as water quality
157 violations²⁷ and water shutoffs⁹ disproportionately affect these communities. Drinking water
158 quality violations often occur at the intersection of race and class where low-income Black and
159 Hispanic communities are at the greatest risk of marginalization²⁸. Our results suggest little
160 progress in addressing these inequities. Increased national awareness of water quality violations
161 following the Flint Water Crisis, emerging local water contaminant exceedances in other cities
162 like Newark, NJ, or other unknown factors may help explain our findings. In particular,
163 increasing reports of lead violations in school drinking water, following increasing requirements
164 to test at the tap, have likely heightened concerns regarding tap water safety³.

165 To address these disparities and distrust, expanded water quality testing, communication,
166 and lead remediation efforts are needed. Simplified water quality reports that are easy to
167 understand coupled with clear communication between utility companies and households are
168 paramount (e.g. http://policyinnovation.org/wp-content/uploads/WaterDataPrize_Report.pdf).
169 Further, targeted interventions for marginalized communities that use trusted community
170 stakeholders can help address environmental injustice and rebuild trust⁹.

171 While recent studies indicate that up to two million Americans lack basic water access¹ -
172 particularly in urban areas²⁹ - our study indicates the broader problem is far more pervasive and
173 growing. In 2017–2018, approximately 20% of US children/adolescents and adults did not drink
174 tap water up from ~13% in 2013-14. Using the NCHS Census population totals¹⁸, this translates
175 into 14.8 million children and 46.6 million adults or 61.4 million people in 2017-18 in the US
176 who did not drink their tap water (**Supplemental Table 2**). Since 2013-14, this represents an
177 increase of 19 million people (~5.6 million children, ~13.4 million adults) not drinking their tap
178 water, which could be considered an epidemic of tap water distrust and disuse.

179 Our results differ somewhat from a recent paper examining trends in tap water intake
180 using NHANES 2011-2016 data, which did not find significant changes in the volume of tap
181 water consumed over time, but did uncover increases in the volume of bottled water consumed³⁰.
182 While our primary variable was based on whether persons consumed their tap water rather than
183 the average volume of water consumed, our results were robust when using a dichotomous
184 variable of any tap water intake from the 24-hour dietary recall. Second, we found that
185 prevalence of any bottled water consumption on a given day was 25% higher in 2017-18 than
186 2013-14, supporting the recent trends of increasing bottled water intake in the US^{12,30,31}. It is
187 critical for public health nutrition to monitor trends in tap and bottled water use beyond average
188 mean intakes because this can provide a window into potential underlying water insecurity.

189 Study limitations include the inability to know why participants did not drink their tap
190 water as follow-up questions were not asked. However, we used two measures of tap water
191 consumption to provide additional internal validity of results and find that shifts to bottled water
192 occurred. We were not able to exclude individuals with no access to tap water, however,
193 NHANES samples the civilian, household population¹⁸ and thus does not sample homeless, who
194 represent a large percentage of those without tap access³². This study is limited by not having
195 geographical data, but its goal was to examine recent national trends in tap water consumption
196 rather than geographic differences. Recent work has demonstrated that structural issues like
197 precarious housing within communities of color increases risk of incomplete plumbing^{29,33}. The
198 question regarding tap water filtration was discontinued following the 2009-2010 survey and
199 thus was not controlled for. However, prior work has demonstrated that use of tap water filters
200 do not have an overall effect on these national estimates of tap water consumption¹². NCHS
201 addressed the declining response rates through the sampling weights, which further took into
202 account non-response and loss of screener to examination¹⁸.

203

204 **Conclusion**

205 Approximately 63% and 40% more Black and Hispanic children and adults did not drink
206 their tap water in 2017-18 compared to 2013-14. Overall, that approximately 61.4 million people
207 in the US did not drink tap water, an increase of 19 million since the Flint Water Crisis,
208 represents a substantial opportunity to improve public health nutrition. Further, bottled water

209 consumption continued to increase in the US from 2011–2018, particularly among Black, Asian,
210 and Hispanic households. This reliance on bottled water levies additional socioeconomic strain.

211 Policies that address the root of environmental injustices that low-income and minority
212 communities face in accessing safe tap water are necessary to halting the growing racial disparity
213 gap observed in tap water consumption²⁵. To design meaningful solutions and rebuild trust more
214 work with communities of color is needed to understand the various factors that contribute to
215 eschewing tap water. The field of public health nutrition should engage with utility companies
216 and community stakeholders to design interventions aimed at ensuring access to clean water and
217 reducing these disparities in tap water distrust.

218

219 References

- 220 1. Roller Z, Gasteyer S, Nelson N, Lai W, Shingne M. Closing the water access gap in the
221 United States: a national action plan. Washington, DC: Dig Deep and US Water Alliance.
222 Retrieved on July. 2019;25:2020.
- 223 2. Hanna-Attisha M, LaChance J, Sadler RC, Champney Schnepf A. Elevated Blood Lead
224 Levels in Children Associated With the Flint Drinking Water Crisis: A Spatial Analysis
225 of Risk and Public Health Response. *American Journal of Public Health*.
226 2015;106(2):283-290.
- 227 3. Cradock AL, Poole MK, Vollmer LY, Barrett JL, Flax CN, Hecht CA. *State approaches*
228 *to testing school drinking water for lead in the United States*. Boston, MA: Prevention
229 Research Center on Nutrition and Physical Activity at the Harvard T.H. Chan School of
230 Public Health;2019.
- 231 4. Pauli BJ. The Flint water crisis. *Wiley Interdisciplinary Reviews: Water*.
232 2020;7(3):e1420.
- 233 5. Jackson DZ. Environmental justice? Unjust coverage of the Flint water crisis.
234 *Shorenstein center on media, politics and public policy*. 2017:1-55.
- 235 6. Butler LJ, Scammell MK, Benson EB. The Flint, Michigan, Water Crisis: A Case Study
236 in Regulatory Failure and Environmental Injustice. *Environmental Justice*. 2016;9(4):93-
237 97.
- 238 7. Gostin LO. Lead in the Water: A Tale of Social and Environmental Injustice. *JAMA*.
239 2016;315(19):2053-2054.
- 240 8. Teodoro MP, Saywitz RR. Water and sewer affordability in the United States: a 2019
241 update. *AWWA Water Science*. 2020;2(2):e1176.
- 242 9. Montag C. Municipal Power and Racial Injustice: Solutions for Water Discrimination.
243 *Natural Resources & Environment*. 2020;34(4):16-20.
- 244 10. Swain M, McKinney E, Susskind L. Water Shutoffs in Older American Cities: Causes,
245 Extent, and Remedies. *Journal of Planning Education and*
246 *Research*.0(0):0739456X20904431.
- 247 11. Jones CP. Invited Commentary: "Race," Racism, and the Practice of Epidemiology.
248 *American Journal of Epidemiology*. 2001;154(4):299-304.
- 249 12. Rosinger AY, Young SL. In-Home Tap Water Consumption Trends Changed Among
250 U.S. Children, but Not Adults, Between 2007 and 2016. *Water Resources Research*.
251 2020;56(7):e2020WR027657.
- 252 13. Rosinger AY, Bethancourt H, Francis LA. Association of Caloric Intake From Sugar-
253 Sweetened Beverages With Water Intake Among US Children and Young Adults in the
254 2011-2016 National Health and Nutrition Examination Survey. *JAMA Pediatrics*.
255 2019;173(6):602-604.
- 256 14. Onufrak SJ, Park S, Sharkey JR, Sherry B. The relationship of perceptions of tap water
257 safety with intake of sugar-sweetened beverages and plain water among US adults.
258 *Public Health Nutrition*. 2014;17(01):179-185.
- 259 15. Javidi A, Pierce G. U.S. Households' Perception of Drinking Water as Unsafe and its
260 Consequences: Examining Alternative Choices to the Tap. *Water Resources Research*.
261 2018;54(9):6100-6113.
- 262 16. Sanders AE, Slade GD. Blood Lead Levels and Dental Caries in U.S. Children Who Do
263 Not Drink Tap Water. *American Journal of Preventive Medicine*. 2018;54(2):157-163.

- 264 17. National Cancer Institute. Dietary Screener Questionnaire in the National Health and
265 Nutrition Examination Survey. Available at:
266 <https://epi.grants.cancer.gov/nhanes/dietscreen/>. Accessed on May 20.
267 18. National Center for Health Statistics. National Health and Nutrition Examination Survey
268 (NHANES): Questionnaires, datasets, and related documentation. 2020;
269 <https://wwwn.cdc.gov/nchs/nhanes/default.aspx>. Accessed August 8, 2020.
270 19. Patel AI, Shapiro DJ, Wang YC, Cabana MD. Sociodemographic Characteristics and
271 Beverage Intake of Children Who Drink Tap Water. *American Journal of Preventive*
272 *Medicine*. 2013;45(1):75-82.
273 20. Graubard BI, Korn EL. Predictive Margins with Survey Data. *Biometrics*.
274 1999;55(2):652-659.
275 21. Muller CJ, MacLehose RF. Estimating predicted probabilities from logistic regression:
276 different methods correspond to different target populations. *Int J Epidemiol*.
277 2014;43(3):962-970.
278 22. U.S. Department of Health and Human Services. U.S. Federal Poverty Guidelines.
279 Washington, D.C 2016
280 23. Korn EL, Graubard BI. *Analysis of health surveys*. Vol 323: John Wiley & Sons; 2011.
281 24. Ingram DD, Malec DJ, Makuc DM, et al. National Center for Health Statistics guidelines
282 for analysis of trends. 2018.
283 25. Patel AI, Schmidt LA. Water Access in the United States: Health Disparities Abound and
284 Solutions Are Urgently Needed. *American Journal of Public Health*. 2017;107(9):1354-
285 1356.
286 26. Rosinger AY, Herrick KA, Wutich AY, Yoder JS, Ogden CL. Disparities in plain, tap
287 and bottled water consumption among US adults: National Health and Nutrition
288 Examination Survey (NHANES) 2007–2014. *Public Health Nutrition*. 2018;21(8):1455-
289 1464.
290 27. McDonald YJ, Jones NE. Drinking Water Violations and Environmental Justice in the
291 United States, 2011–2015. *American Journal of Public Health*. 0(0):e1-e7.
292 28. Switzer D, Teodoro MP. Class, Race, Ethnicity, and Justice in Safe Drinking Water
293 Compliance*. *Social Science Quarterly*. 2018;99(2):524-535.
294 29. Meehan K, Jurjevich JR, Chun NMJW, Sherrill J. Geographies of insecure water access
295 and the housing–water nexus in US cities. *Proceedings of the National Academy of*
296 *Sciences*. 2020;117(46):28700-28707.
297 30. Vieux F, Maillot M, Rehm CD, Barrios P, Drewnowski A. Trends in tap and bottled
298 water consumption among children and adults in the United States: analyses of NHANES
299 2011–16 data. *Nutrition Journal*. 2020;19(1):10.
300 31. Gleick PH. *Bottled and sold: The story behind our obsession with bottled water*. Island
301 Press; 2010.
302 32. Capone D, Cumming O, Nichols D, Brown J. Water and Sanitation in Urban America,
303 2017–2019. *American journal of public health*. 2020;110(10):1567-1572.
304 33. Deitz S, Meehan K. Plumbing Poverty: Mapping Hot Spots of Racial and Geographic
305 Inequality in U.S. Household Water Insecurity. *Annals of the American Association of*
306 *Geographers*. 2019;109(4):1092-1109.

307

308 **Table 1:** Descriptive characteristics of US children/adolescents and adults, NHANES 2011-2018

	Children/adolescents ^{a,c} (n=9,439)	Adults ^{a,c} (n=17,268)
	^b Mean	^b Mean
2011-12	25.9%	24.4%
2013-14	25.3%	25.2%
2015-16	25.0%	25.2%
2017-18	23.8%	25.2%
% Don't drink tap from main water source	15.0%	16.2%
% Didn't drink tap on a given day	48.6%	46.3%
% Drink bottled water on a given day	37.1%	37.5%
% Male	48.5%	51.4%
2-5 years / 20-39 years	22.3%	35.9%
6-11 years / 40-59 years	35.6%	36.8%
12-19 years / 60+ years	42.1%	27.3%
NH white	54.7%	66.6%
NH Black	13.5%	10.8%
Asian	4.0%	5.2%
Hispanic	22.0%	13.9%
% Born in US	95.2%	83.9%
FIPR: ≤130%	34.9%	22.6%
131-350%	37.3%	35.2%
>350%	27.7%	42.2%
Adult education:	--	
Less than high school		13.0%
High school grad	--	22.7%
Some college	--	32.8%
College +	--	31.4%
HH Ref education:		--
Less than high school	17.8%	
High school grad/some college	54.3%	--
College +	27.9%	--

309 ^aUnweighted sample size. ^bWeighted mean percentages for each category. ^cWithout missing
 310 covariate data and valid dietary recall status. Other race/Hispanic origin included in analyses but
 311 not shown. FIPR: Federal income poverty ratio

312

313 **Table 2:** Log-binomial regression models of not drinking any tap water on a given day by survey
 314 cycle, race/ethnicity, and socio-demographics, NHANES 2011-2018 among children/adolescents
 315 and adults.

Independent variables	Adjusted Prevalence ratio of not drinking tap water (95% CI)	Adjusted Prevalence ratio of not drinking tap water (95% CI)
	Children/adolescents	Adults
	Model 1	Model 2
Survey cycle 2011-12	1.06 (0.93 - 1.21)	0.95 (0.86 - 1.05)
2013-2014	1	1
2015-16	1.05 (0.91 - 1.21)	1.02 (0.91 - 1.13)
2017-18	1.14* (0.98 - 1.33)	1.13** (1.01 - 1.25)
Race/ethnicity: NH white	1	1
NH Black	1.38*** (1.22 - 1.56)	1.39*** (1.31 - 1.48)
NH Asian	1.08 (0.93 - 1.26)	0.99 (0.92 - 1.08)
Hispanic	1.40*** (1.25 - 1.56)	1.28*** (1.19 - 1.37)
Other/Mixed	1.08 (0.91 - 1.27)	1.16** (1.02 - 1.32)
Age: ^a 2-5 years / ^b 60+ 6-11 years / 20-39	1 0.95 (0.89 - 1.01)	1 1.11*** (1.05 - 1.17)
12-19 years / 40-59	1.02 (0.95 - 1.09)	1.14*** (1.08 - 1.20)
Nativity status: Born in US	1	1
Born outside US	1.03 (0.94 - 1.14)	1.09*** (1.03 - 1.15)
Male	1	1
Female	1.02 (0.95 - 1.09)	1.00 (0.96 - 1.04)
^c FIPR >350% ≤130%	1.14* (1.00 - 1.30)	1.15*** (1.07 - 1.24)
131-350%	1.08 (0.96 - 1.23)	1.08** (1.02 - 1.15)
Education ^b : College + Less than high school	-- --	1 1.70*** (1.55 - 1.86)
High school grad	--	1.62***


			(1.49 - 1.76)
Some college	--		1.45***
			(1.34 - 1.57)
HH reference education ^a : College + Less than high school	1		--
	1.49***		--
	(1.29 - 1.72)		
High school grad/some college	1.41***		--
	(1.23 - 1.61)		
Observations	10,254		17,829
316	95% CI in parentheses; *** p<0.01, ** p<0.05, * p<0.1;		
317	^a for children; ^b for adults;		
318	^c FIPR: federal income to poverty ratio; household income for children		
319	Note: Data using day one 24-hour dietary recall		
320			

321 **Table 3:** Log-binomial regression models of drinking any bottled water on a given day by survey
 322 cycle, race/ethnicity, and socio-demographics, NHANES 2011-2018 among children/adolescents
 323 and adults.

Independent variables	Adjusted Prevalence ratio of drinking bottled water (95% CI)	Adjusted Prevalence ratio of drinking bottled water (95% CI)
	Children/adolescents	Adults
	Model 1	Model 2
Survey cycle 2011-12	0.94 (0.76 - 1.16)	0.87** (0.77 - 1.00)
2013-2014	1	1
2015-16	1.03 (0.91 - 1.17)	1.08 (0.96 - 1.21)
2017-18	1.26*** (1.12 - 1.41)	1.25*** (1.12 - 1.40)
Race/ethnicity: NH white	1	1
NH Black	1.20** (1.03 - 1.40)	1.56*** (1.43 - 1.69)
NH Asian	1.27*** (1.08 - 1.50)	1.14** (1.01 - 1.28)
Hispanic	1.55*** (1.39 - 1.73)	1.50*** (1.34 - 1.67)
Other/Mixed	0.98 (0.80 - 1.19)	1.25*** (1.06 - 1.47)
Age: ^a 2-5 years / ^b 60+ 6-11 years / 20-39	1 1.11* (0.99 - 1.24)	1 1.23*** (1.13 - 1.34)
12-19 years / 40-59	1.22*** (1.11 - 1.34)	1.22*** (1.12 - 1.34)
Nativity status: Born in US	1	1
Born outside US	1.09* (0.98 - 1.22)	1.26*** (1.17 - 1.36)
Male	1	1
Female	1.17*** (1.07 - 1.27)	1.16*** (1.11 - 1.22)
^c FIPR >350%	1	1
≤130%	0.89* (0.79 - 1.00)	0.90*** (0.84 - 0.96)
131-350%	0.87* (0.76 - 1.00)	0.93** (0.87 - 0.99)
Education ^b : College +	--	1
Less than high school	--	1.33*** (1.21 - 1.46)
High school grad	--	1.24***


			(1.13 - 1.37)
Some college	--		1.29***
			(1.20 - 1.39)
HH reference education ^a : College + Less than high school	1 1.20*		--
	(0.99 - 1.45)		--
High school grad/some college	1.23**		--
	(1.05 - 1.44)		
Observations	10,254		17,829
324	95% CI in parentheses; *** p<0.01, ** p<0.05, * p<0.1;		
325	^a for children; ^b for adults;		
326	^c FIPR: federal income to poverty ratio; household income for children;		
327	Note: Data using day one 24-hour dietary recall		
328			

329 **Figure 1: Log-binomial regression models of not drinking tap water by survey cycle,**
 330 **race/ethnicity, and socio-demographics, NHANES 2011-2018 among (a)**
 331 **children/adolescents and (b) adults.**
 332



333
 334
 335 **Notes:** n=9,439 children/adolescents aged 2-19; n=17,268 adults; models adjusted for all
 336 variables shown in addition to sex, and age (for adults). Full models shown in Supplemental
 337 Table 1.

338 **Figure 2: Covariate-adjusted predicted probability time trends and 95% confidence**
339 **intervals of not drinking tap water by race/ethnicity among (a) children and (b) adults,**
340 **NHANES 2011-2018**
341



342
343 **Notes:** n=9,439 children aged 2-19; n=17,268 adults; models adjusted for nativity status, sex, age
344 group, income level (federal income to poverty ratio), and educational attainment (of household
345 reference for children, self for adults); dashed line at 2013–2014 indicates beginning of Flint,
346 Michigan Water Crisis. Built from models shown in Supplemental Table 1.

347
348

349 **Supplemental Materials for**

350 **Title: Examining Recent Trends in the Racial Disparity Gap in Tap Water Consumption:**

351 **NHANES 2011–2018**

352

353 **Supplemental Table 1:** Log-binomial regression models of not drinking tap water by survey

354 cycle, race/ethnicity, and socio-demographics, NHANES 2011-2018 among children/adolescents

355 and adults.

Independent variables	Adjusted Prevalence ratio of not drinking tap water (95% CI)	Adjusted Prevalence ratio of not drinking tap water (95% CI)
	Children/adolescents	Adults
	Model 1	Model 2
Survey cycle 2011-12	1.00 (0.76 - 1.32)	1.07 (0.84 - 1.37)
2013-2014	1	1
2015-16	1.28* (0.96 - 1.70)	1.17 (0.96 - 1.43)
2017-18	1.63*** (1.25 - 2.12)	1.40*** (1.16 - 1.69)
Race/ethnicity: NH white	1	1
NH Black	2.46*** (1.79 - 3.38)	2.28*** (1.96 - 2.65)
NH Asian	1.84*** (1.35 - 2.52)	1.49*** (1.22 - 1.80)
Hispanic	3.12*** (2.41 - 4.03)	1.89*** (1.56 - 2.29)
Other/Mixed	1.52** (1.04 - 2.20)	1.52*** (1.26 - 1.84)
Age: ^a 2-5 years / ^b 60+ 6-11 years / 20-39	1 0.88** (0.78 - 1.00)	1 1.01 (0.90 - 1.14)
12-19 years / 40-59	0.50*** (0.44 - 0.58)	1.10* (0.99 - 1.22)
Nativity status: Born in US	1	1
Born outside US	1.49*** (1.23 - 1.80)	1.53*** (1.31 - 1.80)
Male	1	1
Female	1.10* (0.99 - 1.22)	1.22*** (1.13 - 1.32)
^c FIPR >350% ≤130%	1 1.42** (1.06 - 1.90)	1 1.23** (1.05 - 1.45)

131-350%	1.33*	1.15*
	(0.98 - 1.81)	(1.00 - 1.34)
Education ^b : College +	--	1
Less than high school	--	2.12***
		(1.77 - 2.53)
High school grad	--	1.93***
		(1.60 - 2.33)
Some college	--	1.61***
		(1.37 - 1.89)
HH reference education ^a : College +	1	--
Less than high school	1.55***	--
	(1.19 - 2.00)	
High school grad/some college	1.42***	--
	(1.10 - 1.84)	
Observations	9,439	17,268

356 95% CI in parentheses; *** p<0.01, ** p<0.05, * p<0.1;

357 ^afor children; ^bfor adults;

358 ^cFIPR: federal income to poverty ratio; household income for children;

359 Note: Data using question inquiring about normal drinking water source in which persons

360 indicated if they did not consume their tap water.

361

362 **Supplemental Table 2:** Estimated amount of people in the US not drinking tap water and
363 change between 2013-2014 and 2017-2018.

364

	2013-2014	2017-2018	Total change
Children/adolescents (2-19 year olds)	9,167,905	14,822,954	5,655,049
Adults (20+ years)	33,184,518	46,553,820	13,369,302
Total	42,352,423	61,376,774	19,024,351

365 Note: Data calculated from adjusted prevalence estimates in this paper multiplied by the NCHS
366 census population totals ([https://wwwn.cdc.gov/nchs/nhanes/responserates.aspx#population-](https://wwwn.cdc.gov/nchs/nhanes/responserates.aspx#population-totals)
367 [totals](https://wwwn.cdc.gov/nchs/nhanes/responserates.aspx#population-totals)) in 2013-14 and 2017-18.

368
369