Direct and indirect effectiveness of mRNA vaccination against SARS-CoV-2 infection

in long-term care facilities in Spain

Susana Monge[1], Carmen Olmedo[2], Belén Alejos, María Fé Lapeña[3], María José Sierra[1]*,

Aurora Limia[2]* and the COVID-19 registries study group.

[1]Centre for the Coordination of Alerts and Health Emergencies, General Directorate of

Public Health, Ministry of Health.

[2]Vaccines division, General Directorate of Public Health, Ministry of Health.

[3]Data analytics department, General Directorate of Digital Health and National Health

Service Information Systems, Ministry of Health.

* These persons contributed equally to this work.

Corresponding author:

Susana Monge

smonge@mscbs.es

Conflicts of interest: authors declare no support from any organisation for the

submitted work; no financial relationships with any organisations that might have an

interest in the submitted work in the previous three years, no other relationships or

activities that could appear to have influenced the submitted work.

Contributions: SM, CO, MJS and AL conceived the idea for the study. SM and CO

designed the analysis which was performed by SM with help from BA, support from

MFL and under the supervision of MJS and AL. All authors participated in the

interpretation of results and critically reviewed the content of the manuscript.

Funding: This work did not receive any specific grant from funding agencies in the

public, commercial, or not-for-profit sectors.

Members of the COVID-19 registries study group: *David Moreno, Manuel Méndez*

Díaz, Ismael Huerta González, Antonia Mª Galmes Truyol, Ana Barreno Estévez,

Valvanuz García Velasco, Mª Jesús Rodríguez Recio, José Sacristán, Montserrat

Martínez Marcos, Eliseo Pastor Villalba , María José Macías Ortiz, Ana García Vallejo,

Sara Santos Sanz, Rocío García Pina , Aurelio Barricarte Gurea, Rosa Sancho Martínez,

Eva María Ochoa, Mauricio Vázquez Cantero, Atanasio Gómez Anés, María Jesús

Pareja Megía, Yolanda Castán, Manuel Roberto Fonseca Álvarez, Antonia Salvà Fiol,

Hilda Sánchez Janáriz, Luz López Arce, María Ángeles Cisneros Martín, José Sacristán,

Frederic Jose Gibernau, Cesar Fernandez Buey, Katja Villatoro Bongiorno, Francisco Javier

Rubio García, Fernando Santos Guerra, Jenaro Astray Mochales, Francisco Javier

Francisco Verdu, Isabel García Romero, Rosa Oriza Bernal, Tomás Gómez Pérez, Salomé

Hijano Villegas, Sergio Román Soto, Virgilio Yagüe Galaup, Mercedes Alfaro Latorre,

Marta Aguilera Guzmán, Belén Crespo Sánchez-Eznarriaga, Montserrat Neira León,

Noemí Cívicos Villa.

Abstract

Objectives To estimate indirect and total (direct plus indirect) effects of COVID-19 vaccination in residents in long-term care facilities (LTCF).

Design Registries-based cohort study including all residents in LTCF ≥65 years offered vaccination between 27 December 2020 and 10 March 2021. Risk of SARS-CoV-2 infection following vaccination was compared with the risk in the same individuals in a period before vaccination. Risk in non-vaccinated was also compared to a period before the vaccination programme to estimate indirect protection. Standardized cumulative risk was computed adjusted by previous documented infection (before the start of follow-up) and daily-varying SARS-CoV-2 incidence and reproductive number.

Participants 573,533 records of 299,209 individuals in the National vaccination registry were selected; 99.0% had ≥1 vaccine-dose, 99.8% was Pfizer/BioNTech (BNT162b2). Residents mean age was 85.9, 70.9% were females. A previous SARS-CoV-2 infection was found in around 25% and 13% of participants, respectively, at the time of vaccine offer and in the reference period.

Main outcome measures Documented SARS-CoV-2 infection identified in the National COVID-19 laboratory registry.

Results Total VE was 57.2% (95% Confidence Interval: 56.1%-58.3%), and was highest ≥28 days after the first vaccine-dose (*proxy* of ≥7 days after the second dose) and for individuals naïve to SARS-CoV-2 [81.8% (81.0%-82.7%)] compared to those with previous infection [56.8% (47.1%-67.7%)]. Vaccination prevented up to 9.6 (9.3-9.9) cases per 10.000 vaccinated per day; 11.6 (11.3-11.9) if naïve vs. 0.8 (0.5-1.0) if previous infection. Indirect protection in the non-vaccinated could only be estimated

for naïve individuals, at 81.4% (73.3%-90.3%) and up to 12.8 (9.4-16.2) infections

prevented per 10.000 indirectly protected per day.

Conclusions Our results confirm the effectiveness of mRNA vaccination in institutionalized elderly population, endorse the policy of universal vaccination in this setting, including in people with previous infection, and suggest that even non-vaccinated individuals benefit from indirect protection.

Key-words: COVID-19; SARS-CoV-2; vaccination; vaccine effectiveness; long-term care facilities; elderly; indirect effects.

Key messages:

- COIVD-19 vaccination reduced the risk of documented SARS-CoV-2 infection in institutionalized elderly by 57.2% (56.1% to 58.3%), which increased to 81.2% (80.2% to 82%) for the fully vaccinated.

- In individuals naïve to SARS-CoV-2 vaccination reduced the risk by up to 81.8% and averted up to 11.6 cases per 10,000 vaccinated persons per day.

- Those with previous infection also benefited from a risk reduction of 57%, which translated in less than 1 infection averted per 10,000 vaccinated persons per day.

- Non-vaccinated individuals living in facilities where the majority (residents and staff) had been vaccinated showed a risk reduction similar to those actually vaccinated.

MAIN TEXT

Introduction

Since the beginning of the COVID-19 pandemic up to March 7 2021, 18,927 residents in

long-term care facilities (LTCF) have died in Spain with confirmed COVID-19, and an

additional 10,492 have died with compatible symptoms [1]. This means a cumulative

mortality rate of 67 per 1,000 residents, accounting only for confirmed infections. This

high vulnerability is due to the higher risk of exposure in dependents living in a closed

institution but also to the higher severity of infection due to advanced age and

presence of comorbidities. Indeed, one on every 5 cases of SARS-CoV-2 infection died

in this setting [1].

COVID-19 vaccination in Spain started on December 27 with the Pfizer/BioNTech

(BNT162b2) vaccine, for which LTCF -both residents and workers- were the first priority

group [2]. The vaccination campaign coincided with the third COVID-19 epidemic wave,

with national 14-day cumulative incidence increasing from less than 250 cases per

100,000 population by the end of 2020 to more than 1,000 by the end of January 2021

[3]. Vaccination started in facilities considered at higher risk, such as those that had

never experienced a COVID-19 outbreak, had higher number of residents or more

difficulties for implementing prevention and control measures. Vaccination teams

visited the facilities and vaccination was universal, including those with previous SARS-

CoV-2 infection. Vaccination was only deferred in people with active infection and,

inconsistently, in people under quarantine. Acceptance has been very high, with 97.8%

of all institutionalized persons (any institution type) having received at least one

vaccine dose, and 88.8% two doses [4].

The Pfizer/BioNTech vaccine has shown an efficacy of 95% in preventing Covid-19 in

randomized clinical trials [5]. However, elderly persons in general, and those

institutionalized in particular, are not represented in randomized studies [6]. Therefore

there is great interest in estimating vaccine effectiveness (VE) in this population

following its widespread vaccination. Moreover, because vaccination coverage was so

high, it is expected that non-vaccinated persons could be indirectly protected if

vaccination reduces infection and transmissibility among vaccinated persons. A few

observational studies focusing on the elderly have been published in the last weeks

[7,8]; one published and two pre-print studies have specifically addressed vaccine

effects in LTCF residents [9,10,11], and none have tried to address the indirect

protection in non-vaccinated individuals in this high-coverage setting.

This study aims to estimate indirect and total (direct plus indirect) effects of

vaccination in residents in LTCF in a high incidence context.

Methods

Data sources

REGVACU is a nation-wide registry of all COVID-19 vaccine-doses administered and

vaccine rejections. Data was extracted on March 15 and the administrative censoring

date was March 10. Individuals ≥65 years of age by December 27, with a valid postal

code, and identified as "resident in elderly homes" according to REGVACU were

selected. SERLAB is a nation-wide registry of all SARS-CoV-2 PCR and rapid antigenic

tests performed. Positive tests within 60 days of a previous positive one were dropped,

as they were considered to belong to the same episode. In LTCF, tests were performed

to symptomatic persons and risk contacts. Incoming residents were also routinely

tested and periodical screenings have also been carried out. Therefore, documented

infections registered in SERLAB may correspond both to symptomatic and

asymptomatic infections, although this circumstance was not recorded in the system.

Residents in REGVACU were cross-matched with SERLAB by person identification

number, date of birth and sex.

Study design

To estimate the total (direct and indirect) effect of vaccination in vaccinated

individuals, the risk of SARS-CoV-2 documented infection in the cohort of individuals

with the first dose administered between December 27 and March 10 was compared to

the risk in the same individuals in a period before the start of the vaccination

programme. A before-after comparison was deemed more appropriate since, due to

the high vaccination coverage at LTCFs, non-vaccinated individuals after December 27

would probably not represent baseline infection risk had the individual not been

vaccinated. Baseline infection risk, on the other hand, is heavily influenced by

community incidence and the vaccination campaign coincided with the third epidemic

wave in Spain. To minimize this effect the second epidemic wave was chosen as

comparison period, starting the follow-up of the non-vaccinated period 87 days before

individual-specific first dose administration date (October 1, at the earliest), with

administrative censoring on December 13, 87 days before March 10 (supplementary

Figure S1).

To estimate the indirect protection of vaccination in not vaccinated individuals, the risk

of SARS-CoV-2 documented infection in the cohort of individuals never vaccinated

between December 27 and March 10 was compared to the risk in the same individuals

87 days before, similarly as previously explained for vaccinated individuals. The follow-

up period started at the earliest date when the vaccine was offered to each individual,

since all residents at the same LTCF were offered vaccination on the same day.

Therefore individuals were ensured to be included on the date that a first vaccine-dose

was administered to most of the co-residents and workers.

The follow-up for all individuals finished at the earliest of a SARS-CoV-2 positive test or

administrative censoring. Unfortunately, no information on the individuals' vital status

was available. Existence of any previous SARS-CoV-2 documented infection on the first

day of follow-up was also registered.

An additional analysis to investigate the possible design-associated bias is presented in

the supplementary material.

The study obtained approval from the research ethics committee at the Instituto de

Salud Carlos III (CEI PI 98_2020). Patients or the public were not involved in the design, or

conduct, or reporting, or dissemination plans of our research. Results of this study are

planned to be disseminated to the broad public.

Data analysis

The standardized cumulative risk of a documented SARS-CoV-2 infection that every

individual had in the sample been either vaccinated or not vaccinated was computed

[12]. To estimate the probability of the event on each follow-up day, conditioned to

remaining event-free up to that day and given the individual covariates, a pooled

logistic regression was fitted adjusting by follow-up day, previous SARS-CoV-2 infection

(before beginning of follow-up), daily-varying 7-day SARS-CoV-2 cumulative incidence

specific to the province, its quadratic term, and the empirical reproduction number for

that province on that date. An interaction between follow-up day and vaccination was

introduced to allow for a time-varying effect of the vaccine. Robust models were built

using individuals as clusters. Standardized cumulative risk curves were derived using

the Kaplan-Meier method. Risk ratios (RR), vaccine effectiveness (VE= 1-RR) and risk

difference (RD) were estimated for the overall period and in four sub-periods after the

administration of the first dose, as proxies of different vaccine protection: (1) 14 days;

(2) 14 to 21 days; (3) 21 to 28 days (proxy of first 7 days after the second dose) and; (4)

>28 days (proxy of fully vaccinated, i.e. ≥7 days after second vaccine dose). Normal

distribution-based confidence intervals were estimated using bootstrapping with 300

repetitions.

Results

Description of participants

Out of 5,068,733 vaccination records from 3,615,403 individuals in REGVACU, 573,533

records from 299,209 individuals were selected; 296,093 (99.0%) had received ≥1

vaccine-dose, of which 99.8% were Pfizer/BioNTech (BNT162b2) and 0.2% Moderna

vaccine; 92.6% of them received a second vaccine-dose in a median of 21 days

(interquartile range: 21-21). Time to vaccination is shown in supplementary figure S2.

Mean age was 85.9 years (standard deviation = 7.8) and 70.9% were females. Selected

individuals were cross-matched with SERLAB; 77,662 (26.0%) had at least one positive

test between March 1, 2020 and March 11, 2021. A SARS-CoV-2 previous infection was

found in 17.5% of participants on the date they started the follow-up for the total

effects study; 22.3% in the vaccinated group and 12.7% in the comparison group (from

87 days before). In the indirect effects analysis, 20.3% had previous infection; 27.7% in

the indirectly protected and 12.9% in the comparison group.

Estimation of vaccine effectiveness in vaccinated persons

This analysis included 16,277,284 and 16,142,536 person-days of follow-up among

vaccinated and non-vaccinated persons, respectively. There were 11,304 and 19,656

documented infections, respectively (supplementary table S1). Detailed information on

the crude estimates and adjusted cumulative risk in each group can be found in the

supplementary material (Figure S3 and Tables S2 and S3).

Vaccine effectiveness for the whole study period was 57.2% (95% Confidence Interval:

56.1% to 58.3%), but it increased after two vaccine doses, and was higher in individuals

without previous SARS-CoV-2 infection; VE was 81.8% (81.0% to 82.7%) for residents

fully vaccinated and with no previous infection, but decreased to 56.8% (47.1% to

67.7%) if previous infection (Table 1, Figure 1). Interestingly, in a separate analysis we

found that previous infection in the reference period was associated to a risk reduction

of 86.6% (85.2%-87.8%), higher than the estimate for complete vaccination.

The estimated number of SARS-CoV-2 infections averted by vaccination (risk

difference) was greatest in the intermediate periods, which coincided with the peak of

the epidemic waves, at 11.6 cases per 10,000 vaccinated persons per day in the group

without previous infection (Table 1). In the group with previous infection, the number

of infections averted was much lower, of around 0.6 – 0.7 per 10,000 vaccinated

persons per day.

Estimation of indirect vaccine effectiveness in non-vaccinated persons

This analysis included 164,520 and 161,388 person-days of follow-up, respectively,

among persons not vaccinated but who had been offered the vaccine at their LTCF

(indirectly protected) and same persons in the reference period (87 days before).

There were 126 and 276 events, respectively (supplementary Table S1). Detailed

information on the crude estimates and the adjusted cumulative risk in each group can

be found in the supplementary material (Figure S3 and Tables S2 and S3).

Indirect protection was estimated at 57.3% (48% to 66.3%) for the whole study-period.

There was no statistically significant reduction in risk in the first 14 days of follow-up

but it increased progressively thereafter, particularly after 28 days (as a proxy of full

immunization of vaccinated persons at the LTCF), when VE reached 79.5% (71.0% to

88.1%) overall and 81.4% (73.3% to 90.3%) for the group with no documented SARS-

CoV-2 infection before the beginning of follow-up (Table 1, Figure 1).

The estimated number of SARS-CoV-2 infections averted by vaccination was similar to

the one found in the vaccinated group for individuals without previous infection, of 11

.0 and 12.8 per 10,000 non-vaccinated persons per day in the intermediate periods

(Table 1).

It was not possible to estimate VE for indirect protection in the group with a previous

SARS-CoV-2 infection since there were only 14 events, confidence intervals virtually

tended to infinite, and the model did not result in credible risk curves.

Discussion

This study on the institutionalized elderly confirms the high benefit of vaccination in

this population, reducing the risk of infection by up to 81.2% and avoiding up to 9.6

cases per 10,000 population per day. The risk reduction was through direct protection

of those vaccinated but also through indirect protection of those who were not-

vaccinated. The vaccine effectiveness increased throughout the study period, likely

showing the progressive immunization of vaccinated persons with increasing time

elapsed since the first-dose and after the receipt of the second dose. While VE was

higher for individuals naïve to SARS-CoV-2, those with previous infection also benefited

from vaccination, even the absolute gain in number of infections averted was low,

possibly due to an already lower baseline risk in this group.

Immunesenescence and factors related to chronic conditions, together with

malnutrition, are known to impair immunity required for an effective vaccine response

[13], and lower neutralizing antibodies response to Pfizer/BioNTech vaccine in people

≥65 years has been reported [6,14]. However, our estimates resulted fairly similar to

those of observational studies in younger adult population and are consistent with

other studies showing high VE in the elderly from the general population. A cohort of

health care workers in the UK found a VE of 70% 21 days after the first dose and of

85% 7 days after the second dose of Pfizer/BioNTech [15]. A slightly higher estimate, of

94.1%, is given by a pre-print with data from Israel [16]. Other observational studies

have explored VE in older age groups. In a registries-based study from Israel, in

persons aged ≥70 years, VE was found to be 44%, 64% and 98% at 14-20 days post-

vaccination, 21-27 days post-vaccination and ≥7 days after the second vaccine-dose,

respectively, which were similar to the results for younger age groups [17]. Bernal et al

have reported vaccine effects to start 10-13 days after vaccination with

Pfizer/BioNTech and reach 61% in people aged ≥70 years and 70% in people aged ≥80

years ≥28 days post-vaccination, and 89% 14 days after the second vaccine-dose [8].

A study in LTCF in Connecticut experiencing COVID-19 outbreaks found a 63%

protection with partial vaccination (between 14 and 28 days of the first dose), close to

our estimates, with unchanged results after excluding those with previous infection

[9]. However, two other existing studies focusing on LTCF have reported lower VE. A

Danish study in pre-print [10], has found no protective effect of a first vaccine-dose, a

52% reduction in days 0-7 after the second dose and 64% beyond day 7. A recently

released pre-print manuscript from the VIVALDI study in the UK has found no

protection conferred by vaccination with the Pfizer/BioNTech vaccine in the first 28

days after the first dose [11]. Nevertheless, VE between days 28 to 47 was between

56% and 62% [11], in a similar range of the effect found in this study for the period 22-

28 days (61.9%). Early results from British Columbia have estimated 80% reduction in

risk 2-3 weeks after the first vaccine-dose [18]. Of note, our work included both

symptomatic and asymptomatic infections, pointing that risk was probably reduced for

both type of endpoints to an unknown degree. As an illustration, in national COVID-19

surveillance, 39% of all notified infections since 10 May 2020 in people ≥65 years of

age were asymptomatic.

A considerable 22% of all participants in our study had a previous documented SARS-

CoV-2 infection, although there are possibly a high number of infections that were not

documented, especially during the first epidemic wave in March-April 2020. Several

studies have documented a high immune response to a first COVID-19 vaccine-dose in

people with previous infection [19,20,21]. The results of this study add to previously

existing literature that, even though the effect was greater in naïve subjects to SARS-

CoV-2, those with previous infection also benefited from a risk reduction of 57%,

although it translated in less than 1 infection averted per 10,000 population per day.

Results from the indirect protection analysis support the hypothesis that vaccination

may reduce transmissibility of SARS-CoV-2 and result in herd immunity. Previous

studies have shown decreased viral load in vaccinated patients, including those in LTCF

[9, 22], and a study from Scotland found a 30% lower risk of SARS-CoV-2 in household

members of vaccinated health-care workers, although the reduction in SARS-CoV-2

transmission from vaccinated individuals could be double that estimate, since

household members could also have been infected in the community [23]. A recent

ecological study from Israel has shown that increasing vaccine coverage provides cross-

protection to unvaccinated individuals in the community [24]. In our study, non-

vaccinated individuals living in facilities where the majority (residents and staff) had

been vaccinated showed a risk reduction similar to those actually vaccinated.

However, the magnitude of protection may be overestimated, since non-vaccinated

individuals could correspond more frequently to persons with previous infection, even

if not documented. This could be controlled in individuals with documented infection

but not in an unknown number with non-documented infection or diagnosed with

serology. Also, indirect protection was measured in a context of very high vaccine

coverage, difficult to attain in a non-institutional setting; therefore our results may not

have generalisability to the community setting.

Some limitations to study results could relate to the before-after comparison. Even

though we tried to minimize it, residual confounding due to higher incidence during

the third epidemic wave and possibly, to the relaxation of the isolation of LTCF during

the Christmas season, with higher number of day-outs and visits, may be present and

could underestimate the protection of the vaccine. This underestimation of the effect

of the vaccine could maybe explain that the effect of natural infection in the non-

vaccinated group was found higher than the effect of the vaccine. The conservative

direction of the possible bias is shown by the bias indicator analysis (supplementary

material).

In conclusion, our results confirm the effectiveness of vaccination in institutionalized

elderly population, endorse the policy of universal vaccination in this setting, including

in people with previous infection, and suggest that even non-vaccinated individuals

benefit from indirect protection. Further questions include the duration of protection

in this population and according to previous infection, and the severity of infection,

which could not be measured in this study.

Acknowledgements

We would like to acknowledge the contribution of the General Directorate of Digital Health and National Health Service Information Systems and the General Sub-Directorate for Health Information (Ministry of Health, Spain) for their work in managing the registries in which this study was based, as well as in providing all the necessary IT support. We specially thank Miguel Hernán for his methodological advice and support in the design of the analysis as well as Luis Sanguiao for his support in implementing parallel computing. We acknowledge as well all the work of the professionals involved in the vaccination program and in managing the vaccination and laboratory registries from the Autonomous Communities.

References

[1] Instituto de Mayores y Servicios Sociales (Imserso). Actualización nº 2. Enfermedad por coronavirus (COVID-19) en Centros Residenciales. 7/3/2021. Available at: https://www.imserso.es/InterPresent2/groups/imserso/documents/binario/inf_resid_20210312.pdf (Accessed 31 March 2021)

[2] Grupo de Trabajo Técnico de Vacunación COVID-19, de la Ponencia de Programa y Registro de Vacunaciones 2 de diciembre 2020. Estrategia de vacunación frente a COVID19 en España. 2 de diciembre 2020. Available at: https://www.mscbs.gob.es/profesionales/saludPublica/prevPromocion/vacunaciones/covid19/docs/COVID-19_EstrategiaVacunacion.pdf (Accessed 31 March 2021)

 [3] Centro de Coordinación de Alertas y Emergencias Sanitarias. Ministry of Health of Spain. Actualización nº 335. Enfermedad por el coronavirus (COVID-19). 18.03.2021. Available at: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov/documentos/Actualizacion_335_COVID-19.pdf (Accessed 31 March 2021)

[4] GIV COVID-19. Gestión Integral de la vacunación COVID. Ministry of Health of Spain. Informe de actividad 18 de marzo. Available at: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov/documentos/Informe_GIV_comunicacion_20210318.pdf (Accessed 31 March 2021)

 [5] Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. N Engl J Med. 2020 Dec 31;383(27):2603-2615. doi: 10.1056/NEJMoa2034577.

[6] Prendki V, Tau N, Avni T, Falcone M, Huttner A, Kaiser L, at al. A systematic review

assessing the under-representation of elderly adults in COVID-19 trials. BMC Geriatr.

2020 Dec 20;20(1):538. doi: 10.1186/s12877-020-01954-5.

[8] Müller L, Andrée M, Moskorz W, et al. Age-dependent immune response to the

Biontech / Pfizer BNT162b2 COVID-19 vaccination. MedRxiv [Preprint] 2021. doi:

10.1101/2021.03.03.21251066.

[7] Bernal JL, Andrews N, Gower C, et al. Early effectiveness of COVID-19 vaccination

with BNT162b2 mRNA vaccine and ChAdOx1 adenovirus vector vaccine on

symptomatic disease, hospitalisations and mortality in older adults in the UK : a test

negative case control study. MedRxiv [Preprint] 2021.

DOI:10.1101/2021.03.01.21252652.

[9] Britton A, Jacobs Slifka K, Edens Ch et al. Effectiveness of the Pfizer-BioNTech

COVID-19 Vaccine Among Residents of Two Skilled Nursing Facilities Experiencing

COVID-19 Outbreaks — Connecticut, December 2020–February 2021. MMWR 2021;70:

Early Release March 15.

[10] Moustsen-Helms IR, Emborg HD, Nielsen J, Nielsen KF, Krause TG, Mølbak H, et al.

Vaccine effectiveness after 1st and 2nd dose of the BNT162b2 mRNA Covid-19 Vaccine

in long-term care facility residents and healthcare workers – a Danish cohort study.

MedRxiv [Preprint] 2021. doi: 10.1101/2021.03.08.21252200.

[11] Shrotri M, Krutikov M, Palmer T, Giddings R, Azmi B,Subbarao S, et al. Vaccine

effectiveness of the first dose of ChAdOx1 nCoV-19 and BNT162b2 against SARS-CoV-2

infection in residents of Long-Term Care Facilities (VIVALDI study). MedRxiv [Preprint]

2021. doi: 10.1101/2021.03.26.21254391.

[12] Hernán MA, Robins JM (2020). Causal Inference: What If. Boca Raton: Chapman &

Hall/CRC. Available at: https://www.hsph.harvard.edu/miguel-hernan/causal-

inference-book/ (Accessed 31 March 2021).

[13] Fulop T, Pawelec G, Castle S, Loeb M. Immunosenescence and vaccination in

nursing home residents. Clin Infect Dis. 2009 Feb 15;48(4):443-8. doi: 10.1086/596475.

[14] Walsh EE, Frenck RW, Falsey AR, *Kitchin N, Absalon J, Gurtman A, et al. Safety

and Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. New England

Journal of Medicine. 2020;383(25):2439-50.

[15] Hall VJ, Foulkes S, Saei A, Andrews N, Oguti B, Charlett et al. Effectiveness of

BNT162b2 mRNA Vaccine Against Infection and COVID-19 Vaccine Coverage in

Healthcare Workers in England, Multicentre Prospective Cohort Study (the SIREN

Study). Available at SSRN: https://ssrn.com/abstract=3790399 (Accessed 31 March

2021)

[16] Haas EJ, Frederick JA, McLaughlin JM, Anis E, Singer SR, Khan F, et al. Nationwide

Vaccination Campaign with BNT162b2 in Israel Demonstrates High Vaccine

Effectiveness and Marked Declines in Incidence of SARS-CoV-2 Infections and COVID-

19 Cases, Hospitalizations, and Deaths. Available at:

https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3811387 (Accessed 31 March

2021)

[17] Dagan N, Barda N, Kepten E, Miron O, Perchik S, Katz MA, et al. BNT162b2 mRNA

Covid-19 Vaccine in a Nationwide Mass Vaccination Setting. N Engl J Med. 2021 Feb 24.

doi: 10.1056/NEJMoa2101765. Online ahead of print. PMID: 33626250

[18] Early findings show the first vaccine dose reduced the risk of COVID-19 by 80 per

cent or more. Press release. February 19, 2021. Available at:

http://www.bccdc.ca/about/news-stories/news-releases/2021/early-findings-show-

the-first-vaccine-dose-reduced-the-risk-of-covid-19-by-80-per-cent-or-more (Accessed

31 March 2021)

[19] Manisty C, Otter AD, Treibel TA, McKnight Á, Altmann DM, Brooks T, et al.

Antibody response to first BNT162b2 dose in previously SARS-CoV-2-infected

individuals. The Lancet. 2021;397(10279):1057-8.

[20] Saadat S, Tehrani ZR, Logue J, *Newman M, Frieman MB, Harris AD, et al. Single

Dose Vaccination in Healthcare Workers Previously Infected with SARS-CoV-2. medRxiv

[Preprint] 2021. doi: 10.1101/2021.01.30.21250843.

[21] Camara C Lozano-Ojalvo D, Lopez-Granados E. Differential effects of the second

SARS-CoV-2 mRNA vaccine dose on T cell immunity in naïve and COVID-19 recovered

individuals. Biorxiv [Preprint] 2021. doi: 10.1101/2021.03.22.436441.

[22] Petter E, Mor O, Zuckerman N, Oz-Levi D, Younger A, Aran D, Erlich Y. Initial real

world evidence for lower viral load of individuals who have been vaccinated by

BNT162b2. MedRxiv [Preprint] 2021. doi: 10.1101/2021.02.08.21251329 .

[23] Shah AS, Gribben C, Bishop J, Hanlon P, Caldwell D, Wood R, et al. Effect of

vaccination on transmission of COVID-19: an observational study in healthcare workers

and their households. MedRxiv [Preprint] 2021. doi: 10.1101/2021.03.11.21253275.

[24] Milman O, Noga Aharony IY, Katz R, Herzel E, Ben-Tov A, Kuint J, et al. SARS-CoV-2

infection risk among unvaccinated is negatively associated with community-level

vaccination rates. MedRxiv [Preprint] 2021. doi: 10.1101/2021.03.26.21254394.

Table 1. Vaccine effectiveness (VE) and risk difference (RD) in residents of elderly long-term care facilities according to evidence of previous

infection and time since first vaccinated (as a proxy of number of vaccine - doses and days since last dose).

		VACCINE EFFECTIVENES (%)			RISK DIFFERENCE (per 10,000 persons per day)		
		Overall	No previous infection	Previous infection	Overall	No previous infection	Previous infection
EFFECTS IN THE VACCINATED	Full period	57.2% (56.1% to 58.3%)	57.6% (56.6% to 58.6%)	36.3% (27.9% to 45.5%)	-6.26 (-6.45 to -6.06)	-7.37 (-7.58 to -7.16)	-0.64 (-0.86 to -0.44)
	Days 0-14	28.5% (26.4% to 30.7%)	28.9% (26.9% to 31%)	9.6% (-6.9% to 26.8%)	-5.06 (-5.52 to -4.57)	-6.05 (-6.56 to -5.54)	-0.25 (-0.72 to 0.23)
	Days 15-21	51.0% (49.7% to 52.3%)	51.9% (50.7% to 53.1%)	25.5% (15.1% to 36.6%)	-9.62 (-9.97 to -9.23)	-11.59 (-12.01 to -11.19)	-0.66 (-1.00 to -0.32)
	Days 22-28	61.9% (60.8% to 63%)	62.9% (61.9% to 64%)	34.6% (25.7% to 44.1%)	-9.65 (-9.92 to -9.35)	-11.59 (-11.92 to -11.28)	-0.76 (-1.03 to -0.5)
	Days ≥29	81.2% (80.2% to 82%)	81.8% (81.0% to 82.7%)	56.8% (47.1% to 67.7%)	-5.59 (-5.76 to -5.41)	-6.47 (-6.66 to -6.28)	-0.75 (-0.98 to -0.53)
INDIRECT EFFECTS	Full period	57.3% (48% to 66.3%)	58.7% (49.4% to 68.5%)	NA	-8.13 (-10.13 to -5.98)	-10.08 (-12.62 to -7.52)	NA
	Days 0-14	18.8% (-1.7% to 39.9%)	18.2% (-3.1% to 39.8%)	NA	-3.31 (-7.29 to 0.79)	-3.79 (-8.54 to 1.14)	NA
	Days 15-21	43.6% (31.3% to 55.5%)	45% (32.8% to 57.1%)	NA	-8.8 (-11.95 to -5.48)	-11.02 (-14.88 to -6.99)	NA
	Days 22-28	55.8% (45.2% to 65.9%)	57.8% (47.5% to 68.2%)	NA	-10.06 (-12.66 to -7.30)	-12.81 (-16.16 to -9.39)	NA
	Days ≥29	79.5% (71.0% to 88.1%)	81.4% (73.3% to 90.3%)	NA	-9.22 (-11.56 to -6.73)	-11.46 (-14.39 to -8.6)	NA

430 NA: The model did not result in plausible bias-free risk curves, therefore no estimations were drawn.

	Figure 1. Cumulative incidence of documented SARS-CoV-2 infection in residents in long-term care facilities estimated from adjusted hazards

	models.