SARS-CoV-2 Detection from the Built Environment and Wastewater and Its Use for Hospital

Surveillance

Aaron Hinz[1,6], Lydia Xing[2], Evgueni Doukhanine[3], Laura A. Hug[4], Rees Kassen[1], Banu Ormeci[5],

Richard J. Kibbee[5], Alex Wong[6], Derek MacFadden*[2,7], Caroline Nott*[2, 7]

*Co-PIs

1. Department of Biology, University of Ottawa, Ottawa, ON Canada

2. Division of Infectious Diseases, Department of Medicine, University of Ottawa, ON

Canada

3. DNA Genotek, Inc., Ottawa, ON, Canada

4. Department of Biology, University of Waterloo, ON Waterloo

5. Department of Civil and Environmental Engineering, Carleton University, Ottawa, ON

Canada

6. Department of Biology, Carleton University, Ottawa, ON Canada

7. The Ottawa Hospital Research Institute, Ottawa, ON Canada

Corresponding Author: Aaron Hinz, Department of Biology, University of Ottawa, 30 Marie-

Curie, Ottawa, ON K1N6N5. Email: ahinz@uottawa.ca

ABSTRACT

Background. SARS-CoV-2 causes significant morbidity and mortality in health care settings.

Our understanding of the distribution of this virus in the built healthcare environment and

wastewater, and relationship to disease burden, is limited.

Methods. We performed a prospective multi-center study of environmental sampling of SARS-

CoV-2 from hospital surfaces and wastewater and evaluated their relationships with regional

and hospital COVID-19 burden. We developed and validated a qPCR-based approach to

surface sampling, and swab samples were collected weekly from different locations and

surfaces across two tertiary care hospital campuses for a 10-week period during the pandemic,

along with wastewater samples.

Results. Over a 10-week period, 963 swab samples were collected and analyzed. We found 61

(6%) swabs positive for SARS-CoV-2, with the majority of these (n=51) originating from floor

samples. Wards that actively managed patients with COVID-19 had the highest frequency of

positive samples (p<0.01). Detection frequency in built environment swabs was significantly

associated with active cases in the hospital throughout the study (p<0.025). Wastewater viral

signal changes appeared to predate change in case burden.

Conclusions. Environment sampling for SARS-CoV-2, in particular from floors, may offer a

unique and resolved approach to surveillance of COVID-19.

Key Words: SARS-CoV-2, COVID-19, environmental screening, burden, wastewater

BACKGROUND

COVID-19 has caused substantial morbidity and mortality worldwide since it was first identified at the end of 2019. Despite the development of effective vaccines and the discovery of multiple therapeutic agents[1,2], the SARS-CoV-2 virus and agent of COVID-19 continues to transmit across all corners of the globe. As of Spring 2021, it is estimated that over 100 million people have been infected with SARS-CoV-2 from the start of the pandemic, with over 2.5 million deaths attributable to COVID-19[3].

Hospitals were recognized early on as a potentially high-risk environment for acquiring COVID-19 infection for patients and healthcare workers, with healthcare facility outbreaks and nosocomial acquisition of COVID-19 common and well described[4]. While the respiratory droplet route is considered to be a primary means for transmission of SARS-CoV-2 in hospitals, it is still not well established if there is appreciable transmission through contact with the built environment[4,5], and to what extent this could contribute to spread.

Laboratory experiments indicate that the virus can remain viable on stainless steel and plastics for approximately 6 hours and SARS-CoV-2 RNA can be detected from wastewater[6]. Preliminary studies have shown a range from no to extensive contamination of hospital ward surfaces, including computer mice, trashcans, and door handles[7]. In addition to identifying the potential routes for fomite-based transmission, surveying the distribution of SARS-CoV-2 throughout the hospital environment over time can provide insight into hospital burden and potentially outbreak identification and monitoring. New approaches for early outbreak detection will be particularly important to reduce institutional morbidity and mortality[8]. The built environment presence of SARS-CoV-2 could act as a spatially resolved indirect measure of symptomatic and asymptomatic carriage/infection within patients and healthcare workers

(HCWs). In a similar fashion, wastewater surveillance may offer a less spatially resolved but

potentially anticipatory mechanism for hospital COVID-19 surveillance and monitoring[9,10].

In this study, we conducted a longitudinal assessment of the prevalence of SARS-CoV-2 on

commonly contacted hospital surfaces as well as from hospital wastewater and compared these

markers with active COVID-19 case prevalence at two large tertiary care hospitals and in the

surrounding community.

METHODS

SARS-CoV-2 control samples

Control samples of heat-inactivated SARS-CoV-2 (ATCC VR-1986HK; lot 70035039) and

SARS-CoV-2 RNA (ATCC VR-1986D; lot 70035624) were obtained from Cedarlane

Laboratories. The copy numbers of the control samples were reported as 4.73×10^3 genome

copies/µL for the RNA and 3.75×10^5 genome copies/µL for the heat-inactivated virus.

Swabbing method and validation

Surfaces were sampled with P-208 Environmental Surface Collection Prototype kit from DNA

Genotek, consisting of a flocked swab and a semi-lytic nucleic acid stabilization solution for

post-collection swab immersion. The swabbing method was validated in laboratory experiments

using surface materials spiked with heat-inactivated SARS-CoV-2. We applied 10^5 copies of the

virus to sheets of acrylic, stainless steel, and vinyl upholstery by pipetting 150 µl of virus

suspension in 30-40 small spots within a 2.25 in by 2.25 in square. After a 2.5 hour drying

period, swabs were pre-wetted with stabilization solution, surfaces were swabbed for 30

seconds, and swabs were stored in stabilization solution in collection vials. As controls, sterile

swabs were placed in collection vials containing virus spiked directly into the stabilization

solution. Following one day of storage at room temperature, RNA was extracted and analyzed

by qPCR. The amount of virus in each sample was estimated using the virus standard curve

that relates Cq values to input genome copies. The percent viral recovery was determined by

dividing the amount of virus estimated for the surface swabbed samples by the amount of virus

estimated in the control samples. Significant differences in recovery efficiencies across surfaces

were determined by ANOVA with post hoc comparisons using Tukey's honestly significant

difference (HSD) test.

RNA extractions and RT-qPCR

Presence of SARS-CoV-2 RNA was determined by quantitative reverse-transcriptase PCR (RT-

qPCR) of RNA extracted from swab sample stabilization solution. RNA was extracted from

samples using the MagMAX Viral/Pathogen II (MVP II) Nucleic Acid Isolation Kit (Thermo-Fisher

Scientific) according to the manufacturer's protocol, with reagent volumes modified

proportionally for 300 µl sample input volumes. Extractions were performed in batches of up to

96 in deep-well plates. Each batch included positive (heat-killed SARS-CoV-2) and negative

(stabilization solution only) control samples to monitor extraction efficacy and potential cross-

contamination. None of the control samples yielded false positive or false negative results.

Quantitative PCR was performed with primers and a TaqMan probe targeting the N gene of

SARS-CoV-2 (Table 1). The probe contained a 6-carboxyfluorescein (FAM) fluorescent dye at

121 the 5' end and the minor groove binder non-fluorescent quencher (MGB-NFQ) at the 3' end.

122 Each qPCR reaction consisted of 5 µl of extracted RNA, 300 nM of each primer, 200 nM of

123 probe, and 5 µl of 4X TaqPath 1-Step Multiplex Master Mix (No ROX) (Applied Biosystems) in a

124 final volume of 20 µl. Thermal cycling, fluorescence quantification, and quantitative cycle (Cq)

125 determination were performed on a Bio-Rad CFX Connect with the following cycling conditions:

126 25 °C for 2 min, 50 °C for 15 min, 95 °C for 2 min, and 45 cycles of denaturation (95 °C for 3 s)

127 and elongation (55 °C for 30 s).

129 **Estimates for Limits of Detection and Quantification**

130 Analytical sensitivity and precision of the qPCR assays were determined by estimating the limits

131 of detection and quantification of the SARS-CoV-2 RNA and virus standards. Serial 10-fold

132 dilutions of the SARS-CoV-2 RNA standard were analyzed by qPCR with 14-20 replicates per

133 dilution. For the virus standard, RNA was extracted from serial dilutions of heat-inactivated virus

134 (7 replicate extractions at each dilution) prior to qPCR analysis.

136 The qPCR results were analyzed as described by Klymus et al.[12] using the Limit of Detection

137 (LOD) Calculator (R script available at: https://github.com/cmerkes/qPCR_LOD_Calc). The

138 standard curves were modeled as linear regressions based on the middle two quartiles of Cq

139 values for each standard concentration exhibiting >50% detection. The limit of detection was

140 defined as the lowest input amount that was detected in >95% of replicates, and the limit of

141 quantification was defined as the lowest input amount yielding reproducible Cq values with a

142 coefficient of variation less than 0.35. For the RNA standards, the LOD with 95% confidence

143 intervals and limit of quantitation (LOQ) were determined by a curve-fitting approach as

described[12]. The virus LOD was estimated using a discrete threshold (*i.e.*, the lowest tested

input amount with >95% detection).

Built Environment Sample Collection

We prospectively collected built environment swabs from two large tertiary-care, academic

hospital campuses (Campus A and Campus B) from Ottawa, Ontario, Canada over a 10-week

period between September 28 and December 6, 2020. We obtained between 40-50 samples

from each campus weekly, generally maintaining location and surface type longitudinally,

though we did not attempt to sample the exact same item longitudinally in order to improve our

study generalizability. For example, a computer keyboard sampled in week 1 on Unit A, may be

different from a computer keyboard sampled on Unit A in week 2. At both campuses, we

collected samples from 1 COVID-dedicated unit, 1 ICU (COVID-designated area), 1 Emergency

Department (ED) area, non-COVID medical and surgical units (n=3 at Campus A, n=4 at

Campus B), 1 ambulatory unit (dialysis), public/non-patient-care spaces within the hospital (e.g.

main elevators), and exterior hospital grounds (e.g. parking garage). A designated HCW space

was also sampled at Campus A. Built environment samples were collected by wetting a P-208

kit flocked swab with the stabilization media, and sweeping the swab tip across an

approximately 2" x 2" area (dependent upon the item/surface size) for approximately 30

seconds. The swab was then transported in the P-208 stabilization media, and stored at

ambient temperature to be processed/analyzed within 1 month of collection. Three study

authors (LX, DM, CN) collected the samples during the study period, and swab technique was

harmonized between them at onset of sample collection. A run-in week was performed prior to

week 1, in order to test sample collection, logistics, and sample processing, but is not included

in this analysis as a number of specimen sites were initially not included for this week, making

inter-week comparisons challenging.

Specific (non-critical care) wards in the hospital were designated for the ongoing inpatient care

of COVID patients, and these were designated as 'COVID Wards'. However, we classified both

the COVID Wards as well as the ICU and ED as 'COVID Risk Units', because they were

anticipated to be managing a large number of SARS-CoV-2 infected patients.

Hospital Practices

The visitor policy at the start of the study period limited visitation to 1 visitor/patient/day;

however, at the start of study week 4, visitors were further restricted to essential caregivers

only. Environmental cleaning was in keeping with provincial best practice and occurred daily in

both public and clinical areas, with all contact points disinfected with Vert-2-Go© everyday

disinfectant (quaternary-ammonium based product) and floors mopped with Vert-2-Go© Oxy

floor cleaner. The only deviation from this was once weekly cleaning of the Campus A parking

garage until sometime between study weeks 5-7, then once daily thereafter. Infection

Prevention and Control (IPAC) policies were also consistent with provincial best practice[6,13] .

Throughout the study period, all staff, visitors, and patients were screened for COVID-19

symptoms and risk factors (e.g. exposures, international travel) on entry to the hospital, in

accordance with Ontario Ministry of Health guidance[14]. Universal masking for all staff and

visitors was introduced prior to the study period. Universal eye protection (visor or goggles) for

staff working in clinical areas was introduced at the start of study week 1.

Wastewater Sample Collection

Raw sewage grab samples from Campus A were taken with a 5L bucket on a rope between

11:10am and 12:30pm each sampling day, which occurred once every 1 to 2 weeks during (and

immediately prior to) the study period. The samples were swirled in the bucket and quickly

poured into pre-labelled sterile 1L Nalgene bottles which were immediately placed into a cooler

with ice packs for transport back to the lab and processed within 1 hr. Temperature of the

samples, measured at the time of sampling with an infrared thermometer, for Oct-Dec were

between 12 and 22 ºC. A 200ml aliquot was concentrated to 0.25-1.25ml using the Centricon®

Plus-70 (MilliporeSigma) centrifugal ultrafilter with a cut-off of 10 kDa[15]. The resulting

concentrate was extracted using the RNeasy PowerMicrobiome Kit (QIAGEN) according to the

manufacturer's instructions with an elution of 100µl.

Primers and probes, and Real-Time Reverse Transcriptase (RT RT-PCR) methods developed

by the US CDC were used with the Bio-Rad CFX96 Touch Real-Time PCR Detection System.

The RT RT-PCR thermal cycling protocol used to amplify the target region within the SARS-

CoV-2 nucleocapsid gene, N2, was carried out at 50 °C for 30 minutes, followed by 95 °C for 15

minutes and 50 cycles of 95 °C for 15 and 60 °C for 30 seconds. Reaction mixture for RT RT-

PCR, 5 µl RNA template, 4 µl of QIAGEN OneStep RT-PCR 5x Buffer, 2 µl dNTP mix (10 mM

each), 2 µl Enzyme Mix, Primers and Probes were used at 500nM/reaction, and 2 µl of 4 mg/ml

BSA. Each reaction volume was adjusted to a final volume of 20 µl with RNase-free water. Each

sample was run in duplicate. An MS-2 bacteriophage matrix spike whole process control was

used to assess the efficiency of the concentration process and RNA extraction as well as

confirm RT RT-PCR protocols. RT RT-PCR inhibition control using MS2 bacteriophage spiked

into raw sewage RNA extracts and molecular grade water was used to assess the performance

of the RNA-extraction, RT-PCR and to detect the presence of inhibitors. Quantification of the N2

gene target in the raw sewage sample concentrates was done using standard curves created

with a 2019-nCoV_N_Positive Control plasmid (IDT, Inc.) Non-template controls (NTC) were run

for each mastermix preparation with molecular grade water as the template. For each week

under observation, we averaged the detectable copies/ml from both N1 and N2 gene targets.

Hospital and Citywide COVID-19 Cases and Outbreaks

Daily active patient COVID-19 case counts and locations, including admitted and non-admitted

Emergency Department cases, were obtained from Infection Prevention and Control (IPAC).

Cases were those confirmed by RT-PCR. We generated a measure of weekly 'active' COVID-

19 cases by taking the mean of these daily active cases for each week (typically data only

available for weekdays). Healthcare worker (HCW) cases were not included in the analysis, as

HCWs were more likely than patients to spend time on multiple units and move throughout the

hospital. Also, once a HCW became symptomatic or had a positive COVID-19 test result, they

would no longer remain within the hospital. Hospital outbreak data were obtained in real time

from IPAC and verified post-study on the publicly available Ottawa Public Health (OPH) COVID-

19 dashboard[16]. An outbreak was defined as per the Ontario Ministry of Health[17] as, "Two or

more laboratory-confirmed COVID-19 cases (patients and/or staff) within a specified area

(unit/floor/service) within a 14-day period where both cases could have reasonably acquired

their infection in the hospital". New and total active COVID-19 cases in Ottawa were obtained

daily from the OPH COVID-19 dashboard, and a weekly 'active' caseload was calculated as the

mean of the daily active cases across a given week.

Data Analysis

To evaluate the characteristics associated with positive swab results, we performed a linear

regression model using specimen characteristics as predictors of built environment sample

positivity (total or floor specific specimens). Specimen characteristics were coded as categorical

variables, including unit COVID Risk type, object/site, material, study week. The outcome

(sample positivity) was coded as a binary variable (1=detected, 0=not-detected). A linear model

was used to predict the outcome variable as it provided improved interpretability of coefficients

and also model convergence stability[18].

We also evaluated the strength of associations between (i) proportion of hospital built

environment swabs positive for SARS-CoV-2 and (ii) proportion of hospital floor swabs positive

for SARS-CoV-2, and hospitalized active COVID-19 cases. We used a negative-binomial

regression model with active cases as the outcome. We did not quantitatively evaluate the

association between wastewater and active cases because of the limited number of wastewater

data points.

For the majority of the analyses we analyzed the two campuses in combination, as they reflect a

single health care center with transfer occurring within and between units in both campuses. We

also chose not to evaluate case burden by specific unit because, in general, as soon as a case

was identified on a given non-COVID risk unit, the patient would be transferred to a specific

COVID Risk unit either for cohorting or due to severity of illness (e.g. ICU).

All statistical analyses were performed using R (Version 4.0.3). Research ethics board approval

was obtained through the Ottawa Health Science Network Research Ethics Board (OHSN-REB)

for this study.

RESULTS

Limit of Detection and Swab Validation

We detected and quantified SARS-CoV-2 RNA by qPCR analysis using the N1 primer and

probe set targeting the nucleocapsid gene[11]. A standard curve relating the quantification cycle

(C_q) values to viral genome copy number was produced by analyzing a 10-fold dilution series of

a SARS-CoV-2 RNA standard (Fig. 1A). We used a model-fitting approach to determine the

LOD, defined as a detection rate greater than 95%, and LOQ defined as a C_q coefficient of

variation less than 0.35[12]. Reactions containing at least 3 genome copies were reliably detected

under our qPCR conditions, whereas precise quantification required at least 150 copies (C_q <

30) (Fig. 1A).

Analysis of the built environment swab samples involved a nucleic acid extraction step prior to

qPCR analysis. We evaluated the extraction method by qPCR analysis of RNA extracted from

samples of a heat-inactivated SARS-CoV-2 virus standard (Fig. 1B). The LOD for the virus

standard (28 copies per reaction) was ~10-fold higher than that of the RNA standard, which can

be due to RNA loss during the extraction procedure, presence of PCR inhibitors in the eluant, or

inaccuracies in the reported titers of the standards. Nevertheless, the procedure was sensitive

and reproducible, and the results yielded a linear standard curve for converting C_q values to viral

copies while accounting for inaccuracies introduced by sample processing.

The built environment sampling approach relied on recovery of virus particles and nucleic acid

from surfaces. In preliminary tests, we determined how efficiently our swabbing technique

recovered virus from surface types representative of the built environment: acrylic, stainless

steel, and vinyl. The surfaces were spiked with equal amounts of heat-inactivated virus, allowed

to dry, and swabbed by the procedure used for built environment sampling. Following incubation

of the swabs in stabilization solution and qPCR analysis of extracted RNA, we estimated the

amount of recovered virus using the virus standard curve and calculated the percent of virus

recovered from each surface (Fig. 2). The mean recovery efficiencies ranged from 28 to 42%

across the three surfaces, values similar to those reported in a previous study of SARS-CoV-2

surface swabbing[19].

Built Environment Screening

Over a 10-week period from September 28th to December 6th 2020 we systematically

performed 963 built environment swabs across 2 tertiary care hospital sites. SARS-CoV-2 was

detected from 6% of swabs (61/963), and was most commonly found from floor samples, with a

detection prevalence of 27% (51/188). SARS-CoV-2 was less frequently identified from elevator

buttons, benches, items (*e.g.*, hand sanitizer pumps), and computer keyboards.

COVID Risk units, namely COVID treatment wards, intensive care units, and emergency rooms,

were the most common locations to find SARS-CoV-2 in the hospital environment, with

detection prevalence of 18%, 8%, and 7% respectively (Table 2, Fig. 3). Non-COVID and

public/HCW spaces tended to have a low frequency of SARS-CoV-2 detection from the

environment. During the course of this study, there were 2 outbreaks involving study units, at

Campus B in the ED (1 patient and 2 HCW) and at Campus A on Ward 3 (5 patients and 1

HCW). A third outbreak occurred on a non-COVID inpatient unit at Campus A but this unit was

not sampled in the study.

Across the 10-week study period, SARS-CoV-2 detection prevalence paralleled total active

cases across the two hospital campuses. Viral prevalence gradually declined over time, but with

spikes in hospitalized cases being clearly reflected in built environment burden (Figure 4A), as

well as an increase reflecting the start of hospitalizations in the second wave of the pandemic

within the city. Built environment burden also paralleled citywide cases, but with less

concordance (Figure 4B). Campus B tended to have SARS-CoV-2 positivity largely restricted to

COVID Risk units, but Campus A seemed to have more activity in non-COVID risk units/public

spaces. A multi-variable logistic regression model identified that increasing study week was

associated significantly with reduced detection of SARS-CoV-2 ($p<0.05$), and that COVID Risk

units and Campus A were significantly associated with increased SARS-CoV-2 detection (Table

3, Table 4).

Hospital floors were the most common surface from which SARS-CoV-2 was detected. We

looked specifically at prevalence on floors and its relationship with cases within the hospital and

citywide (Figure 5). Restricting analysis to only floors showed they parallel the findings seen

with all samples (Figures 4), with stronger correlation with active hospital cases.

Hospital wastewater SARS-CoV-2 viral copies per ml (Figure 6) followed a similar trajectory as

built environment sampling but decline in wastewater levels appeared to predate both the built

environment sampling as well as Campus A and citywide case burden.

We also evaluated the strength of association between (i) all built environment swab positivity

and (ii) floor swab positivity with hospital case burden (Table 4). We found that floor swabs best

described the hospital case burden, having a measure of association with the greatest

precision, and the model with the lowest AIC (Table 4).

DISCUSSION

In this prospective longitudinal study at two large tertiary care hospital campuses, we

demonstrate that SARS-CoV-2 RNA can be readily identified from a variety of hospital surfaces

as well as hospital wastewater. Moreover, both the relative frequency of isolation on surfaces

and the viral titre in wastewater parallel the burden of infection within the hospital. Wards that

specifically managed COVID-19 patients had consistently higher levels of detection compared

to other areas, and frequency of detection paralleled case counts over time. For the single ward-

based outbreak that occurred during the study period, there was detectable viral RNA at the

involved location from the onset and until resolution.

Prior studies have demonstrated the amplification of SARS-CoV-2 RNA from the hospital

environment[7] including high touch surfaces outlined in this paper (*e.g.*, elevator buttons,

computer keyboards, alcohol hand pumps). Floors have been identified as a common location

for identifying SARS-CoV-2 in hospitals, and our results reinforce this finding[7]. Moreover, we

found that performing surveillance from floors provided the best marker for predicting hospital

burden of infection.

To date, fomites have not been identified as a major contributor to transmission of SARS-CoV-2

in the hospital or other settings[5]. While our PCR-based approach for SARS-CoV-2 RNA

detection cannot identify whether viral particles present in our samples were viable or not, our

results do indicate that identifying any RNA from high touch surfaces is relatively rare, further

supporting these as unlikely pathways for transmission in hospitals. Floors, however, were a

common location for SARS-CoV-2 RNA in our study. It is likely they represent a sink for viral

particles which originate from the respiratory droplets/aerosol from patients, health workers, or

visitors, and it is unlikely that floors represent a reservoir for transmission[7].

Interestingly, the non-COVID ward, Ward 3 on Campus A, that experienced the outbreak had

frequent detection of SARS-CoV-2 during the outbreak period, with general absence before and

after. This suggests that built environment sampling may offer a way to potentially identify occult

outbreaks or transmission occurring within patients/visitors/healthcare workers or for monitoring

outbreak status. The outbreak that occurred in the ED at Campus B did have a brief but non-

sustained signal, and we believe this is related to the transient nature of patients (and flux of

staff) within the ED. Patients are flowing from the ED itself to either discharge or transfer to

other sites within the hospital, thus it is unlikely to sustain a patient-based outbreak within that

area. The reason for higher detection of SARS-CoV-2 at Campus A is likely driven by more

frequent detection at non-COVID risk unit sites. The reason for this is unclear and may relate to

both patient and visitor populations.

Where built environment sampling can provide a resolved view of COVID-19 burden in the

hospital, hospital wastewater can provide institution-wide surveillance with the possibility for

anticipatory (predictive) signals. Qualitatively, wastewater appeared to anticipate burden by 1-2

weeks, and is on the order of magnitude reported in the literature[20], though given the limited

number of time points, we were not able to confirm this statistically. A growing number of

studies have shown the potential benefit of wastewater monitoring in both regional as well as

facility-based surveillance[20,21], and our study links this surveillance as complementary to built

environment screening. Combined approaches may ultimately be applicable to other respiratory

viruses that cause significant institutional outbreaks, including influenza, though further

validation is needed.

Our study has some limitations which have not already been noted. Firstly, our study was

performed during a period when there was relatively low activity, with regional case counts

ranging from approximately 2 to 7 cases per 100,000 population per week, and it is possible that

in higher burden settings there could be saturation of ward-based built environment detection

which could limit the correlation with burden or outbreaks. Secondly, outbreaks were relatively

rare in our hospital, owing in part to the low regional burden, and this limits our ability to

evaluate the ability of built environment screening to consistently detect outbreaks. Lastly, our

approaches only detect viral RNA and not viable virus. However, our findings benefit from the

generalizability of our approaches which use standard PCR techniques and could be easily and

quickly performed in most settings worldwide.

In conclusion, we have demonstrated that the detection of SARS-CoV-2 from wastewater and

built environment samples, in particular from floors, parallels hospital burden. Built environment

sampling may offer a spatially-resolved method for surveillance of hospital COVID-19 cases

which can be paired with wastewater monitoring for a comprehensive picture of facility-wide

burden of infection. Further prospective studies are needed to evaluate this approach.

Acknowledgements. We would like to thank Dr. Kathryn Suh and Dr. Raphael Saginur for their

support and advice during this study. We would also like to thank the City of Ottawa Wastewater

Collection Branch, and in particular Benjamin Musyoka and Sandra Gay, for accomodating the

collection of wastewater samples.

Author's contributions. All authors contributed to the design of the study. LX, DM, CN

collected built environment samples for the study and performed data entry. RJK and AH

processed samples for the study. DM and CN contributed to data analyses. All authors

contributed to the writing of the manuscript.

Financial support. This work was supported by The Ottawa Hospital Academic Medical

Organization (TOHAMO), Alliance Grant # 554478 - 20 from the Natural Sciences and

Engineering Research Council (NSERC), Carleton University Rapid Response Research Grant,

and the Jarislowsky Foundation.

Potential conflicts of interest. Author ED works for DNA Genotek which provided sampling

swabs in-kind for this study in an unrestricted fashion. DNA Genotek had no control over the

findings, interpretations, or conclusions published in this paper. All other authors have no

relevant conflicts to disclose.

REFERENCES

1. Siemieniuk, R. A. C. *et al.* Drug treatments for covid-19: living systematic review and network meta-analysis. *BMJ* **370**, (2020).

2. Baden, L. R. *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N. Engl. J. Med.* **384**, 403–416 (2021).

3. The Visual and Data Journalism Team. Covid map: Coronavirus cases, deaths, vaccinations by country. *BBC News* https://www.bbc.com/news/world-51235105 (2021).

4. Islam, M. S. *et al.* Current knowledge of COVID-19 and infection prevention and control strategies in healthcare settings: A global analysis. *Infect. Control Hosp. Epidemiol.* **41**, 1196–1206 (2020).

5. Mondelli, M. U., Colaneri, M., Seminari, E. M., Baldanti, F. & Bruno, R. Low risk of SARS-CoV-2 transmission by fomites in real-life conditions. *Lancet Infect. Dis.* (2020) doi:10.1016/S1473-3099(20)30678-2.

6. van Doremalen, N. *et al.* Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N. Engl. J. Med.* **382**, 1564–1567 (2020).

7. Guo, Z.-D. *et al.* Aerosol and Surface Distribution of Severe Acute Respiratory Syndrome Coronavirus 2 in Hospital Wards, Wuhan, China, 2020. *Emerg. Infect. Dis.* **26**, 1583–1591 (2020).

8. Smith, D. R. M. *et al.* Optimizing COVID-19 surveillance in long-term care facilities: a modelling study. *BMC Med.* **18**, 1–16 (2020).

9. Westhaus, S. *et al.* Detection of SARS-CoV-2 in raw and treated wastewater in Germany - Suitability for COVID-19 surveillance and potential transmission risks. *Sci. Total Environ.* **751**, 141750 (2021).

10. Gonçalves, J. *et al.* Detection of SARS-CoV-2 RNA in hospital wastewater from a low COVID-19 disease prevalence area. *Sci. Total Environ.* **755**, (2021).

11. CDC. Labs. https://www.cdc.gov/coronavirus/2019-ncov/lab/rt-pcr-panel-primer-probes.html

(2020).

12. Klymus, K. E. *et al.* Reporting the limits of detection and quantification for environmental

DNA assays. *Environmental DNA* **2**, 271–282 (2020).

13. Public Health Ontario. https://www.publichealthontario.ca/-/media/documents/b/2020/bp-

novel-respiratory-infections.pdf?la=en.

14. Ontario Ministry of Health. http://health.gov.on.ca/en/pro/programs/publichealth/

coronavirus/docs/2019_reference_doc_symptoms.pdf.

15. Medema, G., Heijnen, L., Elsinga, G., Italiaander, R. & Brouwer, A. Presence of SARS-

Coronavirus-2 RNA in sewage and correlation with reported COVID-19 prevalence in the

early stage of the epidemic in the Netherlands. *Environ. Sci. Technol. Lett.* **7**, 511–516

(2020).

16. Daily COVID-19 Dashboard. https://www.ottawapublichealth.ca/en/reports-research-and-

statistics/daily-covid19-dashboard.aspx (2021).

17. Ontario Ministry of Health. http://www.health.gov.on.ca/en/pro/programs/publichealth/

coronavirus/docs/2019_acute_care_guidance.pdf.

18. Gomila, R. Logistic or linear? Estimating causal effects of experimental treatments on

binary outcomes using regression analysis. *J. Exp. Psychol. Gen.* (2020)

doi:10.1037/xge0000920.

19. Parker, C. W. *et al.* End-to-End Protocol for the Detection of SARS-CoV-2 from Built

Environments. *mSystems* **5**, (2020).

20. SARS-CoV-2 RNA in wastewater anticipated COVID-19 occurrence in a low prevalence

area. *Water Res.* **181**, 115942 (2020).

21. Sharif, S. *et al.* Detection of SARs-CoV-2 in wastewater, using the existing environmental

surveillance network: An epidemiological gateway to an early warning for COVID-19 in

communities. *medRxiv* 2020.06.03.20121426 (2020).

Table 1. qPCR primer and probe sequences

Name	Sequence (5' to 3')*	Reference
N1-Forward	GACCCCAAAATCAGCGAAAT	USCDC[11]
N1-Reverse	TCTGGTTACTGCCAGTTGAATCTG	USCDC[11]
N1-Probe	6FAM-ACCCCGCATTACGTTTGGTGGACC-MGBNFQ	USCDC[11]

480 *6FAM, 6-carboxyfluorescein; MGBNFQ, Minor Groove Binder non-fluorescent quencher

482 **Table 2.** Built environment swab SARS-CoV-2 detection by location and characteristics.

	Campus A		Campus B		Total	
	Detected (N=39)	Not-Detected (N=451)	Detected (N=22)	Not-Detected (N=451)	Detected (N=61)	Not-Detected (N=902)
Unit COVID Risk Type						
COVID Risk Unit	17 (43.6%)	150 (33.3%)	17 (77.3%)	126 (27.9%)	34 (55.7%)	276 (30.6%)
Non-COVID Unit	9 (23.1%)	221 (49.0%)	5 (22.7%)	266 (59.0%)	14 (23.0%)	487 (54.0%)
Public/HCW Area	13 (33.3%)	80 (17.7%)	0 (0%)	59 (13.1%)	13 (21.3%)	139 (15.4%)
Object/Site						
Air (Control)	0 (0%)	10 (2.2%)	0 (0%)	10 (2.2%)	0 (0%)	20 (2.2%)
Computer Keyboard	0 (0%)	75 (16.6%)	1 (4.5%)	77 (17.1%)	1 (1.6%)	152 (16.9%)
Door Handle	0 (0%)	88 (19.5%)	0 (0%)	97 (21.5%)	0 (0%)	185 (20.5%)
Elevator Buttons	4 (10.3%)	46 (10.2%)	2 (9.1%)	57 (12.6%)	6 (9.8%)	103 (11.4%)
Exterior Bench	1 (2.6%)	9 (2.0%)	0 (0%)	0 (0%)	1 (1.6%)	9 (1.0%)
Floor	32 (82.1%)	59 (13.1%)	19 (86.4%)	78 (17.3%)	51 (83.6%)	137 (15.2%)
Hand Sanitizer	1 (2.6%)	49 (10.9%)	0 (0%)	59 (13.1%)	1 (1.6%)	108 (12.0%)
Other Item	1 (2.6%)	37 (8.2%)	0 (0%)	10 (2.2%)	1 (1.6%)	47 (5.2%)
Table	0 (0%)	3 (0.7%)	0 (0%)	2 (0.4%)	0 (0%)	5 (0.6%)
Telephone Receiver	0 (0%)	75 (16.6%)	0 (0%)	61 (13.5%)	0 (0%)	136 (15.1%)
Material						
Air (Control)	0 (0%)	10 (2.2%)	0 (0%)	9 (2.0%)	0 (0%)	19 (2.1%)
Metal	2 (5.1%)	120 (26.6%)	2 (9.1%)	154 (34.1%)	4 (6.6%)	274 (30.4%)
Plastic/Vinyl	37 (94.9%)	318 (70.5%)	20 (90.9%)	287 (63.6%)	57 (93.4%)	605 (67.1%)
Wood	0 (0%)	3 (0.7%)	0 (0%)	1 (0.2%)	0 (0%)	4 (0.4%)
Week						
1	7 (17.9%)	41 (9.1%)	4 (18.2%)	44 (9.8%)	11 (18.0%)	85 (9.4%)
2	6 (15.4%)	43 (9.5%)	2 (9.1%)	45 (10.0%)	8 (13.1%)	88 (9.8%)
3	6 (15.4%)	40 (8.9%)	3 (13.6%)	45 (10.0%)	9 (14.8%)	85 (9.4%)
4	8 (20.5%)	39 (8.6%)	2 (9.1%)	46 (10.2%)	10 (16.4%)	85 (9.4%)
5	3 (7.7%)	47 (10.4%)	3 (13.6%)	43 (9.5%)	6 (9.8%)	90 (10.0%)
6	3 (7.7%)	47 (10.4%)	2 (9.1%)	44 (9.8%)	5 (8.2%)	91 (10.1%)
7	3 (7.7%)	47 (10.4%)	3 (13.6%)	45 (10.0%)	6 (9.8%)	92 (10.2%)
8	2 (5.1%)	48 (10.6%)	1 (4.5%)	46 (10.2%)	3 (4.9%)	94 (10.4%)
9	1 (2.6%)	49 (10.9%)	0 (0%)	47 (10.4%)	1 (1.6%)	96 (10.6%)
10	0 (0%)	50 (11.1%)	2 (9.1%)	46 (10.2%)	2 (3.3%)	96 (10.6%)

485 **Table 3.** Swab characteristics and measure of association with SARS-CoV-2 detection.

Characteristic	Beta	95% CI[1]	p-value
Study Week	-0.01	-0.02, -0.01	<0.001
Material			
Metal	—	—	
Air (Control)	0.12	-0.32, 0.56	0.6
Plastic/Vinyl	0.07	0.00, 0.14	0.065
Wood	-0.02	-0.50, 0.46	>0.9
Specific Site			
Air (Control)	—	—	
Computer Keyboard	0.05	-0.38, 0.47	0.8
Door Handle	0.10	-0.33, 0.53	0.6
Elevator Buttons	0.18	-0.25, 0.61	0.4
Exterior Bench	0.20	-0.25, 0.66	0.4
Floor	0.31	-0.11, 0.74	0.15
Hand Sanitizer	0.05	-0.37, 0.48	0.8
Other Item	0.05	-0.38, 0.47	0.8
Table	0.05	-0.55, 0.65	0.9
Telephone Receiver	0.04	-0.39, 0.47	0.9
Unit COVID Risk Type			
Non-COVID Unit	—	—	
COVID Risk Unit	0.08	0.05, 0.11	<0.001
Public/HCW Area	0.02	-0.02, 0.06	0.3
Campus			
Campus B	—	—	
Campus A	0.03	0.00, 0.06	0.033

[1] CI = Confidence Interval

488 **Table 4.** Univariate coefficients with 95% confidence intervals (CI), p-values, and Akaike

489 information criterion (AIC) for prediction of hospital case burden from (i) proportion of built

490 environment swabs positive and (ii) proportion of floor swabs positive.

Variable	Coefficient	95% CI	P-value	AIC
All Built Environment Swabs	5.8	1-11	0.02	56.5
Floor Swabs	1.7	0.5-3.1	0.01	54.9

Figure 1. Limit of Detection analysis of SARS-CoV-2 RNA and virus standards. Serial dilutions

of (A) SARS-CoV-2 RNA standard and (B) nucleic acid extractions of heat-inactivated virus

were analyzed by qPCR targeting the N1 gene. Mean Cq values and standard deviations are

plotted for 14-20 replicates (RNA standard) or 7 replicates (viral extractions). Grey circles

indicate replicates that were not detected (ND). The limit of detection (red line with shaded 95%

confidence interval) and limit of quantification (blue line) for the RNA standard were estimated

by a model-fitting approach. The LOD for the viral extractions was estimated by a discrete

threshold of >95% positivity. Equations, R^2 values, and amplification efficiency (E) are shown for

each standard curve.

Figure 2. Swabbing recovery of heat-inactivated SARS-CoV-2 virus from spiked surfaces. Test

surfaces were spotted with 10^5 copies of the virus standard, allowed to dry, and swabbed for 30

seconds. Swabs were stored in nucleic acid stabilization solution overnight, RNA was extracted,

and recovered genome copies were quantified by qPCR analysis. Virus recovery was

determined by dividing the copies estimated for each sample by the copies estimated for a "no

swabbing" control. The mean and standard error for three replicates is shown. Letters indicate

significant differences between groups, determined by one-way ANOVA followed by Tukey's

post hoc test.

Figure 3. Time series of built environment swabs positive for SARS-CoV-2 by COVID risk

location. *Blank (white) cells represent weeks where no swab was collected for a given location.*

(A)

(B)

Figure 4. Proportion of hospital built environment swabs positive for SARS-CoV-2 (bars) and [A]

mean hospital COVID-19 active cases (line) or [B] mean citywide COVID-19 active cases (line).

(A)

(B)

Figure 5. A. Proportion of hospital floor swabs positive for SARS-CoV-2 (bars) and [A] mean

hospital COVID-19 active cases (line) or [B] mean citywide COVID-19 active cases (line).

536   (A)

538   (B)

540   **Figure 6.** Campus A wastewater SAR-CoV-2 copies per ml (bars) and [A] mean citywide

541   COVID-19 active cases (line) or [B] mean hospital COVID-19 active cases (line). There was no

542   sample collection on weeks 7, 8 and 10.