IL-1α Mediates Tissue Specific Inflammation and Severe Respiratory Failure In

Covid-19: Clinical And Experimental Evidence

Running heading: IL-1 bioactivity in COVID-19

Georgios Renieris[1], Eleni Karakike[1], Zhongli Xu[2], Theologia Gkavogianni[1],

Dionysia- Eirini Droggiti[1], Katja Schubert[2], Dionysios Kafousopoulos[1],

Marlen Hermann[2], Mihai G. Netea[3,4], Jesper Eugen-Olsen[5], John Simard[6],

Evangelos J. Giamarellos-Bourboulis[1]

[1] 4th Department of Internal Medicine, National and Kapodistrian University of

Athens, Medical School, 124 62 Athens, Greece

[2]InflaRx GmbH, 07745 Jena, Germany

[3]Immunology and Metabolism, Life & Medical Sciences Institute, University of

Bonn, 53115 Bonn, Germany

[4]Department of Internal Medicine and Center for Infectious Diseases, Radboud

University, 6500 Nijmegen, the Netherlands

[5]Department of Clinical Research, Copenhagen University Hospital Hvidovre,

2650 Hvidovre, Denmark

[6]XBiotech, 78744 Austin, Texas, USA

Correspondence

E. J. Giamarellos-Bourboulis, MD, PhD

4th Department of Internal Medicine, ATTIKON University General Hospital

1 Rimini Street 124 62,

Athens, Greece

Tel: +30 210 58 31 994/ Fax: +30 210 53 26 446

E-mail: egiamarel@med.uoa.gr

ABSTRACT

Background: Acute respiratory distress syndrome (ARDS) in COVID-19 has been associated with dysregulated immune responses leading to catastrophic inflammation. The activation pathways remain to be fully elucidated. We investigated the ability of circulating to induce dysregulated immune responses.

Materials & Methods: Calprotectin and high mobility group box 1 (HMGB1) were associated with ARDS in 60 COVID-19 patients. In a second cohort of 40 COVID-19 patients calprotectin at hospital admission was associated with serum levels of soluble urokinase plasminogen activator receptor (suPAR). A COVID-19 animal model was developed by intravenous injection of plasma from healthy volunteers or patients with COVID-19 ARDS into C57/BL6 mice once daily for 3 consecutive days. In separate experiments, mice were treated with a) the IL-1 receptor antagonist Anakinra or vehicle and b) Flo1-2a anti-murine anti-IL-1α monoclonal antibody or the specific anti-human IL-1α antibody XB2001, or isotype controls. Mice were sacrificed on day 4. Cytokines and myeloperoxidase (MPO) in tissues were measured.

Results: Calprotectin, but not HMGB1, was elevated ARDS. Higher suPAR readouts indicated higher calprotectin levels. CHallenge of mice with COVID-19 plasma led to inflammatory reactions in murine lung and intestines as evidenced by increased levels of TNFα, IL-6, IFNγ and MPO. Anakinra treatment brought these levels down. Similar decrease was found in mice treated with Flo1-2a but not with XB2001.

Conclusion: Circulating alarmins, specifically calprotectin, of critically ill COVID-19 patients induces tissue-specific inflammatory responses through an IL-1α mediated mechanism. This could be attenuated through inhibition of IL-1 receptor or specific inhibition of IL-1α.

75   **Keywords**: calprotectin, tumor necrosis factor, interleukin-6, interleukin-1α, suPAR,

76   acute respiratory distress syndrome (ARDS), severe respiratory failure, COVID-19

INTRODUCTION

A little over one year since the onset of the COVID-19 pandemic, more than 2.3 million deaths have been attributed to the virus (https://www.who.int/emergencies/diseases/novel-coronavirus-2019). Despite all efforts to understand the disease, efforts are ongoing to elucidate what drives progression in COVID-19 cases from lower respiratory tract infections (LRTI) to acute respiratory distress syndrome (ARDS). We have suggested that the deterioration from pneumonia to ARDS is the result of a complex immune dysregulation, which in about 25% of patients appears to be related to a macrophage activation like syndrome; while in the remaining 75% of patients the mechanism seems to involve an interleukin (IL)-6 receptor dependent dysregulation, with an associated lymphopenia and a subset of cytokine over-producing monocytes that show decreased human leukocyte antigen (HLA)-DR expression (1).

Recent data suggests that SARS-CoV-2 stimulates vicious pro-inflammatory responses inducing NLRP3 inflammasome activation in monocytes and macrophages (2). In the open-label phase II study SAVE (3), patients with LRTI caused by SARS-CoV-2 and showing plasma concentrations of the biomarker suPAR (soluble urokinase plasminogen activator receptor) ≥ 6 ng/ml were treated with the recombinant human IL-1 receptor antagonist Anakinra. Anakinra is able to block both IL-1α and IL-1β bioactivity at the level of their common receptor (4) and prevent the development of ARDS.

The results of the SAVE trial led us to hypothesise that LRTI caused by SARS-CoV-2 can progress to ARDS. In this hypothesis, as a result of infection by SARS-CoV-2, patients over-produce IL-1α and other danger-associated molecular patterns (DAMPs) from their lung parenchyma. DAMPs trigger the release of IL-1β

and in combination with IL-1α evoke strong innate immune pro-inflammatory

responses which lead to ARDS (1). In order to test our hypothesis, we used a three-

step approach. In the first step, we measured and found higher concentrations of the

DAMP calprotectin in the plasma of patients with ARDS than patients without ARDS.

In the second step, we measured and found that DAMP concentrations in the plasma

of patients with suPAR more than 6 ng/ml were significantly higher than that in

patients with suPAR less than 6 ng/ml. In the third step, we studied a COVID-19-like

animal model, where we prevented the development of strong pro-inflammatory

responses through inhibition of IL-1 signalling via Anakinra or anti-IL-1α antibodies.

PATIENTS & METHODS

Clinical study

Plasma samples from a cohort of 60 patients with and without ARDS that was

determined to be caused by SARS-CoV (approval 30/20 by the National Ethics

Committee of Greece; approval IS 021-20 by the National Organization for

Medicines of Greece) and from 40 participants in the SAVE trial (suPAR-guided

Anakinra treatment for Validation of the risk and Early management of severe

respiratory failure by COVID-19; EudraCT number 2020-001466-11; National Ethics

Committee approval 38/20; National Organization for Medicines approval ISO 28/20;

ClinicalTrials.gov registration NCT04357366) were analyzed.

Patients were enrolled after written informed consent was provided by

themselves or by first-degree relatives in case of patients unable to consent. All

patients were adults of either sex with positive molecular testing of SARS-CoV-2

from respiratory secretions and with LRTI defined by the presence of diffuse

infiltrates by chest X-ray or computed tomography of the chest. Exclusion criteria

were a) HIV-1 infection and b) neutropenia defined as less than 1,000

neutrophils/mm^3. Blood was sampled on the day of hospital admission in the

emergencies. For patients admitted with ARDS, blood was collected within 24 hours

from mechanical ventilation (MV). Control blood samples were collected from healthy

volunteers matched for age and gender.

The following clinical variables were recorded: i) demographics; ii) severity

scores, namely Acute Physiology and Chronic Health Evaluation (APACHE) II score,

Charlson's Comorbidity Index (CCI) and Sequential Organ Failure Assessment

(SOFA) score; (iii) absolute blood cell counts, biochemistry, and blood gases; (iv)

and progression into ARDS during the entire hospital stay. ARDS was defined as a

ratio of partial oxygen pressure to fraction of inspired oxygen (PaO_2 / FiO_2) below

150 mmHg necessitating mechanical ventilation.

Five millilitres of whole blood was collected into a tube containing

ethylenediaminetetraacetic acid (EDTA) and centrifuged. Calprotectin (S100A8/A9);

high-mobility group box 1 (HMGB1); suPAR and ferritin were measured by enzyme-

linked immunosorbent assays (ELISA) (InflaRx, Jena, Germany and Creative

Diagnostics, Shirley, NY, USA; suPARnostic, ViroGates, Lyngby, Denmark and

ORGENTEC Diagnostika GmbH, Mainz, Germany, respectively); and C-reactive

protein (CRP) was measured using a nephelometric assay (Behring, Berlin,

Germany). The lowest limits of detection were: 0.5 ng/ml for HMGB1; 1.1 ng/ml for

suPAR; 0.2 mg/l for CRP; and 75 ng/ml for ferritin. RNA isolation of SARS-CoV-

2 was performed with the QIAamp Viral RNA Mini Kit (Qiagen, Hilden, Germany).

Reverse transcription of the viral RNA into cDNA and enrichment of the sequences

of interest via Real-time Polymerase Chain Reaction (PCR) was performed in the

Rotor-Gene Q instrument (Qiagen) using the SARS-CoV-2 reagent Viasure (CerTest

Biotec SL, Zaragoza, Spain).

Animal model of COVID-19 Pathogenic Inflammation

Animal experiments were conducted in the Unit of Animals for Medical and

Scientific purposes of the University General Hospital "Attikon" (Athens, Greece). All

experiments were licensed from the Greek veterinary directorate under the protocol

number 471955/06-07-2020.

We studied 80 male and female C57Bl6 mice (7-8 weeks old). Mice were

allowed to acclimate for seven days before beginning of the experiments. They were

housed in typical mouse cages, up to 5 mice per cage on 12-h dark/light cycle and

allowed free access to standard dry rodent diet and water. Analgesia was achieved

with paracetamol suppositories to avoid interactions with the immune system.

In order to develop a model of COVID-19-like inflammatory illness, mice were

treated intravenously (iv) with 100 µl of plasma from patients with ARDS due to

SARS-CoV-2 infection for 3 consecutive days; control mice were treated with 100 µl

of plasma from healthy volunteers for 3 consecutive days. On the fourth day, mice

were sacrificed by subcutaneous (sc) injection of 300 mg/kg ketamine, followed by

cervical dislocation. Under sterile conditions a midline abdominal incision was

performed. Then, segments of liver, lower lobe of the right lung, right kidney, ileum

(ca. 1 cm distally to the pyloric sphincter) and colon (ca. 1 cm proximally of the anal

ring) were excised and collected into sterile tubes with 1 ml NaCl 0.9%. The

intestinal samples were thoroughly flushed with NaCl 0.9% for removal of feces

before storage. The samples were weighed and homogenized.

Segments of spleens were gently squeezed and passed through a sterile filter

(250 mm, 12-13 cm, AlterChem Co, Athens, Greece) for the collection of

splenocytes. Peritoneal cells were retrieved by peritoneal lavage with 10 ml of cold

PBS containing 2 mmol/l of EDTA. After three serial washings, spleencells and

peritoneal cells were counted on Neubauer plates with trypan blue for exclusion of

dead cells. A total of 5×10^6 cells/ml were incubated in sterile 24-well plates in

RPMI-1640, supplemented with 2 mM glutamine, 10% fetal bovine serum, 100 U/ml

of penicillin G and 0.1 mg/ml of streptomycin, in the absence or presence of 10 ng/ml

lipopolysaccharide (LPS) of *Escherichia coli* O55:B5 or 5×10^5 cfu/ml heat killed

*Candida αlbicans (*HKCA*)*. After 24 hours or 5 days of incubation at 37°C in 5% CO_2,

the plates were centrifuged, and the supernatants were collected. Concentrations of

tumor necrosis factor alpha (TNFα), interferon gamma (IFNγ) and IL-6, IL-17A and

IL-22 in supernatants from tissue samples and spleencells were measured in

duplicate by an enzyme immunosorbent assay (ThermoFisher Scientific,

Massachussets, USA) according to the manufacturer's instructions. The lowest

detections limits were as follows: for TNFα 19 pg/ml; for IFNγ 16 pg/ml; for IL-6 10

pg/ml; for IL-17A 8 pg/ml; and for IL-22 16 pg/ml.

Myeloperoxidase (MPO) activity in all collected tissues was determined.

Tissue segments were homogenized with T-PER® (ThermoFisher Scientific) and

centrifuged at 10,000 rpm at 4°C. Then the homogenates were incubated in wells of

a 96-well plate at 37°C with 4.2 mM tetramethylbenzidine (Serva, Heidelberg,

Germany), 2.5 mM citrate, 5 mM NaH_2PO_4 and 1.18 mM H_2O_2 pH 5.0 at a final

volume of 150 μl. After 5 minutes the reaction was terminated by adding 50 ml 0.18

M H_2SO_4. Absorbance was read at 450 nm against blank wells. Results were

adjusted for tissue sample protein content on Bradford assay (Sigma-Aldrich) and

they were expressed as MPO units/mg protein/g.

Experiments were repeated in mice receiving in addition to i.v. plasma, a) 100

µl sc of 10 mg/kg Anakinra (Swedish Orphan Biovitrum, Stockholm, Sweden) or 100

µl of 0.9% NaCl once daily for 3 days; and b) 100 µl intraperitoneally (ip) of 1000

µg/ml (12) anti-murine IL-1α antibody Flo1-2a or human IL- 1α antibody XB2001, or

murine isotype control (XBiotech, TX, USA).

In a separate experiment, mice treated with COVID-19 plasma for three days

were challenged on the fourth day ip with 1×10^7 cfu/ml *E. coli* or *Acinetobacter*

baumannii. Survival was recorded every 12 hours for 7 days.

Statistics

Categorical data were presented as frequencies and quantitative variables as

mean ± SE. Comparisons between groups were done using the Fisher exact test for

categorical data. Comparison for quantitative data was performed using the Mann-

Whitney U test for two group comparisons and the one-way ANOVA with the

Bonferroni correction for multiple group comparison. Correlations were performed

using the Spearman's rank of order. Any p value below 0.05 was considered

statistically significant.

RESULTS

Forty consecutive patients with COVID-19 who did not develop ARDS during

their hospital stay and 20 consecutive patients with ARDS were analyzed. Their

demographics are shown in Table 1. Samples of 10 healthy volunteers were

analyzed as healthy controls. Serum calprotectin levels were significantly higher

among patients with COVID-19 and ARDS (Figure 1A). HMGB1 concentrations did

not differ between patients with COVID-19 with and without ARDS (Figure 1B).

Elevated calprotectin levels COVID-19-related ARDS raised the question whether

calprotectin is released early among patients and may serve as a marker for patients

who will progress to ARDS.

suPAR was previously identified as an early predictor of ARDS progression in

patients with at least 6 ng/ml serum levels in the SAVE trial (3). To study of the early

increase of suPAR denotes increases of calcprotectin, calprotectin was measured in

40 patients screened for inclusion in SAVE trial. 20 patients were with suPAR

concentrations less than 6 ng/ml and 20 patients showed suPAR concentrations ≥ 6

ng/ml (Table 2). None of the patients with suPAR less than 6 ng/ml developed ARDS;

30% of those with suPAR ≥ 6 ng/ml progressed into ARDS. Remarkably, patients with

suPAR concentrations ≥ 6 ng/ml had significantly higher concentrations of

calprotectin (Figure 1C). A significant positive correlation between calprotectin and

suPAR was thus demonstrated (Figure 1D). These findings demonstrated that both

suPAR and calprotectin are reliable prognostic biomarkers in predicting the

development of severe respiratory failure in COVID-19. These findings also

suggested that DAMPs like calprotectin, released at early stages of clinical infection

are involved in mediating the progression of pneumonia into ARDS.

In order to investigate the mechanism by which calprotectin induces the

hyperinflammation related to COVID-19 associated respiratory failure, we sought to

design a COVID-19 murine model. Mice were challenged with plasma either from

healthy volunteers (HV) or from patients with ARDS of COVID-19. Plasma of COVID-

19 patients led to organ-specific increase of cytokine concentrations. More precisely,

TNFα and MPO activity were specifically increased in the lung, ileum and colon of

mice challenged with COVID-19 plasma; IL-6 was upregulated in the lung and colon;

and IFNγ was upregulated in the lung (Figure 2A to D). These findings demonstrated

a compartmentalized inflammatory response in mice after challenging with COVID-

19 patient plasma.

DAMPs, like calprotectin, act on monocytes and macrophages and tend to

stimulate intense IL-1 responses. To provide evidence on the contribution of IL-1

stimulation, mice challenged with plasma from HV or COVID-19 patients were

treated with N/S or with Anakinra; the latter blocks both human and murine IL-1α and

IL-1β. Treatment with Anakinra significantly reduced tissue concentrations of TNFα,

IL-6, IFN-γ and MPO activity (Figure 3A to 3I). However, the observed inflammation

may have been mediated through human IL-1α present in the plasma of COVID-19

patients. To further clarify this, mice challenged with COVID-19 plasma were treated

with the anti-murine specific anti-IL-1α antibody Flo1-2a and with the human specific

anti-human IL-1α antibody XB2001, respectively. Treatment with Flo1-2a

significantly reduced tissue concentrations of TNFα, IL- 6, IFNγ and MPO activity in

contrast to treatment with XB2001 (Figure 4A to 4I). This suggests that the host-

derived mouse IL-1α and not any injected human IL-1α was necessary to induce the

compartmentalized inflammation.

Treatment with Anakinra led to an increase in ex vivo production capacity of

IFNγ and TNFα, but not of IL-6, IL-17A or IL-22 from splenocytes following

stimulation with HKCA (Figure 5A to 5F and Figure S1A to S1D). HKCA targets

lymphocyte responses from splenocytes and the cytokine responses are compatible

with increase of Th1 responses following treatment with Anakinra.

The number of transcripts of core SARS-CoV-2 genes *ORF1ab* and *N* were

indirectly provided by the cycles (Ct) of PCR positivity in plasma samples that were

used for the induction of the COVID-19-like hyperinflammation in the murine model.

No correlation was found between Cts and the levels of TNFα, IL-6, IFNγ and MPO in

the lung of COVID-19 patients with ARDS due to COVID-19 (Figure S2). This suggests

that the hyperinflammation is independent from the viral load. In order to mimic hospital

acquired secondary bacterial infections among COVID-19 patients in murine, mice

treated for three consecutive days with COVID-19 plasma were further challenged

with *A. baumannii or E. coli.* Survival was significantly lower after challenging with *A.*

baumannii (Figure 3E)*.* Observations coming from our prospective cohort of 134

ARDS patients revealed higher incidence of secondary infections by A. *baumannii*

when compared to other Gram-negative bacteria (Table S1, Figure S3). This further

supports the immunological similarities of the developed animal model to the human

infection.

DISCUSSION

This study presented evidence that severe respiratory failure in COVID-19

patients is possibly triggered by DAMPs, most likely calprotectin, which are released

early after the interaction of SARS-CoV-2 with host cells. It may well be the case that

calprotectin is this specific DAMP since circulating levels of HMGB1, another DAMP,

do not follow those of calprotectin. DAMPs lead to a compartmentalized inflammation

through enhancement of the production of pro-inflammatory cytokines IL-1α and IL-

1β.

It has been clearly demonstrated that an impaired immune response is a core

characteristic of patients who develop ARDS due to SARS-CoV-2 (5). The

associated great increase in the mortality of these patients has turned the focus on

potential therapeutic targets to disrupt the immune dysregulation. This necessitates a

better understanding of the pathogenesis of COVID-19.

Calprotectin, a heterodimer of S100A8 and S100A9, is a protein secreted by

leukocytes. The upregulation of calprotectin expression has been associated with

various inflammatory processes such as inflammatory bowel disease, rheumatoid

arthritis, spondyloarthritis and cardiovascular disease in type 2 diabetes mellitus (6-

9). Additionally, calprotectin is crucial for neutrophil accumulation and inflammation

in the lung parenchyma during bacterial community acquired pneumonia (10).

Recently, high levels of calprotectin in samples of COVID-19 patients were

correlated with disease severity and were associated with increased levels of

inflammatory cytokines and chemokines and myelopoiesis of abnormal myeloid cells

(11). In this study, we suggest calprotectin as a potential first element in the pathway

leading to ARDS in COVID-19 patients. Indeed, we found greater levels in patients

with ARDS compared to patients without ARDS. On the contrary, HMGB1, an

additional alarmin, which has been associated with sepsis pathogenesis and

outcome (12), did not significantly differ among patients with and without ARDS due

to COVID-19. Moreover, suPAR which is an early predictor of respiratory

deterioration in COVID-19 (3, 13) was indicative of patients with higher calprotectin

levels, suggesting that increased suPAR can be the biomarker of DAMP-induced

pathways in COVID-19.

SARS-CoV-2 uses the cellular surface protein angiotensin-converting enzyme

2 (ACE2) to bind and enter cells. However, the mouse ACE2 does not effectively

bind the viral spike protein (14). Therefore wild-type mice cannot be directly used as

a model to gain insight in COVID-19 pathophysiology. Mice over-expressing ACE2

have been used as a strategy to tackle this problem (15). While it is accepted that

the interaction of the spike protein of SARS-CoV-2 with human angiotensin

converting enzyme 2 (ACE2) is the first step for viral invasion in host cells of the

upper and lower respiratory tract (16), the ability of the virus to directly infect other

human tissues, where ACE2 is also expressed, such as the small intestine, kidneys,

and the heart, is a matter of dispute (17). In addition, we also need to understand the

pure immunological mechanisms of hyperinflammation in COVID-19. This is

introducing the need of COVID-19 models like the one developed after animal

challenge with human plasma. The model simulates the inflammatory reaction in

human SARS-CoV-2 infection since it provides compartmentalized

hyperinflammation of the lung.

Calprotectin acts through the Toll like receptor 4 (TLR4) and the receptor for

advanced glycation end-products (RAGE) and may prime for intracellular pro-IL-1β

(18) which is subsequently cleaved by SARS-CoV-2 activated NLRP3 inflammasome

(2) into IL-1β. Anakinra prevents the inflammatory action of both IL-1α and IL-1β by

blocking their cellular receptor. IL-1α released during cell death is characterized by

an autocrine function enabling an amplifying loop of inflammation (19). Moreover, we

have shown that Anakinra enhances the Th1 response in mice and probably

improves viral clearance.

Our findings provide evidence that progression from LRTI by SARS-CoV-2 to

ARDS is a process which is mediated by DAMPs, like IL-1α and calprotectin,

released following cell destruction. The early detection of the presence of circulating

DAMPs and blockade through IL-1a inhibition may represent a viable therapeutic

strategy. This is the hallmark of the efficacy of Anakinra - guided by the biomarker

suPAR - which led to a significant decrease in the incidence of ARDS in the SAVE

trial (3).

Competing interests

J. Eugen-Olsen is a cofounder, shareholder and CSO of ViroGates A7S, Denmark

and is named inventor on patents on suPAR owned by Copenhagen University

Hospital Hvidovre, Denmark. He is granted by the Horizon 2020 European Grant

RISKinCOVID.

J. Simard is the CEO and founder of XBiotech.

M.G. Netea is a scientific founder of TTxD, and received research grants from GSK

and ViiV Healthcare.

E.J. Giamarellos-Bourboulis has received honoraria from AbbVie USA, Abbott CH,

InflaRx GmbH, MSD Greece, XBiotech Inc. and Angelini Italy; independent

educational grants from AbbVie, Abbott, Astellas Pharma Europe, AxisShield,

bioMérieux Inc, InflaRx GmbH, and XBiotech Inc.; and funding from the FrameWork

7 program HemoSpec (granted to the National and Kapodistrian University of

Athens), the Horizon2020 Marie-Curie Project European Sepsis Academy (granted

to the National and Kapodistrian University of Athens), and the Horizon 2020

European Grant ImmunoSep (granted to the Hellenic Institute for the Study of

Sepsis).

Funding

371 The study was funded in part by the Hellenic Institute for the Study of Sepsis and in

372 part by the Horizon 2020 grant RISKinCOVID. The funders had no role in the design,

373 collection, analysis and interpretation of data, writing, and decision to publish.

REFERENCES

1. Giamarellos-Bourboulis EJ, Netea MG, Rovina N, Akinosoglou K, Antoniadou A, Antonakos N, Damoraki G, Gkavogianni T, Adami ME, Katsaounou P, Ntaganou M, Kyriakopoulou M, Dimopoulos G, Koutsodimitropoulos I, Velissaris D, Koufargyris P, Karageorgos A, Katrini K, Lekakis V, Lupse M, Kotsaki A, Renieris G, Theodoulou D, Panou V, Koukaki E, Koulouris N, Gogos C, Koutsoukou A. Complex immune dysregulation in COVID-19 patients with severe respiratory failure. *Cell Host Microbe.* 2020;27:992-1000.e3.

2. Rodrigues TS, de Sá KSG, Ishimoto AY, Becerra A, Oliveira S, Almeida L, Gonçalves AV, Perucello DB, Andrade WA, Castro R, Veras FP, Toller-Kawahisa JE, Nascimento DC, de Lima MHF, Silva CMS, Caetite DB, Martins RB, Castro IA, Pontelli MC, de Barros FC, do Amaral NB, Giannini MC, Bonjorno LP, Lopes MIF, Santana RC, Vilar FC, Auxiliadora-Martins M, Luppino-Assad R, de Almeida SCL, de Oliveira FR, Batah SS, Siyuan L, Benatti MN, Cunha TM, Alves-Filho JC, Cunha FQ, Cunha LD, Frantz FG, Kohlsdorf T, Fabro AT, Arruda E, de Oliveira RDR, Louzada-Junior P, Zamboni DS. Inflammasomes are activated in response to SARS-CoV-2 infection and are associated with COVID-19 severity in patients. J Exp Med. 2021;218:e20201707.

3. Kyriazopoulou E, Panagopoulos P, Metallidis S, Dalekos GN, Poulakou G, Gatselis N, Karakike E, Saridaki M, Loli G, Stefos A, Prasianaki D, Georgiadou S, Tsachouridou O, Petrakis V, Tsiakos K, Kosmidou M, Lygoura V, Dareioti M, Milionis C, Papanikolaou IC, Akinosoglou K, Myrodia DM, Gravvani A, Stamou A, Gkavogianni T, Katrini K, Marantos T, Trontzas IP, Syrigos K, Chatzis L, Chatzis S, Vechlidis N, Avgoustou C, Chalvatzis S, Kyprianou M, van der Meer JWM,

Eugen-Olsen J, Netea MG, Giamarellos-Bourboulis EJ. An open label trial of Anakinra to prevent respiratory failure in COVID-19. *eLife*. 2021;10:e66125.

4. de Luca A, Smeekens SP, Casagrande A, Iannitti R, Conway KL, Gresnigt MS, Begun J, Plantinga TS, Joosten LA, van der Meer JW, Chamilos G, Netea MG, Xavier RJ, Dinarello CA, Romani L, van de Veerdonk FL. IL-1 receptor blockade restores autophagy and reduces inflammation in chronic granulomatous disease in mice and in humans. *Proc Natl Acad Sci*. 2014;111:3526–31.

5. Bohn MK, Hall A, Sepiashvili L, Jung B, Steele S, Adeli K. Pathophysiology of COVID-19: Mechanisms underlying disease severity and progression. *Physiology*. 2020;35:288-301.

6. Fukunaga S, Kuwaki K, Mitsuyama K, Takedatsu H, Yoshioka S, Yamasaki H, Yamauchi R, Mori A, Kakuma T, Tsuruta O, Torimura T. Detection of calprotectin in inflammatory bowel disease: Fecal and serum levels and immunohistochemical localization. *Int J Mol Med*. 2018;41:107-118.

7. Azramezani Kopi T, Shahrokh S, Mirzaei S, Asadzadeh Aghdaei H, Amini Kadijani A. The role of serum calprotectin as a novel biomarker in inflammatory bowel diseases: a review study. *Gastroenterol Hepatol Bed Bench*. 2019;12:183-189.

8. Jarlborg M, Courvoisier DS, Lamacchia C, Martinez Prat L, Mahler M, Bentow C, Finckh A, Gabay C, Nissen MJ; physicians of the Swiss Clinical Quality Management (SCQM) registry. Serum calprotectin: a promising biomarker in rheumatoid arthritis and axial spondyloarthritis. *Arthritis Res Ther*. 2020;22:105.

9. Oosterwijk MM, Bakker SJL, Nilsen T, Navis G, Laverman GD. Determinants of increased serum calprotectin in patients with type 2 diabetes mellitus. *Int J Mol Sci*. 2020;21:8075.

10. Kotsiou OS, Papagiannis D, Papadopoulou R, Gourgoulianis KI. Calprotectin in

lung diseases. *Int J Mol Sci*. 2021;22:1706.

11. Silvin A, Chapuis N, Dunsmore G, Goubet AG, Dubuisson A, Derosa L, Almire C,

Hénon C, Kosmider O, Droin N, Rameau P, Catelain C, Alfaro A, Dussiau C,

Friedrich C, Sourdeau E, Marin N, Szwebel TA, Cantin D, Mouthon L, Borderie D,

Deloger M, Bredel D, Mouraud S, Drubay D, Andrieu M, Lhonneur AS, Saada V,

Stoclin A, Willekens C, Pommeret F, Griscelli F, Ng LG, Zhang Z, Bost P, Amit I,

Barlesi F, Marabelle A, Pène F, Gachot B, André F, Zitvogel L, Ginhoux F,

Fontenay M, Solary E. Elevated calprotectin and abnormal myeloid cell subsets

discriminate severe from mild COVID-19. *Cell*. 2020;182:1401-1418.e18.

12. Karakike E, Adami ME, Lada M, Gkavogianni T, Koutelidakis IM, Bauer M,

Giamarellos-Bourboulis EJ, Tsangaris I. Late peaks of HMGB1 and sepsis

outcome: evidence for synergy with chronic inflammatory disorders. *Shock*.

2019;52:334-339.

13. Rovina N, Akinosoglou K, Eugen-Olsen J, Hayek S, Reiser J, Giamarellos-

Bourboulis EJ. Soluble urokinase plasminogen activator receptor (suPAR) as an

early predictor of severe respiratory failure in patients with COVID-19 pneumonia.

Crit Care. 2020;24:187.

14. Wan, Y., Shang, J., Graham, R., Baric, R. S. & Li, F. Receptor recognition by

novel coronavirus from Wuhan: an analysis based on decade-long structural

studies of SARS coronaviru**s.** *J Virol*. 2020;94:e00127-20.

15. Muñoz-Fontela C, Dowling WE, Funnell SGP, Gsell PS, Riveros-Balta AX,

Albrecht RA, Andersen H, Baric RS, Carroll MW, Cavaleri M, Qin C, Crozier I,

Dallmeier K, de Waal L, de Wit E, Delang L, Dohm E, Duprex WP, Falzarano D,

Finch CL, Frieman MB, Graham BS, Gralinski LE, Guilfoyle K, Haagmans BL,

Hamilton GA, Hartman AL, Herfst S, Kaptein SJF, Klimstra WB, Knezevic I,

Krause PR, Kuhn JH, Le Grand R, Lewis MG, Liu WC, Maisonnasse P, McElroy

AK, Munster V, Oreshkova N, Rasmussen AL, Rocha-Pereira J, Rockx B,

Rodríguez E, Rogers TF, Salguero FJ, Schotsaert M, Stittelaar KJ, Thibaut HJ,

Tseng CT, Vergara-Alert J, Beer M, Brasel T, Chan JFW, García-Sastre A, Neyts

J, Perlman S, Reed DS, Richt JA, Roy CJ, Segalés J, Vasan SS, Henao-

Restrepo AM, Barouch DH. Animal models for COVID-19. *Nature.* 2020;586:509-

515.

16. Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor

usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol.*

2020;5:562-569.

17. Bost P, Giladi A, Liu Y, Bendjelal Y, Xu G, David E, Blecher-Gonen R, Cohen M,

Medaglia C, Li H, Deczkowska A, Zhang S, Schwikowski B, Zhang Z, Amit I.

Host-viral infection maps reveal signatures of severe COVID-19 patients. *Cell.*

2020;181:1475-1488.e12.

18. Vogl T, Stratis A, Wixler V, Völler T, Thurainayagam S, Jorch SK, Zenker S,

Dreiling A, Chakraborty D, Fröhling M, Paruzel P, Wehmeyer C, Hermann S,

Papantonopoulou O, Geyer C, Loser K, Schäfers M, Ludwig S, Stoll M,

Leanderson T, Schultze JL, König S, Pap T, Roth J. Autoinhibitory regulation of

S100A8/S100A9 alarmin activity locally restricts sterile inflammation. *J Clin*

Invest. 2018;128:1852-66.

19. Mantovani A, Dinarello CA, Molgora M, Garlanda C. Interleukin-1 and related

cytokines in the regulation of inflammation and immunity. *Immunity.* 2019;50:778-

795.

475
476

Table 1. Baseline clinical and laboratory characteristics of patients according to the development of ARDS due to pneumonia by the SARS-CoV-2 coronavirus

	No ARDS (n=40)	ARDS (n=20)	p-value
Age (years, mean ± SD) [#]	57.3 ±14.85	66.60 ± 11.45	0.020[#]
Male gender (n, %)	33 (82.5)	17 (85.0)	0.559[▯]
CCI (mean ± SD) [#]	1.85 ± 1.66	3.55 ± 2.56	0.012[#]
APACHE II score (mean ± SD) [#]	5.50 ± 3.11	10.05 ± 5.07	0.001[#]
SOFA (mean ± SD) [#]	1.87 ± 1.66	3.35 ± 1.06	0.001[#]
Main comorbidities (n, %)			
Type 2 diabetes mellitus	6 (15.0)	6 (30.0)	0.189[▯]
Congestive heart failure	1 (2.5)	2 (10.0)	0.255[▯]
Coronary heart disease	1 (2.5)	4 (20.0)	0.038[▯]
Chronic renal disease	1 (2.5)	0 (0)	0.667[▯]
COPD	4 (10.0)	3 (15.0)	0.676[▯]
Laboratory values (mean ± SD)			
White blood cells (mm^3)	6098.46 ± 2102.44	8758.00 ± 3992.08	0.003[#]
Absolute neutrophil counts (mm^3)	4535.52 ± 2110.35	7437.75 ± 3762.13	0.001[#]
Absolute lymphocyte counts (mm^3)	1065.98 ± 419.86	810.04 ± 426.46	0.025[#]
Platelets (x 10^3, mm^3)	191.2 ± 85.6	178.3 ± 54.9	0.389[#]
C-reactive protein (mg/l)	124.55 ± 100.73	206.38 ± 85.96	0.014[#]
Fibrinogen (mg/dl)	510.59 ± 172.91	611.20 ± 199.78	0.262[#]
Ferritin (ng/ml)	1136.05 ± 1022.05	1288.83 ± 573.13	0.381[#]
Procalcitonin (ng/ml)	0.25 ± 0.39	0.38 ± 0.46	0.280[#]
D-dimers (ng/ml)	599.76 ± 364.72	763.00 ± 702.42	0.464[#]
AST (U/l)	41.32 ± 26.62	64.80 ± 108.08	0.502[#]
ALT (U/l)	39.33 ± 34.24	52.80 ± 74.30	0.822[#]
Creatinine (mg/dl)	0.89 ± 0.18	1.29 ± 1.16	0.006[#]
Total bilirubin (mg/dl)	0.59 ± 0.54	5.25 ± 18.34	0.211[#]

Abbreviations: SD: standard deviation; APACHE: acute physiology and chronic health evaluation; CCI: Charlson's comorbidity index; SOFA: sequential organ failure; COPD: chronic obstructive pulmonary disease; AST: aspartate aminotransferase; ALT: alanine aminotransferase; ARDS: acute respiratory distress syndrome
[▯]Comparison by Fischer exact test; [#]Comparison by the Mann- Whitney U test.

Table 2. Baseline clinical and laboratory characteristics of patients with pneumonia by SARS-CoV-2 coronavirus according to the measured levels of circulating soluble urokinase plasminogen activator receptor (suPAR)

	suPAR < 6 ng/ml (n=20)	suPAR ≥ 6 ng/ml (n=20)	p-value
Age (years, mean ± SD) [#]	46.25 ±17.63	68.30 ± 9.27	6×10^{-5}[#]
Male gender (n, %)	13 (65.0)	13 (65.0)	1.000[ꝉ]
CCI (mean ± SD) [#]	3.56 ± 1.62	5.13 ± 0.99	0.022[#]
APACHE II score (mean ± SD) [#]	7.28 ± 3.43	6.69 ± 5.25	0.953[#]
SOFA (mean ± SD) [#]	0.86 ± 1.07	2.61 ± 1.20	0.004[#]
ARDS (n, %)	0 (0.0)	6 (30)	3×10^{-6}[ꝉ]
Main comorbidities (n, %)			
Type 2 diabetes mellitus	1 (5.0)	4 (20.0)	0.342[ꝉ]
Congestive heart failure	0 (0)	1 (5.0)	0.500[ꝉ]
Coronary heart disease	1 (5.0)	3 (15.0)	0.658[ꝉ]
COPD	0 (0)	1 (5.0)	0.506[ꝉ]
Laboratory values (mean ± SD)			
White blood cells (mm^3)	6007.14 ± 1500.55	7578.88 ± 3201.64	0.220[#]
Absolute neutrophil counts (mm^3)	3573.30 ± 1723.14	5833.80 ± 2988.62	0.055[#]
Absolute lymphocyte counts (mm^3)	2233.86 ± 1872.39	1203.26 ± 614.70	0.198[#]
Platelets (x 10^3, mm^3)	261.8 ± 133.8	212.1 ± 68.2	0.574[#]
C-reactive protein (mg/l)	35.69 ± 53.43	89.19 ± 82.45	0.083[#]
Fibrinogen (mg/dl)	352.00 ± 59.24	612.22 ± 187.67	0.003[#]
Ferritin (ng/ml)	283.41 ± 244.73	882.90 ± 595.75	0.024[#]
Procalcitonin (ng/ml)	0.06 ± 0.05	0.36 ± 0.36	0.133[#]
D-dimers (ng/ml)	421.67 ± 324.10	1407.79 ± 3251.55	0.207[#]
AST (U/l)	22.57 ± 6.68	46.44 ± 29.49	0.041[#]
ALT (U/l)	27.14 ± 14.37	38.67 ± 27.78	0.458[#]
Creatinine (mg/dl)	0.92 ± 0.27	0.97 ± 0.34	0.883[#]
Total bilirubin (mg/dl)	0.259 ± 0.11	1.54 ± 2.06	0.001[#]

Abbreviations: SD: standard deviation; APACHE: acute physiology and chronic health evaluation; CCI: Charlson's comorbidity index; SOFA: sequential organ failure; ARDS: acute respiratory distress syndrome; COPD: chronic obstructive pulmonary disease; AST: aspartate aminotransferase; ALT: alanine aminotransferase.
[ꝉ]Comparison by Fischer exact test; [#]Comparison by the Mann- Whitney U test.

FIGURE LEGENDS

Figure 1 Calprotectin is increased in patients with ARDS due to COVID-19

A-B) Concentrations of calprotectin (S100A8/A9) and high mobility group box 1

(HMGB1) were measured in the plasma from healthy volunteers; from patients with

COVID- 19 who did not develop ARDS during follow-up on the day of hospital

admission; and from patients with ARDS by COVID- 19 on the first day of ARDS.

Comparison by the Mann Whitney U test; ns non- significant; ▯ $p < 0.05$; ▯▯ $p < 0.01$; ▯▯▯

$p < 0.001$; ▯▯▯▯; $p < 0.0001$.

Panels C and D) suPAR was measured in 40 patients screened for participation in the

SAVE trial (3). None of the patients with suPAR less than 6 ng/ml developed acute

respiratory distress syndrome (ARDS), whereas 6 patients (30%) with suPAR 6 ng/ml or

more developed ARDS.

C) Concentrations of calprotectin in 20 patients with suPAR less than 6 ng/ml and in 20

patients with suPAR 6 ng/ml or more. Comparison by the Mann Whitney U test; ▯ $p <$

0.05.

D) Correlation of levels of suPAR with plasma calprotectin on day of admission

Spearman rank correlation coefficients (r_s), interpolation lines and p- values are

provided.

Figure 2 A COVID- like animal model of compartmentalized hyperinflammation

C57Bl6 mice were injected intravenously (i.v.) with plasma of healthy volunteers or

patients with ARDS due to COVID-19 for three consecutive days. Mice were sacrificed

on day 4. Tissue concentrations of A) tumor necrosis factor alpha (TNFα); B) Interleukin

(IL)-6; C) Interferon gamma (IFNγ) and D) myeloperoxidase (MPO) activity were

determined. Comparison by the Mann Whitney U test; ns non-significant; ▢ p< 0.05; ▢▢

p< 0.01; ▢▢▢▢ p< 0.0001.

E) C57Bl6 mice were injected intravenously (i.v.) with plasma of healthy volunteers or

patients with ARDS due to COVID-19 for three consecutive days. On day 4 mice were

challenged intraperitoneally (i.p.) with *E. coli* or *A. baumannii*. Survival comparison by

the log-rank test and the respective p- value are provided.

Figure 3 IL-1 inhibition attenuates the compartmentalized hyperinflammation in a

COVID-like murine model

In a COVID- like infection model, C57Bl6 mice were challenged intravenously (i.v.) with

plasma of healthy volunteers or patients with ARDS due to COVID-19 for three

consecutive days. In separate experiments, on each day of plasma challenge, mice

were treated with Anakinra, which inhibits human and murine IL-1, or vehicle. Mice were

sacrificed on day 4. A-C) Tumor necrosis factor alpha (TNFα); D-E) Interleukin (IL) -6; F)

Interferon gamma (IFNγ) and G-I) myeloperoxidase (MPO) activity was determined in

tissues. Comparison by the Mann Whitney U test; ns non-significant; ▢ p< 0.05; ▢▢ p<

0.01.

Figure 4 Murine IL-1α drives the hyperinflammation caused by SARS-CoV-2

In a COVID- like infection model, C57Bl6 mice were challenged intravenously (i.v.) with

plasma of patients with ARDS due to COVID-19 for three consecutive days. In separate

experiments, on each day of plasma challenge, mice were treated with Flo1-2a anti-

murine IL-1α antibody or XB2001 human IL- 1α antibody or murine isotype control

for Flo1-2a. Mice were sacrificed on day 4. A-C) Tumor necrosis factor alpha (TNFα);

D-E) Interleukin (IL) -6; F) Interferon gamma (IFNγ) and G-I) myeloperoxidase (MPO)

activity was determined in tissues. Comparison by the Mann Whitney U test; ns non-

significant; ▯ $p < 0.05$; ▯▯ $p < 0.01$; ▯▯▯ $p < 0.001$.

Figure 5 IL-1 inhibition primes Th1 responses in COVID-19

A-C) In a COVID- like infection model, C57Bl6 mice were treated intravenously (i.v.) with

plasma of healthy volunteers or patients with ARDS due to COVID-19 for three

consecutive days. Mice were sacrificed on day 4. Splenocytes were isolated and

incubated with heat killed *C. albicans* for the production of interferon-gamma (IFNγ),

tumour necrosis factor-alpha (TNFα) and interleukin (IL)-6.

D-F) In a COVID- like infection model, C57Bl6 mice were treated intravenously (i.v.) with

plasma of patients with ARDS due to COVID-19 with or without treatment with the IL-1

receptor inhibitor Anakinra for three consecutive days.

Mice were sacrificed on day 4. Splenocytes were isolated and incubated with heat killed

C. albicans for the production of IFNγ, TNFα and IL-6. Comparison by the Mann

Whitney U test; ns non-significant; ▯ $p < 0.05$; ▯▯ $p < 0.01$.