Children develop strong and sustained cross-reactive immune responses against spike protein following SARS-CoV-2 infection

Alexander C. Dowell[1], Megan S. Butler[1†], Elizabeth Jinks[1†], Gokhan Tut[1†], Tara Lancaster[1], Panagiota Sylla[1], Jusnara Begum[1], Hayden Pearce[1], Graham S. Taylor[1], Eleni Syrimi[1], Eliska Spalkova[1], Sandra Davies[1], Frances Baawuah, MRCP,[2] Joanne Beckmann, FRCPCH(UK),[3] Ifeanyichukwu Okike, FRCPCH(UK),[2,4] Shazaad Ahmad, MRCPath,[5] Joanna Garstang, MBCHB,[6] Andrew J Brent, FRCP,[7,8] Bernadette Brent, MD,[7] Georgina Ireland, MPH,[2] Felicity Aiano, MSc,[2] Zahin Amin-Chowdhury, BSc,[2] Samuel Jones, PhD,[2] Ray Borrow, FRCPath,[9] Ezra Linley, PhD,[9] John Poh, PhD,[2] Vanessa Saliba, MD,[2] Gayatri Amirthalingam, MFP,[2] Kevin E Brown, FRCPath,[2] Mary E Ramsay, FFPH.[2] Jianmin Zuo[1], Paul Moss[1]*, Shamez Ladhani[2,10]*

†*These author contributed equally

1. Institute of Immunology & Immunotherapy, Collage of Medical and Dental Sciences, University of Birmingham, Birmingham, B15 2TT, UK
2. Public Health England, 61 Colindale Avenue, London NW9 5EQ, UK
3. East London NHS Foundation Trust, 9 Allie Street, London E1 8DE, UK
4. University Hospitals of Derby and Burton NHS Foundation Trust, Uttoxeter New Road, Derby DE22 3NE, UK
5. Manchester University NHS Foundation Trust, Oxford Road, Manchester M13 9WL, UK
6. Birmingham Community Healthcare NHS Trust, Holt Street, Aston B7 4BN, UK
7. Oxford University Hospitals NHS Foundation Trust, Old Road, Oxford OX3 7HE
8. University of Oxford, Wellington Square, Oxford OX1 2JD, UK
9. Public Health England, Manchester Royal Infirmary, Manchester, United Kingdom
10. Paediatric Infectious Diseases Research Group, St. George's University of London, London, UK

Abstract

SARS-CoV-2 infection is generally mild or asymptomatic in children but the biological basis for this is unclear. We studied the profile of antibody and cellular immunity in children aged 3-11 years in comparison with adults. Antibody profiles in children were strong with high titres against spike protein and receptor binding domain (RBD). SARS-CoV-2 seroconversion in children strongly boosted antibody responses against seasonal beta-coronaviruses, partly through cross-recognition of the S2 domain, indicating a broad humoral response that was not seen in adults. T cell responses against spike were also >2-fold higher in children compared to adults and displayed a strong Th1 cytokine profile. SARS-CoV-2 spike-reactive cellular responses were present in more than half the seronegative children, indicating pre-existing cross-reactive responses or sensitization against SARS-CoV-2. Importantly, all children retained high antibody titres and cellular responses for more than 6 months after infection whilst relative antibody waning was seen in adults. Children thus distinctly generate robust, cross-reactive and sustained immune responses after SARS-CoV-2 infection with focussed specificity against spike protein. These observations demonstrate several novel features of SARS-CoV-2-specific immune responses in children and may provide insights into relative clinical protection in this group. Such information on the profile of natural infection will help to guide the introduction of vaccination regimens into the paediatric population.

Introduction

The SARS-CoV-2 pandemic has resulted in over 2.8 million deaths to date and the most significant determinant of outcome is age at the time of primary infection (Booth, Reed et al. 2021). In children, SARS-CoV-2 infection is generally asymptomatic or mild and contrasts with high rates of hospitalisations and deaths among older adults (Viner, Mytton et al. 2021). This natural history is unusual for respiratory viruses such as influenza where children are the main drivers of infection and transmission. As such, there is interest in understanding the profile of the immune response to SARS-CoV-2 in children. Studies of SARS-CoV-2-specific immunity in children have been limited to date but initial findings have reported reduced magnitude of both antibody and cellular responses in comparison to adults, and an absence of nucleocapsid-specific antibody responses during or early post-infection (Pierce, Preston-Hurlburt et al. 2020, Tosif, Neeland et al. 2020, Weisberg, Connors et al. 2021). One unique feature of SARS-CoV-2 infection in children is the development of a rare complication, known as Pediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2 (PIMS-TS) also known as paediatric multisystem inflammatory syndrome (MIS-C), which shares features with Kawasaki disease, toxic shock syndrome or both (Consiglio, Cotugno et al. 2020, Flood, Shingleton et al. 2021). MIS-C develops approximately 2-6 weeks after infection with median age at 9 years of age (Hoang, Chorath et al. 2020). The immunological basis for this condition remains unclear but is characterised by diffuse endothelial involvement and broad autoantibody production (Gruber, Patel et al. 2020).

One potential determinant of differential immune responses to SARS-CoV-2 across the life course may be the timing of exposure to the four additional endemic human coronaviruses (HCoV). These comprise the beta-coronaviruses OC-43 and HKU-1, which have 38% and 35% amino acid homology with SARS-CoV-2, as well as the more distantly related alpha-coronaviruses NL-63 and 229E, each with around 31% homology ((NCBI)). All these coronaviruses cause frequent mild childhood infections and antibody seroconversion occurs typically before the age of 5 years. Infection with one of the Alpha or Beta viruses provides short term immunity against re-infection from coronaviruses and believed to represent transient cross-reactive immunity within the subtypes (Dijkman, Jebbink et al. 2008, Dijkman, Jebbink et al. 2012, Huang, Garcia-Carreras et al. 2020). As such, recent HCoV infection might pre-sensitize children against SARS-CoV-2 infection and may explain cross-reactive SARS-CoV-2-neutralising antibodies in some seronegative children (Ng, Faulkner et al. 2020). Immune responses against HCoV are retained throughout life but do not provide sterilising immunity (Huang, Garcia-Carreras et al. 2020). Consequently, recurrent infections are common, generating concern that a similar pattern will be observed for SARS-CoV-2 infection.

Covid-19 vaccines are now being implemented widely in adults and vaccine trials in children are currently underway, with the aim of increasing protection across the whole population. It is, therefore, imperative to understand the baseline profile of SARS-CoV-2 specific immune responses in children to inform vaccination strategy. Here we provide a comprehensive characterisation of the convalescent humoral and cellular immune response in a cohort of 92 primary school-aged children compared with 155 adults taking part in SARS-CoV-2 surveillance in the schools (sKIDs) study (Ladhani, Baawuah et al. 2021). We demonstrate a markedly different profile of immune response after SARS-CoV-2 infection in children compared to adults. These findings have potential implications for understanding immune responses to infection in children and may provide insights into PIMS-TS. Furthermore, these can also be used to guide and interpret the introduction of COVID-19 vaccination for children.

Methods

Sample collection

Public Health England (PHE) initiated prospective SARS-CoV-2 surveillance in primary schools across the UK after they reopened following the easing of national lockdown in June 2020. The protocol for the COVID-19 Surveillance in School KIDs (sKIDs) is available online (https://www.gov.uk/guidance/covid-19-paediatric-surveillance) (Ladhani, Baawuah et al. 2021). Surveillance comprised two arms, one involving weekly swabbing of primary school students and staff for SARS-CoV-2 infection (from June to mid-July 2020) and the other comprising swabbing and blood sampling taken in 3 rounds: beginning (1-19 June) and end (3-23 July) of the summer half-term when primary schools were partially re-opened, and after full reopening of all schools in September 2020, at the end of the autumn term (23 November-18 December). Samples for extended humoral and cellular analysis were taken in round 3.

For each known SARS-CoV-2 seropositive individual an age-matched (nearest age in years for students, nearest 10 years for teachers) and gender-matched participant also underwent blood sampling. In total 155 adults and 91 children had sufficient blood sample for serology and cellular responses (Table 1).

PBMC and Plasma Preparation

Blood tubes were spun at 300g for 10mins prior to removal of plasma which was then spun again at 800g 10mins and stored at -80°C. Remaining blood was diluted 1:1 with RPMI and PBMC isolated on a Sepmate (Stemcell) density centrifugation tube, washed with RPMI and rested in RPMI+10% FBS overnight at 37°C.

MSD Serology assay

Quantitative IgG antibody titres were measured against trimeric spike (S) protein, nucleocapsid (N) and other coronavirus using MSD V-PLEX COVID-19 Coronavirus Panel 2 (N05368-A1), responses to other respiratory viruses were measured using MSD V-PLEX COVID-19 Respiratory Panel 1 (N05358-A1). Multiplex MSD Assays were performed according to manufacturer instructions. Briefly 96-well plates were blocked. Following washing, samples diluted 1:5000 in diluent, as well as reference standard and internal controls, were added to the wells. After incubation plates were washed and detection antibody added. Plates were washed and were immediately read using a MESO TM QuickPlex SQ 120 system. Data were generated by Methodological Mind software and analysed with MSD Discovery Workbench (v4.0) software. Assay Cut-offs in respect of pre-pandemic plasma samples

from healthy donors are shown in (Extended data 1). Cut-offs used are Spike; 350 AU/ml, RBD and Nucleocapsid; 600 AU/ml, NTD; 15 AU/ml.

Total IgG/A/M anti Spike SARS-CoV-2 ELISA

A total GAM anti-SARS-CoV-2 Spike ELISA kit (Cook, Faustini et al. 2021) was purchased from Binding Site (Birmingham, UK). ELISA was performed following the manufacturer's instructions. Optical density (OD) was compared to a known calibrator and expressed as a ratio to the calibrator. Samples with a ratio above 1.0 were considered sero-positive.

Cross-reactive antibody blocking

Plasma samples were pre-diluted 1:10 with PBS then pre-absorbed by adding an equal volume of either recombinant Spike S1 domain (10569-CV-100) or Spike S2 (10594-CV-100) (R&D systems) at a concentration of 500ug/ml, in PBS, or PBS alone (mock). Samples were incubated at 37°C for 30min. Samples were then diluted to a final dilution of 1:5000 in MSD diluent and run on a MSD V-PLEX COVID-19 Respiratory Panel 2 plate in duplicate.

RBD-ACE-2 interaction neutralisation.

The concentration of antibodies which inhibited interaction between RBD and ACE-2 was determined using a SARS-CoV-2 neutralisation assay (Biolegend), following the manufacturer's instructions. Briefly, plasma or positive control antibody were pre-incubated with biotinylated-Fc-chimera-S1-RBD protein prior to addition of bead bound ACE-2. Binding of RBD to ACE2 was then measured by the addition of streptavidin-PE. Samples were run on a BD LSR-II flow cytometer and analysed using Legendplex v8.0 software (Biolegend). Results were related to a known RBD neutralising antibody standard and displayed as ng/ml.

IFNγ ELISpot

Pepmixes pool containing 15-mer peptides overlapping by 10aa from either SARS-CoV-2 Spike S1 or S2 domains and a combined pool of Nucleoprotien (N) and Membrane (M) and Envelope (E) were purchased from Alta Biosciences (University of Birmingham, UK). Overlapping pepmix from Influenza A Matrix Protein 1; California/08/2009(H1N1) ID:C3W5Z8 and Aichi/2/1968 H3N2 ID:Q67157, were purchased from JPT technologies, and combined as a relevant control.

T cell responses were determined using a IFNγ ELISpot Pro kit (Mabtech), as previously described (Zuo, Dowell et al. 2020). Briefly, standard fresh PBMC were rested over-night prior to assay, 0.25-0.3x10^6 PBMC were added in duplicate per well containing either pep-mix, anti-CD3 (positive) or DMSO (negative) control. Samples were incubated for 16-18hrs. Supernatant was harvested and stored at -

80°C. Plates were developed following the manufacturers instructions and read using an AID plate reader (AID). Cut off values were previously determined (Zuo, Dowell et al. 2020).

Cytokine measurement

Supernatants from donors with a detectable response in overnight Elispot cultures were assessed using a Legendplex Th profile 12-plex kit (BIolegend), following the manufactures instructions. Data was analysed using Legendplex v8.0 Software (Biolegend).

Intracellular Cytokine Staining

Cryopreserved PBMC were thawed and rested overnight. Cells were then stimulated with a combined Spike-S1, S2 peptide pool at a final concentration of 1µg/ml per peptide, or DMSO (Mock). After 1hr eBiosciecne Protein transport inhibitor cocktail (Thermofisher Scientific) was added and cells incubated for a further 5 hrs. eBioscience Cell stimulation cocktail (Thermofisher Scientific) was used as a positive control. Following stimulation cells were washed (PBS+0.1%BSA), and surface stained at 4°C, 30min. Cells were then washed and fixed in 2% paraformaldehyde. Following washing brilliant staining buffer was added (BD bioscience) and a final concentration of 0.4% Saponin added. Cells were stained intracellularly at room temperature for 30min. Cells were then washed and run on a BD Symphony A3 flow cytometer (BD Biosciences). Antibody details are provided in Extended method table 1.

Data visualisation and statistics

Statistical tests, including normality tests, were performed as indicated using Graphpad Prism v9 software. Only results found to be significant ($p < 0.05$) are displayed.

Results

Children develop strong and broad antibody responses after SARS-CoV-2 infection

Blood samples were obtained from 91 children and 154 adults, including 35 children and 81 adults known to be seropositive in previous rounds of testing. The median age of the children was 7 years (range, 3-11) whilst that of adults was 41 years (range, 20-71) in adults. SARS-CoV-2 antibody profile was assessed using the MSD V-plex serology platform to determine serological responses against Spike, Receptor Binding Domain (RBD), N-terminal domain (NTD) and Nucleocapsid (N). In total, 47% of children and 59% of adults were found to be seropositive (Extended Data Table 1).

Seropositive children and adults demonstrated broadly similar antibody responses against the viral proteins. Geometric mean antibody titres against all four regions, were, however, higher in children, most notably against the NTD and RBD domains, which showed 2.3-fold and 1.7-fold increases respectively although these did not reach statistical significance. (Figure 1A). In contrast to previous reports (Weisberg, Connors et al. 2021) we observed a coordinated response against nucleoprotein, with a 1.3-fold increased antibody titre compared to adults (Figure 1 A&B).

SARS-CoV-2 infection strongly boosts antibodies responses against HCoV in children

Pre-existing immune responses against seasonal coronaviruses might act to modulate clinical outcome following primary SARS-CoV-2 infection, and cross-reactive neutralising antibodies have been reported in SARS-CoV-2-seronegative children (Ng, Faulkner et al. 2020). Consequently, we compared antibody levels against the four HCoV in SARS-CoV-2 in seronegative and seropositive children and adults.

A 1.2 to 1.4-fold increase in the titre against HCoV was evident in SARS-CoV-2 seropositive adults compared to the seronegative group. In contrast, antibody levels against all four viruses were markedly higher in SARS-CoV-2 seropositive children, with 2.3, 1.9, 1.5, and 2.1 fold higher antibody levels compared to the seronegative group. These were significant for OC-43 and HKU-1 (p=0.0071, p=0.0024 Brown-Forsythe and Welch ANOVA, with Dunnett's T3 multiple comparison test) (Figure 2A).

In order to assess if this effect was specific to HCoV or represented a more general effect of SARS-CoV-2 infection on antibody responses against heterologous infection, we also examined antibody titres against influenza subtypes and respiratory syncytial virus (RSV) in relation to SARS-CoV-2 serostatus. No significant change in antibody titre against these viruses was seen in either children or adults (Figure 2B). These data show that SARS-CoV-2 infection in children strongly and specifically boosts humoral responses against HCoV.

SARS-CoV-2-specific antibodies in children can cross-react with the S2 domain of betacoronaviruses

Given the large increase in HCoV-specific antibody titre following SARS-CoV-2 infection in children, we were interested to assess to what extent this was cross-reactive against SARS-CoV-2 or could represent a HCoV-specific response. As such, we used recombinant S1 or S2 domains from SARS-CoV-2 to pre-absorb plasma samples prior to assessment of antibody levels to both SARS-CoV-2 and the four HCoV subtypes.

As expected, both the S1 and S2 domains significantly reduced antibody titres against total spike ($p<0.0001$, p0.0031 respectively, Friedman test with Dunn's multiple comparisons test). The S1 domain but not the S2 domain, absorbed RBD and NTD-specific antibodies against SARS-CoV-2 whilst no influence was observed in relation to nucleocapsid-specific binding for either domain (Figure 3). Of note, the S1 domain did not significantly reduce antibody binding to any of the 4 HCoV subtypes indicating little evidence for cross-reactive antibodies against this domain. The S2 domain, however, selectively reduced antibody binding to the two HCoV betacoronaviruses, OC43 and HKU1 ($p <0.0001$ & $p=0.0031$, respectively by repeated measure one-way ANOVA with Holm-Sidak's multiple comparison test). No such effect was observed in relation to binding to the alphacoronaviruses NL-63 and 229e (Figure 3).

These data show that the enhanced antibody response to HCoV in children after SARS-CoV-2 infection is based on a combination of antibodies that are cross-reactive against SARS-CoV-2 and those that are specific for HCoV.

Children develop strong cellular immune responses against spike protein after SARS-CoV-2 infection

We next assessed the magnitude and profile of the cellular immune response against SARS-CoV-2 in children and adults. Elispot analysis against overlapping peptide pools from spike and a combination of nucleocapsid and membrane and envelope (N/M) was performed on samples from 57 children and 93 adults, of which 37 and 64 were seropositive, respectively.

As expected, Elispot responses were common in SARS-CoV-2 seropositive donors, with 86% (32/37) of seropositive children and 69% (45/64) of seropositive adults showing a positive Elispot response to Spike.

The magnitude of the cellular response against spike was 2.1-fold higher in children, with median values of 533 spots/million compared to 195 in adults ($p=0.0003$, , Brown-Forsythe and Welch Anova, with Dunnett's T 3 multiple comparisons test) (Figure 4 A). Cellular responses against the N/M pool were relatively lower in children compared to spike such that the S:N/M ratio was markedly elevated in children, at 4.7 vs 1.8 in adults ($p=0.0007$, , Brown-Forsythe and Welch Anova, with Dunnett's T 3 multiple comparisons test) (Figure 4 B). The proportion of responders against each pool was also

different, with 98% of children with a positive Elispot responding to spike but only 45% responding to the N/M pool. In adults, the proportions were 88% and 64%, respectively (Figure 4 C).

Supernatant samples from the Elispot were then analysed using a multi-analyte bead assay to compare the profile of cytokine production by SARS-CoV-2-specific T cells from children and adults. Samples from adult donors showed a strong IL-2 response with lower levels of IL-10 production. In contrast, IL-2 levels in samples from children were very low (Figure 4D) indicating differential functional response compared with adults. Indeed analysis of three children by flow cytometry indicated that CD8+ IL-2-TNF+IFNγ+ T cells constitute the bulk of the spike-specific T cell response in children (Extended data 2).

Of note, robust cellular responses were also observed in 60% (12/20) seronegative children, all of whom were seronegative by three different serology platforms. Cellular responses of variable but lower magnitude were also present in 34% (10/29) of sero-negative adults (Figure 4 A). Two children and one adult had anti-nucleocapsid antibodies only on the MSD platform (Extended data Table 2.), but this insufficient to provide definitive sero-status. These cellular responses in seronegative donors were markedly spike-specific, with elevated S:N/M ratios of 7.1 and 5 in children and adults (Figure 4 B), respectively, and may indicate pre-existing cross-reactive immunity. Indeed 7 of 12 seronegative children and 6/10 seronegative adults with a positive ELISpot also had increased antibody levels to one or more HCoV (Extended data Table 2) potentially indicating recent HCoV infection. Alternatively, these responses might represent cellular sensitisation to SARS-CoV-2 protein in the absence of sero-conversion.

Children and Adults maintain immune responses against SARS-CoV-2 for at least 6 months

We were next interested to assess the longevity of immune responses within a subgroup of 35 children and 81 adults who had seroconverted at least 6 months prior to analysis. Importantly, all children retained humoral immunity whilst 7% (6/81) of previously seropositive adults failed to show significant antibody responses. Children also maintained higher antibody titres against Spike and RBD which were 1.8-fold higher than adults (Figure 5A&B).

The RBD region of spike is the predominant target of neutralising antibodies and as children had strong RBD-specific antibody responses, we next examined relative inhibition of virus binding in an ACE2 binding assay (n=12 children; n=15 adults). A higher, but non-significant, increase in the mean level of neutralising antibody was seen between adults and children (Figure 5C). Indicating that both groups maintain antibodies capable of neutralisation of RBD binding to ACE2, although this may be higher in children, consistent with higher RBD antibody levels (Figure 5A).

Cellular immune responses were detectable in 84% of children and 79% of adults at least 6 months after infection. The magnitude of the spike-specific response remained higher in children than in adults (p=0.032) whereas responses to the N/M pool were seen in only 31% of children compared to 68% of adults (Figure 5 D&E). As such, the ratio of the S:N/M response in children over time diverged further from adults at this later time point (p<0.0001) (Figure 5 F).

These data show children generate durable cellular and humoral immune responses following natural SARS-Cov-2 infection within an increasing focus against the spike protein.

Discussion

Age is the primary determinant of the clinical severity of SARS-CoV-2 infection and a life course assessment of virus-specific immunity is essential to understand pathogenesis and design vaccine strategies in children. Our detailed analysis of adaptive immune memory identifies a number of novel features in young children. A key finding was that the magnitude of the adaptive immune response to SARS-CoV-2 is higher in children compared to adults. This is somewhat different to previous reports which have reported lower antibody titres in children. It has also been reported that children do not mount effective antibody responses against nucleocapsid in the early post-infection period. We did observe antibody responses against both spike and nucleocapsid in children but it was also clear that immune responses are much more strongly focussed against spike. This is likely because the spike protein - and RBD in particular - is a major virulence factor for SARS-CoV-2 and antibodies against RBD correlate with neutralising activity *in vitro*.(Ju, Zhang et al. 2020) At the same time, nucleocapsid is a much more abundant protein within the virion and it is possible that the magnitude of the N-specific response is a reflection of peak or aggregate viral load. Levels of virus within the upper airways at the time of primary infection are equivalent in children and adults (Madera, Crawford et al. 2021), but relative changes over the course of infection are not known. Enhanced innate immune responses in children may play an important role in limiting systemic replication, which may explain the higher rates of asymptomatic and transient, mild illness in children exposed to the virus compared to adults (Neeland, Bannister et al. 2021). One striking feature was that SARS-CoV-2 infection in children doubled the antibody titres against all four alpha and beta HCoV subtypes. This pattern was not seen in adults where increased humoral responses were modest, even to the most closely related beta-coronavirus (Anderson, Goodwin et al. 2020, Westerhuis, Aguilar-Bretones et al. 2020). Of note, increased antibody titres against HoCV have also been observed after SARS-CoV-1 infection (Chan, Cheng et al. 2005). Using protein domain pre-absorption, we identified that a proportion of this response represented a SARS-CoV-2-specific humoral response that was cross-reactive with the S2 domain of the two more closely-related betacoronaviruses. The S2 domain is more highly conserved than S1 between CoV and this pattern is compatible with preferential targeting of structurally-conserved epitopes by HCoV-specific antibodies in children (Khan, Rahman et al. 2021). Some studies have shown these to have neutralising activity against SARS-CoV-2 (Shrock et al 2020, Ng et al 2020). Paediatric SARS-CoV-2 infection, however, also boosts HCoV-specific antibody responses that are not directly cross-reactive, as demonstrated by increased titres against alphacoronaviruses that could not be pre-absorbed. This may potentially reflect weakly cross-reactive B cell clones, potentially activated through T cell cross recognition, and is reminiscent of antibody boosting against H3 haemagglutinin after H1N1 infection in children who have had previous H3N2 infection (Meade, Kuan et al. 2020). As

such, there may be a lower affinity threshold for boosting of related but heterologous immune responses in children which might potentially represent an evolutionary adaptation to expand the memory pool early in life. It is interesting to speculate whether the observation that SARS-CoV-2 infection triggers a broadly reactive humoral response in children might provide insight into the pathophysiology of PIMS-TS, where intense B cell activation and auto-antibody production leads to an inflammatory syndrome. Of note, lack of pre-existing immunity to HCoV is a risk factor for development of PIMS-TS and may potentially indicate a protective influence from a stronger prior cross-reactive memory B cell pool (Consiglio, Cotugno et al. 2020).

There is currently very little information regarding the cellular immune response to SARS-CoV-2 in children. Interestingly, we find that the virus-specific T cell response is stronger in children than in adults. Furthermore, the specificity mirrors that of the humoral response in that responses against the spike protein are markedly increased compared to nucleocapsid and envelope proteins.

Virus-specific T cell responses in children also showed a differential cytokine response with markedly reduced production of IL-2 and evidence of increased IL-10 production. This may suggest a more highly differentiated profile in comparison to adults, in line with the enhanced magnitude of the response. Indeed, IL-2-IFNg+TNF+ CD8 T cell responses represented the majority of the Spike-specific T cell pool observed in three child at 6 months after primary infection. As such further characterisation of the T cell response in children is required.

A further noteworthy feature was that virus-specific T cell responses were detectable in more than half the seronegative children. It is possible that these represent HCoV-specific T cell responses that cross react against SARS-CoV-2 peptides (Mateus, Grifoni et al. 2020, Nelde, Bilich et al. 2021), as previously observed for antibody responses that are also focussed against spike protein (Ng et al 2020). HCoV infections are more common in young children and might potentially lead to generation of effector cells that were detected in the SARS-CoV-2 ELISpot. As such, this profile of cross-reactive antibody and cellular responses in children may help to explain the excellent clinical outcomes in this group. Alternatively, it is possible that these responses represent genuine SARS-CoV-2-specific T cells that have been generated following virus exposure in the absence of antibody sero-conversion. This pattern has been reported in health care workers with high levels of viral exposure (Reynolds, Swadling et al. 2020) and it is possible that such conditions are also seen in primary schools where enforcement of social distancing is challenging.

Many participants were seropositive 6 months prior to analysis and our data extend previous findings in adults (Zuo, Dowell et al. 2020, Dan, Mateus et al. 2021) to show sustained immunity over this time period. Moreover, we also found that children maintained stronger antibody and cellular immune responses at this timepoint compared to adults with no loss of humoral response compared to 7% in

adults. These data provide encouragement that immunity generated in childhood may provide longer term protection in the light of increasing concerns that SARS-CoV-2 will become an endemic infection (Shaman and Galanti 2020). In addition they may indicate a need for less frequent booster vaccination in children.

In conclusion we show that children display a novel and characteristic profile of adaptive immune response against SARS-CoV-2. This is likely to contribute to the strong clinical protection in this age group but may also provide insight into the characteristic immuno-pathology that may develop. Furthermore they will help to guide the introduction and interpretation of vaccine deployment in the paediatric population.

Acknowledgement

We would like to express our gratitude to the staff, parents and especially children, for their participation in this study, without whom this work would not have been possible.

		Number	Age (years) Median & Range	Infection >6mth[+]
Adults	**All**	154	40 (20-71)	
	Seropositive	91	39 (20-71)	81
	Seronegative	63	43 (26-65)	
Children	**All**	91	7 (3-11)	
	Seropositive	43	8 (5-11)	35
	Seronegative	48	7 (3-11)	

[+] Seropositive at first assessment (June 2020)

Table 1. Demographics and serostatus of study participants.

Figure 1

Children (red) and adults (blue) induce strong coordinated antibody responses to SARS-CoV-2

A) SARS-CoV-2 antibody levels determined by MSD assay, serostatus was determined based upon Spike serology and used to divide the cohorts into seropositive (red/blue) and seronegative (light red/blue). Dotted lines represent cut off values for serostatus. Fold change indicates the difference between the geometric mean titres in seropositive children and adults. Bars indicate geometric mean, with 95% CI.

B) The level of anti-spike and anti-nucleocapsid were compared within the whole cohorts showing coordinated responses.

Figure 2.
Antibody responses against the seasonal beta-coronaviruses are back-boosted in children following SARS-CoV-2 infection.

Antibody titres to the seasonal coronavirus (upper)and other respiratory viruses (lower) by MSD in children (red) and adults (blue) based on serostatus. Fold change indicates the difference between the geometric mean titres in seropositive children and adults. Bars indicate geometric mean, with 95% CI. Only significant differences are shown. Brown-Forsythe and Welch ANOVA, with Dunnett's T3 multiple comparison test.

Figure 3   **Cross reactive antibodies against Spike 2 domain contribute to increased HCoV-specific antibody responses in SARS-CoV-2 seropositive children**

Plasma from SARS-CoV-2 seropositive children was assessed for binding to the spike protein of the four HCoV or the Spike, RBD, NTD or nucleocapsid regions of SARS-CoV-2. Plasma was either applied neat (control) or following pre-absorption with either recombinant S1 (Spike 1 Block) or S2 (Spike 2 Block) domain protein. S1 pre-absorption strongly reduced binding to SARS-CoV-2 spike whilst S2 pre-absorption reduced binding to OC43 and HKU1. Repeated measure one-way ANOVA with Holm-Sidak's multiple comparison test, or Friedman test with Dunn's multiple comparisons test as appropriate

Figure 4.

Children induce strong spike T cell responses, and show evidence of pre-existing responses.

A) SARS-CoV-2 T cell response in children (red) and adults (blue) was determined by IFNγ ELISpot using pepmixes containing overlapping peptides to Spike, Nucleoprotein and viral Membrane (N/M), or Influenza, and shown in respect of spike serostatus.
B) The level of spike response was compared to N/M response and displayed as a ratio in seropositive and seronegative adults and children, as indicated. Bars indicate mean, Brown-Forsythe and Welch Anova, with Dunnett's T 3 multiple comparisons test.
C) The proportions of each cohort scored as responding to SARS-CoV-2 is indicated
D) Supernatant from overnight ELISpot cultures showing responses were analyses for additional cytokines as shown (children red, adults blue). Bars indicate mean±SD.

Figure 5.

Strong Spike responses are maintained in children at least six months after infection. Antibody and T cell responses in children (red) and adults (blue) who were seropositive at first testing indicating infection greater than six months prior.

A) Antibody titres as determined by MSD assay, bars indicate geometric mean± 95%CI. Dotted lines indicate seropositive cut-offs. Fold change indicates change in geometric mean titre. Black dots indicate individuals who were seronegative for spike but retained a nucleocapsid antibody response.

B) Proportions of individuals retaining antibody responses to antigens as indicated.

C) The level of antibodies which neutralise interaction of RBD with the ACE2 protein were determined in children (red) and adults (blue), ten pre-pandemic adult samples were included as a control (black). Dotted line indicates results below limit of detection. Bars indicate geometric mean ±95% CI.

D) IFNγ ELISPOT results in children (red) and adults (blue). Black dots indicate individuals who lacked a spike response but retained a nucleocapsid response. , Brown-Forsythe and Welch Anova, with Dunnett's T 3 multiple comparisons test.

E) The proportions of each cohort scored as responding to SARS-CoV-2 is indicated

F) The level of spike response was compared to N/M response and displayed as a ratio in seropositive and seronegative adults and children, as indicated. Bars indicate mean. Un-paired t-test.

Extended Data 1. MSD assay validation

Validation of cut-offs was tested using plasma samples from nine children (red) and 50 adults (blue) taken prior to CoVID-19. Dotted lines indicate cut-off used.

Extended Data 2. IL-2-TNF+IFNγ+ CD8 T cells dominate the Spike-specific T cell response in children

PBMC were stimulated for 6hrs in the presence of total-spike peptide pool and then analysed by flow cytometry to assess intracellular cytokine production. A) Gating strategy for analysis. B) Representative example of ICS staining from one child at six months post SARS-CoV-2 infection showing strong TNF+IFNγ+ CD8+ T cell response representing 0.45% of the global CD8+ T cell repertoire.

		Children		Adult		Pre-CoVID	
		Seropositive*	Seronegative*	Seropositive*	Seronegative*	Children	Adult
MSD serology – Geometric mean ±SD (AU/ml)[1]	SARS-CoV-2 Spike	11589±4.097 (43/43)	57.43±2.178 (0/48)	8593±4.173 (91/91)	61.14 ±2.047 (0/63)	44.34±3.719 (0/9)	13.22±4.893 (0/50)
	SARS-CoV-2 Nucleoprotein	5914±3.735 (40/43, 93%)	160.4±3.625 (7/48, 15%)	5321±5.116 (85/90, 94%)	216.6±5.038 (17/64, 27%)	68.71±2.764 (0/9)	76.238±6.238 (6/50, 12%)
	SARS-CoV-2 RBD	4527±3.673 (40/43, 93%)	70.80±2.435 (1/48, 2%)	2845±4.755 (79/91, 87%)	106.0±2.604 (2/64, 5%)	49./68±3.496 (0/9)	6.085±4.865 (0/50)
	SARS-CoV-2 NTD	234.9±3.436 (41/43, 95%)	2.119±2.630 (1/47, 2%)	106.9±6.290 (79/91, 87%)	2.271±3.265 (2/60, 3%)	1.757±4.179 (1/9, 11%)	1.196±1.99 (0/11)
	OC-43	36938±2.780	19296±2.531	38644±2.151 (91)	30526±1.974 (64)		
	HKU-1	8014±2.346 (43)	3544±3.181 (48)	12810±2.402 (91)	9090±2.698 (64)		
	NL-63	2861±2.482 (43)	1899±2.599 (48)	3112±1.914 (91)	2643±2.304 (64)		
	229e	11327±5.498 (43)	4845±6.982 (48)	22822±2.273 (91)	17032±2.951 (64)		
ELIspot – mean±SD sfc/10[6] PBMC[2]	Spike (S1+S2)	532.9±591.0 (32/37, 86%)	212.7±242.6 (12/20, 60%)	195.0±280.7 (45/64, 69%)	111.7±244.1 (10/29, 34%)		
	N/M	33.81±43.27 (16/37, 44%)	4.172±8.574 (2/20, 10%)	75.91±147.9 (40/64, 63%)	19.59±83.46 (2/29, 7%)		
	Flu	110.0±160.9 (37)	159.2±243.0 (18)	147.6±369.9 (52)	164.4±275.1 (24)		
+6mth Infection[3]							
MSD serology	SARS-CoV-2 Spike	11450±4.428 (34/35, 97%)		6102±7.131 (73/81, 90%)			
	SARS-CoV-2 Nucleoprotein	6086±2.26 (35/35, 100%)		4611±5.45 (75/81, 93%)			
	SARS-CoV-2 RBD	4366±(3.624) (33/35, 94%)		2482±5.029 (69/81, 85%)			
ELIspot	Spike (S1+S2)	530.8±600.7 (23/27, 85%)		176.1±228.9 (37/52, 71%)			
	N/M	35.41±42.51 (12/27, 44%)		86.44±161.7 (35/52, 67%)			
	Neutralising antibody (ng/ml)	278.9±1.485 (12)		247.4±1.608 (15)			n.d. (10)

Extended Data Table 1. Immunological Characteristics of Seropositive and Seronegative Children and Adults

Adult/Child	Age	ELIspot (sfc/10^6 PBMC)			MSD Serology – AU/ml								sKIDS round 1	sKIDS round 2	IgG,A,M ELISA
		Spike	N/M	Flu	COV-2-Spike	COV-2-RBD	Cov-2 NTD	COV2-Nucleocapsid	OC43 Spike	HKU1 Spike	NL63 Spike	229e Spike			
Child	5-11	**110**	4	172	5	285	2	50	28273	*12195*	2925	*87935*	Negative	Negative	Negative
Child	5-11	**243**	0	0	12	10	1	546	20067	1650	2012	432	Negative	Negative	Negative
Child	5-11	**418**	0	766	39	14	0	29	*31263*	*16729*	3684	*18035*	Negative		Negative
Child	5-11	**70**	0	796	73	225	2	**603**	7360	5213	1263	*60299*	Negative	Negative	Negative
Child	5-11	**142**	0	230	40	25	0	59	23156	5741	4402	8572	Negative	Negative	Negative
Child	5-11	**274**	0	276	127	97	4	56	*53420*	3992	*7148*	*14452*			Negative
Child	5-11	**330**	6	54	134	171	3	241	12221	1481	4213	*37470*	Negative	Negative	Negative
Child	5-11	**168**	0	4	220	47	2	43	*29509*	5580	675	*12144*	Negative	Negative	Negative
Child	5-11	**742**	2	36	25	37	1	93	6215	2197	1425	2828	Negative	Negative	Negative
Child	5-11	**776**	**32**	16	42	40	2	51	2637	268	305	153	Negative	Negative	Negative
Child	5-11	**484**	8	136	20	58	1	42	2610	2239	1072	6684	Negative		Negative
Child	5-11	**387**	**24**	104	109	36	1	**1465**	9255	*8513*	4582	5311	Negative	Negative	Negative
Staff	40-49	**148**	0	8	56	323	5	51	43504	11602	2468	28990	Negative		Negative
Staff	30-39	**62**	0	430	69	114	2	46	*106800*	*26036*	*7039*	*37074*	Negative	Negative	Negative
Staff	40-49	**496**	2	1246	64	132	0	299	30002	9771	2877	19008	Negative	Negative	Negative
Staff		**37**	1	19	84	75	3	105	43170	3686	4180	12688	Negative	Negative	Negative
Staff	40-49	**160**	0	392	225	136	7	**804**	23610	5171	826	29936	Negative	Negative	Negative
Staff	30-39	**1124**	16	374									Negative	Negative	Negative
Staff	60-69	**524**	2	320	56	39	1	150	*62856*	*25802*	2758	*33779*	Negative	Negative	Negative
Staff	30-39	**84**	4		65	55	2	278	*63770*	9017	3795	24533	Negative		Negative
Staff	50-59	6	**32**	96	62	64	12	60	30596	6158	*5140*	2549	Negative	Negative	Negative
Staff	30-39	8	**452**	64	33	395	8	22	25250	1613	692	*46524*	Negative	Negative	Negative
Staff	50-59	**428**	2		46	151	1	49	29970	*18828*	4699	*35577*	Negative		Negative

Extended data table 2. Immune characteristics of Elispot Responding Seronegative donors. Bold values indicate values above cut-off. ***Bold-italicised*** indicates values +50% higher than expected.

Antigen	Clone	Fluorophore	Supplier	Code
eBioscience Fixable Viability Dye	-	eFluor 780	ThermoFisher Scientific	65-0865-14
CD14	HCD14	APC-Cy7	Biolegend	325620
CD19	HIB19	APC-Cy7	Biolegend	302218
CD3	UCHT1	Fitc	Biolegend	300406
CD4	RPAT4	Per-CP-Cy5.5	Biolegend	300529
CD8	SK1	BV510	Biolegend	344732
IL-2	MQ117H12	Pe-Cy7	Biolegend	500326
IL-4	MP4 25D2	APC	Biolegend	500813
IL-6	MQ2 13A5	PE-Dazzle 594	Biolegend	501122
IL-10	Jes3-9D7	PE	Biolegend	501404
IL-17A	BL168	BV421	Biolegend	512322
IFNγ	4S.B3	AF700	Biolegend	502520
TNF	Ab11	BV711	Biolegend	502940

Table 1. Flow cytometry antibodies

References

(NCBI), N. C. f. B. I. "Protein BLAST." Retrieved Febuary, 2021, from
https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastp&PAGE_TYPE=BlastSearch&LINK_LOC=blasthome.

Anderson, E. M., E. C. Goodwin, A. Verma, C. P. Arevalo, M. J. Bolton, M. E. Weirick, S. Gouma, C. M. McAllister, S. R. Christensen, J. Weaver, P. Hicks, T. B. Manzoni, O. Oniyide, H. Ramage, D. Mathew, A. E. Baxter, D. A. Oldridge, A. R. Greenplate, J. E. Wu, C. Alanio, K. D'Andrea, O. Kuthuru, J. Dougherty, A. Pattekar, J. Kim, N. Han, S. A. Apostolidis, A. C. Huang, L. A. Vella, U. P. C. P. Unit, E. J. Wherry, N. J. Meyer, S. Cherry, P. Bates, D. J. Rader and S. E. Hensley (2020). "Seasonal human coronavirus antibodies are boosted upon SARS-CoV-2 infection but not associated with protection." medRxiv : the preprint server for health sciences: 2020.2011.2006.20227215.

Booth, A., A. B. Reed, S. Ponzo, A. Yassaee, M. Aral, D. Plans, A. Labrique and D. Mohan (2021). "Population risk factors for severe disease and mortality in COVID-19: A global systematic review and meta-analysis." PLOS ONE **16**(3): e0247461.

Chan, K. H., V. C. C. Cheng, P. C. Y. Woo, S. K. P. Lau, L. L. M. Poon, Y. Guan, W. H. Seto, K. Y. Yuen and J. S. M. Peiris (2005). "Serological responses in patients with severe acute respiratory syndrome coronavirus infection and cross-reactivity with human coronaviruses 229E, OC43, and NL63." Clinical and diagnostic laboratory immunology **12**(11): 1317-1321.

Consiglio, C. R., N. Cotugno, F. Sardh, C. Pou, D. Amodio, L. Rodriguez, Z. Tan, S. Zicari, A. Ruggiero, G. R. Pascucci, V. Santilli, T. Campbell, Y. Bryceson, D. Eriksson, J. Wang, A. Marchesi, T. Lakshmikanth, A. Campana, A. Villani, P. Rossi, N. Landegren, P. Palma and P. Brodin (2020). "The Immunology of Multisystem Inflammatory Syndrome in Children with COVID-19." Cell **183**(4): 968-981.e967.

Cook, A. M., S. E. Faustini, L. J. Williams, A. F. Cunningham, M. T. Drayson, A. M. Shields, D. Kay, L. Taylor, T. Plant, A. Huissoon, G. Wallis, S. Beck, S. E. Jossi, M. Perez-Toledo, M. L. Newby, J. D. Allen, M. Crispin, S. Harding and A. G. Richter (2021). "Validation of a combined ELISA to detect IgG, IgA and IgM antibody responses to SARS-CoV-2 in mild or moderate non-hospitalised patients." J Immunol Methods **494**: 113046.

Dan, J. M., J. Mateus, Y. Kato, K. M. Hastie, E. D. Yu, C. E. Faliti, A. Grifoni, S. I. Ramirez, S. Haupt, A. Frazier, C. Nakao, V. Rayaprolu, S. A. Rawlings, B. Peters, F. Krammer, V. Simon, E. O. Saphire, D. M. Smith, D. Weiskopf, A. Sette and S. Crotty (2021). "Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection." **371**(6529): eabf4063.

Dijkman, R., M. F. Jebbink, N. B. El Idrissi, K. Pyrc, M. A. Müller, T. W. Kuijpers, H. L. Zaaijer and L. van der Hoek (2008). "Human coronavirus NL63 and 229E seroconversion in children." Journal of clinical microbiology **46**(7): 2368-2373.

Dijkman, R., M. F. Jebbink, E. Gaunt, J. W. A. Rossen, K. E. Templeton, T. W. Kuijpers and L. van der Hoek (2012). "The dominance of human coronavirus OC43 and NL63 infections in infants." Journal of clinical virology : the official publication of the Pan American Society for Clinical Virology **53**(2): 135-139.

Flood, J., J. Shingleton, E. Bennett, B. Walker, Z. Amin-Chowdhury, G. Oligbu, J. Avis, R. M. Lynn, P. Davis, T. Bharucha, C. E. Pain, D. Jyothish, E. Whittaker, B. Dwarakanathan, R. Wood, C. Williams, O. Swann, M. G. Semple, M. E. Ramsay, C. E. Jones, A. V. Ramanan, N. Gent and S. N. Ladhani (2021). "Paediatric multisystem inflammatory syndrome temporally associated with SARS-CoV-2 (PIMS-TS): Prospective, national surveillance, United Kingdom and Ireland, 2020." The Lancet Regional Health - Europe **3**: 100075.

Gruber, C. N., R. S. Patel, R. Trachtman, L. Lepow, F. Amanat, F. Krammer, K. M. Wilson, K. Onel, D. Geanon, K. Tuballes, M. Patel, K. Mouskas, T. O'Donnell, E. Merritt, N. W. Simons, V. Barcessat, D. M. Del Valle, S. Udondem, G. Kang, S. Gangadharan, G. Ofori-Amanfo, U. Laserson, A. Rahman, S. Kim-Schulze, A. W. Charney, S. Gnjatic, B. D. Gelb, M. Merad and D. Bogunovic (2020). "Mapping Systemic Inflammation and Antibody Responses in Multisystem Inflammatory Syndrome in Children (MIS-C)." Cell **183**(4): 982-995.e914.

Hoang, A., K. Chorath, A. Moreira, M. Evans, F. Burmeister-Morton, F. Burmeister, R. Naqvi, M. Petershack and A. Moreira (2020). "COVID-19 in 7780 pediatric patients: A systematic review." EClinicalMedicine **24**: 100433.

Huang, A. T., B. Garcia-Carreras, M. D. T. Hitchings, B. Yang, L. C. Katzelnick, S. M. Rattigan, B. A. Borgert, C. A. Moreno, B. D. Solomon, L. Trimmer-Smith, V. Etienne, I. Rodriguez-Barraquer, J. Lessler, H. Salje, D. S. Burke, A. Wesolowski and D. A. T. Cummings (2020). "A systematic review of antibody mediated immunity to coronaviruses: kinetics, correlates of protection, and association with severity." Nature Communications **11**(1): 4704.

Ju, B., Q. Zhang, J. Ge, R. Wang, J. Sun, X. Ge, J. Yu, S. Shan, B. Zhou, S. Song, X. Tang, J. Yu, J. Lan, J. Yuan, H. Wang, J. Zhao, S. Zhang, Y. Wang, X. Shi, L. Liu, J. Zhao, X. Wang, Z. Zhang and L. Zhang (2020). "Human neutralizing antibodies elicited by SARS-CoV-2 infection." Nature **584**(7819): 115-119.

Khan, T., M. Rahman, F. A. Ali, S. S. Y. Huang, M. Ata, Q. Zhang, P. Bastard, Z. Liu, E. Jouanguy, V. Béziat, A. Cobat, G. K. Nasrallah, H. M. Yassine, M. K. Smatti, A. Saeed, I. Vandernoot, J.-C. Goffard, G. Smits, I. Migeotte, F. Haerynck, I. Meyts, L. Abel, J.-L. Casanova, M. R. Hasan and N. Marr (2021). "Distinct antibody repertoires against endemic human coronaviruses in children and adults." JCI insight **6**(4): e144499.

Ladhani, S. N., F. Baawuah, J. Beckmann, I. O. Okike, S. Ahmad, J. Garstang, A. J. Brent, B. Brent, J. Walker, N. Andrews, G. Ireland, F. Aiano, Z. Amin-Chowdhury, L. Letley, J. Flood, S. E. I. Jones, R. Borrow, E. Linley, M. Zambon, J. Poh, V. Saliba, G. Amirthalingam, J. Lopez Bernal, K. E. Brown and M. E. Ramsay (2021). "SARS-CoV-2 infection and transmission in primary schools in England in June-December, 2020 (sKIDs): an active, prospective surveillance study." Lancet Child Adolesc Health.

Madera, S., E. Crawford, C. Langelier, N. K. Tran, E. Thornborrow, S. Miller and J. L. DeRisi (2021). "Nasopharyngeal SARS-CoV-2 viral loads in young children do not differ significantly from those in older children and adults." Scientific Reports **11**(1): 3044.

Mateus, J., A. Grifoni, A. Tarke, J. Sidney, S. I. Ramirez, J. M. Dan, Z. C. Burger, S. A. Rawlings, D. M. Smith, E. Phillips, S. Mallal, M. Lammers, P. Rubiro, L. Quiambao, A. Sutherland, E. D. Yu, R. da Silva Antunes, J. Greenbaum, A. Frazier, A. J. Markmann, L. Premkumar, A. de Silva, B. Peters, S. Crotty, A. Sette and D. Weiskopf (2020). "Selective and cross-reactive SARS-CoV-2 T cell epitopes in unexposed humans." Science **370**(6512): 89-94.

Meade, P., G. Kuan, S. Strohmeier, H. E. Maier, F. Amanat, A. Balmaseda, K. Ito, E. Kirkpatrick, A. Javier, L. Gresh, R. Nachbagauer, A. Gordon and F. Krammer (2020). "Influenza Virus Infection Induces a Narrow Antibody Response in Children but a Broad Recall Response in Adults." **11**(1): e03243-03219.

Neeland, M. R., S. Bannister, V. Clifford, K. Dohle, K. Mulholland, P. Sutton, N. Curtis, A. C. Steer, D. P. Burgner, N. W. Crawford, S. Tosif and R. Saffery (2021). "Innate cell profiles during the acute and convalescent phase of SARS-CoV-2 infection in children." Nature Communications **12**(1): 1084.

Nelde, A., T. Bilich, J. S. Heitmann, Y. Maringer, H. R. Salih, M. Roerden, M. Lübke, J. Bauer, J. Rieth, M. Wacker, A. Peter, S. Hörber, B. Traenkle, P. D. Kaiser, U. Rothbauer, M. Becker, D. Junker, G. Krause, M. Strengert, N. Schneiderhan-Marra, M. F. Templin, T. O. Joos, D. J. Kowalewski, V. Stos-Zweifel, M. Fehr, A. Rabsteyn, V. Mirakaj, J. Karbach, E. Jäger, M. Graf, L.-C. Gruber, D. Rachfalski, B. Preuß, I. Hagelstein, M. Märklin, T. Bakchoul, C. Gouttefangeas, O. Kohlbacher, R. Klein, S. Stevanović, H.-G. Rammensee and J. S. Walz (2021). "SARS-CoV-2-derived peptides define heterologous and COVID-19-induced T cell recognition." Nature Immunology **22**(1): 74-85.

Ng, K. W., N. Faulkner, G. H. Cornish, A. Rosa, R. Harvey, S. Hussain, R. Ulferts, C. Earl, A. G. Wrobel, D. J. Benton, C. Roustan, W. Bolland, R. Thompson, A. Agua-Doce, P. Hobson, J. Heaney, H. Rickman, S. Paraskevopoulou, C. F. Houlihan, K. Thomson, E. Sanchez, G. Y. Shin, M. J. Spyer, D. Joshi, N. O'Reilly, P. A. Walker, S. Kjaer, A. Riddell, C. Moore, B. R. Jebson, M. Wilkinson, L. R. Marshall, E. C. Rosser, A. Radziszewska, H. Peckham, C. Ciurtin, L. R. Wedderburn, R. Beale, C. Swanton, S. Gandhi, B. Stockinger, J. McCauley, S. J. Gamblin, L. E. McCoy, P. Cherepanov, E. Nastouli and G. Kassiotis

(2020). "Preexisting and de novo humoral immunity to SARS-CoV-2 in humans." **370**(6522): 1339-1343.

Pierce, C. A., P. Preston-Hurlburt, Y. Dai, C. B. Aschner, N. Cheshenko, B. Galen, S. J. Garforth, N. G. Herrera, R. K. Jangra, N. C. Morano, E. Orner, S. Sy, K. Chandran, J. Dziura, S. C. Almo, A. Ring, M. J. Keller, K. C. Herold and B. C. Herold (2020). "Immune responses to SARS-CoV-2 infection in hospitalized pediatric and adult patients." **12**(564): eabd5487.

Reynolds, C. J., L. Swadling, J. M. Gibbons, C. Pade, M. P. Jensen, M. O. Diniz, N. M. Schmidt, D. K. Butler, O. E. Amin, S. N. L. Bailey, S. M. Murray, F. P. Pieper, S. Taylor, J. Jones, M. Jones, W.-Y. J. Lee, J. Rosenheim, A. Chandran, G. Joy, C. Di Genova, N. Temperton, J. Lambourne, T. Cutino-Moguel, M. Andiapen, M. Fontana, A. Smit, A. Semper, B. O'Brien, B. Chain, T. Brooks, C. Manisty, T. Treibel, J. C. Moon, M. Noursadeghi, D. M. Altmann, M. K. Maini, Á. McKnight and R. J. Boyton (2020). "Discordant neutralizing antibody and T cell responses in asymptomatic and mild SARS-CoV-2 infection." **5**(54): eabf3698.

Shaman, J. and M. Galanti (2020). "Will SARS-CoV-2 become endemic?" **370**(6516): 527-529.

Tosif, S., M. R. Neeland, P. Sutton, P. V. Licciardi, S. Sarkar, K. J. Selva, L. A. H. Do, C. Donato, Z. Quan Toh, R. Higgins, C. Van de Sandt, M. M. Lemke, C. Y. Lee, S. K. Shoffner, K. L. Flanagan, K. B. Arnold, F. L. Mordant, K. Mulholland, J. Bines, K. Dohle, D. G. Pellicci, N. Curtis, S. McNab, A. Steer, R. Saffery, K. Subbarao, A. W. Chung, K. Kedzierska, D. P. Burgner and N. W. Crawford (2020). "Immune responses to SARS-CoV-2 in three children of parents with symptomatic COVID-19." Nature Communications **11**(1): 5703.

Viner, R. M., O. T. Mytton, C. Bonell, G. J. Melendez-Torres, J. Ward, L. Hudson, C. Waddington, J. Thomas, S. Russell, F. van der Klis, A. Koirala, S. Ladhani, J. Panovska-Griffiths, N. G. Davies, R. Booy and R. M. Eggo (2021). "Susceptibility to SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults: A Systematic Review and Meta-analysis." JAMA Pediatr **175**(2): 143-156.

Weisberg, S. P., T. J. Connors, Y. Zhu, M. R. Baldwin, W.-H. Lin, S. Wontakal, P. A. Szabo, S. B. Wells, P. Dogra, J. Gray, E. Idzikowski, D. Stelitano, F. T. Bovier, J. Davis-Porada, R. Matsumoto, M. M. L. Poon, M. Chait, C. Mathieu, B. Horvat, D. Decimo, K. E. Hudson, F. D. Zotti, Z. C. Bitan, F. La Carpia, S. A. Ferrara, E. Mace, J. Milner, A. Moscona, E. Hod, M. Porotto and D. L. Farber (2021). "Distinct antibody responses to SARS-CoV-2 in children and adults across the COVID-19 clinical spectrum." Nature Immunology **22**(1): 25-31.

Westerhuis, B. M., M. Aguilar-Bretones, M. P. Raadsen, E. de Bruin, N. M. A. Okba, B. L. Haagmans, T. Langerak, H. Endeman, J. P. C. van den Akker, D. A. M. P. J. Gommers, E. C. M. van Gorp, B. H. G. Rockx, M. P. G. Koopmans and G. P. van Nierop (2020). "Severe COVID-19 patients display a back boost of seasonal coronavirus-specific antibodies." 2020.2010.2010.20210070.

Zuo, J., A. Dowell, H. Pearce, K. Verma, H. Long, J. Begum, F. Aiano, Z. Amin-Chowdhury, B. Hallis, L. Stapley, R. Borrow, E. Linley, S. Ahmad, B. Parker, A. Horsley, G. Amirthalingam, K. Brown, M. Ramsay, S. Ladhani and P. Moss (2020). "Robust SARS-CoV-2-specific T-cell immunity is maintained at 6 months following primary infection." 2020.2011.2001.362319.