**A cross-sectional study of workplace measures against COVID-19 and psychological**

**distress among Japanese workers**

Yoshino Yasuda[1], Tomohiro Ishimaru[1], Seiichiro Tateishi[2], Hisashi Eguchi[3], Mayumi Tsuji[4],

Akira Ogami[5], Koji Mori[6], Shinya Matsuda[7], and Yoshihisa Fujino[1] for the CORoNaWork

project

**Author contributions:** Y.F. was the chairperson of the study group. Y.Y. and Y.F.

conceived the research questions. Y.F. conducted the statistical analysis. Y.Y., T.I. and Y.F.

drafted the manuscript. All the authors designed the research protocol and developed the

questionnaire. All the authors reviewed and approved the final manuscript.

**Authors' affiliations:**

[1]Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan, Kitakyushu, Japan

[2] Department of Occupational Medicine, School of Medicine, University of Occupational and

Environmental Health, Japan, Kitakyushu, Japan Department of Psychiatry, University of

Occupational and Environmental Health, Japan, Kitakyushu, Japan

[3] Department of Mental Health, Institute of Industrial Ecological Sciences, University of

Occupational and Environmental Health, Japan, Kitakyushu, Japan

[4] Department of Environmental Health, School of Medicine, University of Occupational and

Environmental Health, Japan, Kitakyushu, Japan

[5] Department of Work Systems and Health, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan, Kitakyushu, Japan

[6] Department of Occupational Health Practice and Management, Institute of Industrial

Ecological Sciences, University of Occupational and Environmental Health, Japan,

Kitakyushu, Japan

[7] Department of Preventive Medicine and Community Health, School of Medicine,

University of Occupational and Environmental Health, Japan, Kitakyushu, Japan

**Corresponding author:** Yoshihisa Fujino[1], M.D., M.P.H., Ph.D.

Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan

1-1, Iseigaoka, Yahatanishiku, Kitakyushu, 807-8555, Japan

Tel: +81-93-691-7401

Email: zenq@med.uoeh-u.ac.jp

**Abstract**

**Objectives:** This study examined the relationship between the status of infection control efforts against COVID-19 in the workplace and workers' mental health using a large-scale Internet study.

**Methods:** This cross-sectional study was based on an Internet monitoring survey conducted during the third wave of the COVID-19 epidemic in Japan. Of the 33,302 people who participated in the survey, 27,036 were included in the analyses. Participants answered whether or not each of 10 different infection control measures were in place at their workplace (e.g. wearing masks at all times during working hours). A Kessler 6 (K6) score of ≥5 was defined as mild psychological distress. The odds ratios (ORs) of psychological distress associated with infection control measures at the workplace were estimated using a multilevel logistic model nested in the prefectures of residence.

**Results:** The OR of subjects working at facilities with 4 or 5 infection control measures was 1.15 (95% confidence interval [CI]: 1.02-1.29, p=0.025), that in facilities with 2 or 3 infection control measures was 1.32 (95% CI: 1.15-1.50, p<0.001), and that in facilities with 1 or no infection control measures was 1.48 (95% CI: 1.30-1.68, p<0.001) compared to subjects whose workplaces had ≥6 infection control measures.

**Conclusion:** Our findings suggest that proactive COVID-19 infection control measures can influence the mental health of workers.

**Keywords:** COVID-19, Japan, Occupational Health, Psychological Distress, Prevention and Control

**Introduction**

A new type of coronavirus infection (COVID-19) was confirmed in China in December 2019 and spread rapidly around the world. On April 7, 2020, a state of emergency was declared in Tokyo and six other prefectures, followed by a nationwide declaration on April 16. Although the number of infected people temporarily decreased in Japan, a second wave arrived in August of the same year, and a third wave was noted in December to January 2021.

The spread of COVID-19 has transformed people's lifestyles. To stop the spread of the infection, events and gatherings, including the 2020 Tokyo Olympics, were postponed or cancelled, and restrictions on going out, as well as online schooling and telecommuting were encouraged. The government recommends that residents avoid the "three Cs" of "closed, crowded, and close" in their daily lives by maintaining as much distance between people as possible, going outdoors rather than staying indoors when interacting, and avoiding talking directly to one another whenever possible.[1,2]

In addition to the above measures, various measures are also being taken to prevent infection in the workplace, including maintaining physical distance, wearing masks, basic hygiene (e.g. hand washing), daily health checks, and telework. Further, infection control measures are being implemented not only in public facilities but also in the private sector. Measures to ensure social distance in the workplace, such as spacing out seats, are increasingly common. Even in the hospitality industry, where wearing a mask has historically been frowned upon, the donning of a face mask for health purposes is now routine. Many companies have also introduced programs to allow employees to telecommute, some on a permanent basis. According to a survey,[3] the number of companies in Tokyo practicing telework has increased markedly, from 24% in March 2020 to 57% in January 2021.

However, with the spread of COVID-19, mental health problems have become an emerging public health issue. Previous studies have reported increased anxiety and mental

burden among residents of areas that experienced lockdowns.[4] Increased rates of mental

health problems, such as depression, worsening of existing mental illnesses, suicide, and

alcohol dependence, have also been reported.[5–7] In addition, encouraging avoidance of the

three Cs is also suspected to have had a negative impact on the mental health of the public, as

such measures necessarily interfere with public interaction, communication, and socialization.

Further, anxiety is likely to be heightened by the spread of unconfirmed or even outright false

information and the continuous state of fear and worry.[8] However, while authorities have

recommended a variety of measures, the extent of infection prevention measures actually

implemented varies among companies. Depending on the industry, some measures may be

difficult to implement.

In general, anxiety about infectious diseases such as influenza, HIV, and Middle East

Respiratory Syndrome is known to affect mental health,[9–11] and the same holds true for

COVID-19.[12] A recent review reported that occupational factors and workplace environment

were associated with workers' mental health.[13] To our knowledge, however, no report has yet

examined how the status of infection control efforts against COVID-19 in the workplace

influences workers' mental health.

We hypothesized that the implementation of good infection control efforts in the

workplace would have a positive impact on workers' mental health. Here, we examined the

relationship between the status of infection control efforts against COVID-19 in the

workplace and workers' mental health using a large-scale Internet study.

**Material and Methods**

*Study design and subjects*

This cross-sectional study was based on an Internet monitoring survey conducted on

December 22nd to 26th, 2020, when the third wave of the COVID-19 epidemic began in

Japan. Details of the protocol have already been reported.[14] In brief, data were collected from

workers who had employment contracts at the time of the survey, allocated by prefecture,

occupation, and sex. Of the 33,302 people who participated in the survey, 27,036 were

included in the study, excluding fraudulent responses.

This study was approved by the ethics committee of the University of Occupational

and Environmental Health, Japan (reference No. R2-079). Informed consent was obtained via

the website.

*Evaluation of infection control measures in the workplace*

Participants were asked to answer yes or no concerning whether the following measures had

been implemented in their workplace: refraining from and restrictions on business trips;

refraining from and restrictions on visitors; refraining from or requesting a limit on the

number of people at social gatherings and dinners; refraining from or limiting face-to-face

internal meetings; wearing masks at all times during working hours; installing partitions and

revising the workplace layout; recommendation of performing daily temperature checks by

workers themselves at home; encouraging telecommuting; prohibiting workers from eating at

their own desk; and requesting employees not come to work when they were not feeling well.

*Assessment of psychological distress*

The Kessler 6 (K6) was used to assess psychological distress.[15] The validity of the Japanese

version of the K6 has been confirmed.[16,17] The K6 was developed to screen for mental

disorders, such as depression and anxiety, and is also used as an indicator of the degree of

such mental problems, including psychological stress. The K6 is a series of six questions,

with scores ranging from 0 (never) to 4 (always), depending on how often the topics of the

questions were experienced in the past 30 days. The higher the total score, the greater the

possibility of depression or anxiety disorder, with a score of ≥5 indicating the possibility of

some kind of depression or anxiety problem and a score of ≥13 indicating a suspected

depression or anxiety disorder. In the present study, a K6 score of ≥5 was used as the cut-off

value for mild psychological distress.

*Other covariates*

Participants noted the following about themselves in an online form: age, sex, prefecture of

residence, marital status (married, unmarried, bereaved/divorced), job type (mainly desk

work, mainly involving interpersonal communication, and mainly labor), number of

employees in the workplace, educational background, equivalent household income

(household income divided by the square root of the household size), smoking status, and

alcohol consumption (6–7 days a week, 4–5 days a week, 2–3 days a week, less than 1 day a

week, hardly ever).

*Statistical analyses*

The odds ratios (ORs) of psychological distress associated with infection control measures at

the workplace were estimated using a multilevel logistic model nested in the prefecture of

residence. The multivariate model was adjusted for sex, age, education, equivalent household

income, job type, number of employees at the workplace, smoking status, and alcohol

consumption. The incidence rate of COVID-19 by prefecture was also used as a prefecture-

level variable. We further estimated the multivariate ORs of psychological distress associated

with the number of infection control measures at the workplace.

A p-value of <0.05 was considered statistically significant. All analyses were

conducted using Stata (Stata Statistical Software, Release 16; StataCorp LLC, College

Station, TX, USA).

**Results**

Table 1 shows the basic characteristics according to the number of infection control measures implemented at the workplace. Of the 27036 participants, 14852 (55%) indicated that they had ≥6 infection control measures implemented in their workplace. In contrast, 4177 (15%) of the participants reported having ≤1 infection control measures implemented in their workplace. Subjects with desk jobs had more infection control measures implemented than those in other lines of work. Further, the greater the number of employees, the more infection control measures were implemented.

Table 2 shows the number of infection control efforts put in place and the details of each effort. The most commonly implemented infection control measures were "wearing masks at all times during working hours" (79%) and "refraining from or requesting a limit on the number of people at social gatherings and dinners" (71%). In contrast, relatively few companies had implemented "requesting employees not come to work when they are not feeling well" (9%) and "prohibiting workers from eating at their own desk" (17%).

Table 3 shows the association of the number of workplace measures against COVID-19 and severe psychological distress. As the number of measures increased, the OR for psychological distress decreased. Compared to subjects whose workplaces had implemented 6 or more infection control measures, the OR of subjects with 4 or 5 measures was 1.15 (95% CI: 1.02-1.29, p=0.025), 1.32 (95% CI: 1.15-1.50, p<0.001) for 2 or 3 measures, and 1.48 (95% CI: 1.30-1.68, p<0.001) for 1 or no measures.

Table 4 shows the association of each infection control measure and psychological distress. Most of the measures were significantly associated with a reduction in psychological distress, with ORs ranging from approximately 0.80 to 0.90, with statistical significance. In particular, "requesting employees not come to work when they are not feeling well" was associated with a decreased OR for psychological distress (OR=0.60, 95% CI: 0.54-0.66,

184    p<0.001), whereas "prohibiting workers from eating at their own desk" was associated with

185    increased psychological distress (OR=1.12, 95% CI:1.00-1.25, p=0.050).

**Discussion**

This study demonstrated an association between psychological distress and the implementation of 10 common COVID-19 infection prevention measures recommended in the workplace. In addition, our study revealed that the more infection control measures implemented in the workplace, the lower the workers' mental burden.

Each individual preventive measure was found to be associated with low psychological distress. Particularly, the policy of "requesting employees not come to work when they are not feeling well" decreased the psychological distress. This measure is thought to be directly effective for minimizing damage in the event of infection around workers and linked to the mental stability of workers because it reduces the risk of infection to others when they themselves are infected. Sickness presenteeism is reportedly influenced by organizational culture, employment instability, and socioeconomic status.[18–20] Such a clear statement of company policy would discourage workers with a poor socioeconomic status against concealing their fever and coming to work, thereby preventing infection.

In contrast, psychological distress was increased for the measure "prohibiting workers from eating at their own desk", possibly reflecting the psychological burden caused by reducing communication with peers during a time of refreshment. These results suggest that communication is becoming an issue during the COVID-19 pandemic and that efforts to consciously balance COVID-19 infection control and communication in the workplace are needed.[21]

The present study revealed that the COVID-19 infection control measures implemented by companies varied. These differences in among companies are attributed to various factors, including the difficulty of adopting some measures in certain industries, the influence of the sense of risk among companies and employers, and the costs associated with the implementation of measures. The feasibility of infection control measures also differs

depending on the work environment. For example, in workplaces that do not have break

rooms for taking meals, prohibiting workers from eating at their own desks may be difficult.

In workplaces where employees have fixed starting and working hours, crowding of changing

rooms at certain times may be unavoidable. The ability to implement such measures varies

depending on the budget available, the size of the company, and the presence of an individual

in charge of promoting safety and health.

Our study also found that the more infection control measures implemented in the

workplace, the less substantial workers' mental burden. We assumed that companies with

more infection control measures in place would have stronger company governance. The

results suggest that the implementation of proactive infection control helps reduce

psychological distress. Infection anxiety concerning COVID-19 reportedly affects mental

health,[5,7,8,12,22] which is consistent with findings for other infectious diseases, such as hepatitis

and HIV.[23,24] These results suggest that infection control measures alleviate workers' anxiety

about infection. Proactive infection prevention measures in the workplace are expected to

reduce workers' anxiety about infection and thereby have a positive impact on workers'

mental health.[25]

Several possible reasons may explain why active infection control in the workplace is

associated with psychological distress in workers. As mentioned above, infection prevention

measures have a direct effect on reducing infection anxiety. In addition, the implementation

of appropriate COVID-19 control measures based on correct information can itself serve as a

form of infection prevention education for workers.[26] When an organization faces a crisis,

whether or not the employer provides appropriate support affects the mental health of the

workers.[22] A clear COVID-19 infection prevention policy in the workplace can build a

relationship of trust between the employer and workers and consequently reduce workers'

psychological distress.[13]

Several limitations associated with the present study warrant mention. First, this study

was a cross-sectional study, so causality is unclear. More psychologically anxious people

may have been prone to underestimate the efficacy of the infection control measures in the

workplace. However, since this study inquired about the existence of visible and physical

measures, we believe that the likelihood of wrong answers was low. Second, we did not

confirm the time when the countermeasures were implemented. This study investigated the

situation as of December 2020, when the infection was at its peak in Japan. Therefore, the

degree of the subjects' anxiety and their reaction to their workplace's efforts may have

differed between the time when the infection rate was relatively low and the time of the

survey, when the rate was relatively high. Third, we did not ask about the subjects'

workplaces or industries in detail, but some of these factors likely affected the results, such as

the size of the workplace space (small office vs. a large space, such as a retail store) or how

often the subjects were in contact with unspecified numbers of people. Finally,

establishments that have implemented many measures are more likely to be companies that

are enthusiastic about mental health measures and consider their employees' well-being on a

regular basis, so the mental burden may have already been low at baseline; however, this

aspect was not evaluated.

This study suggests that proactive COVID-19 infection control measures can lead to

improved mental health care for workers. Proactive infection prevention measures in the

workplace are expected to reduce workers' anxiety about infection and have a positive impact

on workers' mental health. The implementation of COVID-19 infection control measures in

the workplace is recommended not only as a way to prevent infection but also as a new

mental health measure.

**Acknowledgements**

This study was supported and partly funded by the University of Occupational and

Environmental Health, Japan; General Incorporated Foundation (Anshin Zaidan); The

Development of Educational Materials on Mental Health Measures for Managers at Small-

sized Enterprises; Health, Labour and Welfare Sciences Research Grants; Comprehensive

Research for Women's Healthcare (H30-josei-ippan-002); Research for the Establishment of

an Occupational Health System in Times of Disaster (H30-roudou-ippan-007), scholarship

donations from Chugai Pharmaceutical Co., Ltd., the Collabo-Health Study Group, and

Hitachi Systems, Ltd.

The current members of the CORoNaWork Project, in alphabetical order, are as

follows: Dr. Yoshihisa Fujino (present chairperson of the study group), Dr. Akira Ogami, Dr.

Arisa Harada, Dr. Ayako Hino, Dr. Hajime Ando, Dr. Hisashi Eguchi, Dr. Kazunori Ikegami,

Dr. Kei Tokutsu, Dr. Keiji Muramatsu, Dr. Koji Mori, Dr. Kosuke Mafune, Dr. Kyoko

Kitagawa, Dr. Masako Nagata, Dr. Mayumi Tsuji, Ms. Ning Liu, Dr. Rie Tanaka, Dr.

Ryutaro Matsugaki, Dr. Seiichiro Tateishi, Dr. Shinya Matsuda, Dr. Tomohiro Ishimaru, and

Dr. Tomohisa Nagata. All members are affiliated with the University of Occupational and

Environmental Health, Japan.

**Disclosure**

**Ethical approval:** This study was approved by the ethics committee of the University of

Occupational and Environmental Health, Japan (reference No. R2-079).

**Informed Consent:** Informed consent was obtained through an online website.

**Registry and the Registration No. of the study/Trial:** N/A

**Animal Studies:** N/A

**Conflicts of Interest:** The authors declare no conflicts of interest associated with this

manuscript.

**References**

1. Shimizu K, Wharton G, Sakamoto H, Mossialos E. Resurgence of covid-19 in Japan.
*BMJ*. 2020;370:m3221.

2. Sarwar A, Imran M. Prioritizing Infection Prevention and Control Activities for SARS-
CoV-2 (COVID-19): A Multi-Criteria Decision-Analysis Method. *J Healthc Leadersh*.
2021;13:77-84.

3. Tokyo Metroplolitan Government. [Results of an emergency survey on telework
adoption rates]. Published May 11, 2020. Accessed December 25, 2020.
https://www.metro.tokyo.lg.jp/tosei/hodohappyo/press/2020/05/12/10.html

4. Hyland P, Shevlin M, McBride O, et al. Anxiety and depression in the Republic of
Ireland during the COVID-19 pandemic. *Acta Psychiatr Scand*. 2020;142(3):249-256.

5. Shi L, Lu Z-A, Que J-Y, et al. Prevalence of and Risk Factors Associated With Mental
Health Symptoms Among the General Population in China During the Coronavirus
Disease 2019 Pandemic. *JAMA Netw Open*. 2020;3(7):e2014053.

6. Yao H, Chen J-H, Xu Y-F. Patients with mental health disorders in the COVID-19
epidemic. *Lancet Psychiatry*. 2020;7(4):e21.

7. Niedzwiedz CL, Green MJ, Benzeval M, et al. Mental health and health behaviours
before and during the initial phase of the COVID-19 lockdown: longitudinal analyses of
the UK Household Longitudinal Study. *J Epidemiol Community Health*.
2021;75(3):224-231.

8. Gao J, Zheng P, Jia Y, et al. Mental health problems and social media exposure during
COVID-19 outbreak. *PLoS One*. 2020;15(4):e0231924.

9. Starks TJ, Rendina HJ, Breslow AS, Parsons JT, Golub SA. The psychological cost of
anticipating HIV stigma for HIV-negative gay and bisexual men. *AIDS Behav*.
2013;17(8):2732-2741.

10.  Kunzler AM, Stoffers-Winterling J, Stoll M, et al. Mental health and psychosocial

support strategies in highly contagious emerging disease outbreaks of substantial public

concern: A systematic scoping review. *PLoS One*. 2021;16(2):e0244748.

11.  Aiello A, Khayeri MY-E, Raja S, et al. Resilience training for hospital workers in

anticipation of an influenza pandemic. *J Contin Educ Health Prof*. 2011;31(1):15-20.

12.  Torales J, O'Higgins M, Castaldelli-Maia JM, Ventriglio A. The outbreak of COVID-19

coronavirus and its impact on global mental health. *Int J Soc Psychiatry*.

2020;66(4):317-320.

13.  Giorgi G, Lecca LI, Alessio F, et al. COVID-19-Related Mental Health Effects in the

Workplace: A Narrative Review. *Int J Environ Res Public Health*. 2020;17(21).

doi:10.3390/ijerph17217857

14.  Fujino Y, Ishimaru T, Eguchi H, et al. Protocol for a nationwide Internet-based health

survey in workers during the COVID-19 pandemic in 2020. *bioRxiv*. Published online

February 5, 2021. doi:10.1101/2021.02.02.21249309

15.  Kessler RC, Andrews G, Colpe LJ, et al. Short screening scales to monitor population

prevalences and trends in non-specific psychological distress. *Psychol Med*.

2002;32(6):959-976.

16.  Sakurai K, Nishi A, Kondo K, Yanagida K, Kawakami N. Screening performance of

K6/K10 and other screening instruments for mood and anxiety disorders in Japan.

*Psychiatry Clin Neurosci*. 2011;65(5):434-441.

17.  Furukawa TA, Kawakami N, Saitoh M, et al. The performance of the Japanese version

of the K6 and K10 in the World Mental Health Survey Japan. *Int J Methods Psychiatr*

*Res*. 2008;17(3):152-158.

18.  Widera E, Chang A, Chen HL. Presenteeism: a public health hazard. *J Gen Intern Med*.

2010;25(11):1244-1247.

19. Leineweber C, Westerlund H, Hagberg J, Svedberg P, Alexanderson K. Sickness

presenteeism is more than an alternative to sickness absence: results from the

population-based SLOSH study. *Int Arch Occup Environ Health*. 2012;85(8):905-914.

20. Webster RK, Liu R, Karimullina K, Hall I, Amlôt R, Rubin GJ. A systematic review of

infectious illness Presenteeism: prevalence, reasons and risk factors. *BMC Public Health*.

2019;19(1):799.

21. Abrams EM, Greenhawt M. Risk Communication During COVID-19. *J Allergy Clin*

*Immunol Pract*. 2020;8(6):1791-1794.

22. Hamouche S. COVID-19 and employees' mental health: stressors, moderators and

agenda for organizational actions. *Emerald Open Research*. 2020;2:15.

doi:10.35241/emeraldopenres.13550.1

23. Morrison MF, Petitto JM, Ten Have T, et al. Depressive and anxiety disorders in women

with HIV infection. *Am J Psychiatry*. 2002;159(5):789-796.

24. Wada K, Smith DR, Ishimaru T. Reluctance to care for patients with HIV or hepatitis B

/ C in Japan. *BMC Pregnancy Childbirth*. 2016;16:31.

25. Havaei F, Ma A, Staempfli S, MacPhee M. Nurses' Workplace Conditions Impacting

Their Mental Health during COVID-19: A Cross-Sectional Survey Study. *Healthcare*

*(Basel)*. 2021;9(1). doi:10.3390/healthcare9010084

26. Coelho CM, Suttiwan P, Arato N, Zsido AN. On the Nature of Fear and Anxiety

Triggered by COVID-19. *Front Psychol*. 2020;11:581314.

Table 1. Characteristics of the subjects according to the number of workplace measures against COVID-19

| Factor | Number of workplace measures against COVID-19 (n=27036) | | | |
|---|---|---|---|---|
| | 0-1 | 2-3 | 4-5 | ≥6 |
| Number of subjects | 4177 | 3404 | 4603 | 14852 |
| Age, mean (SD) | 48.6 (9.9) | 47.0 (10.3) | 46.8 (10.6) | 46.6 (10.7) |
| Sex, male | 2489 (59.6%) | 1726 (50.7%) | 2222 (48.3%) | 7377 (49.7%) |
| Marital status, married | 2006 (48.0%) | 1751 (51.4%) | 2462 (53.5%) | 8810 (59.3%) |
| Job type | | | | |
|    Mainly desk work | 1954 (46.8%) | 1475 (43.3%) | 2119 (46.0%) | 7920 (53.3%) |
|    Jobs mainly involving interpersonal communication | 870 (20.8%) | 929 (27.3%) | 1342 (29.2%) | 3786 (25.5%) |
|    Mainly labor | 1353 (32.4%) | 1000 (29.4%) | 1142 (24.8%) | 3146 (21.2%) |
| Equivalent income (million JPY) | | | | |
|    50-249 | 1453 (34.8%) | 933 (27.4%) | 990 (21.5%) | 2334 (15.7%) |
|    250-374 | 1197 (28.7%) | 1101 (32.3%) | 1368 (29.7%) | 3884 (26.2%) |
|    375-489 | 838 (20.1%) | 727 (21.4%) | 1116 (24.2%) | 3944 (26.6%) |
|    ≥490 | 689 (16.5%) | 643 (18.9%) | 1129 (24.5%) | 4690 (31.6%) |
| Education | | | | |
|    Junior high school | 111 (2.7%) | 83 (2.4%) | 71 (1.5%) | 103 (0.7%) |
|    High school | 1475 (35.3%) | 1043 (30.6%) | 1213 (26.4%) | 3222 (21.7%) |
|    Vocational school/college, university, graduate school | 2591 (62.0%) | 2278 (66.9%) | 3319 (72.1%) | 11527 (77.6%) |
| Current smoker | 1289 (30.9%) | 990 (29.1%) | 1219 (26.5%) | 3506 (23.6%) |
| Alcohol consumption, 6–7 days a week | 1057 (25.3%) | 755 (22.2%) | 927 (20.1%) | 2935 (19.8%) |
| Number of employees in the workplace | | | | |
|    1-29 | 2394 (57.3%) | 1248 (36.7%) | 977 (21.2%) | 1546 (10.4%) |
|    30-99 | 1046 (25.0%) | 1203 (35.3%) | 1572 (34.2%) | 3119 (21.0%) |
|    100-999 | 451 (10.8%) | 594 (17.5%) | 1152 (25.0%) | 4956 (33.4%) |
|    ≤1000 | 286 (6.8%) | 359 (10.5%) | 902 (19.6%) | 5231 (35.2%) |
| Severe psychological distress (K6≥13) | 464 (11.1%) | 355 (10.4%) | 429 (9.3%) | 1212 (8.2%) |

Table 2. Implemented workplace measurements against COVID-19

| | Number of workplace measures against COVID-19 | | | | Total |
|---|---|---|---|---|---|
| | 0-1 | 2-3 | 4-5 | ≥6 | |
| | n=4177 | n=3404 | n=4603 | n=14852 | N=27036 |
| Refraining from and restriction of business trips | 56 (1.3%) | 400 (11.8%) | 1479 (32.1%) | 12725 (85.7%) | 14660 (54.2%) |
| Refraining from or restriction on visitors | 26 (0.6%) | 189 (5.6%) | 867 (18.8%) | 11217 (75.5%) | 12299 (45.5%) |
| Refraining from or requesting a limit on the number of people at social gatherings and dinners | 149 (3.6%) | 1200 (35.3%) | 3340 (72.6%) | 14525 (97.8%) | 19214 (71.1%) |
| Refraining from or limiting face-to-face internal meetings | 8 (0.2%) | 250 (7.3%) | 1418 (30.8%) | 13001 (87.5%) | 14677 (54.3%) |
| Wearing masks at all times during working hours | 551 (13.2%) | 2315 (68.0%) | 3970 (86.2%) | 14387 (96.9%) | 21223 (78.5%) |
| Installation of partitions and consideration of workplace layout (e.g. altering desk layout or adjusting flow lines) | 83 (2.0%) | 748 (22.0%) | 2316 (50.3%) | 12587 (84.7%) | 15734 (58.2%) |
| Recommended daily temperature checks by workers themselves at home | 183 (4.4%) | 1349 (39.6%) | 2892 (62.8%) | 12825 (86.4%) | 17249 (63.8%) |
| Encouragement of telecommuting | 114 (2.7%) | 255 (7.5%) | 539 (11.7%) | 6886 (46.4%) | 7794 (28.8%) |
| Prohibiting eating at a worker's own desk | 12 (0.3%) | 87 (2.6%) | 292 (6.3%) | 4206 (28.3%) | 4597 (17.0%) |
| Requesting employees not to come to work when they are not feeling well | 464 (11.1%) | 355 (10.4%) | 429 (9.3%) | 1212 (8.2%) | 2460 (9.1%) |

Table 3. Association of the number of workplace measures against COVID-19 and severe psychological distress (K6≥13)

| number of workplace measures against COVID-19 | Age-sex adjusted | | | | Multivariate* | | | |
|---|---|---|---|---|---|---|---|---|
| | OR | 95% CI | | p | OR | 95% CI | | p |
| 0-1 | 1.53 | 1.37 | 1.72 | p<0.001 | 1.48 | 1.30 | 1.68 | p<0.001 |
| 2-3 | 1.34 | 1.18 | 1.52 | p<0.001 | 1.32 | 1.15 | 1.50 | p<0.001 |
| 4-5 | 1.16 | 1.04 | 1.31 | 0.010 | 1.15 | 1.02 | 1.29 | 0.025 |
| ≥6 | reference | | | | reference | | | |

* The multivariate model included sex, age, education, equivalent household income, occupation, number of business establishments, smoking status, and alcohol consumption. The incidence rate of COVID-19 by prefecture was also used as a prefecture-level variable.

Table 4. Association of workplace measures against COVID-19 and severe psychological distresss (K6≥13)

| | Age-sex adjusted | | | Multivariate* | | |
|---|---|---|---|---|---|---|
| | OR | 95% CI | p | OR | 95% CI | p |
| Refraining from and restrictions on business trips | 0.78 | 0.72 0.85 | <0.001 | 0.83 | 0.76 0.91 | <0.001 |
| Refraining from and restrictions on visitors | 0.85 | 0.78 0.92 | <0.001 | 0.90 | 0.83 0.99 | 0.027 |
| Refraining from or requesting a limit on the number of people at social gatherings and dinners | 0.75 | 0.68 0.81 | <0.001 | 0.80 | 0.72 0.88 | <0.001 |
| Refraining from or limiting face-to-face internal meetings | 0.77 | 0.70 0.83 | <0.001 | 0.80 | 0.73 0.87 | <0.001 |
| Wearing masks at all times during working hours | 0.82 | 0.74 0.91 | <0.001 | 0.86 | 0.77 0.96 | 0.006 |
| Installation of partitions and consideration of workplace layout (e.g. altering desk layout or adjusting flow lines) | 0.79 | 0.72 0.86 | <0.001 | 0.81 | 0.74 0.89 | <0.001 |
| Recommended daily temperature checks by workers themselves at home | 0.82 | 0.75 0.90 | <0.001 | 0.84 | 0.76 0.92 | <0.001 |
| Encouragement of telecommuting | 0.85 | 0.77 0.93 | 0.001 | 0.89 | 0.80 0.99 | 0.025 |
| Prohibiting workers from eating at their own desk | 1.08 | 0.97 1.20 | 0.172 | 1.12 | 1.00 1.25 | 0.050 |
| Requesting employees not come to work when they are not feeling well | 0.60 | 0.55 0.65 | <0.001 | 0.60 | 0.54 0.66 | <0.001 |

* The multivariate model included sex, age, education, equivalent household income, occupation, number of business establishments, smoking status, and alcohol consumption. The incidence rate of COVID-19 by prefecture was also used as a prefecture-level variable.