Global Burden of Disease worldwide cohort analysis of

dietary and other risk factors for cardiovascular diseases:

lipid hypothesis versus fat-soluble vitamin hypothesis

David K Cundiff (0000-0002-3206-9665), independent researcher,[1, 3] Chunyi Wu (0000-0002-

2186-3433), Research Epidemiologist/Statistician,[2, 3]

1 Long Beach, California, USA

2 Area Specialist Lead in Epidemiology and Statistics, Michigan Medicine, Ann Arbor,

Michigan, USA

3 Volunteer collaborators with the Institute of Health Metrics and Evaluation, Seattle,

Washington, USA

Correspondence to David K Cundiff davidkcundiff@gmail.com Phone: 01-562-438-8805

Word count: abstract 416, Text: 3884

Key Points

Question: Is the lipid hypothesis valid and/or is the fat-soluble vitamin hypothesis valid?

Findings: This worldwide cohort study suggested the lipid hypothesis is valid in cohorts with high intakes of animal foods and added fats. The fat-soluble vitamin hypothesis is valid in worldwide cohorts.

Meaning: That both the lipid hypothesis and fat-soluble vitamin hypothesis are valid suggests that a yet not precisely defined moderate amount of dietary animal foods and added fats contributes to the lowest cardiovascular risk.

Abstract

Importance Despite most academic thought leaders and the USDA Dietary Guidelines endorsing the lipid hypothesis, debate about whether dietary saturated fat and cholesterol cause cardiovascular disease has continued.

Objective To test validities of the lipid hypothesis and our "fat-soluble vitamin hypothesis"

Design Using Global Burden of Disease (GBD 2017) data on cardiovascular disease deaths/100k/year, ages 15-69 years old in male and female cohorts (CVD) and dietary and other risk factors, we formatted and population weighted data from 195 countries.

Setting Worldwide

Participants The formatted 7846 rows of data (cohorts) represented about 1 million people each, about 7.8 billion people in 2020.

Interventions We correlated CVD with dietary and other risk factors worldwide and in appropriate subsets.

Exposures Dietary and other risk factors

Main Outcome measures CVD versus dietary and other risk factors worldwide and in subsets.

Results

After empirical exploration, we defined a "fat-soluble vitamin variable" (FSVV) as the sum of

kilocalories/day (KC/d) of processed meat + red meat + fish + milk + poultry + eggs + (saturated

fatty acids (SFA) + polyunsaturated fatty acids (PUFA) + trans fatty acids (TFA)) * 0.46. Low

density lipoprotein cholesterol (LDL-c) correlated strongly positively with FSVV worldwide

(r=0.780, 95% CI 0.771 to 0.788, p<0001, n=7846 cohorts), so we considered FSVV our marker

variable to test the lipid hypothesis and our fat-soluble vitamin hypothesis. FSVV correlated

positively with CVD in high FSVV cohorts (FSVV≥567.3 KC/d: r=0.523, 95% CI 0.476 to

0.567, p<0001, n=974 cohorts, worldwide mean FSVV=285.4 KC/d). FSVV correlated

negatively with CVD worldwide (r= -0.329, 95% CI -0.349 to -0.309, $p<0.0001$). The analysis

dataset and associated SAS codes are available to check the accuracy of the tables in this

analysis: https://data.mendeley.com/datasets/g6b39zxck4/3. SAS statisticians who check the

analysis and report their findings as comments on this preprint, will be eligible to become paper

contributors by contacting lead author (davidkcundiff@gmail.com).

Conclusion and Relevance

Since FSVV correlated positively with CVD only in high FSVV cohorts, the lipid hypothesis

was supported only in GBD cohorts (and by inference individuals) with high FSVV intake.

These worldwide data supported the fat-soluble vitamins hypothesis because FSVV correlated

negatively with CVD, meaning the more fat-soluble vitamin containing animal foods and added

fat for gut absorption the less the CVD. A moderate intake of FSVV foods, not yet precisely

quantified, is predicted to contribute to reducing CVD. This GBD cohort data analysis

70   methodology could supplement prospective observational studies of individuals to be used in

71   developing food policy and education strategies for improving public health.

73   **Trial Registration:** N/A

Introduction

The scientific validity of the Dietary Guidelines for Americans for 2015-2020, including

guidelines based on the lipid hypothesis, was challenged by Journalist Nina Teicholz in the

BMJ.[1] The Center for Science in the Public Interest called for the *BMJ* to retract the article.

The peer reviewers the *BMJ* selected to adjudicate the far-reaching dispute concluded that,

"Teicholz's criticisms of the methods used by Dietary Guidelines for Americans Committee are

within the realm of scientific debate."[2] Influential Stanford University meta researcher, Dr.

John Ioannidis, called for radical reform of all nutritional epidemiology methodologies used to

influence food/agricultural policies and to produce dietary guidelines for clinicians and the

public.[3] Currently, no methodology for relating cardiovascular disease events to food intake

and other risk factors has been generally accepted as rigorous, replicable, and scientifically valid.

This novel statistical analysis of cardiovascular disease early death related to diet and other risk

factors will attempt to answer Dr. Ioannidis' call.

We define our fat-soluble vitamin hypothesis as, "suboptimal dietary intake of fat-

soluble vitamins and added fat for gut absorption increases cardiovascular disease

risk." Fat-soluble vitamins include vitamins A, D, E, and K (K1=phylloquinones and K2=

menaquinones). Vitamin A and its beta-carotene precursor are important in preventing congenital

heart disease.[4, 5] Vitamin E protects the vasculature against oxidative stress, including lipid

peroxidation and the production of atherogenic forms of LDLc—a part of the pathogenesis of

cardiovascular disease.[6] Vitamin K2 (menaquinones from animal foods and fermented plant

products) inhibits arterial calcification, a major factor in the pathogenesis of atherosclerosis and

cardiovascular disease.[7] Regarding vitamin D, a meta-analysis of prospective studies

showed an inverse correlation between cardiovascular disease and circulating

25(OH)-vitamin D ranging from 20 to 60 nmol/L.[8]

The four fat-soluble vitamins come primarily from animal foods. Vitamin K2 levels

are high in aged cheeses, other fermented dairy products,[9] organ meats (e.g.,

bovine and pork liver), chicken, and egg yolks.[10] The highest vitamin K2 levels

per 100g serving are in fermented legumes, particularly soy (e.g., natto and miso

(Japan),[11, 12] soybean Douchi, soybean Meitauza, and soybean curd (China and Taiwan),[13]

and Cheonggukjang, a fermented soybean stew (South Korea)).[14]

This paper will use population weighted, formatted, worldwide GBD 2017 data from the Institute

of Health Metrics and Evaluation (IHME) to assess the impact of dietary, metabolic, and other

risk factors on cardiovascular disease early deaths.

Methods

As volunteer collaborators with the IHME, we acquired and utilized raw GBD worldwide ecological data (GBD 2017: 195 countries and 365 subnational locations, n=1120 male and female cohorts). Data consisted of the rate of cardiovascular disease early deaths, metabolic risk factors, dietary risk factors and covariates, and other risk factors of male and female cohorts 15-49 years old and 50-69 years old from each year 1990-2017. GBD worldwide citations of over 12,000 surveys constituting ecological data inputs for this analysis are available online from IHME.[15] The main characteristics of IHME GBD data sources, the protocol for the GBD study, and all risk factor values have been published by IHME GBD data researchers and discussed elsewhere.[16-19] These include detailed descriptions of categories of input data, potentially important biases, and the methodologies of analysis. We did not clean or pre-process any of the GBD data. GBD cohort risk factor and health outcome data from the IHME had no missing records. Food risk factors came from IHME as gram/day (g/d) consumed per capita. GBD dietary covariate data originally came from Food and Agriculture Organization surveys of animal and plant food commodities available per capita in countries worldwide (potatoes, corn, rice, sweet potatoes, poultry, and eggs)—as opposed to consumed per capita.[15]

Supplementary Table 1 lists the relevant GBD risk factors, covariates, and other available variables with definitions of those risk factor exposures.[20] Once IHME staff time constraints due to COVID-19 data collection and modeling are over, the updated GBD 2019 data with all the variables as the GBD 2017 data we used for this analysis may be obtained by volunteer researchers collaborating with IHME.[21]

Study design and population

For cardiovascular disease early deaths and all risk factors, we averaged the values for ages 15-

49 years old together with 50-69 years old for each male and female cohort for each year.

Finally, for each male and female cohort, data from all 28 years (1990-2017) of the means of the

rate of cardiovascular disease early deaths and risk factor exposures were averaged using the

computer software program R. We defined the rate of cardiovascular disease early deaths as

"deaths/100k/year of males and females 15-69 years old": abbreviated "CVD."

To weigh the data according to population, internet searches (mostly Wikipedia) yielded the

most recent population estimates for countries and subnational states, provinces, and regions.

The 1120 GBD cohorts available were population weighted by a software program in R resulting

in an analysis dataset with 7846 population weighted cohorts representing about 7.8 billion

people in 2020. Each male or female cohort in the population-weighted analysis dataset

represented approximately 1 million people (range: < 100,000 to 1.5 million). For example, India

with about 1.234 billion people had 617 rows of the same data for males and 617 rows for

females. Maldives, with about 445,000 people, had a single row of data for males and another for

females. Without population-weighting the data, cohorts in India and Maldives each would have

had one row for males and one row for females in the analysis dataset, invalidating the analysis

results. World population data from the World Bank or the Organisation for Economic Co-

operation and Development could not be used because they did not include all 195 countries or

any subnational data.

Supplementary Table 2 details how omega-3 fatty acid g/d was converted to fish g/d using data

on the omega-3 fatty acid content of frequently eaten fish from the National Institutes of Health

Office of Dietary Supplements (USA).[22] As shown in Supplementary Table 3, we converted all

of the animal and plant food data, including alcohol and sugary beverage consumption, from g/d

to kilocalories/day (KC/d). For the g/d to KC/d conversions, we used the Nutritionix track

app,[23] which tracks types and quantities of foods consumed. Saturated fatty acids (SFA: 0-1

portion of the entire diet) was not available with GBD 2017 data, so GBD saturated fatty acids

risk factor data from GBD 2016 was used. Polyunsaturated fatty acid (PUFA) and trans fatty acid

(TFA) GBD risk factor data from 2017 (0-1 portion of the entire diet) were also utilized, but

monounsaturated fat data were not available. These fatty acid data expressed as 0-1 portion of

the entire diet were converted to KC/d by multiplying by the total KC/d available per capita for

each cohort.

170   Outcome variable and covariates

The principal outcome variable, CVD, was a combination variable consisting of the

deaths/100k/year from ischemic heart disease, stroke, hypertensive heart disease, rheumatic heart

disease, non-rheumatic valvular disease, subarachnoid haematoma, myocarditis, alcoholic

cardiomyopathy, endocarditis, aortic aneurysm, and atrial fibrillation. Given the multiple dietary

and other influences on the causation of CVD, we did not attempt to differentiate between risk

factors and covariates.

Statistical methods

To determine the strengths of the risk factor correlations with CVD of population weighted worldwide cohorts (n=7846 cohorts) or subgroups of cohorts (e.g., highest and lowest CVD cohorts, etc.), we utilized Pearson correlation coefficients: r, 95% confidence intervals (CIs), and p values. A subgroup of the cohorts with the lowest CVD values was determined as 500 pair of male and female cohorts (mean of male and female CVD values) totaling 1000 total cohorts (an arbitrary number). Using the FSVV median value in that low CVD subgroup, we formed a subgroup with FSVV greater than or equal to the median FSVV. This allowed us to test the lipid hypothesis in cohorts with the highest intakes of animal foods and added fats.

To identify confounding in the univariate correlations, we used partial correlations, holding constant the suspected confounder variables.

We used SAS and SAS OnDemand for Academics software 9.4 (SAS Institute, Cary, NC) for the data analysis.

Sensitivity analyses

We performed subset analyses on the 1000 cohorts with the lowest levels of CVD and on cohorts with FSVV levels greater than or equal to the median FSVV KC/d value of the 1000 cohorts with the lowest levels of CVD.

Results

To test the fat-soluble vitamin hypothesis, we summed the KC/d of processed meat, red meat, fish, milk (including all dairy products), poultry, eggs, and added fat (SFA, PUFA, and TFA) to create a fat-soluble vitamins variable (FSVV). In determining the portion of SFA, PUFA, and TFA added in addition to the SFA, PUFA, and TFA in the animal and plant foods, an adjustment factor (fatty acids * 0.46), adapted from the website "Our World in Data," differentiated the fatty acids in individual foods (54% of the total) from the added fatty acids (46% of the total).[24] This adjustment prevented double counting fatty acids KC/d and allowed a reasonably accurate determination of FSVV KC/d.

Table 1 shows the 43 dietary, metabolic, and other risk factors potentially contributing to CVD. See Supplementary Table 1 for definitions of the risk factor and co-variate variables. LDLc negatively correlated with CVD worldwide (r=-0.279, 95% CI -0.299 to -0.258, p<0001, Table 1), meaning the higher the LDLc mmol/L the lower the CVD. Figure 1 gives the worldwide plot of LDLc versus CVD, showing widely scattered data points with negative slope.

Similar to LDLc, FSVV negatively correlated with CVD worldwide (r=-0.329, 95% CI -0.349 to -0.309, p<0001, Table 1), meaning more FSVV KC/d (processed meats KC/d + red meats KC/d + fish and seafood KC/d + milk KC/d + poultry KC/d + eggs KC/d + added fatty acids) correlated with reduced CVD.

Table 1 also shows that six of the nine plant foods negatively correlated with CVD. Potatoes (KC/d available, a covariate derived from total potato KC available/population, including highly

processed potatoes) positively correlated with CVD (r=0.050, 95% CI 0.028 to 0.073, p<0001).

Sugary beverages also correlated positively with CVD (r=0.113, 95% CI 0.091 to 0.135,

p<0001). Rice and legumes were not significantly correlated with CVD.

Subset analyses

Table 2 includes 500 male and female pairs of cohorts (total=1000 cohorts, an arbitrary number,

representing approximately 1 billion people out of about 7.8 billion people worldwide) with the

lowest CVD values (CVD≤ 293.0 compared with mean CVD=543.7 worldwide).

Table 3 shows FSVV related risk factors (processed meat, red meat, fish, milk, poultry, eggs, and

added (SFA+PUFA+TFA)) for the 34 countries in Table 2 in descending order by CVD.

Table 4 gives all the CVD risk factors for seven representative countries out of the 34 countries

in Tables 2 and 3 (CVD≤ 293.0).

Table 5 shows CVD and risk factor statistics for cohorts with FSVV≥567.3 KC/d (i.e., at or

above the median FSVV of Table 2 countries). In this subset of 974 cohorts, of which 500 were

included in the low CVD subset (Table 2), FSVV strongly positively correlated with CVD

(r=0.523, 95% CI 0.476 to 0.567, p<0001).

Although FSVV correlated strongly with LDLc worldwide (r=0.780, 95% CI 0.771 to 0.788,

p<0001, n=7846 cohorts, Table 1), FSVV negatively correlated with LDLc in cohorts with

FSVV ≥ 567.3 KC/d (r=-0.571, 95% CI -0.612 to -0528, p<0001, n= 974 cohorts). Also with

FSVV ≥ 567.3 KC/d, LDLc and CVD negatively correlated (r=-0.254, 95% CI -0.312 to -0.194,

p<0001, n= 974 cohorts). Because of the inconsistency of the FSVV versus LDLc relationship

with very high FSVV KC/d levels, we considered LDLc confounded at high FSVV levels. We

therefore used FSVV, rather than LDLc, to test for the lipid hypothesis in high FSVV cohorts.

The average alcohol intake was 81.0 KC/d worldwide, and alcohol KC/d correlated negatively

with CVD (r=-0.061, 95% CI -0.083 to -0.039, p<0001, Table 1). However, in high FSVV

cohorts (FSVV≥567.3 KC/d, mean FSVV=692.6 KC/d and mean alcohol=155.0 KC/d), alcohol

positively correlated with CVD (r=0.475, 95% CI 0.425 to 0.522, p<0001, n=974 cohorts, Table

5).

The only fat-soluble vitamin in the GBD database, vitamin A deficiency incidence/100k in

children ≤ 5 years old, correlated positively with CVD worldwide (r=0.210, 95% CI 0.189 to

0.231, p<0001, Table 1).

Dietary fiber was not significantly correlated with CVD worldwide (r=0.019, 95% CI -0.003 to

0.041, p=0.09, Table 1). Once it was adjusted for FSVV, alcohol, sugary beverages, and potatoes

by partial correlation analysis; dietary fiber correlated negatively with CVD (r=-0.052, 95% CI -

0.074 to -0.030, p<0.0001).

Physical activity positively correlated with CVD worldwide (r=0.160, 95% CI 0.139 to 0.182,

p<0.0001, Table 1). Once it was adjusted for FSVV, smoking, and sex by partial correlation

analysis; physical activity correlated negatively with CVD (r=-0.127, 95% CI -0.149 to -0.104,

p<0.0001).

In cohorts, mostly in developed countries, with FSVV ≥567.3 KC/d discontinuing breast feeding

correlated positively with CVD (r=0.268, 95% CI 0.208 to 0.325, p<0.0001, Table 5).

Secondhand smoking negatively correlated with CVD worldwide (r=-0.225, 95% CI -0.246 to -

0.204, p<0.0001, Table 1). After adjusting secondhand smoking for smoking, sublingual tobacco

use, household smoke, ambient air pollution, and sex by partial correlation analysis; secondhand

smoke correlated positively with CVD (r=0.048, 95% CI 0.025 to 0.070, p<0.0001).

The metabolic risk factors, BMI, FPG, and LDLc, all correlated negatively with CVD worldwide

(r=-0.240, 95% CI -0.261 to -0.219, p<0.0001, r=-0.178, 95% CI -0.200 to -0.157, p<0.0001, and

r=-0.279, 95% CI -0.299 to -0.258, p<0.0001, respectively, Table 1). Once they were

adjusted for FSVV, alcohol, sugary beverages, and potatoes by partial correlation analysis; they

all correlated positively with CVD (r=0.028, 95% CI 0.006 to 0.050, p=0.0144, r=0.076, 95% CI

0.054 to 0.098, p<0.0001, r=0.170, 95% CI 0.148 to 0.191, p<0.0001, respectively). In the

cohorts with FSVV≥567.3 KC/d, BMI and FPG positively correlated with CVD (r=0.484, 95%

CI 0.434 to 0.530, p<0.0001 and r=0.091, 95% CI 0.028 to 0.153, p=0.0046, respectively).

Discussion

In high FSVV cohorts, (FSVV≥ 567.4 KC/d), why LDL-c correlated negatively with FSVV (r=-0.571 95% CI -0.612 to -0.528, *p*<0.0001, n=974 cohorts) but positively correlated with LSVV worldwide (r=0.780, 95% CI 0.771 to 0.788, p<0001) is not entirely clear. However, multiple studies have documented that LDLc may be influenced by genetic, racial, or ethnic differences that do not necessarily correspond with risk of CVD.[25, 26] Additionally, LDLc lowering medications in wide use especially in high FSVV developed countries may also confound the relationship of LDLc with both LSVV and CVD risk in this subset cohort analysis.

Since there are high FSVV and low FSVV individual people widely distributed worldwide, prospective observational studies of individuals in high FSVV areas, rather than GBD cohort studies, could better demonstrate positive correlations between LDLc (or total cholesterol) and CVD. This may explain why the Seven Countries Study[27] of Ancel Keys and the Framingham Heart Study,[28] both in high FSVV regions, supported the lipid hypothesis. However, the Prospective Urban Rural Epidemiology (PURE) Study, with individual data on 135,335 patients from three high-income, 11 middle-income, and four low-income countries was clearly at variance with the lipid hypothesis.[29] The PURE Study found that higher saturated fat intake was associated with lower risk of stroke (quintile 5 *vs* quintile 1, HR 0.79, (95% CI 0.64 to 0.98), p_{trend}=0.0498).

This global data analysis supports the fat-soluble vitamin hypothesis or possibly another mechanism that accounts for FSVV KC/d being negatively correlated with CVD worldwide (r=-0.308, 95% CI -0.358 to -0.257, p<0001, Table 1).

Given the negative correlations of CVD with both fish and eggs in both tables 1 and 5, the

reduction in FSVV might best come from meat, poultry, and added fats. Although milk

(including other dairy products) correlated positively with CVD in Table 5 (r=0.185, 95% CI

0.123 to 0.245, p<0001), this probably related to Japan and Peru having the lowest and fifth

lowest CVD in Table 3 (Japan: mean CVD=169.2, mean milk=29.0 KC/d, Peru: mean

CVD=197.7, mean milk=15.3 KC/d). Without Japan and Peru in the analysis (n=834 cohorts),

milk correlated negatively with CVD (r=-0.071, 95% CI -0.138 to -0.030, p=0.0405). Of the 34

low CVD countries in Table 3, 20 had high milk intakes (mean milk ≥68.8 KC/d versus mean

milk worldwide=25.0 KC/d) and these high milk intake cohorts all had relatively high per capita

intakes of cheese.[30, 31] Fermented, full-fat dairy products have high levels of vitamin K2,[9,

10] which have been found to be important in preventing calcification of arteries and

atherosclerosis.[32]

Table 4 shows that the Japanese, with low milk KC/d intake (mean milk=29.0 KC/d), had the

world's lowest CVD (mean CVD=169.2) while crucial risk factors were high (e.g., smoking:

mean male/female smoking=26.8%, salt intake (mean sodium= 6.01 g/d, systolic blood pressure

(mean SBP= 137.6 mm/Hg). Additionally, relative to fermented cheeses, the content of vitamin

K2 in fish is low to negligible.[10] Instead, Japanese consume high quantities of vitamin K2 in

the form of fermented soy products like natto and miso.[11, 12] As noted in the introduction,

people in Taiwan and South Korea (also listed among low CVD countries) also have high intakes

of fermented legumes.

Potatoes KC/d available positively correlated with CVD both worldwide (r=0.050, 95% CI 0.028

to 0.073, p< 0.0001, Table 1) and in cohorts with FSVV≥ 567.3 KC/d (r=0.109, 95% CI 0.046 to

0.170, p< 0.0001, Table 5). Notably, half, or more of the potatoes consumed worldwide are in

the form of highly processed food products.[33] Recent large prospective observational studies

have found higher consumption of ultra-processed foods associated with an increased risk of

cardiovascular disease incidence and CVD mortality.[34, 35] Data from 79 high- and middle-

income countries show that ultra-processed products dominate the food supplies of high-income

countries and that their consumption is now rapidly increasing in middle-income countries.[36]

The finding that sodium negatively correlated with CVD worldwide (r=-0.213, 95% CI -0.234 to

-0.192, p<0.0001) is contrary to an analysis of sodium intake versus cardiovascular deaths in 66

countries by Mozaffarian, et. al who attributed 9.5% of cardiovascular deaths worldwide to high

sodium intake.[37] However, O'Donnell and colleagues found the relationship of sodium to

cardiovascular disease to be J shaped curve and suggested that the lowest cardiovascular risk is

with moderate sodium intake in the 3-5 g/d range.[38]

The positive correlation of physical activity with CVD was surprising (Table 1). However, (1)

countries with higher FSVV KC/d levels also had less physical activity (FSVV versus physical

activity: r=-0.366, 95% CI: -0.385 to -0.347, *p*< 0.0001), (2) females had less physical activity

than males (mean physical activity males=5061 METs/week, mean physical activity

females=4356 METs/week), along with less CVD (mean CVD males=656.2, mean CVD

females=428.6), and (3) males smoked tobacco more than females (mean smoking (0-1)

males=0.339 or 33.9%, mean smoking (0-1) females=0.070 or 7%). When FSVV, sex, and

smoking were held constant, physical activity correlated negatively with CVD worldwide (r= -

0.126, 95% CI -0.148 to -0.104, p<0.0001).

The moderately strong positive correlation of child ≤5 years old severe underweight (>2 SD

below the mean for height) with CVD worldwide (r=0.306, 95% CI 0.285 to 0.325, p<0.0001,

Table 1) suggested a relationship of infant/child malnutrition with later CVD. Babies surviving

the so called "Dutch famine" toward the end of the second world war (1944-45) have been

shown to have higher subsequent heart disease than earlier or later cohorts in Holland.[39] This

suggested that in utero and early infancy severe malnutrition may subsequently contribute

substantially to CVD, especially in developing countries.

Prematurely stopping breast feeding positively correlated with CVD in the countries with FSVV

≥567.3 KC/d (r=0.268 95% CI: 0.208 to 0.325, p=0.0027, Table 5). In those countries,

prematurely stopping breast feeding was more common than in developing countries (range (0-

1): 0.129 to 0.268, mean=0.196). A review of observational studies, in developed countries, of

breast feeding related to metabolic risk factors for CVD suggested that breast feeding was

associated with increased insulin sensitivity and decreased systolic blood pressure in later life.

Breast feeding also has metabolic benefits for the mother.[40]

According to The World Health Organisation, ambient air pollution (particulate

matter ≤2·5 micrometers diameter (PM $_{2.5}$)) causes cardiovascular and respiratory diseases and

cancers.[41] The WHO considers ≥10 μg/m^3 of PM$_{2.5}$ particles a health hazard. In this

analysis, the ambient air pollution (PM $_{2.5}$) in the countries with FSVV≥567.3 KC/d

averaged 12.7 $\mu g/m^3$ and mean CVD=273.7 from 1990 to 2017 (Table 5). Worldwide,

comparable numbers were mean CVD=543.7 and air pollution (PM $_{2.5}$) mean=44.7 $\mu g/m^3$.

These IHME GBD data support the link of smoking tobacco to CVD (r=0.298, 95% CI

0.278 to 0.318, p<0.0001, Table 1). However, given the multiple dietary and other

contributors, the CVD population attributable risk for smoking is probably far below

the World Health Organization estimate that tobacco accounts for 20% of coronary

artery disease mortality.[42] The mean incidence of smoking worldwide=20.5%

(Table 1). In the 34 countries with the lowest CVD, the mean smoking

incidence=23.3% (n=974 cohorts). Smoking was clearly a major risk factor for CVD,

but these data suggest that diet was more influential

The positive correlation of blood lead levels with CVD worldwide (r=0.180, 95% CI 0.159 to

0.201, p<0.0001, Table 1) is consistent with reports that lead toxicity leads to hypertension.[43]

Also, environmental lead exposure has been linked with all-cause mortality,

cardiovascular disease mortality, and ischaemic heart disease mortality in the

US.[44]

This analysis showed that kidney disease correlated positively with CVD (r=0.194, 95% CI

0.173 to 0.215, p<0.0001, Table 1). A systematic GBD review of kidney disease revealed that

chronic kidney disease resulted in an estimated 1.2 million deaths in 2017, of which a large

portion were from CVD.[45] Other CVD risk factors may also lead to kidney disease (e.g.,

systolic hypertension, types 1 and 2 diabetes).

Limitations of this study

Our study was subject to all the limitations discussed in previous GBD publications.[46, 47] These included gaps, biases, and inconsistencies in data sources as well as limitations in the methods of data processing and estimation. Having comprehensive data on dietary inputs is key to more accurate and reliable analyses. The GBD data on animal foods, plant foods, alcohol, sugary beverages, and fatty acids were not comprehensive and comprised only 1191.4 KC/d per person on average worldwide. Subnational data on all risk factors were available on only four countries. Because the data formatting and statistical methodology were new, this was necessarily a post hoc analysis and no pre-analysis protocol was possible. This GBD data analysis should be repeated with the most recently released GBD 2019 data when it becomes available to volunteer collaborators.

Generalizability

This new methodology presented for probing the GBD raw data to find correlations between dietary and other risk factors and health outcomes (e.g., CVD) can be applied to any of the risk factors and health outcomes available with the new IHME GBD 2019 data.

Conclusion

The lipid hypothesis (e.g., dietary SFA and cholesterol from animal foods causing CVD) was consistent with the strong positive correlation of FSVV with CVD in high FSVV cohorts but not worldwide. Global data analysis supported the fat-soluble vitamin hypothesis, because FSVV negatively correlated with CVD worldwide. Defining the optimal FSVV that will reduce CVD risk while not increasing other diseases (e.g., cancers and obesity) will require more research. Supplements of fat-soluble vitamins to reduce the FSVV required to minimize CVD should be studied, especially in developing countries with less access to animal foods.

This methodology of analysing IHME GBD data should augment systematic literature reviews in developing policy recommendations, clinical practice guidelines, and public health recommendations. GBD data analysis can bring more rigor, precision, and consensus to the field of population health, especially in the areas of dietary risk factors.

Figure 1. CVD versus LDLc scatter plot

440
441
442

Table 1. Dietary and other risks related to CVD worldwide (n=7846 cohorts)

CVD and CVD risk factors	Mean	SD	Min	Max	r	95% CI	95% CI	p
Cardiovascular disease deaths/100k/year ages 15-69 (CVD)	543.66	288.01	73.47	1844				
CVD mean m/f	543.69	246.13	135.46	1727	0.855	0.849	0.860	<.0001
LDL cholesterol mmol/L	2.35	0.40	1.27	3.25	-0.279	-0.299	-0.258	<.0001
Fat-soluble vitamin variable (FSVV)	285.36	193.31	58.78	932.18	-0.329	-0.349	-0.309	<.0001
Processed meat KC/d	5.33	9.72	0.20	68.77	-0.204	-0.226	-0.183	<.0001
Red meat KC/d	50.27	45.13	3.21	235.95	-0.232	-0.253	-0.211	<.0001
Fish KC/d	9.99	36.52	0.40	370.36	-0.203	-0.224	-0.181	<.0001
Milk KC/d	25.04	27.05	1.06	146.82	-0.192	-0.214	-0.171	<.0001
Poultry KC/d available	44.32	50.08	1.06	411.87	-0.289	-0.309	-0.268	<.0001
Eggs KC/d available	19.36	14.71	0.79	69.64	-0.390	-0.408	-0.371	<.0001
Added Saturated fatty acids KC/d	87.67	29.41	32.56	221.29	-0.239	-0.260	-0.219	<.0001
Added PUFAs KC/d	37.30	33.78	1.35	175.40	-0.316	-0.336	-0.296	<.0001
Added Trans fatty acids KC/d	6.09	6.28	0.91	35.77	-0.104	-0.126	-0.082	<.0001
Alcohol KC/d	81.03	57.33	4.25	429.81	-0.061	-0.083	-0.039	<.0001
Sugary beverages KC/d	298.36	152.38	72.91	1472.00	0.113	0.091	0.135	<.0001
Potatoes KC/d available	84.04	74.60	3.07	533.88	0.050	0.028	0.073	<.0001
Corn KC/d available	34.72	48.28	0.16	305.17	-0.062	-0.084	-0.040	<.0001
Fruits KC/d	40.21	22.50	3.58	161.39	-0.355	-0.374	-0.336	<.0001
Vegetables KC/d	79.76	43.12	9.48	304.17	-0.107	-0.128	-0.085	<.0001
Nuts and seeds KC/d	8.41	8.36	0.05	102.99	-0.277	-0.297	-0.256	<.0001
Whole grains KC/d	55.65	30.93	1.14	235.10	-0.194	-0.216	-0.173	<.0001
Legumes KC/d	51.74	32.23	0.51	194.70	-0.024	-0.046	-0.002	0.0319
Rice KC/d available	141.86	116.34	1.42	461.80	0.007	-0.015	0.029	0.5417
Sweet potatoes KC/d available	22.76	35.95	0.02	364.74	-0.167	-0.189	-0.146	<.0001
Total KC/d available	2574	418	1579	3898	-0.203	-0.224	-0.181	<.0001
Vit A deficiency children/100k/yr	23205	10939	1267	50969	0.210	0.189	0.231	<.0001
Sodium g/d	4.45	2.34	1.33	9.21	-0.214	-0.235	-0.193	<.0001
Calcium g/d	0.301	0.179	0.081	1.044	-0.169	-0.191	-0.148	<.0001
Dietary fiber g/d	9.21	3.15	2.72	22.68	0.019	-0.003	0.041	0.0932
Physical activity METs	4714	1368	1609	7669	0.160	0.139	0.182	<.0001
Child underweight >2SD	0.186	0.171	0.004	0.535	0.302	0.282	0.322	<.0001
Stop breast feeding <6 months	0.119	0.055	0.016	0.242	-0.302	-0.322	-0.282	<.0001
Ambient pollution PM $_{0.25}$	44.73	26.46	4.38	95.54	0.153	0.131	0.174	<.0001
Smoking rate (0-1)	0.205	0.176	0.003	0.640	0.297	0.277	0.317	<.0001
Secondhand smoking (0-1)	0.376	0.155	0.164	0.796	-0.225	-0.246	-0.204	<.0001
Sublingual tobacco rate (0-1)	0.068	0.095	0.001	0.419	0.284	0.264	0.304	<.0001
Blood lead level mcg/dl	5.01	1.01	1.22	8.37	0.180	0.159	0.201	<.0001
Household air pollution (0-1)	0.482	0.325	0.000	0.996	0.179	0.158	0.201	<.0001
Kidney disease stage III (0-1)	0.056	0.028	0.015	0.154	0.194	0.173	0.215	<.0001
Type 1 DM early deaths	10.37	9.39	0.55	112.49	0.340	0.320	0.359	<.0001
Type 2 DM early deaths	17.50	15.65	0.63	269.67	0.227	0.205	0.247	<.0001
BMI kg/M^2	21.77	2.29	17.95	29.39	-0.240	-0.261	-0.219	<.0001
Fasting plasma glucose mmol/L	4.30	0.35	3.32	5.58	-0.178	-0.200	-0.157	<.0001
Systolic BP mm Hg	133.91	4.32	123.41	147.89	0.195	0.174	0.216	<.0001
Socio-demographic index (0-1)	0.543	0.174	0.112	0.896	-0.337	-0.357	-0.317	<.0001
Sex male 1 and female 2	1.50	0.50	1.00	2.00	-0.395	-0.414	-0.376	<.0001

447 **Table 2. CVD lowest and highest 1000 cohorts (≈1 billion people)**

CVD and risk factors	CVD lowest ≈1 billion people (n=1000 cohorts)				CVD highest ≈1 billion people (n=1000 cohorts)			
	Mean	SD	Min	Max	Mean	SD	Min	Max
Cardiovascular Disease deaths/100k/year	227.81	95.25	73.47	422.36	1045	356.90	430.20	1844
CVD male/female mean	227.81	42.52	135.46	292.97	1045	200.47	802.24	1727
LDL cholesterol mmol/L	2.808	0.256	1.598	3.247	2.376	0.380	1.35	3.20
Fat-soluble vitamin variable (FSVV)	531.4	180.20	143.63	912.69	235.66	99.68	80.53	503.98
Processed meat KC/d	15.30	12.20	0.61	61.09	3.70	4.95	0.22	18.12
Red meat KC/d	90.72	48.06	12.03	235.95	36.82	29.40	7.07	172.00
Fish KC/d	49.54	92.63	2.85	370.36	4.56	2.69	0.78	12.97
Milk KC/d	55.64	35.39	7.70	146.82	25.61	16.78	1.06	75.87
Poultry KC/d available	90.44	47.27	4.74	240.56	30.23	22.39	2.78	100.04
Eggs KC/d available	36.63	11.25	4.77	69.64	14.73	13.63	1.51	46.81
Added Saturated fatty acids KC/d	113.40	36.53	59.52	221.29	87.22	24.03	33.03	205.85
Added PUFAs KC/d	69.80	35.78	15.77	175.40	25.48	14.54	2.73	84.90
Added Trans fatty acids KC/d	9.91	9.25	1.65	34.90	7.31	8.62	1.21	35.77
Alcohol KC/d	119.72	66.67	4.25	296.70	45.83	46.24	5.77	340.37
Sugary beverages KC/d	320.29	251.94	72.91	1472	303.30	63.68	180.80	685.42
Potatoes KC/d available	81.34	51.14	9.50	177.30	98.34	96.77	3.94	533.88
Corn KC/d available	48.19	75.26	1.79	287.14	31.77	37.30	0.20	305.17
Fruits KC/d	64.84	16.49	23.05	128.61	31.23	13.93	3.58	70.11
Vegetables KC/d	108.02	44.60	9.48	214.26	83.83	56.67	14.43	198.51
Nuts and seeds KC/d	15.90	10.50	0.66	39.80	4.81	2.91	0.05	17.91
Whole grains KC/d	64.71	38.62	1.61	173.46	28.30	33.01	1.14	156.75
Legumes KC/d	47.88	25.99	2.95	133.26	31.55	18.93	0.51	103.17
Rice KC/d available	78.10	97.95	7.58	349.08	75.93	107.38	1.42	461.80
Sweet potatoes KC/d available	3.92	5.14	0.04	27.51	5.84	20.20	0.02	364.74
Total KC/d available	2972	409	1948	3572	2627	409	1579	3254
Vit A deficiency children/100k/yr	14181	8723	1400	44100	20559	11395	1722	50969
Sodium g/d	3.92	1.22	1.33	6.70	3.30	0.76	1.33	5.96
Calcium g/d	0.506	0.189	0.183	1.044	0.321	0.150	0.089	0.792
Dietary fiber g/d	10.58	2.90	5.15	17.95	9.53	3.32	3.87	18.83
Physical activity METs	3802	1164	2162	7607	4273	1497	1838	7607
Child underweight >2SD	0.038	0.043	0.004	0.242	0.186	0.141	0.011	0.411
Stop breast feeding <6 mo	0.176	0.032	0.071	0.218	0.116	0.044	0.036	0.208
Ambient pollution PM $_{0.25}$	17.60	8.27	4.38	38.42	40.81	22.19	7.90	77.69
Smoking rate (0-1)	0.217	0.121	0.012	0.448	0.212	0.190	0.005	0.627
Secondhand smoking (0-1)	0.328	0.097	0.164	0.569	0.397	0.158	0.164	0.780
Sublingual tobacco rate (0-1)	0.010	0.015	0.001	0.111	0.060	0.072	0.001	0.238
Blood lead level mcg/dl	4.08	1.00	1.22	6.85	4.74	1.28	1.92	8.37
Household air pollution (0-1)	0.103	0.182	0.001	0.839	0.404	0.353	0.002	0.993
Kidney disease stage III (0-1)	0.042	0.020	0.015	0.111	0.078	0.024	0.035	0.139
Type 1 DM early deaths	4.79	3.88	0.59	20.65	16.38	15.31	2.62	112.49
Type 2 DM early deaths	15.69	18.23	0.75	87.92	23.65	24.88	0.89	269.67
BMI kg/M^2	23.74	1.60	19.61	27.06	22.46	2.15	17.95	26.08
Fasting plasma glucose mmol/L	4.56	0.25	3.54	5.12	4.28	0.42	3.38	5.18
Systolic BP mm Hg	133.26	4.41	123.41	142.15	136.86	4.39	124.59	147.89
Socio-demographic index (0-1)	0.757	0.125	0.351	0.896	0.497	0.183	0.186	0.824
Sex male 1 and female 2	1.50	0.50	1.00	2.00	1.50	0.50	1.00	2.00

Table 3. CVD related to fat-soluble vitamin related risk factors in low CVD countries

CVD lowest Countries (n=1000 cohorts)	n co-horts	CVD ascending order	FSVV	Processed meat	Red meat	Fish	Milk	Poultry	Eggs	Added SFA +PUFA +TFA
Japan	158	169.19	618.69	19.19	58.72	260.54	29.04	63.93	52.23	135.04
France	64	174.00	644.78	13.59	135.07	11.17	95.75	99.86	43.30	246.04
Switzerland	8	175.04	558.41	11.50	129.01	6.75	110.90	55.57	31.20	213.47
Andorra	2	185.55	763.18	16.58	150.84	16.44	99.23	189.92	47.55	242.62
Peru	32	197.74	183.12	1.01	16.92	3.82	15.34	35.04	12.03	98.95
Spain	46	206.25	597.93	11.65	139.92	10.28	82.99	113.88	43.35	195.86
Italy	60	206.44	587.57	15.78	126.36	9.15	89.54	84.16	36.39	226.20
Iceland	2	206.51	609.62	11.97	112.73	11.67	85.37	63.57	29.51	294.81
Australia	24	208.03	698.26	9.79	164.14	14.15	97.05	153.33	24.92	234.89
Canada	36	222.11	624.98	25.31	126.28	18.48	76.13	147.46	34.94	196.39
Israel	8	228.32	662.00	5.95	59.50	15.35	68.79	240.56	40.54	231.30
South Korea	50	232.63	357.23	5.97	82.88	5.96	15.72	43.10	26.89	176.71
Belgium	12	235.98	636.39	12.28	122.36	11.19	93.44	85.00	33.28	278.86
Netherlands	18	236.61	629.56	15.72	133.06	8.66	114.16	72.70	41.67	243.59
Taiwan	24	237.37	587.36	3.01	106.58	9.15	15.95	132.38	34.87	285.41
Mexico	108	237.55	413.25	11.29	63.03	8.69	34.29	85.85	39.92	170.19
Panama	4	244.31	383.83	2.75	61.32	9.44	20.72	95.43	12.96	181.23
Sweden	10	247.21	563.83	29.31	119.77	11.80	124.13	45.42	35.04	198.37
New Zealand non Mauri	4	247.32	699.81	9.31	158.86	11.56	88.74	128.14	40.24	262.95
Norway	32	248.72	605.30	48.14	109.37	10.10	100.96	41.65	31.52	263.56
England	20	253.09	534.56	19.21	96.29	10.70	87.80	89.30	32.90	198.36
Costa Rica	4	260.73	408.42	2.63	51.74	8.12	46.55	78.05	28.17	193.14
Luxembourg	2	262.53	578.00	14.65	163.17	11.79	91.99	71.48	27.60	197.32
Chile	18	267.50	404.12	19.11	90.33	9.65	31.93	97.29	19.70	136.10
Guatemala	16	269.85	231.05	1.89	20.05	5.28	13.32	53.94	30.46	106.12
Kenya	12	270.04	150.04	1.41	25.83	3.09	18.89	4.74	4.77	91.31
Ecuador	16	270.64	375.35	6.87	56.88	7.34	35.51	59.92	14.66	194.16
USA	94	274.00	825.54	36.45	136.35	15.79	90.90	190.90	41.10	314.05
Denmark	6	275.10	600.48	14.46	137.75	13.38	98.01	74.42	45.96	216.51
Austria	8	277.67	621.00	8.95	194.65	9.26	98.68	70.00	39.96	199.50
Portugal	10	278.50	524.81	5.16	124.77	8.77	79.43	94.26	24.18	188.25
Nicaragua	6	282.04	214.24	1.52	14.93	4.07	20.66	41.76	17.41	113.90
Tehran (Iran)	16	283.29	299.65	1.23	22.63	5.36	19.23	65.90	18.89	166.41
Thailand	70	289.91	247.43	0.70	32.01	6.95	7.85	57.07	31.10	111.75
Total cohorts	1000									

Table 4. Representative examples of low CVD countries (n=1000 cohorts) including all risk factors

CVD and CVD risk factors	Japan[a]	France[b]	Peru[c]	Mexico[d]	Panama[e]	Guatemala[f]	Ecuador[g]
n cohorts	158	64	32	108	4	16	16
CVD/100k/year ages 15-69	169.19	174.00	197.74	237.55	244.31	269.85	270.64
Fat-soluble vitamin variable	618.69	644.78	183.12	413.25	383.83	231.05	375.35
LDL cholesterol mmol/L	2.81	3.08	2.37	2.59	2.61	2.20	2.46
Processed meat KC/d	19.19	13.59	1.01	11.29	2.75	1.89	6.87
Red meat KC/d	58.72	135.07	16.92	63.03	61.32	20.05	56.88
Fish KC/d	260.54	11.17	3.82	8.69	9.44	5.28	7.34
Milk KC/d	29.04	95.75	15.34	34.29	20.72	13.32	35.51
Poultry KC/d available	63.93	99.86	35.04	85.85	95.43	53.94	59.92
Eggs KC/d available	52.23	43.30	12.03	39.92	12.96	30.46	14.66
Added Saturated fatty acids KC/d	77.73	172.79	68.34	89.13	101.32	60.62	133.54
Added PUFAs KC/d	53.78	67.38	24.47	52.14	72.29	41.90	51.69
Added Trans fatty acids KC/d	3.54	5.88	6.13	28.92	7.62	3.60	8.93
Alcohol KC/d	183.16	118.64	55.79	58.39	27.44	28.79	59.77
Sugary beverages KC/d	94.78	323.88	284.76	847.81	921.54	964.08	283.57
Potatoes KC/d available	42.29	126.40	142.55	25.79	32.01	16.68	67.56
Corn KC/d available	27.58	23.85	33.54	242.41	50.86	206.19	11.76
Fruits KC/d	44.43	59.56	52.27	65.68	52.92	39.61	107.07
Vegetables KC/d	149.32	101.07	46.17	60.72	27.81	54.98	31.79
Nuts and seeds KC/d	9.26	16.94	1.02	6.27	1.92	12.72	0.68
Whole grains KC/d	76.05	16.16	58.94	142.06	83.46	68.32	43.78
Legumes KC/d	71.84	17.07	45.78	76.99	38.35	73.84	21.99
Rice KC/d available	124.52	12.09	93.09	14.07	141.66	13.46	118.02
Sweet potatoes KC/d available	12.09	0.04	7.94	0.69	8.67	1.71	0.49
Total KC/d available	2590	3406	1948	3015	2288	2330	2273
Vit A deficiency children/100k/yr	8151	1643	19372	23939	11463	17947	20185
Sodium g/d	6.01	3.21	3.14	2.62	2.97	1.83	3.16
Calcium g/d	0.425	0.707	0.299	0.370	0.293	0.220	0.314
Dietary fiber g/d	13.43	8.68	9.44	15.51	5.90	12.54	6.25
Physical activity METs	3460	2795	3774	3833	4935	6142	3740
Child underweight >2SD	0.045	0.012	0.069	0.050	0.042	0.193	0.096
Stop breast feeding <6 months	0.171	0.208	0.108	0.171	0.140	0.087	0.130
Ambient pollution PM $_{0.25}$	13.10	13.87	28.99	24.28	13.24	28.61	17.86
Smoking rate (0-1)	0.268	0.289	0.075	0.149	0.073	0.077	0.080
Secondhand smoking (0-1)	0.353	0.320	0.238	0.344	0.178	0.224	0.206
Sublingual tobacco rate (0-1)	0.012	0.003	0.004	0.002	0.002	0.001	0.004
Blood lead level mcg/dl	2.65	3.86	5.17	5.34	4.80	5.89	4.51
Household air pollution (0-1)	0.002	0.010	0.454	0.213	0.193	0.627	0.080
Kidney disease stage III (0-1)	0.038	0.028	0.050	0.078	0.068	0.075	0.055
Type 1 DM early deaths	1.25	2.91	4.84	12.64	7.71	12.03	6.92
Type 2 DM early deaths	2.03	3.70	11.37	60.97	18.07	40.38	27.44
BMI kg/M^2	21.89	23.47	23.29	24.82	22.98	21.97	25.04
Fasting plasma glucose mmol/L	4.77	4.46	4.01	4.53	4.31	4.12	4.26
Systolic BP mm Hg	137.56	135.11	125.19	130.31	133.66	131.15	126.85
Socio-demographic index	0.828	0.822	0.590	0.608	0.644	0.431	0.575
Likely mechanism of low CVD (see discussion for explanation)	FSVV, Plants	FSVV	Smoking, metabolic	FSVV, Plants	Physical activity, smoking	Physical activity, smoking	FSVV, smoking

[a] Japan not only had high FSVV but also had relatively high intake of GBD risk factor plant
foods (fruits, vegetables, nuts and seeds, whole grains, and legumes: 350.9 KC/d).

[b] France was characteristic of 23 other low CVD countries in that intake of milk products was
high (France milk=95.75 KC/d and in the other 23 countries' milk>68 KC/d, Table 3 compared
with worldwide mean milk=25.04 KC/d).

[c] The low CVD of Peru might be attributed to a relatively low smoking rate (7.5%) and good
metabolic statistics (mean BMI=23.3 Kg/M^2, mean FPG=4.01 mmol/L, mean LDLc=2.37
mmol/L, mean SBP=125.2 mm/Hg).

[d] While Mexico had a lower FSVV than most low CVD countries (Mexico FSVV=413.3 KC/d),
like Japan it had relatively high intake of important plant foods (fruits, vegetables, nuts and
seeds, whole grains, and legumes: totaling 351.7 KC/d).

[e] Panama, with moderate FSVV=383.8 KC/d, had relatively high physical activity (Panama
physical activity=4935 METs) and low smoking rate (male/female mean smoking=7.3%).

[f] While Guatemala had relatively low FSVV (FSVV=231.1 KC/d) and low GBD risk factor plant
food intake (fruits, vegetables, nuts and seeds, whole grains, and legumes: 249.5 KC/d), it had
relatively high physical activity (physical activity=6142 METs) and a low smoking rate
(male/female mean smoking=7.7%).

[g] Ecuador had an intermediatory FSVV (FSVV=375.4 KC/d) and relatively low intake of
important plant foods (fruits, vegetables, nuts and seeds, whole grains, and legumes: totaling
205.3 KC/d). However, Ecuador had a low rate of smoking (male/female mean smoking=8.0%).
and a low SBP (mean SBP=126.9 mm/Hg).

Table 5. CVD and risk factors for countries with FSVV ≥567.27 KC/d (n=974 cohorts)

CVD and CVD risk factors	Mean	SD	Min	Max	r	95% CI low	95% CI high	P
Cardiovascular deaths/100k/year ages 15-69	273.72	143.95	73.47	913.18				
LDL cholesterol mmol/L	2.91	0.14	2.36	3.20	-0.254	-0.312	-0.194	<.0001
Fat-soluble vitamin variable	692.58	111.88	527.67	932.18	0.523	0.476	0.567	<.0001
Processed meat KC/d	25.01	15.30	2.16	68.77	0.459	0.408	0.507	<.0001
Red meat KC/d	122.16	41.41	43.65	235.95	0.655	0.618	0.690	<.0001
Fish KC/d	49.08	89.24	7.60	370.36	-0.259	-0.317	-0.199	<.0001
Milk KC/d	78.58	26.14	14.60	146.82	0.185	0.123	0.245	<.0001
Poultry KC/d available	129.97	60.21	41.65	411.87	0.366	0.311	0.419	<.0001
Eggs KC/d available	40.80	7.25	14.59	69.64	-0.242	-0.301	-0.182	<.0001
Saturated fatty acids KC/d	299.88	66.83	129.39	481.07	0.193	0.132	0.253	<.0001
PUFAs KC/d	211.86	83.06	80.53	381.31	0.382	0.327	0.435	<.0001
Trans fatty acids KC/d	25.17	17.82	4.23	66.89	0.224	0.163	0.283	<.0001
Alcohol KC/d	155.02	65.23	11.92	429.81	0.475	0.425	0.522	<.0001
Sugary beverages KC/d	231.66	121.92	72.91	769.89	0.332	0.274	0.386	<.0001
Potatoes KC/d available	106.39	40.47	9.50	224.86	0.109	0.046	0.170	0.00
Corn KC/d available	19.45	9.41	1.79	47.39	0.086	0.024	0.148	0.01
Fruits KC/d	66.91	15.93	34.83	161.39	-0.014	-0.077	0.049	0.66
Vegetables KC/d	117.42	34.24	42.27	304.17	-0.127	-0.188	-0.064	<.0001
Nuts and seeds KC/d	21.98	8.91	0.85	102.99	0.177	0.115	0.237	<.0001
Whole grains KC/d	52.11	20.93	1.61	92.45	0.121	0.059	0.183	0.00
Legumes KC/d	40.76	25.29	2.95	133.26	0.040	-0.023	0.102	0.21
Rice KC/d available	38.82	46.75	6.50	179.63	-0.138	-0.199	-0.076	<.0001
Sweet potatoes KC/d available	3.63	3.99	0.04	17.63	-0.128	-0.189	-0.066	<.0001
Total KC/d available	3219	292	2516	3898	0.249	0.189	0.307	<.0001
Vit A deficiency children/100k/yr	9316	6877	1400	28081	-0.105	-0.166	-0.042	0.00
Sodium g/d	3.98	0.96	2.22	6.70	-0.040	-0.102	0.023	0.21
Calcium g/d	0.645	0.129	0.353	1.044	0.396	0.342	0.448	<.0001
Dietary fiber g/d	10.62	1.75	6.30	18.15	0.241	0.181	0.299	<.0001
Physical activity METs	3523	765	1781	5494	0.578	0.535	0.618	<.0001
Child underweight >2SD	0.015	0.013	0.004	0.058	-0.197	-0.257	-0.136	<.0001
Stop breast feeding <6 months	0.196	0.017	0.129	0.219	0.268	0.208	0.325	<.0001
Ambient pollution PM $_{0.25}$	12.67	7.13	4.38	87.22	0.005	-0.058	0.068	0.87
Smoking rate (0-1)	0.233	0.084	0.021	0.444	0.233	0.172	0.291	<.0001
Secondhand smoking (0-1)	0.309	0.070	0.201	0.586	-0.306	-0.362	-0.248	<.0001
Sublingual tobacco rate (0-1)	0.013	0.020	0.001	0.125	0.579	0.536	0.620	<.0001
Blood lead level mcg/dl	3.93	0.77	1.22	5.72	0.451	0.400	0.500	<.0001
Household air pollution (0-1)	0.012	0.026	0.000	0.201	-0.011	-0.073	0.052	0.74
Kidney disease stage III (0-1)	0.036	0.012	0.016	0.109	-0.028	-0.090	0.035	0.39
Type 1 DM early deaths	3.93	2.67	0.55	15.97	0.573	0.529	0.613	<.0001
Type 2 DM early deaths	9.54	7.52	0.75	54.22	0.622	0.582	0.659	<.0001
BMI kg/M^2	24.87	1.78	21.40	29.39	0.484	0.434	0.530	<.0001
Fasting plasma glucose mmol/L	4.73	0.23	3.84	5.58	0.091	0.028	0.153	0.00
Systolic BP mm Hg	133.639	4.362	126.863	142.147	-0.148	-0.209	-0.086	<.0001
Socio-demographic index (0-1)	0.838	0.053	0.592	0.896	-0.052	-0.114	0.011	0.11
Sex male 1 and female 2	1.50	0.50	1.00	2.00	-0.745	-0.772	-0.716	<.0001

Data Availability: The formatted database on which this analysis is based is posted on the

Mendeley data repository (https://data.mendeley.com/datasets/g6b39zxck4/3):

Cundiff, David and Wu, Chunyi (2021), "IHME GBD CVD risk factors", Mendeley Data, V3,

doi: 10.17632/g6b39zxck4.3. Data and SAS codes are available to check the accuracy of the

tables in this analysis: https://data.mendeley.com/datasets/g6b39zxck4/3. SAS Statisticians who

check the analysis and report their findings as comments on this preprint, will be eligible to

become paper contributors by contacting the lead author: davidkcundiff@gmail.com).

487•   **Funding:** This research received no grant from any funding agency in the public, commercial or

488   not-for-profit sectors. The Bill and Melinda Gates Foundation funded the acquisition of the data

489   for this analysis by the IHME. The data were provided to the authors as volunteer collaborators

490   with IHME.

491•

492   **Competing interests:** The authors declare no competing interests.

493•   **Competing interests' statement:** Both authors have completed the ICMJE uniform disclosure

494   form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the

495   submitted work; no financial relationships with any organisations that might have an interest in

496   the submitted work in the previous three years; no other relationships or activities that could

497   appear to have influenced the submitted work.

498•

499•   **Contributors**

500•   DKC acts as guarantor; conceived and designed the study, acquired and analysed the GBD data

form IHME, interpreted the study findings, drafted the manuscript, critically reviewed and edited

the manuscript and tables, and approved the final version for publication.

CW designed software programs in R to format and population weighted the data,

aided with the SAS statistical analysis, critically reviewed the manuscript, and

approved the final version for publication.

The lead author (DKC) affirms the manuscript is an honest, accurate, and transparent account of

the study being reported; that no important aspects of the study have been omitted; and that any

discrepancies from the study as planned (and, if relevant, registered) have been explained.

Author access to data: As volunteer collaborators with the Institute of Health Metrics and

Evaluation, we received about 1.4 gigabytes of raw data on noncommunicable disease deaths and

43 relevant risk factors.

Acknowledgments

We thank Scott Glenn and Brent Bell from IHME who supplied us with the GBD risk factor

exposure data.

Ethical approval: Studies based solely on data from IHME GBD database do not need approval

from any bioethics committee or other agency.

Checklist: This report followed the STrengthening the Reporting of OBservational studies in

Epidemiology (STROBE) guidelines for reporting global health estimates.[48]

References

1. Teicholz N. The scientific report guiding the US dietary guidelines: is it scientific? BMJ. 2015;351:h4962. doi: 10.1136/bmj.h4962.
2. Teicholz N. The scientific report guiding the US dietary guidelines: is it scientific? BMJ. 2015;351. doi: 10.1136/bmj.h4962.
3. Ioannidis JA. The challenge of reforming nutritional epidemiologic research. JAMA. 2018;320(10):969-70. doi: 10.1001/jama.2018.11025.
4. Zile MH. Vitamin A-not for your eyes only: requirement for heart formation begins early in embryogenesis. Nutrients. 2010;2(5):532-50. PubMed PMID: 22254040.
5. Xing X, Tao F. [Advance of study on vitamin A deficiency and excess associatied with congenital heart disease]. Wei Sheng Yan Jiu. 2008;37(6):754-6. PubMed PMID: 19239019.
6. Saremi A, Arora R. Vitamin E and cardiovascular disease. Am J Ther. 2010;17(3):e56-65. PubMed PMID: 19451807.
7. Beulens JWJ, Booth SL, van den Heuvel EGHM, Stoecklin E, Baka A, Vermeer C. The role of menaquinones (vitamin K2) in human health. British Journal of Nutrition. 2013;110:1357–68.
8. Wang L, Song Y, Manson JE, Pilz S, März W, Michaëlsson K, et al. Circulating 25-Hydroxy-Vitamin D and Risk of Cardiovascular Disease: A Meta-Analysis of Prospective Studies. Circulation: Cardiovascular Quality and Outcomes. 2012;5(6):819-29. doi: 10.1161/circoutcomes.112.967604.
9. Fu X, Harshman SG, Shen X, Haytowitz DB, Karl JP, Wolfe BE, et al. Multiple Vitamin K Forms Exist in Dairy Foods. Curr Dev Nutr. 2017;1(6):e000638-e. doi: 10.3945/cdn.117.000638. PubMed PMID: 29955705.
10. Walther B, Chollet M. Vitamin K2 - Vital for Health and Wellbeing: IntechOpen; 2017. Available from: https://www.intechopen.com/books/vitamin-k2-vital-for-health-and-wellbeing/menaquinones-bacteria-and-foods-vitamin-k2-in-the-diet.
11. Kamao M, Suhara Y, Tsugawa N, Uwano M, Yamaguchi N, Uenishi K, et al. Vitamin K content of foods and dietary vitamin K intake in Japanese young women. J Nutr Sci Vitaminol (Tokyo). 2007;53(6):464-70. doi: 10.3177/jnsv.53.464. PMID: 18202532.
12. Kaneki M HS, Hosoi T, Fujiwara S, Lyons A, Crean SJ, Ishida N, Nakagawa M, Takechi M, Sano Y, Mizuno Y, Hoshino S, Miyao M, Inoue S, Horiki K, Shiraki M, Ouchi Y, Orimo H. Japanese fermented soybean food as the major determinant of the large geographic difference in circulating levels of vitamin K2: possible implications for hip-fracture risk. Nutrition. 2001;17(4):315-21. doi: 10.1016/s0899-9007(00)00554-2.
13. Tamang JP, Cotter PD, Endo A, Han NS, Kort R, Liu SQ, et al. Fermented foods in a global age: East meets West. Comprehensive Reviews in Food Science and Food Safety. 2020;19(1):184-217. doi: https://doi.org/10.1111/1541-4337.12520.
14. Joseph M. Fermented Soy Products: A Guide To 12 Traditional Foods: Nutrition Advance; 2020 [April 11, 2021].
15. Global Burden of Disease Study 2017 (GBD 2017) Data Input Sources Tool Seattle, Washington: Institute of Health Metrics and Evaluation; [updated 2019August 10, 2019]. Available from: http://ghdx.healthdata.org/gbd-2017/data-input-sources.
16. PROTOCOL FOR THE GLOBAL BURDEN OF DISEASES, INJURIES, AND RISK FACTORS STUDY (GBD) Version 3.0; Issue 26 Seattle, Washington: Institute for Health

Metrics and Evaluation; [updated February 2018August 2, 2019]. Available from:
http://www.healthdata.org/sites/default/files/files/Projects/GBD/GBD_Protocol.pdf.
17.   Naghavi M, Abajobir AA, Abbafati C, Abbas KM, Abd-Allah F, Abera SF, et al. Global,
regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a
systematic analysis for the Global Burden of Disease Study 2016. The Lancet. 2017;390
(10100):1151-210.
18.   Flaxman AD, Lee YY, Vos T, Blore J, Degenhardt L, Peterson HM, et al. An Integrative
Metaregression Framework for Descriptive Epidemiology. Seattle, WA: University of
Washington Press; 2015. Available from: http://www.guilles.website/wp-
content/uploads/2017/11/Metaregression-AFlaxman.pdf.
19.   GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk
assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of
risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of
Disease Study 2017. . The Lancet. 2018;392(10159):1923-94.
20.   GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk
assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of
risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of
Disease Study 2017: Table of risk factor definitions. The Lancet. 2018;392(10159):1923-94.
21.   Call for Collaborators. Seattle, Washington: Institute for Health Metrics and Evaluation
at the University of Washington; 2019 [updated 2019April 25, 2019]. Available from:
http://www.healthdata.org/gbd/call-for-collaborators.
22.   Omega 3 Fatty Acids: Fact Sheet for Health Professionals. In: US Department of Health
and Human Services, editor.: The Office of Dietary Supplements; 2018.
23.   Nutritionix Track App: Syndigo LLC [updated 2019April 25, 2019]. Available from:
https://www.nutritionix.com/.
24.   Ritchie H, Roser M. Diet Compositions. Our World in Data. 2017.
25.   Pu J, Romanelli R, Zhao B, Azar KMJ, Hastings KG, Nimbal V, et al. Dyslipidemia in
special ethnic populations. Cardiol Clin. 2015;33(2):325-33. doi: 10.1016/j.ccl.2015.01.005.
PubMed PMID: 25939303.
26.   Frank ATH, Zhao B, Jose PO, Azar KMJ, Fortmann SP, Palaniappan LP. Racial/ethnic
differences in dyslipidemia patterns. Circulation. 2014;129(5):570-9. Epub 11/05. doi:
10.1161/CIRCULATIONAHA.113.005757. PubMed PMID: 24192801.
27.   Keys A, Menotti A, Karvonen MJ, Aravanis C, Blackburn H, Buzina R, et al. The diet
and 15-year death rate in the seven countries study. Am J Epidemiol. 1986;124(6):903-15.
PubMed PMID: 3776973.
28.   Mahmood SS, Levy D, Vasan RS, Wang TJ. The Framingham Heart Study and the
epidemiology of cardiovascular disease: a historical perspective. Lancet (London, England).
2014;383(9921):999-1008. Epub 09/29. doi: 10.1016/S0140-6736(13)61752-3. PubMed PMID:
24084292.
29.   Dehghan M, Mente A, Zhang X, Swaminathan S, Li W, Mohan V, et al. Associations of
fats and carbohydrate intake with cardiovascular disease and mortality in 18 countries from five
continents (PURE): a prospective cohort study. The Lancet. 2017;390(10107):2050-62. doi:
10.1016/S0140-6736(17)32252-3.
30.   Countries Who Eat The Most Cheese: World Atlas; 2021 [updated 2021March 28, 2021].
Available from: https://www.worldatlas.com/articles/countries-who-consume-the-most-

cheese.html#:~:text=The%20top%20cheese%20consumer%20is,kilograms%20of%20cheese%2
0per%20capita.
31.	Cheese Consumption Per Capita in New Zealand Czech Republic: Helgi Library, Source:
Faostat; [updated Oct. 18, 2018April 9, 2021]. Available from:
https://www.helgilibrary.com/indicators/cheese-consumption-per-capita/new-zealand/.
32.	Khalil Z, Alam B, Akbari AR, Sharma H. The Medical Benefits of Vitamin K(2) on
Calcium-Related Disorders. Nutrients. 2021;13(2):691. doi: 10.3390/nu13020691. PubMed
PMID: 33670005.
33.	POTATO PROCESSING AND USES Lima, Peru: International Potato Center; [updated
2020May 28, 2020]. Available from: https://cipotato.org/potato/potato-processing-uses/.
34.	Juul F, Vaidean G, Lin Y, Deierlein Andrea L, Parekh N. Ultra-Processed Foods and
Incident Cardiovascular Disease in the Framingham Offspring Study. Journal of the American
College of Cardiology. 2021;77(12):1520-31. doi: 10.1016/j.jacc.2021.01.047.
35.	Srour B, Fezeu LK, Kesse-Guyot E, Allès B, Méjean C, Andrianasolo RM, et al. Ultra-
processed food intake and risk of cardiovascular disease: prospective cohort study (NutriNet-
Santé). BMJ. 2019;365:l1451. doi: 10.1136/bmj.l1451.
36.	Monteiro CA, Moubarac JC, Cannon G, Ng SW, Popkin B. Ultra-processed products are
becoming dominant in the global food system. Obes Rev. 2013;14 Suppl 2:21-8. PubMed PMID:
24102801.
37.	Mozaffarian D, Fahimi S, Singh GM, Micha R, Khatibzadeh S, Engell RE, et al. Global
Sodium Consumption and Death from Cardiovascular Causes. New England Journal of
Medicine. 2014;371(7):624-34. doi: 10.1056/NEJMoa1304127. PubMed PMID: 25119608.
38.	O'Donnell M, Mente A, Yusuf S. Sodium Intake and Cardiovascular Health. Circ Res.
2015;116(6):1046-57. doi: doi:10.1161/CIRCRESAHA.116.303771.
39.	Roseboom T, van der Meulen JHP, Osmond C, Barker D, Ravelli A, Schroeder-Tanka J,
et al. Coronary heart disease after prenatal exposure to the Dutch famine, 1944-45. Heart.
2000;84(6):595-8. doi: DOI:10.1136/heart.84.6.595. PubMed PMID: PMC1729504.
40.	Dieterich CM, Felice JP, O'Sullivan E, Rasmussen KM. Breastfeeding and health
outcomes for the mother-infant dyad. Pediatric clinics of North America. 2013;60(1):31-48.
Epub 11/03. doi: 10.1016/j.pcl.2012.09.010. PubMed PMID: 23178059.
41.	Ambient (outdoor) air pollution Geneva, Switzerland: World Health Organization; 2018
[March 24, 2021]. Available from: https://www.who.int/news-room/fact-sheets/detail/ambient-
(outdoor)-air-quality-and-health.
42.	Tobacco responsible for 20% of deaths from coronary heart disease Geneva, Switzerland:
World Health Organization; 2020 [March 2, 2021]. Available from:
https://www.who.int/news/item/22-09-2020-tobacco-responsible-for-20-of-deaths-from-
coronary-heart-disease.
43.	Vaziri ND. Mechanisms of lead-induced hypertension and cardiovascular disease.
American Journal of Physiology-Heart and Circulatory Physiology. 2008;295(2):H454-H65. doi:
10.1152/ajpheart.00158.2008. PubMed PMID: 18567711.
44.	Lanphear BP, Rauch S, Auinger P, Allen RW, Hornung RW. Low-level lead exposure
and mortality in US adults: a population-based cohort study. The Lancet Public Health.
2018;3(4):e177-e84. doi: 10.1016/S2468-2667(18)30025-2.
45.	Bikbov B, Purcell CA, Levey AS, Smith M, Abdoli A, Abebe M, et al. Global, regional,
and national burden of chronic kidney disease, 1990-2017: a systematic analysis for the Global

Burden of Disease Study 2017. The Lancet. 2020;395(10225):709-33. doi: 10.1016/S0140-
6736(20)30045-3.
46.   Global, regional, and national age-sex-specific mortality for 282 causes of death in 195
countries and territories, 1980-2017: a systematic analysis for the Global Burden of Disease
Study 2017. Lancet. 2018;392(10159):1736-88. PubMed PMID: 30496103.
47.   Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, Salama JS, et al. Health effects of
dietary risks in 195 countries, 1990-2017: a systematic analysis for the Global Burden of Disease
Study 2017. The Lancet. 2019;393(10184):1958-72. doi: 10.1016/S0140-6736(19)30041-8.
48.   Stevens GA, Alkema L, Black PRE, Boerma JT, Collins GS, Ezzati PM, et al. Guidelines
for Accurate and Transparent Health Estimates Reporting: the GATHER statement. The Lancet.
2016;388(10062):e19-e23.
671