1   **Health Belief, Planned Behavior, or Psychological Antecedents: What predicts**
2   **COVID-19 Vaccine Hesitancy better among the Bangladeshi Adults?**
3

4   Mohammad Bellal Hossain[1]*, Md. Zakiul Alam[1], Md. Syful Islam[2], Shafayat Sultan[1],
5   Md. Mahir Faysal[1], Sharmin Rima[3], Md. Anwer Hossain[1], Abdullah Al Mamun[1]

ABSTRACT

Background This study aimed to determine the prevalence and investigate the constellations of psychological determinants of the COVID-19 vaccine hesitancy among the Bangladeshi adult population utilizing the health belief model-HBM (perceived susceptibility to and severity of COVID-19, perceived benefits of and barriers to COVID-19 vaccination, and cues to action), the theory of planned behavior-TPB (attitude toward COVId-19 vaccine, subjective norm, perceived behavioral control, and anticipated regret), and the novel 5C psychological antecedents (confidence, constraints, complacency, calculation, and collective responsibility). We compared the predictability of these theoretical frameworks to see which framework explains the highest variance in COVID-19 vaccine hesitancy.

Methods This study adopted a cross-sectional research design. We collected data from a nationally representative sample of 1497 respondents through both online and face-to-face interviews. We employed multiple linear regression analysis to assess the predictability of each model of COVID-19 vaccine hesitancy.

Results We found a 41.1% prevalence of COVID-19 vaccine hesitancy among our study respondents. After controlling the effects of socio-economic, demographic, and other COVID-19 related covariates, we found that the TPB has the highest predictive power (adjusted R^2=0.43), followed by the 5C psychological antecedents of vaccination (adjusted R^2=0.32) and the HBM (adjusted R^2=0.31) in terms of explaining total variance in the COVID-19 vaccine hesitancy among the adults of Bangladesh.

Conclusions This study provides evidence that theoretical frameworks like the HBM, the TPB, and the 5C psychological antecedents can be used to explore the psychological determinants of vaccine hesitancy, where the TBP has the highest predictability. Our findings can be used to design targeted interventions to reduce vaccine hesitancy and increase vaccine uptake.

35
36
37

[1] Department of Population Sciences, University of Dhaka, Dhaka, Bangladesh
[2] Department of Population Science, Jatiya Kabi Kazi Nazrul Islam University, Mymensingh
[3] Ovibashi Karmi Unnayan Program (OKUP), Dhaka
* Correspondence: Professor Mohammad Bellal Hossain, PhD, E-mail: bellal@du.ac.bd

Key Questions

What is already known?

- There is a global-level insurgence of COVID-19 vaccine hesitancy, where the majority of studies come from western, educated, industrialized, rich, and democratic (WEIRD) countries.
- To date, an online survey found that the prevalence of COVID-19 vaccine hesitancy in Bangladesh was 32.5%.
- Few studies from WEIRD countries have adopted the Health Believe Model and/or the Theory of Planned Behavior to explore the predictors of COVID-19 vaccine hesitancy.

What are the new findings?

- This study found a 41.1% prevalence of COVID-19 vaccine hesitancy among a nationally representative sample of Bangladesh.
- After controlling the effects of demographic, socio-economic, and other COVID-19 related covariates, we found that the TPB has the highest predictive power, followed by the 5C psychological antecedents and the HBM in terms of explaining total variance in the COVID-19 vaccine hesitancy among the adults of Bangladesh.

What do the new findings imply?

- Theoretical frameworks like the HBM, the TPB, and the 5C psychological antecedents can be used to explore the multitude of the psychological determinants of vaccine hesitancy, where the TPB has the highest predictability.
- Findings can be used to design targeted interventions to reduce COVID-19 vaccine hesitancy and increase vaccine demand and uptake.

INTRODUCTION

Though vaccines, in the form of successful mass immunization programs, have saved millions of lives and improved health and wellbeing across the world [1], historically, such successes have constantly been challenged by a minority, yet a significant proportion of 'vaccine-hesitant individuals and groups for a variety of environmental, cultural, political and psychological reasons [2–5]. The World Health Organization has identified vaccine hesitancy as one of the top ten global health threats [6], where vaccine hesitancy has been defined as a delay in acceptance or refusal of vaccination despite its availability [5].

The majority of the studies on vaccine hesitancy comes from the western, educated, industrialized, rich, and democratic (WEIRD) countries [7–9]. Thus, very little is known about the contexts of developing, low- and middle-income countries [10]. The world has once again witnessed an insurgence of vaccine hesitancy globally during the pandemic of Coronavirus Disease 2019 (COVID-19)[11]. A systematic review found that many studies reported a COVID-19 vaccine acceptance rate below 60%, ranging from the lowest 23.6% in Kuwait to the highest 97% in Ecuador [9].

Studies recommend that 60-75% immunity at the population level would be necessary to
halt the Coronavirus transmission and community spread [12,13]. Thus, the Government of
Bangladesh aims to achieve an 80% coverage of the COVID-19 vaccination program [14].
However, there is an apparent dearth of research on COVID-19 vaccination behavior in
Bangladesh. To date, an online survey found that the prevalence of vaccine hesitancy was
32.5% among the adult population of Bangladesh aged 18 years and above, though this
study lacks generalizability and was conducted before the vaccine was publicly available [15].
Research on vaccine hesitancy has received less attention in Bangladesh until the recent
pandemic of COVID-19 due to the success stories of expanded programs on immunization
[16,17].

Along with quantifying the prevalence of vaccine hesitancy among the population, it is
crucial to understand the determinants of the individual decision-making process that result
in delay or omission of vaccination [18,19]. Because vaccine hesitancy reduces vaccine demand
and the uptake [5], which can impact vaccination coverage, that may ultimately hinder the
successful control of the COVID-19 pandemic in Bangladesh. This study, therefore, aimed to
determine the prevalence of COVID-19 vaccine hesitancy and investigate the constellations
of psychological determinants of vaccine hesitancy among the adult population of
Bangladesh, utilizing three of the most widely used theoretical frameworks and models-, the
Health Belief Model (HBM) [20,21], the Theory of Planned Behavior (TPB) [22,23] and the novel 5C
psychological antecedents of vaccination [10,24].

Vaccine hesitancy: theories and models to understand vaccination behavior
Vaccine hesitancy is a complex, emerging term in socio-medical literature that recognizes a
continuum between total acceptance and outright refusal of some or all vaccines [5,19]. Hence,
vaccine-hesitant individuals comprise a heterogeneous group in the middle of a continuum
ranging from the total acceptors of vaccine to the complete refusers [4].

The complex nature of motives behind vaccine hesitancy has primarily been explored and
analyzed using the epidemiological triad of *environment* (contextual factors), *agent* (vaccine
and disease), and *host* factors (individual characteristics) [1,19]. Building on this triad, the WHO
Strategic Advisory Group of Experts (SAGE) on immunization drafted a "Model of
determinants of vaccine hesitancy" around three key domains: 1) Contextual influences –
comprising historical, socio-cultural, environmental, health system/institutional, economic
or political factors; 2) Individual and group influences – including influences arising from the
personal perception of and attitude towards the vaccine or influences of the social/peer
environment; and 3) Vaccine and vaccination-specific issues those are associated with the
vaccine characteristics or the vaccination process [4,5].

However, in this study, our focus was explicitly directed towards the psychological
determinants of vaccine hesitancy [18,19], as vaccine hesitancy is predominantly the outcome
of the individual decision-making process, which is influenced by individual's feeling about
the vaccination or a particular vaccine, barriers, and enablers to vaccinate [10,18,19]. Thus, it is
crucial to understand which psychological determinants or drivers determine to delay or
refusal of the vaccination [18,19] so that targeted interventions can be designed to reduce
vaccine hesitancy and increase vaccine demand [5,25]. Therefore, to fulfill that intent, we have
adopted two of the most widely used social cognition models- the HBM [26] and the TPB [27],

and the latest framework of 5C psychological antecedents of vaccination[10] to explore and
predict the COVID-19 vaccine hesitancy among adults of Bangladesh.
The HBM is one of the most widely used models in vaccination behavior, particularly in
influenza [28], swine flu [22], ebola [28], and hepatitis B [29]. Few studies have investigated COVID-19
vaccination behavior utilizing the HBM [20,21,30]. The key argument of HBM is that the
likelihood of an individual adopting a particular health behavior (e.g., getting COVID-19
vaccine) is determined by the perceived susceptibility and severity of illness or disease (e.g.,
COVID-19), along with the belief in the effectiveness of the recommended health behavior
(e.g., COVID-19 vaccination) [26]. The model is comprised of, as applied to COVID-19 and its
vaccine, *perceived severity* of and *perceived susceptibility* to COVID-19, *perceived benefits* of
and *perceived barriers* to getting a COVID-19 vaccine, and *cues to action* which include
implicit or explicit incentives or situations that serve to motivate vaccination, such as,
information from mass media [30].
138
In contrast, the TPB argues that behavior is driven by the intention to carry out the
behavior, which ultimately is determined by an individual's 'belief structure' [27]. As applied to
the context of COVID-19 vaccine, belief structure is comprised of *attitude towards COVID-19*
vaccine (i.e., its perceived necessity, benefit, and effectiveness), *subjective norms* (i.e.,
whether significant others support getting a COVID-19 vaccine), and *perceived behavioral*
control (i.e., to what extent COVID-19 vaccination is perceived within the individual's
control) [31]. However, new components are continually being added to the TPB framework
to increase its usefulness [22]. For example, Gallagher and Povey (2006) found that the
addition of 'anticipated regret' substantially increased the predictive value of TPB in terms
of older adults' intention to get a seasonal influenza vaccination.
149
150 A few studies have combined the HBM and the TPB to identify health-related behaviors and
151 explore the multitude of the psychological determinants that influence individual decision-
152 making, such as receiving influenza vaccine among the general public [22,32] or in the context
153 of COVID-19 vaccination [30,31]. Gerund & Shepherd (2012) found that the TPB explained more
154 variance and produced a better model fit than the HBM [33].
155
Finally, Betsch and colleagues (2018) have incorporated and expanded existing vaccination
behavior measures and proposed a novel framework of 5C psychological antecedents of
vaccination. It includes *confidence* (trust in vaccine effectiveness, safety, and necessity, and
the system that delivers it), *complacency* (perceiving the disease as low risk), *constraints*
(perceived low vaccine availability, affordability, accessibility, and other barriers to
vaccinating), *calculation* (analyzing pros and cons of vaccination), and *collective*
responsibility (willingness to take the vaccine for protecting others via herd immunity).
Compared with other existing models, the 5C psychological antecedents have explained a
greater extent of vaccination variance. However, it has not been tested alongside the HBM
and the TPB [10].
166

METHODS

Study Design and Data Collection
The study adopted a cross-sectional research design. The calculated sample size was 1635,
where the Z-score for 95% confidence interval was 1.96, prevalence (p) of COVID-19 vaccine
hesitancy from a previous study was 0.33[15], the margin of error (e) was 0.03, design effect

(Deff) for sampling variation was 1.6, and a non-response rate (NR) was 10%. However, 112
respondents did not consent to participate in the survey, while another 26 respondents did
not know about the COVID-19 vaccine. After excluding them, the final sample stood 1497
for the analysis.

We collected data from all eight administrative divisions of Bangladesh using probability-
proportionate to each division's population size. Both online and face-to-face interviews
were conducted to collect data. Data were collected using the online platform Google form
from one-third of the respondents. A link to the survey questionnaire was created and sent
to the prospective respondents via e-mail, WhatsApp, and Facebook messenger. All the
respondents to whom the survey link was sent were requested to share the link in their
network to reach more people. The research team members circulated the survey link in
their respective professional and social networks. The online link was valid for three days.
After that, the collected data were downloaded, and divisional distribution was assessed.
We then collected data for the remaining sample size of each division using face-to-face
interviews. For this, we randomly selected two districts from each division. Within each
district, the sample was distributed proportionately according to the rural-urban distribution
of its population. Then, convenience sampling was adopted to accomplish face-to-face
interviews from the population-based households.

The respondent selection criteria for the face-to-face interview were adult people of
18 years and above living in Bangladesh and knowing about the COVID-19 vaccine. Reading
and writing ability and using the Internet were added criteria for selecting the online survey
respondents. The data is now available in the Mendeley open research data repository [34].

Measures

Outcome Variable: COVID-19 Vaccine Hesitancy
We used two 6-points Likert-type items to measure COVID-19 vaccine hesitancy among the
respondents: (a*) If you get the chance of getting a COVID 19 vaccine for free, what will you
do?* (with the response of 1= *Surely, I will take it*; 2= *Probably I will take it*; 3= *I will delay
taking it*; 4= *I am not sure what I will do*; 5= *Probably I will not take it*; 6= *Surely, I will not
take it*), and (b) *If your family or friends think of getting COVID 19 vaccine, what will you do?*
(with the response of 1= *Strongly encourage them*; 2=*Encourage them*; 3=*Ask them to delay
getting the vaccine*; 4=I *will not say anything about it*; 5=*Discourage them to take vaccine*;
6=*Forbid them to take vaccine*). We combined these two items and calculated the level of
COVID-19 vaccine hesitancy, where a higher score indicated a higher level of hesitancy
toward the COVID-19 vaccine. The theoretical total score ranges from 2 to 12. The reliability
analysis of the scale demonstrates an excellent internal consistency (Cronbach
Alpha=0.833).

Predictor Variables
The HBM constructs
The HBM constructs consisted of the following components: perceived susceptibility
(included two items, α=0.657), perceived severity (included two items, α=0.612), perceived
benefits (included three items, α=0.841), perceived barriers (included five items, α=0.735),
and cues to action (Table 1). These items were measured on a 5-point Likert scale (1=

Strongly Disagree to 5= Strong Agree). Cues to action included self-reported COVID-19
positive status for self and family members, and sources of vaccine-related knowledge
(social media/ online news portals/blog, and print media).
The TPB constructs
The TPB constructs consisted of four domains: attitude toward vaccine (including six items,
α=0.739), subjective norm, perceived behavioral control, and anticipated regret (Table 1).
Each item of the four domains was assessed on a 5-point scale (1 = *strongly disagree* to 5 =
strongly agree). The mean score of the items under attitude toward vaccine component was
calculated, with a higher average score indicating more negative attitude toward COVID-19
vaccine.
228
The 5C psychological antecedents of vaccination
The 5C psychological antecedents of vaccination consisted of 5 antecedents: confidence
(included three items, α=0.808), complacency (included three items, α=0.753), constraints
(included a single item), the calculation (included three items, α=0.909), and collective
responsibility (included two items, α=0.612) (Table 1). Each item of the five antecedent
domains was assessed on a 5-point scale (1 = *strongly disagree*; 5 = *strongly agree*). Except
for the constraints domain, mean scores of items under each domain were computed, with
a higher average score indicating the corresponding domain's stronger agreement.
237
238 **Table 1: Items used to measure HBM, TPB, and 5C psychological antecedents**

Statements	Cronbach α
The Health Belief Model [26]	
Perceived Susceptibility	**0.657**
I am worried about the likelihood of getting infected by COVID-19	
I am at high risk of COVID-19 because of my health conditions	
Perceived Severity	**0.612**
I will be very sick if I get infected by COVID-19	
I am very concerned that I could die from COVID-19	
Perceived Benefits	**0.841**
I think vaccination is good because it will make me less worried about COVID-19	
I believe vaccination will decrease my risk of getting infected by COVID-19	
I think the complications of COVID-19 will decrease if I get vaccinated and then get infected with the Coronavirus.	
Perceived Barriers	**0.700**
I am worried that the possible side effects of the COVID-19 vaccination would interfere with my usual activities	
I am concerned about the efficacy of the COVID-19 vaccine	
I have a concern that I may receive faulty/fake COVID-19 vaccine	
It concerns me that the development of a COVID-19 vaccine is too rushed to test its safety properly	
I am concerned about the long-term side effects of the COVID-19 vaccination	
Cues to Action	**NA**
Respondents got infected with Coronavirus	
A family member got Infected with Coronavirus	
Social media (e.g., Facebook) or online news portals/blog as a source of knowledge about the COVID-19 vaccine	
Printed newspaper as a source of knowledge about COVID-19 vaccine	
The Theory of Planned Behavior [27]	
Attitude toward Vaccine	0.739
I think the COVID-19 vaccine probably will not work	
I do not trust the COVID-19 vaccine	
I think the COVID-19 vaccine is unnecessary	

Statements	Cronbach α
I think it is not important to get a vaccine to protect people from the COVID-19	
I do not need a COVID-19 vaccine because I am healthy and at low risk for infection	
I do not need a COVID-19 vaccine because even if I get infected, I will not become seriously ill	
Subjective Norm	NA
I believe my family members will support me to get vaccinated against COVID-19	
Perceived Behavioral Control	NA
If I want, I can register for COVID 19 vaccination	
Anticipated Regret	NA
If I do not get a COVID-19 vaccine and end up getting Coronavirus, I will regret not getting the vaccination	
The 5C Psychological Antecedents of Vaccination [10]	
Confidence	0.808
I am completely confident that COVID 19 vaccines are safe	
I am completely confident that COVID 19 vaccines are effective	
Regarding COVID 19 vaccines, I am confident that public authorities decide in the best interest of the community	
Constraints	NA
Everyday work stress may prevent me from getting vaccinated	
Complacency	0.753
I think it is unnecessary to receive vaccinations as it cannot prevent COVID-19	
I believe my immune system is powerful; it will protect me from COVID-19	
I believe COVID-19 is not much a severe disease that I should get vaccinated against it	
Calculation	0.909
When I think about getting vaccinated against COVID 19, I weigh the benefits and risks to make the best decision possible	
When I think about getting vaccinated against COVID 19, I will first consider whether it is effective or not	
Before I get COVID-19 vaccinated, I need to know about this vaccine in details	
Collective Responsibility	0.612
I will take COVID 19 vaccine because, in that way, I can protect people with a weaker immune system	
I think vaccination against COVID 19 is a collective action to prevent the spread of diseases	

Other Covariates

We also collected data on the following independent variables: age, sex, religion, marital
status, educational attainment, place of residence, geographic region, occupation, number
of family members, household income, knowledge about COVID-19 vaccine, knowledge
about vaccination process, and behavioral practice to prevent COVID-19 (Table 2).

Knowledge about COVID-19 vaccine: We assessed the knowledge related to the COVID-19
vaccine using four Likert-type items: *COVID-19 vaccine is safe for pregnant women, COVID-*
19 vaccine is safe for children under 18 years old, COVID-19 vaccine have a very mild side
effect, and *side effects COVID-19 vaccine do not last longer than 2 days.* The response
options against these items were "strongly disagree = 1", "disagree = 2", "neither agree nor
disagree = 3", "agree = 4", and "strongly agree = 5". The total score of these items ranged
between 4 to 20, with a higher score indicating higher knowledge. Reliability analysis of the
scale suggested an acceptable internal consistency (Cronbach alpha, α=0.643).

Knowledge about vaccination process: Knowledge about the COVID-19 vaccination process
was measured by the following six binary response (yes=1, no=0) questions: (1) do you know
that you need to consult with a doctor to receive the COVID-19 vaccination?; (2) do you

know that you need to register online to receive the COVID-19 vaccination?; (3) do you
know that you can receive the COVID-19 vaccination only from selected health facilities?; (4)
do you know that you cannot receive the COVID-19 vaccination from a pharmacy?; (5) do
you know that health care workers will not provide the COVID-19 vaccine at your doorstep?;
and (6) do you know the correct doses of the COVID 19 vaccine? The reliability analysis
showed good internal consistency among these six questions (α=0.765). The total score
ranged between 0 to 6, with a higher score indicating better knowledge about the COVID-19
vaccination process.

Behavioral practice to prevent COVID-19: The level of preventive behavioral practices related
to COVID-19 was measured using three Likert-type items: *I avoid crowds as much as possible*
to prevent my risk of getting COVID-19, I always wear a mask when I am outside of my home
or around other people, and *I am conscious about using sanitizer or hand wash*. The
response options against these items were "never = 1", "sometimes = 2", "often = 3", and
"regularly = 4". The total score ranged between 3 to 12, with a higher score indicating a
higher level of preventive practices. The reliability analysis showed good internal
consistency in this scale (α=0.857).

Statistical Analysis
We employed multiple linear regression analysis to assess the selected model's
predictability to COVID-19 vaccine hesitancy after checking the assumptions and multi-
collinearity. We produced three models. In each model, significant demographic, socio-
economic, knowledge, and practice of COVID-19 related variables were controlled to predict
the used model. Akaike Information Criterion (AIC), Amemiya Prediction Criterion (APC),
Mallows's Prediction Criterion (MPC), and Schwarz Bayesian Criterion (SBC) were used to
assess model performance.

Patient and Public Involvement
No public or patients were involved in the process of this research.

Human Subject's Protections
The present study was carried out following the Declaration of Helsinki. The respondents
were informed about the aims, objectives, potential scopes, and implications of this study's
findings and were requested to participate voluntarily.

RESULTS
Background Characteristics of the Respondents
The demographic, socio-economic, and other background characteristics of the respondents
are summarized in Table 2. The respondents' average age was 33.7 years, with a standard
deviation (SD) of 12.9. The highest proportion (28.9%) of respondents was from 18–24
years. About 47% of the respondents were women, while most of the respondents (86.9%)
were Muslim. About two-thirds of the respondents (61.6%) were married, while 20.6% of
the respondents had less than a secondary education level. About two-thirds of the
respondents (64.3%) were from rural areas, while 31.9% were from Dhaka Division. One-
third (31.6%) of the respondents were students and unemployed. The mean number of
household members was 4.9, while the mean household income was BDT 37627 (Table 2).

Table 2 Sample characteristics of the respondents

Variables	Estimates, n (%)
Age in years	
18-24	432 (28.9)
25-30	362 (24.2)
31-39	254 (17.0)
40-49	236 (15.8)
50+	213 (14.2)
Average Age, Mean (SD)	33.7 (12.9)
Sex	
Women	692 (46.2)
Men	805 (53.8)
Religion	
Others	196 (13.1)
Muslim	1301 (86.9)
Marital status	
Never-married	575 (38.4)
Ever-married	922 (61.6)
Education	
No education	129 (8.6)
Primary	179 (12.0)
Secondary and higher secondary	448 (29.9)
Graduate	400 (26.7)
Masters and MPhil/PhD	341 (22.8)
Place of residence	
Rural area	963 (64.3)
Urban area (other than city corporation)	179 (12.0)
City Corporation	355 (23.7)
Administrative division	
Barishal	114 (7.6)
Chattogram	253 (16.9)
Dhaka	478 (31.9)
Khulna	137 (9.2)
Mymensingh	108 (7.2)
Rajshahi	180 (12.0)
Rangpur	114 (7.6)
Sylhet	113 (7.5)
Occupation	
Government, private, & NGO sector job	202 (13.5)
Professional (teacher, engineer, lawyer, doctor, nurse, paramedics, pharmacist)	277 (18.5)
Housewife	348 (23.2)
Students and unemployed	473 (31.6)
Agriculture and day Laborer	102 (6.8)
Others	95 (6.3)
Household members, Mean (SD)	4.9 (1.98)
Household income in BDT, Mean (SD)	37627.2 (81295.9)
Knowledge about COVID-19 vaccine, Mean (SD)	11.4 (2.2)
Knowledge about vaccination process, Mean (SD)	2.8 (2.0)
Behavioral practice to prevent COVID-19, Mean (SD)	8.8 (2.7)
Total	**1497 (100.0)**

Prevalence of COVID-19 Vaccine Hesitancy

Nearly 43% of respondents reported that they would surely take the COVID-19 vaccine if it
were available for free, whereas another 17.7% of respondents would probably take it

(Figure 1). Thus, a total of 60.6% of the respondents were potential COVID-19 *vaccine*
acceptors. However, 39.4% of respondents were found vaccine-hesitant, where nearly 7%
were complete refusers of the COVID-19 vaccine (Figure 1). A similar level of COVID-19
vaccine hesitancy was found among the respondents when asked about their opinion about
their family members or relatives' hypothetical vaccination decisions. A total of 60.3% of the
respondents mentioned that they would strongly encourage or encourage their family
members or relatives if they decided to take the COVID-19 vaccine (Figure 1). Thus, rest
39.7% of respondents were vaccine-hesitant in terms of their family members and relatives'
decision to vaccinate, where 3% would forbid them to take the COVID-19 vaccine (Figure 1).
The mean score of the hesitancy scale was 4.93 with an SD of 2.68. The accuracy of the
hesitancy scale was 41.1% (4.93/12*100), indicating that 41.1% of the respondents had
hesitancy to accept the COVID-19 vaccine.

Figure 1 Prevalence (%) of COVID-19 vaccine hesitancy in Bangladesh
324

Predictors of COVID-19 Vaccine Hesitancy in Bangladesh

After controlling the effects of demographic, socio-economic, and other COVID-19 related
covariates, our first model shows that the HBM constructs explained 31% of the variance in
COVID-19 vaccine hesitancy (adjusted R^2=0.31) (Table 3). Among the HBM constructs,
perceived susceptibility, perceived severity, perceived benefits, and perceived barriers were
the significant predictors of COVID-19 vaccine hesitancy. An increase in perceived
susceptibility of COVID-19 tended to reduce vaccine hesitancy (β=-0.06, p<0.05). Similarly,
increased perceived severity of COVID-19 also reduced vaccine hesitancy (β=-0.11, p<0.01).
However, among the HBM constructs, perceived benefits had the largest standardized co-
efficient. In other words, a one-unit increase in perceived benefits of getting the COVID-19
vaccine reduced the vaccine hesitancy by 0.33 units (β=-0.33, p<0.01). On the other hand,
perceived barriers to getting the COVID-19 vaccine tended to increase vaccine hesitancy (β=
0.30, p<0.01). However, the cues to action construct were found insignificant in predicting
COVID-19 vaccine hesitancy, except the component 'Social media (e.g., Facebook) or online

news portals/blog as a source of knowledge about the COVID-19 vaccine'. In other words,
respondents who heard about the COVID-19 vaccine from social media (e.g., Facebook) or
online news portals were less vaccine-hesitant (β= -0.05, p<0.05).
343

344 **Table 3 Multiple linear regression analysis to predict COVID-19 vaccine hesitancy**

Variables	Adjusted R^2	Beta (SE)
Model 1: Constant+ Health Belief Model constructs (n=1497)	0.31	
Perceived susceptibility		-0.06 (0.04)**
Perceived severity		-0.11 (0.04)***
Perceived benefits		-0.33 (0.03)***
Perceived barriers		0.30 (0.02)***
Cues to action		
Respondents got infected with Coronavirus		
Yes		-0.03 (0.27)
No (RC)		
A family member got Infected with Coronavirus		
No		0.04 (0.21)*
Yes (RC)		
Social media (e.g., Facebook) or online news portals/blog as a source of knowledge about the COVID-19 vaccine		
Yes		-0.05 (0.14)**
No (RC)		
Printed newspaper as a source of knowledge about the COVID-19 vaccine		
Yes		0.01 (0.15)
No (RC)		
Model 2: Constant+ Theory of Planned Behavior constructs (n=1497)	0.43	
Attitude toward COVID-19 vaccine		0.27 (0.02)***
Subjective norm		-0.31 (0.07)***
Perceived behavioral control		-0.05 (0.05)**
Anticipated regret		-0.18 (0.05)***
Model 3: Constant+ 5C Model constructs (n=1497)	0.32	
Confidence		-0.38 (0.03)***
Constraints		0.03 (0.06)
Complacency		0.18 (0.03)***
Calculation		0.09 (0.03)***
Collective responsibility		-0.06 (0.05)**

345 Note: the effects of the following variables were controlled for all models: sex of the respondents,
346 religion, place of residence, administrative division, knowledge about COVID-19 vaccine, knowledge
347 about vaccination process, and behavioral practice to prevent COVID-19; * indicates p-value < 0.10,
348 ** indicates p-value < 0.05, *** indicates p-value < 0.01; Reference category = RC
349

The second model shows that 43% of the variance in COVID-19 vaccine hesitancy was
explained by the TPB (adjusted R^2=0.43) (Table 3). All four components of the TPB were the
significant predictors of COVID-19 vaccine hesitancy. Respondents who had a more negative
attitude toward the COVID-19 vaccine were more vaccine-hesitant (β=0.27, p<0.01). Vaccine
hesitancy tended to decrease with the increase of familial support in favor of vaccination
regarding the TPB's subjective norm (β=-0.31, p<0.01). However, in terms of perceived

behavioral control, respondents who mentioned registering for COVID-19 vaccination was within their control were less vaccine-hesitant (β=-0.05, p<0.05). Finally, an increase in anticipated regret among the respondents resulted in reduced vaccine hesitancy (β=-0.18, p<0.01).

The third model included the 5C psychological antecedents, which explained 32% of the variance in COVID-19 vaccine hesitancy (adjusted R^2=0.32) (Table 3). An increase in COVID-19 vaccine confidence tended to decrease vaccine hesitancy (β=-0.38, p<0.05). The more complacent individuals had a more vaccine hesitancy (β=0.18, p<0.01). Similarly, respondents who were more calculative about the pros and cons of getting vaccinated or needed more information about the vaccine before getting vaccinated were significantly more COVID-19 vaccine-hesitant (β=0.09, p<0.01). Finally, respondents who had a sense of collective responsibility to vaccinate against COVID-19 were significantly less vaccine-hesitant (β=-0.06, p<0.05).

Table 4 presents the model summary and the selection criteria to assess model performance. For each of the three multiple linear regression models to identify psychological determinants of COVID-19 vaccine hesitancy, we compared AIC, APC, MPC, SBC to select the best fitting model. The model with the smallest AIC, APC, MPC, SBC values was the best-fitted model. According to these indicators, the TPB constructs (Model 2) were the best-fitted model (Table 4).

Table 4 Model Summary and Selection Criteria

Model Summary	Model 1: HBM	Model 2: TPB	Model 3: 5c
n	1497	1497	1497
R	0.56	0.66	0.57
R^2	0.32	0.43	0.33
Adjusted R^2	0.31	0.43	0.32
F Change	108.2	205.09	92.69
df	4, 1474	4, 1478	5, 1477
Significance of F Change	<.001	<.001	<.001
Selection Criteria			
Akaike Information Criterion (AIC)	2427.1	2144.6	2399.0
Amemiya Prediction Criterion (APC)	0.70	.58	0.69
Mallows' Prediction Criterion (MPC)	23.0	19.0	20.0
Schwarz Bayesian Criterion (SBC)	2549.2	2245.5	2505.23

DISCUSSION

Given the dearth of research on COVID-19 vaccination behavior in Bangladesh, this study aimed to determine the prevalence of COVID-19 vaccine hesitancy and explore its psychological determinants among Bangladeshi adults. To fulfill that intent, we have utilized three of the most widely used tools in the field of vaccination behavior- the HBM, the TPB, and the novel 5C psychological antecedents of vaccination, and compared their predictability in terms of predicting the COVID-19 vaccine hesitancy.

We found a 41.1% prevalence of the COVID-19 vaccine hesitancy among our study respondents, a significantly higher estimate than found by Ali and colleagues (2021) (32.5%). This higher estimation may, in part, be explained by the fact that Ali and colleagues (2021)

conducted an online survey which may biasedly present the view of 'Netigens' of
Bangladesh [35]. In contrast, in our study, data were collected through online and face-to-face
interviews from a nationally representative sample covering all eight administrative
divisions. In that sense, our study findings provide a more accurate estimate of COVID-19
vaccine hesitancy that is generalizable in the context of the adult population living in
Bangladesh.
397
However, as per the second objective of this study, we found that after controlling the
effects of socio-economic and demographic variables, level of knowledge related to COVID-
19, its vaccine and vaccination process, and level of preventive practices towards COVID-19,
the TPB has the highest predictive power (adjusted R^2=0.43), followed by the 5C model
(adjusted R^2=0.32) and the HBM (adjusted R^2=0.31) in terms of explaining total variance in
COVID-19 vaccine hesitancy among the adults of Bangladesh. This finding is particularly
unique because, to date, no study compared the predictability of these three behavioral
frameworks or models in terms of predicting vaccine hesitancy, especially in Bangladesh.
Moreover, given the newness of the 5C scale [10], the available literature is still scanty that
tests this model's predictive validity [24,36]. However, various other studies that adopted only
the HBM and the TPB validate our findings that TPB constructs are a better predictor of
vaccination behavior [23], specifically in the intention to vaccinate against swine flu [22] human
papillomavirus [33] or in the context of COVID-19 vaccination [20,30,31].
411
According to the findings of the HBM constructs, an increase in perceived benefits of the
COVID-19 vaccine, along with increasing perceived severity of and perceived susceptibility to
COVID-19, significantly reduced the vaccine hesitancy. On the other hand, an increase in
perceived barriers to getting vaccinated acted as a significant vaccine hesitancy promoter.
These findings essentially correspond to other studies related to influenza vaccination [37],
though Lin et al. (2020) found perceived susceptibility was not a significant predictor of the
COVID-19 vaccine hesitancy in China. In terms of cues to action, though COVID-19 infection
status (of both self or family members) were non-significant predictors at a 5 percent level
of significance. We found that social media (e.g., Facebook) or online news portals as the
source of information about the COVID-19 vaccine was the significant predictor, and
respondents who heard about the COVID-19 vaccine from social media (e.g., Facebook) or
online news portals or blogs were less hesitant. Taken together, these findings suggest that
imparting adequate and proper information about the COVID-19 vaccine to the public, along
with solid evidence of the safety, efficacy, and benefits of the COVID-19 vaccine, can be a
crucial strategy to reduce vaccine hesitancy and increase its demand and actual uptake. In
that case, social media and online news portals may act as more effective means than
printed newspapers to disseminate COVID-19 related information, as found in our study.
429
According to the findings of the TPB constructs, an increase of negative attitude towards the
COVID-19 vaccine and a decrease in perceived behavioral control significantly increased the
COVID-19 vaccine hesitancy. Other studies support these findings [23,30], though perceived
behavioral control was found non-significant predictor in other contexts [22,38,39]. However,
subjective norms in the form of family members' support for having COVID-19 vaccination
significantly reduced vaccine hesitancy, corresponding to other studies [23,30]. Consistent with
earlier researches [22,40,41], anticipated regret was a significant predictor of COVID-19 vaccine
hesitancy. This finding implies that an intervention to increase the COVID-19 vaccination

uptake should circulate the message that it is better to get vaccinated than regret later.
Alternative measures should be devised to reduce the barriers related to COVID-19
vaccination, such as online registration to get the COVID-19 vaccine. This provision is
inconvenient, especially for older persons, people living in rural areas, and those who do not
have internet access [42].
443

Finally, according to the 5C model, more substantial confidence and higher collective
responsibility were significantly associated with reduced COVID-19 vaccine hesitancy,
whereas increased complacency and calculation significantly increased the vaccine
hesitancy. These findings are supported by other studies in both COVID-19 and other
contexts [24,36], though calculation and constraints were non-significant predictors of COVID-
19 vaccine hesitancy among the nurses in Hong Kong [24]. In our study, the constraint related
to the COVID-19 vaccination was a non-significant predictor of COVID-19 vaccine hesitancy.
These findings suggest that public confidence in the vaccine and the health system that
delivers the vaccination service are crucial. Widespread misinformation, conspiracy beliefs,
and superstitions regarding the COVID-19 vaccine and its potential health hazards have
been found to diminish public trust [43] that need to be addressed through proper
communication. Extensive information searching about the subjective utility of vaccination,
as evidenced in earlier studies [18,19,44], might have resulted in more vaccine hesitancy as more
calculation increased COVID-19 vaccine hesitancy among the respondents, along with
complacency. These findings warrant the urgency of re-iterating the risk communication and
health benefits message of getting vaccinated to mass people.
460

Strengths and Limitations of the Study

This study is the first to explore the COVID-19 vaccine hesitancy among Bangladeshi adults,
adopting a large and diversified representative sample. Moreover, this study also explored
many psychological antecedents of the COVID-19 vaccine hesitancy utilizing three of the
most widely used theoretical tools- the HBM, the TPB, and the 5C psychological
antecedents, and used multivariate modeling to identify the most salient predictors.
Therefore, this study's findings can help design targeted intervention to reduce the COVID-
19 vaccine hesitancy, which will help the Government of Bangladesh attain the target of
80% vaccination coverage for the COVID-19 vaccine. Another strength of the study is that
our study provides the findings on the prevalence of the COVID-19 vaccine hesitancy and its
predictors while the COVID-19 vaccine was publicly available in Bangladesh. Finally, in terms
of theoretical contribution of the study in the field of vaccination behavior, this study
contributes evidence from a non-WEIRD country that, at one hand, assess the predictive
validity of the novel 5C psychological antecedents of vaccination [10] and, on the other hand,
validates the theoretical supremacy of the TPB over the HBM and the 5C model in predicting
the COVID-19 vaccine hesitancy among the adult population in Bangladesh. However, our
study also has some limitations. This study could not use probability sampling completely.
We tried to draw our sample following the national population distribution in terms of age,
sex, residence, region, marital status, and religion. However, the distribution of education
among the respondents is not comparable to national data. Moreover, this study collected
self-reported data that may suffer from reporting bias. Finally, this research used a cross-
sectional study design which can not establish causality.

CONCLUSION

This study provides evidence that theoretical frameworks like the TPB, the HBM, and the 5C psychological antecedents can explore the psychological determinants that influence a person's vaccination decision-making process. Among the frameworks of determinants, the TPB has the highest predictive power in determining the vaccination decision. These findings can be used to craft targeted interventions to reduce vaccine hesitancy and increase vaccine uptake. Thus, this study's findings will steer Bangladesh's vaccination campaign and those alike to reach the targeted coverage of the COVID-19 vaccination program and, thereby, paving the way for successfully eradicating the never-ending pandemic of COVID-19.

Acknowledgments The authors would like to thank the respondents of this study for their valuable time and inputs. We would also like to convey our thanks to selected students of the Department of Population Sciences, the University of Dhaka, for their contribution in conducting field-level data collection amidst this challenging time of the COVID-19 pandemic.

Contributors MBH conceptualized the study. All the authors designed the study and collected the data. MBH, MAH, and MZA analyzed and interpreted the data and drafted the manuscript. MSI, SS, MMF, SR, and AAM revised the manuscript critically for valuable intellectual content and approved the final version to be published. All authors remained in agreement to be accountable for all aspects of the work.

Funding This study received no funding or whatsoever from any organization or donor agency.

Competing interests The authors declared no conflicts of interest.

Ethical Approval The study was approved by the Bangladesh Medical Research Council (BMRC).

Data availability statement Data associated with this study has been deposited at Mendeley at http://dx.doi.org/10.17632/jzvbvvknkv.1

REFERENCES

1.	WHO. Report of the SAGE working group on vaccine hesitancy. Published online 2014:1-64. https://www.who.int/immunization/sage/meetings/2014/october/1_Report_WORKING_GROUP_vaccine_hesitancy_final.pdf

2.	Porter D, Porter R. The politics of prevention: Anti-vaccinationism and public health in nineteenth-century England. *Med Hist*. 1988;32(3):231-252. doi:10.1017/S0025727300048225

3.	Wolfe RM, Sharp LK. Anti-vaccinationists past and present. *Br Med J*. 2002;325(7361):430-432. doi:10.1136/bmj.325.7361.430

4.   Larson HJ, Jarrett C, Eckersberger E, Smith DMD, Paterson P. Understanding vaccine
hesitancy around vaccines and vaccination from a global perspective: A systematic
review of published literature, 2007-2012. *Vaccine*. 2014;32(19):2150-2159.
doi:10.1016/j.vaccine.2014.01.081

5.   MacDonald NE, Eskola J, Liang X, et al. Vaccine hesitancy: Definition, scope and
determinants. *Vaccine*. 2015;33(34):4161-4164. doi:10.1016/j.vaccine.2015.04.036

6.   Scheres J, Kuszewski K. The ten threats to global health in 2018 and 2019. A welcome
and informative communication of WHO to everybody. *Zdr Publiczne i Zarządzanie*.
2019;17(1):2-8. doi:10.4467/20842627oz.19.001.11297

7.   Lazarus J V., Ratzan SC, Palayew A, et al. A global survey of potential acceptance of a
COVID-19 vaccine. *Nat Med*. 2021;27(2):225-228. doi:10.1038/s41591-020-1124-9

8.   Henrich J, Heine SJ, Norenzayan A. The weirdest people in the world? *Behav Brain Sci*.
2010;33(2-3):61-83. doi:10.1017/S0140525X0999152X

9.   Sallam M. Covid-19 vaccine hesitancy worldwide: A concise systematic review of
vaccine acceptance rates. *Vaccines*. 2021;9(2):1-15. doi:10.3390/vaccines9020160

10. Betsch C, Schmid P, Heinemeier D, Korn L, Holtmann C, Böhm R. Beyond confidence:
Development of a measure assessing the 5C psychological antecedents of vaccination.
PLoS One. 2018;13(12):e0208601. doi:10.1371/journal.pone.0208601

11. Harrison EA, Wu JW. Vaccine confidence in the time of COVID-19. *Eur J Epidemiol*.
2020;35(4):325-330. doi:10.1007/s10654-020-00634-3

12. Anderson RM, Vegvari C, Truscott J, Collyer BS. Challenges in creating herd immunity
to SARS-CoV-2 infection by mass vaccination. *Lancet*. 2020;396(10263):1614-1616.
doi:10.1016/S0140-6736(20)32318-7

13. Billah MA, Miah MM, Khan MN. Reproductive number of coronavirus: A systematic
review and meta-analysis based on global level evidence. *PLoS One*.
2020;15(11):e0242128. doi:10.1371/journal.pone.0242128

14. Tajmim T. Coronavirus Vaccination: Bangladesh stands out in South Asia. The Business
Standard. Published 2021. Accessed April 13, 2021.
https://www.tbsnews.net/coronavirus-chronicle/covid-19-bangladesh/vaccination-
bangladesh-stands-out-south-asia-205324

15. Ali M, Hossain A. What is the extent of COVID-19 vaccine hesitancy in Bangladesh?:
A cross-sectional rapid national survey. *medRxiv*. Published online 2021.
doi:10.1101/2021.02.17.21251917

16. Jamil K, Bhuiya A, Streatfield K, Chakrabarty N. The immunization programme in
Bangladesh: Impressive gains in coverage, but gaps remain. *Health Policy Plan*.
1999;14(1):49-58. doi:10.1093/heapol/14.1.49

17. Sarkar PK, Sarker NK, Doulah S, Bari TIA. Expanded programme on immunization in
Bangladesh: A success story. *Bangladesh J Child Heal*. 2017;39(2):93-98.
doi:10.3329/bjch.v39i2.31540

18. Betsch C, Böhm R, Chapman GB. Using behavioral insights to increase vaccination
policy effectiveness. *Policy Insights from Behav Brain Sci*. 2015;2(1):61-73.
doi:10.1177/2372732215600716

19. Kumar D, Chandra R, Mathur M, Samdariya S, Kapoor N. Vaccine hesitancy:
Understanding better to address better. *Isr J Health Policy Res*. 2016;5(1):1-8.
doi:10.1186/s13584-016-0062-y

20. Wong LP, Alias H, Wong PF, Lee HY, AbuBakar S. The use of the health belief model to
assess predictors of intent to receive the COVID-19 vaccine and willingness to pay.
Hum   Vaccines   Immunother.   2020;16(9):2204-2214.
doi:10.1080/21645515.2020.1790279

21. Lin Y, Hu Z, Zhao Q, Alias H, Danaee M, Wong LP. Understanding COVID-19 vaccine
demand and hesitancy: A nationwide online survey in China. *PLoS Negl Trop Dis*.
2020;14(12):e0008961. doi:10.1371/journal.pntd.0008961

22. Myers LB, Goodwin R. Determinants of adults' intention to vaccinate against
pandemic swine flu. *BMC Public Health*. 2011;11:15. doi:10.1186/1471-2458-11-15

23. Xiao X, Wong RM. Vaccine hesitancy and perceived behavioral control: A meta-
analysis. *Vaccine*. 2020;38(33):5131-5138. doi:10.1016/j.vaccine.2020.04.076

24. Kwok KO, Li KK, WEI WI, Tang A, Wong SYS, Lee SS. Influenza vaccine uptake, COVID-
19 vaccination intention and vaccine hesitancy among nurses: A survey. *Int J Nurs
Stud*. 2021;114:103854. doi:10.1016/j.ijnurstu.2020.103854

25. de Figueiredo A, Simas C, Karafillakis E, Paterson P, Larson HJ. Mapping global trends
in vaccine confidence and investigating barriers to vaccine uptake: A large-scale
retrospective   temporal   modelling   study.   *Lancet*.   2020;396(10255):898-908.
doi:10.1016/S0140-6736(20)31558-0

26. Rosenstock IM. The health belief model and preventive health behavior. *Heal Educ
Behav*. 1977;2(4):354-386. doi:10.1177/109019817400200405

27. Ajzen I. The theory of planned behavior. *Organ Behav Hum Decis Process*.
1991;50(2):179-211. doi:10.1016/0749-5978(91)90020-T

28. Bish A, Yardley L, Nicoll A, Michie S. Factors associated with uptake of vaccination
against pandemic influenza: A systematic review. *Vaccine*. 2011;29(38):6472-6484.
doi:10.1016/j.vaccine.2011.06.107

29. Huynh G, Pham L, Tran T, Cao N, Nguyen T, Bui Q. How knowledge of hepatitis B
disease and vaccine influences vaccination practices among parents in Ho Chi Minh
City, Vietnam. *Asian Pac J Trop Med*. 2021;14(3):122-127. doi:10.4103/1995-
7645.307534

30. Shmueli L. Predicting intention to receive COVID-19 vaccine among the general
population using the Health belief model and the theory of planned behavior model.
medRxiv. Published online 2020. doi:10.1101/2020.12.20.20248587

31. Guidry JPD, Laestadius LI, Vraga EK, et al. Willingness to get the COVID-19 vaccine
with and without emergency use authorization. *Am J Infect Control*. 2021;49(2):137-

142. doi:10.1016/j.ajic.2020.11.018

32.   Yang ZJ. Predicting young adults intentions to get the H1N1 vaccine: An integrated
model. *J Health Commun*. 2015;20(1):69-79. doi:10.1080/10810730.2014.904023

33.   Gerend MA, Shepherd JE. Predicting human papillomavirus vaccine uptake in young
adult women: Comparing the health belief model and theory of planned behavior.
Ann Behav Med. 2012;44(2):171-180. doi:10.1007/s12160-012-9366-5

34.   Hossain MB, Alam MZ, Islam MS, et al. Data on Health Belief, Planned Behavior, or
Psychological Antecedents of COVID-19 vaccine hesitancy among the Bangladeshi
adults. *Mendeley Data*. 2021;V1. doi:10.17632/jzvbvvknkv.1

35.   Paul A, Sikdar D, Hossain MM, et al. Knowledge, attitudes, and practices toward the
novel coronavirus among Bangladeshis: Implications for mitigation measures. *PLoS*
One. 2020;15(9):e0238492. doi:https://doi.org/10.1371/journal.pone.0238492

36.   Neufeind J, Betsch C, Habersaat KB, Eckardt M, Schmid P, Wichmann O. Barriers and
drivers to adult vaccination among family physicians – Insights for tailoring the
immunization program in Germany. *Vaccine*. 2020;38(27):4252-4262.
doi:10.1016/j.vaccine.2020.04.052

37.   Schmid P, Rauber D, Betsch C, Lidolt G, Denker ML. Barriers of influenza vaccination
intention and behavior - A systematic review of influenza vaccine hesitancy, 2005-
2016. *PLoS One*. 2017;12(1):e0170550. doi:10.1371/journal.pone.0170550

38.   Trafimow D, Sheeran P, Conner M, Finlay KA. Evidence that perceived behavioural
control is a multidimensional construct: Perceived control and perceived difficulty. *Br*
J Soc Psychol. 2002;41(1):101-121. doi:10.1348/014466602165081

39.   Povey R, Conner M, Sparks P, James R, Shepherd R. Application of the Theory of
Planned Behaviour to two dietary behaviours: Roles of perceived control and self-
efficacy. *Br J Health Psychol*. 2000;5(2):121-139. doi:10.1348/135910700168810

40.   Sandberg T, Conner M. A mere measurement effect for anticipated regret: Impacts on
cervical screening attendance. *Br J Soc Psychol*. 2009;48(2):221-236.
doi:10.1348/014466608X347001

41.   Gallagher S, Povey R. Determinants of older adults' intentions to vaccinate against
influenza: A theoretical application. *J Public Health (Bangkok)*. 2006;28(2):139-144.
doi:10.1093/pubmed/fdl008

42.   Ahamad MG, Islam AKMN, Talukder B, Ahmed MU. COVID-19 vaccination in
Bangladesh: Challenges on price, misinformation, and trust. *SocArXiv*. Published
online 2021. doi:10.31219/osf.io/kn6ts

43.   COVID: Bangladesh vaccination drive marred by misinformation. Deutsche Welle.
Published January 27, 2021. Accessed April 13, 2021. https://www.dw.com/en/covid-
bangladesh-vaccination-drive-marred-by-misinformation/a-
56360529?fbclid=IwAR0ujYXHNJ8rZjn5yARGlZSnGe0uDRZAbsmmfb0lrRouVtl1qhidYV
hF3XE

44.   Leask J. Target the fence-sitters. *Nature*. 2011;473(7348):443-445.

648