Longitudinal analysis shows durable and broad immune memory after SARS-

CoV-2 infection with persisting antibody responses and memory B and T cells

Kristen W. Cohen[1,*], Susanne L. Linderman[2,*], Zoe Moodie[1], Julie Czartoski[1], Lilin

Lai[4,10,11], Grace Mantus[4,10], Carson Norwood[4,10], Lindsay E. Nyhoff[4,11], Venkata

Viswanadh Edara[4,10,11], Katharine Floyd[4,10,11], Stephen C. De Rosa[1,7], Hasan Ahmed[8],

Rachael Whaley[1], Shivan N. Patel[3], Brittany Prigmore[1], Maria P. Lemos[1], Carl W.

Davis[2], Sarah Furth[1], James O'Keefe[6], Mohini P. Gharpure[2], Sivaram Gunisetty[2], Kathy

A. Stephens[5], Rustom Antia[8], Veronika I. Zarnitsyna[2,9], David S. Stephens[6], Srilatha

Edupuganti[3], Nadine Rouphael[3], Evan J. Anderson[4-6], Aneesh K. Mehta[6], Jens

Wrammert[4,10**], Mehul S. Suthar[4,10,11**], Rafi Ahmed[2,**], and M. Juliana McElrath[1,7,**]

*Co-first authors and **Co-senior authors

Address correspondence to jmcelrat@fredhutch.org or rahmed@emory.edu

[1]Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Research Center,

Seattle, WA, 98109, USA

[2]Emory Vaccine Center and Department of Microbiology and Immunology, Emory

University, Atlanta, GA 30322, USA

[3]Department of Medicine, Division of Infectious Diseases, Hope Clinic of Emory Vaccine

Center, Emory University School of Medicine, Atlanta, GA 30329, USA

[4]Center for Childhood Infections and Vaccines of Children's Healthcare of Atlanta,

Emory University Department of Pediatrics Department of Medicine, Atlanta, GA 30322,

USA

[6]Emory University School of Medicine, Department of Medicine, Atlanta, GA 30322,

USA

[7]Departments of Laboratory Medicine and Medicine, University of Washington, Seattle,

WA 98195, USA

[8]Department of Biology, Emory University, Atlanta, GA 30322, USA

[9]Department of Microbiology and Immunology, Emory University, Atlanta, GA 30322,

USA

[11]Yerkes National Primate Research Center, Atlanta, GA 30329, USA

Funding Acknowledgments:

The research reported in this publication was supported in part by COVID supplements

from the National Institute of Allergy and Infectious Diseases and the Office of the

Director of the National Institutes of Health under award numbers UM1AI068618-14S1

and UM1AI069481-14S1 (MJM); UM1A057266-S1, U19AI057266-17S1 and

U19AI090023 (R. Ahmed); ORIP/OD P51OD011132 (MSS); and T32AI074492 (LEN).

This work was also supported by grants from the Oliver S. and Jennie R. Donaldson

Charitable Trust (R. Ahmed); Paul G. Allen Family Foundation Award #12931 (MJM);

Seattle COVID-19 Cohort Study (Fred Hutchinson Cancer Research Center, MJM); the

Joel D. Meyers Endowed Chair (MJM); An Emory EVPHA Synergy Fund award (MSS

and JW); COVID-Catalyst-I[3] Funds from the Woodruff Health Sciences Center (MSS);

the Center for Childhood Infections and Vaccines (MSS and JW); Children's Healthcare

of Atlanta (MSS and JW), a Woodruff Health Sciences Center 2020 COVID-19 CURE

SUMMARY

Ending the COVID-19 pandemic will require long-lived immunity to SARS-CoV-2. We evaluated 254 COVID-19 patients longitudinally from early infection and for eight months thereafter and found a predominant broad-based immune memory response. SARS-CoV-2 spike binding and neutralizing antibodies exhibited a bi-phasic decay with an extended half-life of >200 days suggesting the generation of longer-lived plasma cells. In addition, there was a sustained IgG+ memory B cell response, which bodes well for a rapid antibody response upon virus re-exposure. Polyfunctional virus-specific CD4+ and CD8+ T cells were also generated and maintained with an estimated half-life of 200 days. Interestingly, the CD4+ T cell response equally targeted several SARS-CoV-2 proteins, whereas the CD8+ T cell response preferentially targeted the nucleoprotein, highlighting the importance of including the nucleoprotein as a potential vaccine antigen. Taken together, these results suggest that broad and effective immunity may persist long-term in recovered COVID-19 patients.

Keywords:

SARS-CoV-2, COVID-19, Immune memory, Antibody, Kinetics, Neutralization, CD4+ T cells, CD8+ T cells, B cells, Spike, RBD, Endemic coronaviruses

INTRODUCTION

The COVID-19 pandemic caused by the rapid spread of SARS-CoV-2, a novel

betacoronavirus, continues to cause significant morbidity and mortality. The induction of

effective early immune control of SARS-CoV-2 and durable immune memory is critical

to prevent severe disease and to protect upon re-exposure. SARS-CoV-2 infection

induces polyclonal humoral and cellular responses targeting multiple viral proteins

described in cross-sectional and longitudinal studies (Sette and Crotty, 2021). More

comprehensive, quantitative analyses with extensive serial sampling in larger numbers

of COVID-19 patients are limited, and could resolve some conflicting views about the

durability of humoral immunity. Importantly, defining the frequency, immune function

and specificity of the antibodies, memory B and T cell responses among COVID-19

patients, and identifying when they appear and how long they persist can provide

understanding of the integral components for long-lived immunity to SARS-CoV-2 and

potentially other human coronaviruses that emerge in the future (Stephens and

McElrath, 2020).

We initiated two prospective COVID-19 patient cohorts in Seattle and Atlanta during

the first surge of the pandemic to investigate long-term immunity to SARS-CoV-2.

Among 254 COVID-19 patients enrolled and frequently sampled, we identify binding

and neutralizing antibodies to SARS-CoV-2 as well as antigen-specific B and T cells

elicited early after infection, define their specificities, quantify the extent of antibody

boosting of cross-reactive responses to other coronaviruses, and further characterize

the decay rate and durability of these immune parameters over 250 days. We employ

highly standardized or validated assays that are also being used to evaluate immunity in

recent and ongoing clinical vaccine trials (Anderson et al., 2020; Doria-Rose et al.,

2021; Sadoff et al., 2021). This indepth longitudinal study demonstrates that durable

immune memory persists in most COVID-19 patients, including those with mild disease,

and serves as a framework to define and predict long-lived immunity to SARS-CoV-2

after natural infection. This investigation will also serve as a benchmark for immune

memory induced in humans by SARS-CoV-2 vaccines.

RESULTS

COVID-19 study population

COVID-19-confirmed patients were recruited into our longitudinal study of SARS-

CoV-2 specific B and T cell memory after infection. A total of 254 patients enrolled at

two sites, Atlanta and Seattle, starting in April 2020 and returned for follow up visits over

a period of 250 days. We were able to collect blood samples at 2-3 time points from 165

patients and at 4-7 time points from another 80 patients, which allowed us to perform a

longitudinal analysis of SARS-CoV-2-specific B and T cell responses on a large number

of infected patients. The demographics and baseline characteristics of this cohort are

described in Table S1. The study group was 55% female and 45% male and between

18-82 years old (median, 48.5 years). Based on World Health Organization (WHO)

guidelines of disease severity, 71% of study participants exhibited mild disease, 24%

had moderate disease, and 5% experienced severe disease.

Antibody responses to SARS-CoV-2 spike protein show a bi-phasic decay with an extended half-life

Binding antibodies to the SARS-CoV-2 full length spike protein, to the receptor binding domain (RBD) and to the N terminal domain (NTD) of the spike protein were assessed in COVID-19 patients (n=222) over a period of 8 months post symptom onset. We included healthy individuals age 18-42 years as negative controls whose longitudinal blood samples were collected before the emergence of the COVID-19 pandemic. These pre-pandemic samples (n=51) were from recipients of either the seasonal inactivated influenza vaccine (n=27, collected from 2014-2018) or the live yellow fever virus (YFV-17D) vaccine (n=24, collected from 2005-2007). The Mesoscale multiplex assay was used to measure IgG, IgA, and IgM antibody responses to SARS-CoV-2 proteins in the COVID-19 patients and in the pre-pandemic healthy controls.

The magnitude of serum IgG antibodies binding to the SARS-CoV-2 spike protein increased in 92% of COVID-19 convalescent participants (n=222) relative to pre-pandemic controls (Figure 1A). The IgG responses to SARS-CoV-2 spike, RBD, and NTD declined over time with half-lives of 126 (95% confidence interval [95%CI] [107, 154]), 116 (95%CI [97, 144]), and 130 (95%CI [110, 158]) days, respectively, as estimated by an exponential decay model (Figure 1A-C and S1A). We also estimated antibody waning using a power law model, which models a scenario in which the rate of antibody decay slows over time. The power law model produced a better fit for the decay of the SARS-CoV-2 spike, RBD, and NTD binding IgG antibodies (ΔAICs>10), suggesting that spike-specific antibodies plateau over time. Because the decay rate

changes over time, the half-life is predicted to change over time as well; therefore, we

used the power law model to estimate the half-lives at 120 days after symptom onset.

The power law estimated half-lives for the IgG antibody responses to spike ($t_{1/2}$=238

days), RBD ($t_{1/2}$=209 days), and NTD ($t_{1/2}$=244 days) were longer than those estimated

by the exponential decay model (Figure S1A, C), indicating that the concentration of

these IgG antibodies may be starting to stabilize. IgA (Figure 1D-F) and IgM (Figure 1G-

I) antibodies reactive to the SARS-CoV-2 spike also increased after SARS-CoV-2

infection but were detected at lower levels and declined faster than the SARS-CoV-2-

reactive IgG antibodies. As expected, spike-binding IgM decayed more rapidly than

spike-binding IgA and IgG. Taken together these results show that antibody responses,

especially IgG antibody, were not only durable in the vast majority of patients in the 250

day period but also that the bi-phasic decay curve suggests the generation of longer

lived plasma cells producing antibody to the SARS-CoV-2 spike protein.

We also examined the antibody response to the SARS-CoV-2 nucleocapsid protein

in these infected patients. As expected, the COVID-19 patients showed higher levels of

antibody to the nucleocapsid protein compared to the pre-pandemic healthy controls

(Figure S2). However, the nucleocapsid-specific antibodies declined with a much

shorter half-life of 63 days (95%CI [58, 70]) compared to the spike protein antibodies

(Figure S1A, C). Also, the nucleocapsid reactive IgG decay rate was best fit by the

exponential model and not the power law model in contrast to what we observed with

the spike IgG antibody decay rate (Figure S1A). Thus, the nucleocapsid reactive IgG

not only declined much faster but also showed less evidence of stabilizing antibody

levels consistent with a response driven disproportionately by short-lived antibody

secreting cells – at least at this stage of the immune response.

Stable and long-lived antibody responses to common human alpha- and

betacoronaviruses in pre-pandemic healthy controls

We were interested in determining if SARS-CoV-2 infection had any effect on the

levels of antibody to the circulating human alpha- and betacoronaviruses. As a prelude

to this question, we first examined antibody levels to the spike protein of the two

circulating alphacoronaviruses (229E and NL63) and the two betacoronaviruses (HKU1

and OC43) in our pre-pandemic samples. As shown in Figure 2, all 51 pre-pandemic

samples had clearly detectable levels of IgG and IgA antibodies to the spike proteins of

the four human coronaviruses. This is the expected result since seropositivity to these

coronaviruses is very high in the adult population. But what was quite interesting was

the remarkable stability of these antibody responses over a 200-day period in the pre-

pandemic serum samples (shown as red lines in Figure 2). These were essentially flat

lines with no decline in the antibody levels and question the prevailing belief that

antibody responses to the endemic coronaviruses are short-lived (Callow et al., 1990;

Edridge et al., 2020; Lavine et al., 2021). While some occasional boosting of these

childhood-acquired coronavirus infections cannot be ruled out, these data showing such

stable antibody titers are best explained by the persistence of long-lived plasma cells in

the bone marrow many years after infection (Amanna et al., 2007; Davis et al., 2020;

Hammarlund et al., 2003; Manz et al., 1997; Slifka et al., 1998).

COVID-19 infection results in increased levels of antibodies to two common

human betacoronaviruses (HKU1 and OC43) and to SARS-CoV-1

We next examined if SARS-CoV-2 infection had any impact on the levels of

antibodies to the other human coronaviruses. We measured IgG, IgA, and IgM antibody

binding to the spike proteins of other known human coronaviruses in the COVID-19

patients (n=222 for IgG and n=190 for IgA and IgM) and compared these data to the 51

pre-pandemic healthy donor samples. In the COVID-19 patients, IgG and IgA antibodies

to the alphacoronaviruses, 229E and NL63, did not show any significant changes

compared to the antibody levels in the pre-pandemic healthy controls (Figure 2A, B, F,

G; Figure S1C, D). In contrast, the IgG and IgA antibodies to betacoronaviruses HKU1

and OC43 were substantially elevated in COVID-19 patients relative to pre-pandemic

controls (Figure 2C, D, H, I; Figure S1C, D; p<0.0001). After this boost, HKU1 and

OC43 IgG antibody levels declined with estimated half-lives of 288 (95%CI [235, 372])

and 212 (95%CI [176, 268]) days, respectively (exponential decay model). IgM levels to

common betacoronaviruses HKU1 and OC43 were low in both pre-pandemic controls

and COVID-19 patients (Figure 2M, N). While pre-existing exposure and antibodies

against HKU1 and OC43 betacoronaviruses are common in adults, pre-existing SARS-

CoV-1 exposure is rare and antibody levels to SARS-CoV-1 spike protein were very low

(essentially negative) in the pre-pandemic healthy controls. However, SARS-CoV-1

spike-reactive antibodies increased significantly after SARS-CoV-2 infection. These

increases were quite striking for IgG (p=0.0038) and also IgA (p=0.0084) and most likely

represent cross-reactive antibodies directed to SARS-CoV-2 spike epitopes that are

conserved between SARS-CoV-2 and SARS CoV-1 (Ellis et al., 2021). These newly

induced cross-reactive IgG antibodies generated after COVID-19 infection declined with

an estimated half-life of 215 days (95%CI [168, 298]) (exponential decay model) (Figure

2). Taken together, these results show that people infected with SARS-CoV-2 may have

also have some heightened immunity against the common human betacoronaviruses

and more importantly against SARS-CoV-1.

Durable neutralizing antibody responses to SARS-CoV-2 in infected patients

Neutralizing antibodies were measured with a live virus focus reduction

neutralization test that uses a recombinant SARS-CoV-2 virus expressing the

fluorescent reporter gene mNeonGreen (FRNT-mNG) (Figure 3A). During the first 250

days post-symptom onset, $FRNT_{50}$ titers varied considerably between individuals and

ranged from <20 to 3726 (Figure 3A). Of the 183 individuals for whom longitudinal

neutralization titers were assayed, 140 (77%) had at least one timepoint with

neutralization titers above the limit of detection (>20). Seventy-five percent (43/57) of

COVID-19 patients generated serum neutralizing antibodies between 30 – 50 days after

symptom onset, and similarly 72% (48/67) had measurable titers between 180 – 263

days after symptom onset. Using an exponential decay model, we evaluated the

kinetics of neutralizing antibody titers after day 42 and estimated a half-life of 150 days

(95%CI [124, 226]). However, similar to the spike reactive IgG binding antibodies, we

hypothesized that the neutralizing antibody rate of decay may actually slow over time

during the recovery period. To address this, we fit a power law to the data. The power

law model fit significantly better than the exponential decay model ($\Delta AIC=9$) and

estimated the half-life of neutralizing antibody responses at 120 days post-symptom

onset to be 254 days (95%CI [183, 400]).

Next, we assessed the relationship between the levels of spike and RBD binding

antibodies and SARS-CoV-2 neutralization. Figures 3B and C show the SARS-CoV-2

spike and RBD binding antibody response kinetics of the 183 participants for whom

neutralization titers were assessed. These exhibited a wide range of antibody binding

levels ranging from non-responders (n=11) who did not elicit antibody titers above those

of pre-pandemic controls (defined as a COVID-19 patient titer below the mean pre-

pandemic antibody titer plus 3 standard deviations, see dashed line on Figure 3B, C) to

those with IgG levels >200,000 AU/ml. Spike and RBD binding IgG levels correlated

significantly with the neutralization titers (Figure 3D, E; $p<0.0001$).

Taken together, our findings show that induction of neutralizing antibodies occurs in

the majority of COVID-19 patients. These neutralizing antibodies can persist over the 8-

9 month period following infection, and show a correlation with spike and RBD binding

IgG.

SARS-CoV-2 spike and RBD-specific memory B cells increase for several months

after infection and then plateau over 8 months

Memory B cells (MBC) are an important component of humoral immunity and

contribute to viral control by generating antibody responses upon re-exposure to the

pathogen. We used full-length spike and RBD antigen probes to quantify the

frequencies of SARS-CoV-2 spike- and RBD-specific MBC in longitudinal PBMC

samples from 111 COVID-19 patients (Figure 4) and from 29 pre-pandemic controls

(Figure S3A, B). Our flow cytometric gating strategy to identify SARS-CoV-2 specific

MBC and classify them as IgG, IgM and IgA MBC isotypes is shown in Figure 4A.

Among the total MBC, the spike IgG+ MBCs were significantly increased in COVID-

19 patients (n=111; Figure 4B) in comparison to pre-pandemic controls (n=29; Figure

S3A) (median increase, 0.73% vs. 0.02%; p<0.0001). After a steep early expansion

over the first 2-3 months, the spike IgG+ MBC persisted in COVID-19 patients with no

decline out to 250 days post symptom onset. These findings (Figure 4B) are supported

by a positive slope (0.004) from the model of the longitudinal spike IgG+ MBC

responses after day 30 (95%CI [0.002, 0.006], p<0.001; Figures S4A, B).

The spike IgM+ MBC appeared within the first two weeks post-symptom onset and

quickly declined (Figure 4C, D). The decay continued after day 30 (slope=-0.007,

95%CI [-0.010, -0.005], p<0.001). One month after symptom onset, 56% of spike MBC

were IgG+, which increased to a peak of 80% at 5-6 months (Figure 4D). Circulating

spike IgA+ MBC were also detectable in many subjects at low frequencies and without

significant change over time (day 30 to 250: slope=0.000, 95%CI [-0.002, 0.002],

p=0.91, Figure 4D).

Since the RBD contains the primary neutralizing epitopes on the spike, we also

used an RBD-specific probe to characterize this subset of spike-specific memory B

cells. Overall, approximately 20% of the spike IgG+ memory B cells targeted the RBD,

which was consistent across subjects and time (Figure 4E, F). As expected, RBD+ IgM+

MBC emerged early in infection and subsequently switched to RBD+ IgG+ MBCs, which

gradually increased during follow-up (day 30 to 250: slope=0.004, 95%CI [0.002, 0.005],

p<0.001, Figure 4E). Thus, the maintenance of circulating spike- and RBD-specific IgG

memory B cells suggests that these cells could be recruited for a rapid secondary

response following re-exposure or after vaccination.

Induction of durable and polyfunctional virus specific memory CD4+ and CD8+ T

cells in infected patients

CD4+ T cells are critical for generation of high affinity antibody responses and can

also have anti-viral effects. In addition, they provide help for CD8+ T cell responses,

which are vital for killing infected cells and mediating viral clearance. Thus, we next

examined virus-specific CD4+ and CD8+ T cell responses longitudinally in COVID-19

patients and uninfected controls using a high-dimensional, multi-parameter *ex vivo*

intracellular cytokine staining (ICS) assay. The assay is sensitive, precise and specific

for detection of antigen-specific T cells expressing multiple cytokines and effector

molecules following a short-term (6 hours) stimulation with peptide pools. Our lab

developed and validated the assay, and we are currently using the method to quantitate

Th1/Th2 CD4+ and CD8+ T cell responses in SARS-CoV-2 vaccine trials. Here, we

assessed T cell responses to the SARS-CoV-2 structural (S, E, M and N) and

accessory proteins (ORF 3a, 6, 7a, 7b, and 8) using overlapping peptide pools that

span the sequences of these proteins.

Among COVID-19 patients, 89% (102/113) mounted CD4+ T cell responses (Figure

5A) recognizing at least one SARS-CoV-2 structural protein that was detectable at one

or more visits. By contrast, SARS-CoV-2 specific CD4+ T cells were rarely detected in

the uninfected control group using this assay (Figure S3C). Antigen-specific CD4+ T

cells expanded over the first month after infection and then gradually declined over

subsequent months. Their estimated half-life was 207 days (95%CI [104, 211]) as

shown in Figure 5A, and these findings are supported by the individual CD4+ T cell

response levels and slopes after day 30 (slope= -0.0033, 95%CI [-0.0017, -0.0066],

p<0.0001) (Figure S4C, D). Of note, we observed a wide range in the total magnitude of

responses, some reaching >1% of circulating CD4+ T cells, and an overall median

frequency of 0.51% (Figures 5A and S5).

To better characterize the development of T cell memory in SARS-CoV-2 infection,

we examined the differentiation profiles of virus-specific T cells longitudinally in COVID-

19 patients. Based on CD45RA and CCR7 expression, SARS-CoV-2-specific CD4+ T

cells were primarily central memory phenotype (CD45RA- CCR7+) and to a lesser

extent effector memory (CCRA-CCR7-); this profile of the memory T cell subsets was

very consistent between subjects and stable over time (Figure 5B). The antigen-specific

CD4+ T cells were Th1-biased with a predominant CXCR3+CCR6- phenotype, and

highly polyfunctional, with simultaneous detection of antigen-specific CD154, IFN-γ, IL-

2, TNF-α and less frequently granzyme B in the early expansion phase (21-60 days post

symptom onset; median, 30 days) (Figure 5C). Interestingly, many of the virus-specific

CD4+ T cells also exhibited this polyfunctionality at the memory time point (>180 days

post symptom onset; median, 203 days) (Figure 5D). Circulating SARS-CoV-2-specific

Th2 (IL-4, IL-5 and IL-13), Th17 (IL-17) or perforin-expressing subsets were not

detected (Figure 5C and D).

Next, we examined the CD8+ T cell responses in COVID-19 patients and found that

69% generated CD8+ T cells recognizing at least one SARS-CoV-2 structural protein

that were detectable at one or more visits (Figure 6A), in contrast to infrequent to rare,

low-level antigen specific responses in the uninfected control donors (Figure S3D).

Expansion of CD8+ T cells occurred over the first month and then frequencies gradually

declined, with a half-life of 196 days (95%CI [92, 417]) and a negative estimated slope

after 30 days of symptom onset (slope=-0.004, 95%CI [-0.002, -0.008], $p < 0.0001$)

(Figure 6A). The median frequency of SARS-CoV-2-specific CD8+ T cells was 0.2%,

indicating a lower overall response magnitude than observed for CD4+ T cells.

However, like the CD4+ T cells, a wide range in magnitudes was observed with many

SARS-CoV-2-specific CD8+ T cell frequencies above 1% and even up to 12% (Figure

6A).

A very different pattern of phenotypic changes were observed with virus-specific

CD8+ T cells compared to what we saw with the CD4+ T cells (Figure 6B vs. Figure

Figure 5B). In contrast to the dominance of the central memory subset with SARS-CoV-

2-specific CD4+ T cells, the vast majority of the virus-specific CD8+ T cells showed an

effector memory phenotype during the early phase of the response. However, this

population of SARS-CoV-2-specific effector memory (CD45RA-CCR7-) contracted over

time (slope=-0.904, $p < 0.0001$; Figure 6B) and simultaneously there was an increase in

the proportion of the TEMRA (CD45RA+CCR7-) subset of virus-specific CD8+ T cells

(slope=0.075, p<0.0001; Figure 6B). A small but stable fraction of SARS-CoV-2-specific

CD8+ T cells expressed a central memory phenotype (slope=0.024, p=ns; Figure 6B).

The SARS-CoV-2-specific CD8+ T cells were highly polyfunctional with the highest

magnitude populations secreting IFN-γ, TNF-α and granzyme B; other dominant

subsets also expressed IL-2 or perforin (Figure 6C, D). This polyfunctional profile was

seen in the expansion phase (median 30 days; Figure 6C) and also at the later time

points (>180 days post symptom onset; median 203 days; Figure 6D). It is important to

note that this pattern of CD8+ T cell differentiation has been described in detail after

vaccination in humans with the live attenuated yellow fever virus vaccine (YFV-17D)

(Akondy et al., 2017). This YFV-17D vaccine generates long-lived and functional virus

specific memory CD8+ T cells that persist in humans for decades (Akondy et al., 2017;

Veit et al., 2018). That the CD8+ T cell differentiation program after COVID-19 infection

resembles what is seen after YFV infection of human suggests that COVID-19 patients

may also generate long-lived CD8+ T cell memory.

CD4+ and CD8+ cells target different SARS-CoV-2 antigen specificities

The majority of COVID-19 patients generated CD4+ T cells that recognized most

SARS-CoV-2 viral structural and accessory proteins, with the highest percentage

responding to S2 (78%) and S1 (69%) (Figure 5E, F). Among the COVID-19 subjects

with positive responses, the proportion of SARS-CoV-2-specific CD4+ T cells reacting to

each peptide pool was evenly distributed (Figure 5F). Thus, CD4+ T cells equally

targeted multiple SARS-CoV-2 proteins.

In contrast to the results seen with CD4+ T cells, SARS-CoV-2-specific CD8+ T

cells showed preferential recognition of the nucleocapsid protein. The dominant CD8+ T

cell response rate was directed to the nucleocapsid (57%), followed by ORFs 7a, 7b,

and/or 8 (25%), S1 (25%), ORFs 3a and/or 6 (16%), S2 (12%) and E and/or M (9%)

(Figure 6E). Also, among the COVID-19 patients with CD8+ T cell responses, there was

a bias with the largest percentage (median, 43%) reacting to the nucleoprocapsid

protein (Figure 6F). While SARS-CoV-2 CD8+ T cell responses rates were much lower

in uninfected controls, when present in a few control donors with lower frequencies,

these were also targeted to the nucleocapsid protein (Figure S3D). A likely explanation

for these findings is that in SARS-CoV-2 infection, antigen-presenting cells *in vivo* may

display a higher proportion of peptides derived from the nucleocapsid protein and hence

more nucleocapsid-specific CD8+ T cells are generated during infection. This has

interesting implications suggesting that nucleocapsid-specific CD8+ T cells might be

more efficient in recognizing virally infected cells.

Age and disease severity are significantly associated with magnitude of SARS-

CoV-2 immune responses

We evaluated whether COVID-19 patient age, disease severity, or sex could

account in part for the heterogeneity observed among the SARS-CoV-2-specific

immune responses as estimated from the individual models (post day 30 for cellular and

post day 42 for antibody responses). We observed that age was significantly associated

with higher immune responses to SARS-CoV-2, independently of any covariation with

disease severity (Figure 7A). Neutralizing antibody titers and IgG antibody responses to

nucleocapsid increased 1.35-fold and 1.25-fold, respectively, with each decade of age

and the same disease severity (95%CIs [1.19, 1.54] and [1.08, 1.43], p values<0.003).

Similarly, increased age positively correlated with increased frequencies of spike and

RBD-specific IgG+ memory B cells, with 1.19 to 1.24-fold higher responses per decade

of age (p values<0.02; Figure 7A), accounting for disease severity. Increased age also

correlated with higher SARS-CoV-2 and S1-specific CD4+ T cell responses (1.16-1.20-

fold increase by decade of age, p values<0.02) and N-specific CD8+ T cell responses

(1.24-fold increase by decade of age, p=0.039) accounting for disease severity (Figure

7A).

Since the cohort included primarily persons with mild to moderate COVID-19, we

had limited ability to assess the relationship of severe disease and SARS-CoV-2

immune responses, especially among the cellular responses. However, we found that

after accounting for age, severe disease (WHO score >4) was associated with higher

IgG antibodies to nucleocapsid, spike, RBD, and NTD (Figure 7B, C), and SARS-CoV-2

neutralization titers (Figure 7D). Severe disease was also associated with 2.30 to 2.46-

fold higher S1, E and/or M, and nucleocapsid-specific CD4+ T cells (all p values<0.05;

Figure 7B). We found no significant relationships between sex and the immune

responses evaluated, apart from 1.66-fold higher IgG NTD responses antibodies among

males compared to females, after accounting for age and disease severity (95%CI

[1.08, 2.55], p=0.022). In all, our analyses suggest that there are synergistic, but also

independent mechanisms driving higher adaptive immune responses in COVID-19

patients who are older and/or who experienced more severe disease.

Early SARS-CoV-2 B and T cell responses correlated with durable spike and RBD

IgG antibody binding and neutralization titers

We assessed correlations between SARS-CoV-2-specific immune responses using

the individual-level models to interpolate the magnitude of responses for each COVID-

19 patient at early (day 30) or later (day 180) convalescent time points (Figure 7E). We

found that durable serum neutralization titers correlated with the magnitude of IgG+

binding antibodies to spike, NTD and RBD at day 180 each (day 180; Spearman

R=0.62, 0.61, and 0.61, respectively; all p values<0.0001). Similarly, the frequency of

RBD+ IgG+ memory B cells at day 30 correlated with the maintenance of RBD+ IgG

antibodies (day 180; Spearman R=0.53, p<0.0001) and neutralization antibody titers

(day 180; Spearman R=0.48, p<0.0001). We also observed that the magnitude of S1-

specific CD4+ T cells at day 30 correlated with durable IgG antibodies against spike

(day 180; Spearman R=0.56, p<0.0001), NTD (Spearman R=0.62, p<0.0001), and RBD

(Spearman R=0.47, p=0.0002) (Figure 7E). These findings are consistent with early

SARS-CoV-2 memory B cells and CD4+ T cells supporting the generation of durable

antibody responses.

DISCUSSION

Establishing immune memory is essential in the defense against SARS-CoV-2

infection. To end the COVID-19 pandemic, it is critical to know how long immunity

against SARS-CoV-2 will persist after infection and whether it will be sufficient to

prevent new infections and severe disease in years to come. Identifying in depth the

adaptive immune components leading to recovery and modeling the trends of each

response was enabled by the longitudinal sampling of a large number of COVID-19

patients. Here, we show that most convalescent COVID-19 patients mount durable

antibody, B and T cells specific for SARS-CoV-2 up to 250 days, and the kinetics of

these responses provide an early indication for a favorable course ahead to achieve

long-lived immunity. Because the cohort will be followed for 2-3 more years, we can

build on these results to define the progression to long-lived immunity against this novel

human coronavirus, which can guide rational responses when future outbreaks occur.

The hallmark of the initial immune defense against SARS-CoV-2 is the emergence

of antibodies recognizing the SARS-CoV-2 spike protein, including the RBD and NTD

components of the S1 subunit, during the early phase of viral replication. These

antibodies are likely secreted from plasmablasts rapidly generated from B cells that are

activated upon their first encounter with the pathogen spike antigen. The brisk rise over

the first month of infection, followed by a fast decline of the circulating spike IgG and IgA

antibodies, is a consistent finding and likely explained by the disappearance of the

short-lived plasmablasts. These events occur even sooner for the spike IgM and

nucleocapsid antibodies.

Some antibodies that bind to specific epitopes on the spike RBD and NTD can block

SARS-CoV-2 infection of respiratory epithelial cells by inhibiting the interactions of the

viral spike with the ACE2 receptor (Ju et al., 2020; Seydoux et al., 2020; Walls et al.,

2020; Zost et al., 2020). Thus, as expected, the early rise and decline of antibodies

neutralizing live SARS-CoV-2 were similar to the kinetics of antibodies binding the spike

and RBD protein. The striking finding is the bi-phasic curve of the spike-specific binding

and neutralizing antibody responses when analyzed with the power law model, which

provides a better fit for the antibody kinetics after the peak response (Zarnitsyna et al.,

2021). With sampling data extended to 250 days, we were able to detect a slowing of

the decay of these functional antibodies toward a plateau level, suggestive of the

generation of longer-lived plasma cells, and durable antibody responses. The

importance of these observations is that following recovery, neutralizing antibodies may

persist, albeit at low levels, and may act as the first line of defense against future

encounters of SARS-CoV-2 and possibly related human coronaviruses.

Another interesting finding of this investigation is the remarkably stable antibody

responses among the pre-pandemic and COVID-19 patients to the common human

coronaviruses that are acquired in children and adults. These data are most consistent

with the generation of long-lived plasma cells and refute the current notion that these

antibody responses to human coronaviruses are short-lived. Moreover, the COVID-19

patients mounted increased IgG antibody responses to SARS-CoV-1, a related

pathogen that none likely had experienced previous exposure. This finding is consistent

with the booster response of SARS-CoV-1 neutralizing antibodies that we recently

observed following SARS-CoV-2 mRNA vaccination (Doria-Rose et al., 2021;

Stamatatos et al., 2021). Taken together, these results may have implications for a

broader strategy for vaccines targeting multiple betacoronaviruses.

The durable antibody responses in the COVID-19 recovery period are further

substantiated by the ongoing rise in both the spike and RBD memory B cell responses

after over 3-5 months before entering a plateau phase over 6-8 months. Persistence of

RBD memory B cell has been noted (Dan et al., 2021; Gaebler et al., 2021; Rodda et

al., 2021). We presume this may be explained by sustained production of memory B

cells in germinal centers of lymph nodes draining the respiratory tract in the early

months, followed by the memory B cell redistribution into the circulation as the germinal

centers begin to recede. Thus, the induction and maintenance of memory B cells and,

over time, long-lived plasma cells, will continue to furnish higher affinity antibodies if re-

exposures occur.

In contrast to spike memory B cell kinetics, SARS-CoV-2-specific CD4+ and CD8+

memory T cells each peak early, within the first month, but then slowly decline over the

next 6-7 months. Central memory Th1-type CD4+ T cells dominate throughout the early

infection and recovery period. However, the CD8+ T cells exhibit a predominant

effector memory phenotype early that transitions to those effector memory cells re-

expressing CD45RA, maintaining expression of antiviral cytokines and effector functions

which have been shown to provide protective immunity against other viral pathogens.

We also provide clear evidence that the CD4+ T cells mount a broader antigen-specific

response across the structural and accessory gene products, whereas the CD8+ T cells

are predominantly nucleocapsid specific and spike-specific responses are substantially

lower in frequency.

Our study demonstrates the considerable immune heterogeneity in the generation

of potentially protective response against SARS-CoV-2, and by focusing on the

dynamics and maintenance of B and T cell memory responses, we were able to identify

features of these early cellular responses that can forecast the durability of a potentially

effective antibody response. The ability to mount higher frequencies of RBD-specific

memory IgG+ B cells early in infection was the best indicator for a durable RBD-specific

IgG antibody and neutralizing antibody response. In addition, higher frequency CD4+ T

cells were associated with stronger spike IgG and neutralizing antibody responses.

However, the induction and peak response of SARS-CoV-2-specific CD8+ T cells

occurs independently to these antibody responses. Interestingly, while it has been

widely reported that age correlates with COVID-19 disease severity, we found that age

and disease severity were independent co-variates associated with the magnitude of

both SARS-CoV-2-specific CD4+ T cell and humoral SARS-CoV-2 immunity, but not

with the magnitude of CD8+ T cell responses. In the case of T cells, whether the T cell

differences are related to the frequencies or specificities of pre-existing coronavirus

CD4+ and CD8+ T cell immunity will require additional future analysis.

The COVID-19 pandemic remains a global public health threat after one year of

overwhelming disruption and loss. Overcoming the challenges to end the pandemic is

accentuated by the recognition that SARS-CoV-2 can undergo rapid antigenic variation

that may lower vaccine effectiveness in preventing new cases and progression to

severe disease (Edara et al., 2021; Mascola et al., 2021; Stamatatos et al., 2021). Our findings

show that most COVID-19 patients induce a wide-ranging immune defense against

SARS-CoV-2 infection, encompassing antibodies and memory B cells recognizing both

the RBD and other regions of the spike, broadly-specific and polyfunctional CD4+ T

cells, and polyfunctional CD8+ T cells. The immune response to natural infection is

highly likely to provide protective immunity even against the SARS-CoV-2 variants

because the CD4+ and CD8+ T cell epitopes will be conserved. Thus, recovered

COVID-19 patients are likely to better defend against the variants than persons who

have not been infected but have been immunized with spike-containing vaccines only.

Nevertheless, vaccine induction of CD8+ T cells to more conserved antigens such as

the nucleocapsid, rather than just to SARS-CoV-2 spike antigens, may add benefit to

more rapid containment of infection as SARS-CoV-2 variants overtake the prevailing

strains.

ACKNOWLEDGMENTS

First, we thank the participants for volunteering their time and effort to participate in

this study. We thank Children's Healthcare of Atlanta, the Georgia Research Alliance,

and the Donaldson Trust for their support. The Emory Children's Center-Vaccine

Research Clinic also thanks Laila Hussaini, Ashley Tippett, Amy Muchinsky, and

Sydney Biccum for their assistance with this study. At the Hope clinic of Emory Vaccine

Center, we thank Rebecca Fineman, Dumingu Nipuni Gomes, Ellie Butler, Michelle

Wiles for their assistance with the study. At the Fred Hutchinson Cancer Research

Center, we thank Roland Strong for providing recombinant SARS-CoV-2 hexapro spike

(S6P) and Leo Stamatatos for providing RBD protein. We also thank Rebecca Putnam,

Todd Haight, Kim Louis, Ro Yoon, Carol Marty, Daryl Morris, Xiaoling Song, Mark

Majeres, Joe Abbott, Omolara Akingba, Josh Donahue, Tu Anh Nguyen, Katharine

Schwedhelm, Carly Sprague, and Terri Stewart for their vital assistance with this study.

AUTHOR CONTRIBUTIONS

DECLARATION OF INTERESTS

546 The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. Longitudinal SARS-CoV-2 spike binding antibody responses.

IgG (A-C), IgA (D-F), and IgM (G-I) antibodies reactive to SARS-CoV-2 spike (A, D, G), spike receptor binding domain (RBD; B, E, H), and the spike N terminal domain (NTD; C, F, I) were measured by an electrochemiluminescent multiplex immunoassay and reported as arbitrary units per ml (AU/ml) as normalized by a standard curve. Longitudinal antibody titers of COVID-19 patients (in blue, n=222 COVID-19+ for IgG; n=190 COVID-19+ for IgA and for IgM) are plotted over days since symptom onset, whereas longitudinal pre-pandemic donor samples (in red, n=51 for IgG, IgA and IgM) were collected in the course of a non-SARS-CoV-2 vaccine study before 2019 and plotted over days since immunization. IgG decay curves and half-lives estimated by an exponential decay model are shown in black, and the decay curves and half-lives at day 120 post symptom onset estimated by a power law model are shown in green.

Figure 2. Longitudinal binding antibody responses to other coronavirus spike proteins.

IgG (A-E), IgA (F-J), and IgM (K-O) antibody responses in sera collected from COVID-19+ patients (in blue, n=222 for IgG; n=190 for IgA and IgM) and pre-pandemic donors (in red, n=51 for IgG, IgA and IgM) that were reactive to 229E spike (A, F, K), NL63 spike (B, G, L), HKU1 spike (C, H, M), OC43 spike (D, I, N), and the SARS-CoV-1 spike protein (E, J, O). Antibody responses were measured by an electrochemiluminescent multiplex immunoassay and reported as arbitrary units per ml (AU/ml) as normalized by a standard curve. IgG decay curves and half-lives estimated by an exponential decay

model are shown in black. There was no significant decline in IgG reactive to endemic

alpha and betacoronaviruses in longitudinal samples collected in healthy donors before

the pandemic (red, A-D).

Figure 3. Neutralizing antibody responses to SARS-CoV-2

(A) *In vitro* serum neutralization antibody titers to SAR-CoV-2 were measured by focus-

reduction neutralization assay in 183 COVID-19 patients. The limit of detection is

indicated with a dashed line at FRNT-mNG$_{50}$ = 20. The half-life estimated by the

exponential decay model (black) is 150 days, whereas the half-life estimated at day 120

using the power law model (green) is 254 days. IgG antibody titers reactive to SARS-

CoV-2 spike (B) and RBD (C) of the matched 183 COVID-19 for whom neutralization

titers were assessed. The geometric mean titer plus 3 standard deviations of pre-

pandemic samples is indicated by a dashed line (B and C). SARS-CoV-2 spike (D) and

RBD (E) reactive IgG levels correlated with neutralization titers at the matched time

point (repeated measures correlation, p<0.0001). The limit of detection is indicated with

a dashed line at FRNT-mNG$_{50}$ = 20.

Figure 4. SARS-CoV-2 spike and RBD specific memory B cells.

(A) Representative memory B cell gating strategy is shown for identification of SARS-

CoV-2 spike and RBD-specific IgD- IgG+, IgD- IgM+ and IgD- IgA+ memory B cells in

PBMCs from a SARS-CoV-2 convalescent participant. The frequency of spike+ (B)

IgG+ and (C) IgM+ memory B cells out of memory B cells (IgD- CD19+ CD20+) is

displayed over time from initial symptom onset among SARS-CoV-2 infected subjects

(n=105 subjects). The dashed line indicates the limit of detection. The bold line

represents the median fitted curve from a linear mixed effects model of post-day 30

responses. (D) The median percent of spike+ memory B cells expressing IgG, IgM or

IgA isotypes was assessed at monthly intervals post-symptom onset. (E) The frequency

of RBD+ IgG+ of memory B cells over time (n=141). (F) The proportion of S+ IgG+

memory B cells that are specific for the receptor binding domain are depicted over time.

Figure 5. CD4+ T cell responses to SARS-CoV-2 antigens.

(A) The sum of background-subtracted CD4+ T cells expressing *ex vivo* IFN-γ, IL-2

and/or CD40L to peptide pools spanning SARS-CoV-2 structural proteins: S1, S2,

envelope (E), membrane (M), nucleocapsid (N), and the following ORFs: 3a, 3b, 6, 7a,

7b, and 8 (n=114) for each individual/timepoint. Each sample that is 'positive' (by

MIMOSA) for at least one SARS-CoV-2 antigen is indicated by a solid circle, whereas

samples that are 'negative' for all of the SARS-CoV-2 antigens at that timepoint are

indicated by open triangles. The bold line represents the median fitted curve from a

nonlinear mixed effects model of post-day 30 responses among those with a positive

response at >= 1 time point; $t_{1/2}$ is the median half-life estimated from the median slope,

with 95%CI [104, 411]. (B) The proportion of SARS-CoV-2-specific CD4+ T cells

expressing a specific memory phenotype over time: central memory (CCR7+ CD45RA-

), effector memory (CCR7- CD45RA-) or T_{EMRA} (CCR7- CD45RA+); restricted to

'positive' responders. Polyfunctionality of SARS-CoV-2-specific CD4+ T cells are shown

at (C) 21-60 days since symptom onset (median, 30 days) and (D) >180 days median

post symptom onset (median, 203 days). Percentages of cytokine-expressing CD4+ T

cells are background subtracted and only subsets with detectable T cells are displayed.

Data shown were restricted to 'positive' responders and a single data point per

individual per time frame. All subsets were also evaluated for expression of IL-4, IL-5,

IL-13, IL-17 and perforin, and were found to be negative. (E) Bar graphs indicate the

proportion of COVID-19 convalescent patients who had a positive CD4+ T cell response

to the individual SARS-CoV-2. peptide pool *ex vivo* stimulations. Some antigens were

combined for stimulation as indicated. (F) For each subject with positive SARS-CoV-2-

specific CD4+ T cells, the proportion of the total SARS-CoV-2 responding CD4+ T cells

that are specific for each stimulation.

Figure 6. CD8+ T cell responses to SARS-COV-2 antigens.

(A) The sum of background-subtracted CD8+ T cells expressing IFN-γ (with or without

other cytokines), in response to peptide pools covering SARS-CoV-2 structural proteins:

S1, S2, envelope (E), membrane (M), nucleocapsid (N), and the following ORFs: 3a, 3b,

6, 7a, 7b, and 8 (n=114) for each individual/timepoint. Each sample that is 'positive'

(MIMOSA) for at least 1 SARS-CoV-2 antigen is indicated by a solid circle, whereas

samples that are 'negative' for all of the SARS-CoV-2 antigens at that timepoint are

indicated by open triangles. The bold black line represents the median fitted curve from

a nonlinear mixed effects model of post-day 30 responses among those with a positive

response to the antigen(s) under consideration at ≥ 1 time point; $t_{1/2}$ shown is the

median half-life estimated from the median slope, with 95%CI [92, 417]. (B) The

proportion of SARS-CoV-2-specific CD8+ T cells by memory phenotype over time:

effector memory (EM; CCR7- CD45RA-), T$_{EMRA}$ (CCR7- CD45RA+) and central memory

(CM; CCR7+ CD45RA-). Analyses were restricted to 'positive' responders.

Polyfunctionality of SARS-CoV-2-specific CD8 T cells at (C) 21-60 days post symptom

onset (median, 30 days) and (D) >180 days median post symptom onset (median, 203

days). Percentages of cytokine expressing CD8+ T cells are background subtracted and

only subsets with detectable T cells are displayed. Data shown were restricted to

positive responders and a single data point per individual per time frame. All CD8+ T

cell subsets were also evaluated for expression of IL-4, IL-5, IL-13, and IL-17 and were

found to be negative. (E) The bar graphs indicate the proportion of COVID-19

convalescent patients who had a 'positive' CD8+ T cell response to the individual

SARS-CoV-2 stimulations. (F) The fraction of the total SARS-CoV-2 responding CD8+ T

cells per subject that are specific for each peptide pool.

Figure 7. Correlations between SARS-CoV-2-specific immune responses and

assessment of covariates.

(A) The forest plot depicts the estimated fold-change in the level of each immune

response per decade of age, with 95% Wald-based CIs and p-values. (B) The forest

plot shows the estimated fold-change in the level of each immune response for severe

(WHO score > 4) vs. non-severe (WHO score ≤ 4) disease, with 95% Wald-based CIs

and p-values. S1 CD8+ T cell responses compared moderate-severe (WHO score > 2)

to mild (WHO score ≤ 2) disease as there were no participants with severe disease with

at least one positive S1 CD8+ T cell response post-day 30. Estimates in (A) and (B) are

from mixed effects models of post-day 30 (B and T cell responses) or post-day 42

(antibody responses) among responders that account for fixed effects of age and

disease severity on the level of immune response. Univariate assessment of disease

severity on the magnitude of (C) spike IgG antibodies and (D) SARS-CoV-2 neutralizing

antibodies at day 120 is shown for mild (WHO score: 0-2), moderate (WHO score: 3-4),

and severe disease (WHO score: 5+); p-values from one-way ANOVA. (E) The heatmap

shows Spearman correlations between critical SARS-CoV-2 memory immune

responses (day 30 B and T cell responses and day 180 antibody responses) with

significance levels: *$p<0.05$, **$p<0.01$, ***$p<0.001$. The tile size and color intensity

correspond to the absolute value of the Spearman rank correlation coefficient, with red

or blue indicating a positive or negative correlation, respectively. Day 30, 42 and 180

immune responses were estimated from mixed effects models of the longitudinal SARS-

CoV-2 binding antibodies, SARS-CoV-2 neutralizing antibodies, CD4+ and CD8+ T

cell responses, and B cell responses.

<u>METHODS DESRIPTIONS</u>

Study Populations

Two longitudinal COVID-19 cohort studies at Fred Hutchinson Cancer Research

Center (Seattle, Washington) and Emory University (Atlanta, Georgia) began after

receiving institutional review board approvals (IRB 10440, IRB 00001080 and

IRB00022371). Adults ≥18 years were enrolled who met eligibility criteria for SARS-

CoV-2 infection and provided informed consent. Study participants provided medical

history of co-morbidities, presentation of SARS-CoV-2 infection onset and disease

course, and peripheral blood at initial and follow up visits for analysis of serum antibody

and cellular immune responses. Additional longitudinal archived sera and PBMC from

pre-pandemic study populations from Emory and Seattle served as controls for the

immune assays.

The Atlanta study population included adult volunteers over the age of 18 who were

diagnosed with COVID-19 by a commercially available SARS CoV-2 PCR assay, rapid

antigen test, or clinical syndrome only (later confirmed with serology) due to limited

SARS-CoV-2 testing during the early period of the pandemic. Ambulatory participants

were recruited through local advertisements, internet-based avenues (such as social

media, list serves), COVID-19 testing sites, and primary care clinics. Hospitalized

patients were identified through SARS-CoV-2 testing. Informed consent was obtained

from all participants prior to conduct of study procedures. Initial acute peripheral blood

samples were collected from hospitalized patients at the time of enrollment.

Convalescent samples from hospitalized patients were collected when the patients were

able to return for a visit to the clinical research site at the next study visit. Serial

peripheral blood samples were collected starting at about 30 days after the onset of

COVID-19 symptoms and/or after PCR positivity for SARS-CoV-2. Thereafter, samples

were collected at 3, 6, and 9 months. The study is ongoing with expected completion of

sample collection from participants in February 2023. Participants were excluded if they

were immunocompromised, HIV positive, had active hepatitis B or C virus infection,

used immunosuppressive drugs for 2 weeks or more in the preceding 3 months,

received blood products or immune globulin 42 days prior to enrollment, received

convalescent COVID-19 plasma, or were pregnant or breast feeding. We report on 110

participants to date, of which 73% were diagnosed by SARS-CoV-2 PCR, the remaining

were diagnosed by rapid antigen test or serology. Demographic features of the

participants are as follows: median age was 48; 45% were male; the majority (80%)

were white, 11% Black/African American, 6% Asian, and 8% were Hispanic/Latinx

ethnicity. The most frequent co-morbid conditions were hypertension, obesity, heart

disease and diabetes mellitus. The most frequent COVID-19 symptoms were

myalgia/fatigue, fever, cough, headache, loss of smell and taste (Table S1).

Hospitalized patients were older, with a median age of 56; a higher percentage were

Black/African American (27%); and 100% had fever.

Longitudinal pre-pandemic sera samples from Emory were collected from

individuals participating in a yellow fever vaccine study from 2014-2016 or an influenza

vaccine study from 2015-2018 (Akondy et al., 2017; Ellebedy et al., 2016). Data were

included for analysis of binding antibody responses and are presented as days post-

irrelevant (yellow fever) vaccination.

The Seattle COVID-19 Cohort study participants were recruited from the Seattle

metropolitan area by social media advertisements, partnership with the local emergency

medical service and by word of mouth. Study participants were screened and enrolled

by the Seattle Vaccine Trials Unit staff. Eligibility criteria included adults at risk for

SARS-CoV-2 infection or those diagnosed with COVID-19 by a commercially available

SARS-CoV-2 PCR assay or blood antibody test and willing to have at least four blood

draws collected over one year. Exclusion criteria included pregnancy and inability to

donate blood.

Informed electronic consent was obtained from all Seattle participants during a

screening phone call with study clinical staff. Interested participants were screened,

consented and medical history and COVID-19 illness onset date and symptoms

collected. Participants undiagnosed with COVID-19 had a nasopharyngeal (NP) swab

collected and tested for SARS-CoV-2 via an FDA-approved PCR test and blood was

collected for SARS-CoV-2 antibody (Abbott) and study assays. Those with either a

positive PCR or antibody test were asked to return for future blood draws. Those who

tested negative were asked to return as controls for the positive cohort and in case they

tested positive in the future. Participants with a positive test prior to study enrollment or

those diagnosed in study were asked to provide blood donation at approximately 7

days, 2 weeks, 1, 2, 3, 4, 6, 9- and 12-months post symptom onset. After completing

one year of study, participants will be given the option of continuing the longitudinal

study for up to two or more years. At each study visit, participant symptoms and medical

history is updated. Those with COVID-19 symptoms after enrollment in all groups are

offered a nasopharyngeal swab PCR SARS-CoV-2 test.

As of October 2020, 805 individuals have contacted the Seattle COVID-19 cohort

study and 425 have enrolled. This includes 281 negative and 144 SARS-CoV-2 positive

participants. Reasons for not enrolling include lack of interest, not meeting the eligibility

criteria, inability to travel to blood draw location and inability to collect study blood. No

participants have terminated from the study. Study enrollment and follow-up remains

ongoing. Samples from SARS-CoV-2 negative subjects were included in B and T cell

assays as 'contemporaneous' negative controls.

Peripheral blood mononuclear cells (PBMC) were obtained from HIV-1 seronegative

donors who were recruited at the Seattle Vaccine Trials Unit before 2019 as part of the

study "Establishing Immunologic Assays for Determining HIV-1 Prevention and Control".

All participants signed informed consent, and the Fred Hutchinson Cancer Research

Center IRB (Seattle, WA, USA) institutional human subjects review committee approved

the protocol prior to study initiation. Pre-pandemic samples from this cohort were used

as assay controls in B and T cell assays.

PBMC processing

PBMC for cellular assays were isolated by density centrifugation and cryopreserved

from ACD-anticoagulated whole blood within eight hours of venipuncture, as described

previously (Cross-Network PBMC SOP Working Group, 2018). Sera were also

processed and cryopreserved within 4 hours after collection.

Antibody binding assay

Antibody binding titers were measured using a multiplex plate coated with the SARS-

CoV-2 spike, SARS-CoV-2 spike receptor binding domain, SARS-CoV-2 spike N

terminal domain, SARS-CoV-2 nucleocapsid, SARS-CoV-1 spike, 229E spike, NL63

spike, HKU1 spike, and OC43 spike proteins (Mesoscale Discovery). Plates were

blocked with 150μl/well with 5% bovine serum albumin in phosphate buffered saline

(PBS) and shaken at 700 RPM at room temperature for at least 30 min. Plates were

washed 3 times with 150μl/well 0.05% Tween-20 in PBS. Serum and plasma samples

were added to the plate at dilutions between 1:500 and 1:50,000 and shaken at 700

RPM at room temperature for 2 hr. Following a wash, plates were incubated with

50ul/well of Sulfo-Tag anti-human IgG, IgA, or IgM detection antibody and shaken at

700RPM at room temperature for 1 hr. After a subsequent wash, 150μl/well of MSD

GOLD read buffer was added to the plate and plates were immediately read on the

MSD instrument to measure light intensity. Antibody levels are reported as arbitrary

units/ml (AU/ml) based on normalization to a standard curve.

Viruses and cells

VeroE6 cells were obtained from ATCC (clone E6, ATCC, #CRL-1586) and cultured in

complete DMEM medium consisting of 1× DMEM (VWR, #45000-304), 10% FBS,

25mM HEPES Buffer (Corning Cellgro), 2mM L-glutamine, 1mM sodium pyruvate, 1×

Non-essential Amino Acids, and 1× antibiotics. The infectious clone SARS-CoV-2

(icSARS-CoV-2-mNG), derived from the 2019-nCoV/USA_WA1/2020 strain, was

propagated in VeroE6 cells (ATCC) and sequenced (Xie et al., 2020).

Focus reduction neutralization test

Neutralization assays with SARS-CoV-2 virus were performed as previously described

(Suthar et al., 2020; Vanderheiden et al., 2020; Xie et al., 2020). Plasma/serum were

serially diluted (three-fold) in serum-free Dulbecco's modified Eagle's medium (DMEM)

in duplicate wells and incubated with 100–200 FFU infectious clone derived SARS-CoV-

2-mNG virus at 37°C for 1 hr (Xie et al., 2020). The antibody-virus mixture was added to

VeroE6 cell (C1008, ATCC, #CRL-1586) monolayers seeded in 96-well blackout plates

and incubated at 37°C for 1 hr. Post-incubation, the inoculum was removed and

replaced with pre-warmed complete DMEM containing 0.85% methylcellulose. Plates

were incubated at 37°C for 24 hr. After 24 hr, methylcellulose overlay was removed,

cells were washed twice with PBS and fixed with 2% paraformaldehyde in PBS for

30 min at room temperature. Following fixation, plates were washed twice with PBS and

foci were visualized on a fluorescence ELISPOT reader (CTL ImmunoSpot S6 Universal

Analyzer) and enumerated using Viridot (Katzelnick et al., 2018). The neutralization

titers were calculated as follows: 1 - (ratio of the mean number of foci in the presence of

sera and foci at the highest dilution of respective sera sample). Each specimen was

tested in two independent assays performed at different times. The FRNT-mNG$_{50}$ titers

were interpolated using a 4-parameter nonlinear regression in GraphPad Prism

8.4.3. Samples with an FRNT-mNG$_{50}$ value that was below the limit of detection were

plotted at 20.

Spike and RBD memory B cell flow cytometry assays

Fluorescent SARS-CoV-2-specific S6P (Hsieh et al., 2020) (provided by Roland Strong,

Fred Hutchinson Cancer Research Center, Seattle, WA) and RBD (provided by

Leonidas Stamatatos, Fred Hutchinson Cancer Research Center, Seattle, WA) probes

were made by combining biotinylated protein with fluorescently labeled streptavidin

(SA). The S6P probes were made at a ratio of 1:1 molar ratio of trimer to SA. Two S6P

probes, one labeled with AlexaFluor488 (Invitrogen), one labeled with AlexaFluor647

(Invitrogen), were used in this panel in order to increase specificity of the detection of

SARS-CoV-2-specific B cells. The RBD probe was prepared at a 4:1 molar ratio of RBD

monomers to SA, labeled with R-phycoerythrin (Invitrogen). Cryopreserved PBMCs

from SARS-CoV-2-convalescent participants and a pre-pandemic SARS-CoV-2-naïve

donor were thawed at 37°C and stained for SARS-CoV-2-specific memory B cells as

described previously (Seydoux et al., 2020) with a flow cytometry panel shown in

Reagents Table 1. Cells were stained first with the viability stain (Invitrogen) in PBS for

15 min at 4°C. Cells were then washed with 2% FBS/PBS and stained with a cocktail of

the three probes for 30 min at 4°C. The probe cocktail was washed off with 2%

FBS/PBS and the samples were stained with the remaining antibody panel and

incubated for 25 min at 4°C. The cells were washed two times and resuspended in 1%

paraformaldehyde/1× PBS for collection on a LSR II or FACSymphony flow cytometer

(BD Biosciences). Data was analyzed in Flow Jo version 9.9.4.

Intracellular cytokine staining (ICS) assay

Flow cytometry was used to examine SARS-CoV-2-specific CD4+ and CD8+ T-cell

responses using a validated ICS assay. The assay was similar to a published report

(Dintwe et al., 2019; Horton et al., 2007; Sadoff et al., 2021) and the details of the

staining panel are included in Reagents Table 2. Peptide pools covering the structural

proteins of SARS-CoV-2 were used for the six-hour stimulation. Peptides matching the

SARS-CoV-2 spike sequence (316 peptides, plus 4 peptides covering the G614 variant)

were synthesized as 15 amino acids long with 11 amino acids overlap and pooled in 2

pools (S1 and S2) for testing (BioSynthesis). All other peptides were 13 amino acids

overlapping by 11 amino acids and were synthesized by GenScript. The peptides

covering the envelope (E), membrane (M) and nucleocapsid (N) were initially combined

into one peptide pool, but the majority of the assays were performed using a separate

pool for N and one that combined only E and M. Several of the open reading frame

(ORF) peptides were combined into two pools: ORF 3a and 6, and ORF 7a, 7b and 8.

All peptide pools were used at a final concentration of 1 µg/ml for each peptide. As a

negative control, cells were not stimulated, only the peptide diluent (DMSO) was

included. As a positive control, cells were stimulated with a polyclonal stimulant,

staphylococcal enterotoxin B (SEB). Cells expressing IFN-γ and/or IL-2 and/or CD154

was the primary immunogenicity endpoint for CD4+ T cells and cells expressing IFN-γ

was the primary immunogenicity endpoint for CD8+ T cells. The overall response to

SARS-CoV-2 was defined as the sum of the background-subtracted responses to each

of the individual pools. A sample was considered positive for CD4+ or CD8+ T cell

responses to SARS-CoV-2 if any of the CD4+ or CD8+ T cell responses to the

individual peptide pool stimulations was positive. Positivity was determined using

MIMOSA (Finak et al., 2014). The total number of CD4+ T cells must have exceeded

10,000 and the total number of CD8+ T cells must have exceeded 5,000 for the assay

data to be included in the analysis.

Statistical methods

Binding and neutralizing antibody responses

Mixed effects exponential and power law models were used to analyze waning of

antibody (day 42 to day 263 post symptom onset). For binding antibody analyses,

antibody (Ab) was natural log transformed, yielding linear equations of the form

$\ln(Ab)=a+b*(day-42)$ and $\ln(Ab)=a+b*\ln(day/42)$ for the exponential and power law

models, respectively, and fit using the lmer function (lme4 package) in R. Models

included population level fixed effects and individual level random effects for intercept

and slope and covariance between the random effects. Simplified models – with random

effects only for intercept – were also fit. Neutralization antibody data were analyzed in

Monolix (Lixoft). For analysis in Monolix, the exponential and power law models were

formulated as ordinary differential equations, $dAb/dt=k*Ab$ and $dAb/dt=k*Ab/t$,

respectively, with antibody at day 42 lognormally distributed and lognormal multiplicative

error. Neutralization titers < 20 were treated as left censored. For comparison of

models, difference in Akaike information criterion (ΔAIC) > 4 was considered statistically

significant. Models (in R and Monolix) were fit using maximum likelihood. To account for

repeated measures, correlations between antibody binding levels and neutralization

titers were calculated using a repeated measures correlation (rmcorr package) in R

(Bakdash and Marusich, 2017).

B cell responses

We considered linear mixed effects models for B cell response, $\mathcal{Y}_{ij}$, as a function of t_{ij},

the j^{th} time since symptom onset for the i^{th} individual, with random effects for intercept

and slope and $t_{ij} > 30$ days for all i, j:

883
$$\log_e \mathcal{Y}_{ij} = \beta_{0i} + \beta_{1i} t_{ij} + \epsilon_{ij}$$

884   where $\beta_{0i} = \beta_0 + b_i$ and $\beta_{1i} = \beta_1 + c_i$ with (b_i, c_i) iid $\sim N_2(\mathbf{0}, \mathbf{\Sigma})$, with

886
$$\mathbf{\Sigma} = \begin{bmatrix} \sigma_b^2 & Cov(b, c) \\ Cov(b, c) & \sigma_c^2 \end{bmatrix}$$

and σ_b^2 and σ_c^2 are the between-person variation in the intercept and slope of log B cell

responses respectively, *Cov(b, c)* is the covariance between the intercept and slope,

and ϵ_{ij} iid $\sim N(0, \sigma^2)$. The random effects, b_i and c_i, are each assumed to be

independent for different individuals and the within-individual errors ϵ_{ij} are assumed to

be independent for different *i, j* and to be independent of the random effects. The

function lme from the R package nlme was used to fit the model.

T cell responses

Longitudinal analyses of CD4+ and CD8+ T cell responses were performed for

individuals with a positive response for at least one time point 30 days after symptom

onset. The MIMOSA (Mixture Models for Single-Cell Assays) (Finak et al., 2014) model

incorporated cell count and cell proportion information to define a positive CD4+/CD8+

T cell response by ICS by comparing peptide pools stimulated cells and unstimulated

negative controls. This method assumed a common distribution for cytokine positive

CD4+/CD8+ T cells in stimulated and unstimulated samples in non-responders,

resulting in paired differences that were zero on average. In contrast, for responders,

the distribution of the proportion of cytokine positive cells for stimulated samples was

assumed to be greater than for unstimulated samples, resulting in paired differences

that were greater than zero on average. The MIMOSA method modeled this structure

through a Bayesian hierarchical mixture model framework. One component (or

distribution) of the model represented the responders, and the other component

modeled the non-responders. The parameters defining these distributions, as well as

the probabilities that each ICS response was either a responder or non-responder, were

estimated from the observed data. This sharing of information across SARS-CoV-2

responders and non-responders increased the sensitivity and specificity to make

positivity calls (Newton et al., 2004). Responses with probability of response > 0.999

were considered positive responders.

We considered nonlinear mixed effects models for T cell response, $\mathcal{Y}_{ij}$, as a function of

t_{ij}, the j^{th} time since symptom onset for the i^{th} individual, with random effects for

intercept and slope and $t_{ij} > 30$ days for all i, j:

917
$$\log_e \mathcal{Y}_{ij} = \beta_{0i} - \exp(\beta_{1i})\, t_{ij} + \epsilon_{ij}$$

918   where $\beta_{0i} = \beta_0 + b_i$ and $exp(\beta_{1i}) = exp(\beta_1 + c_i)$ with (b_i, c_i) iid $\sim N_2(\mathbf{0}, \mathbf{\Sigma})$, with

919
$$\Sigma = \begin{bmatrix} \sigma_b^2 & 0 \\ 0 & \sigma_c^2 \end{bmatrix}$$

and σ_b^2 and σ_c^2 are the between-person variation in the intercept and slope of log T cell

responses respectively, and ϵ_{ij} iid ~ $logNormal(0, \sigma^2)$. The random effects, b_i and c_i,

are each assumed to be independent for different individuals and the within-individual

errors ϵ_{ij} are assumed to be independent for different *i, j* and to be independent of the

random effects. The function nlme from the R package nlme was used to fit the models.

Individual-level estimates at days 30 (T and B cell responses), day 42 (antibody

responses) and day 180 (all responses) were obtained from the mixed effects models

described above. Spearman rank correlations, Wald-based two-sided 95% confidence

intervals and p-values were reported.

Generalized estimating equations (GEE), with an independence working covariance

matrix, were used to confirm the results of the covariate assessments for B and T cell

responses from the mixed effects models. Two-tailed P-values based on the robust

standard error estimates for the covariate coefficients were consistent with the

corresponding two-tailed P-values for the covariate associations from the mixed effects

models.

Reagents Table 1. Reagents used in the B cell staining panel.

Antibody	Manufacturer	Clone
CD3 BV510	BD Biosciences	HIT3a
CD14 BV510	BD Biosciences	MφP9
CD56 BV510	BD Biosciences	NCAM16.2
CD19 BUV395	BD Biosciences	SJ25-C1
CD20 BUV737	BD Biosciences	2H7
CD21 PE-Cy7	BD Biosciences	B-ly4
CD27 BV605	BioLegend	O323
CD38 BB700	BD Biosciences	HIT2
IgA VioBlue	Miltenyi Biotec	IS11-8E10
IgD BV650	BD Biosciences	IA6-2
IgG BV786	BD Biosciences	G18-145
IgM PE/Dazzle 594	BioLegend	MHM-88
R-Phycoerythrin Streptavidin	Invitrogen	N/A
AlexaFluor 488 Streptavidin	Invitrogen	N/A
AlexaFluor 647 Streptavidin	Invitrogen	N/A
LIVE/DEAD Fixable Aqua Stain	Invitrogen	N/A

937 *APC, allophycocyanin; BB, brilliant blue; BUV, brilliant ultraviolet; BV, brilliant violet; Cy, cyanine;*
938 *FITC, fluorescein isothiocyanate; PE R-phycoerythrin; UViD, Live/Dead fiaxable ultraviolet dead cell*
939 *stain*
940

941 **Reagents Table 2. T cell intracellular cytokine staining flow cytometry panel.**

Detector	Fluorochrome	Specificity	Clone	Purpose
B515	FITC	Perforin	B-D48	Function
B610	BB630	IL-5	TRFK5	Function
B610	BB630	IL-13	JES10-5A2	Function
B660	BB660	Ki67	B56	Activation
B710	BB700	IL-4	MP4-25D2	Function
G575	PE	CRTh2	BM16	T helper class
G610	PE-Dazzle594	CD32	FUN-2	Fcg receptor
G660	Pe-Cy5	CXCR3 (CD183)	1C6/CXCR3	T helper class
G710	PE-Cy5.5	FOXP3	PCH101	Treg
G780	PE-Cy7	IL-17a	BL168	Function
R660	APC	IL-2	MQ1-17H12	Function
R710	Alexa 700	Granzyme B	GB11	Function
R780	APC-Fire750	CD3	UCHT1	T cell lineage
U395	BUV395	TNF	MAb11	Function
U450	UViD	Viability	N/A	Viability
U500	BUV496	CD45RA	HI100	Differentiation
U570	BUV563	CD19	SJ25C1	B cells
U660	BUV661	CD14	MFP9	Monocytes
U730	BUV737	CD154	TRAP1	Function
U780	BUV805	CD8	SK1	T cell lineage
V450	V450	IFN-γ	B27	Function
V510	BV480	CD4	SK3	Function
V570	BV570	CD16	3G8	NK, NKT
V610	BV605	CCR7	G034H7	Differentiation
V655	BV650	CD25	M-A251	Treg, activation
V710	BV711	CD64	10.1	Fcg receptor
V750	BV750	CD56	5.1H11	NK, NKT
V780	BV786	CCR6 (CD196)	11A9	T helper class

References

References

Akondy, R.S., Fitch, M., Edupuganti, S., Yang, S., Kissick, H.T., Li, K.W., Youngblood,
B.A., Abdelsamed, H.A., McGuire, D.J., Cohen, K.W*., et al.* (2017). Origin and
differentiation of human memory CD8 T cells after vaccination. Nature *552*, 362-367.

Amanna, I.J., Carlson, N.E., and Slifka, M.K. (2007). Duration of humoral immunity to
common viral and vaccine antigens. N Engl J Med *357*, 1903-1915.

Anderson, E.J., Rouphael, N.G., Widge, A.T., Jackson, L.A., Roberts, P.C., Makhene,
M., Chappell, J.D., Denison, M.R., Stevens, L.J., Pruijssers, A.J*., et al.* (2020). Safety
and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine in Older Adults. *383*, 2427-
2438.

Bakdash, J.Z., and Marusich, L.R. (2017). Repeated Measures Correlation. Front
Psychol *8*, 456.

Callow, K.A., Parry, H.F., Sergeant, M., and Tyrrell, D.A. (1990). The time course of the
immune response to experimental coronavirus infection of man. Epidemiol Infect *105*,
435-446.

Cross-Network PBMC SOP Working Group (2018). Cross-Network PBMC Processing
SOP v6.0. HIV/AIDS Network Coordination (HANC)
https://www.hanc.info/labs/Documents/PBMC%20Documents/HANC-LAB-
P0001_v6.0_2018-04-26_PBMC_SOP.pdf.

Dan, J.M., Mateus, J., Kato, Y., Hastie, K.M., Yu, E.D., Faliti, C.E., Grifoni, A., Ramirez,
S.I., Haupt, S., Frazier, A*., et al.* (2021). Immunological memory to SARS-CoV-2
assessed for up to 8 months after infection. Science *371*.

Davis, C.W., Jackson, K.J.L., McCausland, M.M., Darce, J., Chang, C., Linderman,
S.L., Chennareddy, C., Gerkin, R., Brown, S.J., Wrammert, J*., et al.* (2020). Influenza
vaccine-induced human bone marrow plasma cells decline within a year after
vaccination. Science *370*, 237-241.

Dintwe, O., Rohith, S., Schwedhelm, K.V., McElrath, M.J., Andersen-Nissen, E., and De
Rosa, S.C. (2019). OMIP-056: Evaluation of Human Conventional T Cells, Donor-
Unrestricted T Cells, and NK Cells Including Memory Phenotype by Intracellular
Cytokine Staining. Cytometry A *95*, 722-725.

Doria-Rose, N., Suthar, M.S., Makowski, M., O'Connell, S., McDermott, A.B., Flach, B.,
Ledgerwood, J.E., Mascola, J.R., Graham, B.S., Lin, B.C*., et al.* (2021). Antibody
Persistence through 6 Months after the Second Dose of mRNA-1273 Vaccine for Covid-
19. N Engl J Med.

Edara, V.V., Norwood, C., Floyd, K., Lai, L., Davis-Gardner, M.E., Hudson, W.H.,
Mantus, G., Nyhoff, L.E., Adelman, M.W., Fineman, R*., et al.* (2021). Infection- and
vaccine-induced antibody binding and neutralization of the B.1.351 SARS-CoV-2
variant. Cell Host Microbe *29*, 516-521 e513.

Edridge, A.W.D., Kaczorowska, J., Hoste, A.C.R., Bakker, M., Klein, M., Loens, K.,
Jebbink, M.F., Matser, A., Kinsella, C.M., Rueda, P.*, et al.* (2020). Seasonal coronavirus
protective immunity is short-lasting. Nat Med *26*, 1691-1693.

Ellebedy, A.H., Jackson, K.J., Kissick, H.T., Nakaya, H.I., Davis, C.W., Roskin, K.M.,
McElroy, A.K., Oshansky, C.M., Elbein, R., Thomas, S.*, et al.* (2016). Defining antigen-
specific plasmablast and memory B cell subsets in human blood after viral infection or
vaccination. Nat Immunol *17*, 1226-1234.

Finak, G., McDavid, A., Chattopadhyay, P., Dominguez, M., De Rosa, S., Roederer, M.,
and Gottardo, R. (2014). Mixture models for single-cell assays with applications to
vaccine studies. Biostatistics *15*, 87-101.

Gaebler, C., Wang, Z., Lorenzi, J.C.C., Muecksch, F., Finkin, S., Tokuyama, M., Cho,
A., Jankovic, M., Schaefer-Babajew, D., Oliveira, T.Y.*, et al.* (2021). Evolution of
antibody immunity to SARS-CoV-2. Nature *591*, 639-644.

Hammarlund, E., Lewis, M.W., Hansen, S.G., Strelow, L.I., Nelson, J.A., Sexton, G.J.,
Hanifin, J.M., and Slifka, M.K. (2003). Duration of antiviral immunity after smallpox
vaccination. Nat Med *9*, 1131-1137.

Horton, H., Thomas, E.P., Stucky, J.A., Frank, I., Moodie, Z., Huang, Y., Chiu, Y.L.,
McElrath, M.J., and De Rosa, S.C. (2007). Optimization and validation of an 8-color
intracellular cytokine staining (ICS) assay to quantify antigen-specific T cells induced by
vaccination. J Immunol Methods *323*, 39-54.

Hsieh, C.L., Goldsmith, J.A., Schaub, J.M., DiVenere, A.M., Kuo, H.C., Javanmardi, K.,
Le, K.C., Wrapp, D., Lee, A.G., Liu, Y.*, et al.* (2020). Structure-based design of
prefusion-stabilized SARS-CoV-2 spikes. Science *369*, 1501-1505.

Ju, B., Zhang, Q., Ge, J., Wang, R., Sun, J., Ge, X., Yu, J., Shan, S., Zhou, B., Song,
S.*, et al.* (2020). Human neutralizing antibodies elicited by SARS-CoV-2 infection.
Nature *584*, 115-119.

Katzelnick, L.C., Coello Escoto, A., McElvany, B.D., Chavez, C., Salje, H., Luo, W.,
Rodriguez-Barraquer, I., Jarman, R., Durbin, A.P., Diehl, S.A.*, et al.* (2018). Viridot: An
automated virus plaque (immunofocus) counter for the measurement of serological
neutralizing responses with application to dengue virus. PLoS Negl Trop Dis *12*,
e0006862.

Lavine, J.S., Bjornstad, O.N., and Antia, R. (2021). Immunological characteristics
govern the transition of COVID-19 to endemicity. Science *371*, 741-745.

Manz, R.A., Thiel, A., and Radbruch, A. (1997). Lifetime of plasma cells in the bone
marrow. Nature *388*, 133-134.

Mascola, J.R., Graham, B.S., and Fauci, A.S. (2021). SARS-CoV-2 Viral Variants-
Tackling a Moving Target. JAMA *325*, 1261-1262.

Newton, M.A., Noueiry, A., Sarkar, D., and Ahlquist, P. (2004). Detecting differential
gene expression with a semiparametric hierarchical mixture method. Biostatistics *5*,
155-176.

Rodda, L.B., Netland, J., Shehata, L., Pruner, K.B., Morawski, P.A., Thouvenel, C.D.,
Takehara, K.K., Eggenberger, J., Hemann, E.A., Waterman, H.R*., et al.* (2021).
Functional SARS-CoV-2-Specific Immune Memory Persists after Mild COVID-19. Cell
184, 169-183 e117.

Sadoff, J., Le Gars, M., Shukarev, G., Heerwegh, D., Truyers, C., de Groot, A.M.,
Stoop, J., Tete, S., Van Damme, W., Leroux-Roels, I*., et al.* (2021). Interim Results of a
Phase 1-2a Trial of Ad26.COV2.S Covid-19 Vaccine. N Engl J Med.

Sette, A., and Crotty, S. (2021). Adaptive immunity to SARS-CoV-2 and COVID-19. Cell
184, 861-880.

Seydoux, E., Homad, L.J., MacCamy, A.J., Parks, K.R., Hurlburt, N.K., Jennewein,
M.F., Akins, N.R., Stuart, A.B., Wan, Y.H., Feng, J*., et al.* (2020). Analysis of a SARS-
CoV-2-Infected Individual Reveals Development of Potent Neutralizing Antibodies with
Limited Somatic Mutation. Immunity *53*, 98-105 e105.

Slifka, M.K., Antia, R., Whitmire, J.K., and Ahmed, R. (1998). Humoral immunity due to
long-lived plasma cells. Immunity *8*, 363-372.

Stamatatos, L., Czartoski, J., Wan, Y.H., Homad, L.J., Rubin, V., Glantz, H., Neradilek,
M., Seydoux, E., Jennewein, M.F., MacCamy, A.J*., et al.* (2021). mRNA vaccination
boosts cross-variant neutralizing antibodies elicited by SARS-CoV-2 infection. Science.

Stephens, D.S., and McElrath, M.J. (2020). COVID-19 and the Path to Immunity. JAMA
324, 1279-1281.

Suthar, M.S., Zimmerman, M.G., Kauffman, R.C., Mantus, G., Linderman, S.L., Hudson,
W.H., Vanderheiden, A., Nyhoff, L., Davis, C.W., Adekunle, O*., et al.* (2020). Rapid
Generation of Neutralizing Antibody Responses in COVID-19 Patients. Cell Rep Med *1*,
100040.

Vanderheiden, A., Edara, V.V., Floyd, K., Kauffman, R.C., Mantus, G., Anderson, E.,
Rouphael, N., Edupuganti, S., Shi, P.Y., Menachery, V.D*., et al.* (2020). Development of
a Rapid Focus Reduction Neutralization Test Assay for Measuring SARS-CoV-2
Neutralizing Antibodies. Curr Protoc Immunol *131*, e116.

Veit, O., Domingo, C., Niedrig, M., Staehelin, C., Sonderegger, B., Hequet, D., Stoeckle,
M., Calmy, A., Schiffer, V., Bernasconi, E*., et al.* (2018). Long-term Immune Response
to Yellow Fever Vaccination in Human Immunodeficiency Virus (HIV)-Infected
Individuals Depends on HIV RNA Suppression Status: Implications for Vaccination
Schedule. Clinical infectious diseases : an official publication of the Infectious Diseases
Society of America *66*, 1099-1108.

Walls, A.C., Park, Y.J., Tortorici, M.A., Wall, A., McGuire, A.T., and Veesler, D. (2020).
Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell *181*,
281-292 e286.

Xie, X., Muruato, A., Lokugamage, K.G., Narayanan, K., Zhang, X., Zou, J., Liu, J.,
Schindewolf, C., Bopp, N.E., Aguilar, P.V*., et al.* (2020). An Infectious cDNA Clone of
SARS-CoV-2. Cell Host Microbe *27*, 841-848 e843.

Zarnitsyna, V.I., Akondy, R.S., Ahmed, H., McGuire, D.J., Zarnitsyn, V.G., Moore, M.,
Johnson, P.L.F., Ahmed, R., Li, K., Hellerstein, M*., et al.* (2021). Dynamics and turnover
of memory CD8 T cell responses following yellow fever vaccination. bioRxiv,
2021.2001.2023.427919.

Zost, S.J., Gilchuk, P., Case, J.B., Binshtein, E., Chen, R.E., Nkolola, J.P., Schafer, A.,
Reidy, J.X., Trivette, A., Nargi, R.S*., et al.* (2020). Potently neutralizing and protective
human antibodies against SARS-CoV-2. Nature *584*, 443-449.

1068

SARS-CoV-2

	Spike	RBD	NTD
Exponential :	$t_{1/2}$ = 126 days	$t_{1/2}$ = 116 days	$t_{1/2}$ = 130 days
Power Law :	$t_{1/2}$ = 238 days	$t_{1/2}$ = 209 days	$t_{1/2}$ = 244 days

IgG
n=222
n=51

A   B   C

IgA
n=190
n=51

$t_{1/2}$ = 148 days $t_{1/2}$ = 157 days $t_{1/2}$ = 159 days
$t_{1/2}$ = 324 days $t_{1/2}$ = 385 days $t_{1/2}$ = 284 days

D   E   F

IgM
n=190
n=51

$t_{1/2}$ = 65 days $t_{1/2}$ = 66 days
$t_{1/2}$ = 98 days $t_{1/2}$ = 105 days

G   H   I

COVID-19   Pre-pandemic   Exponential   Power Law

Human Alphacoronaviruses   **Human Betacoronaviruses**

	229E	NL63	HKU1	OC43	SARS-CoV-1
COVID+ :	$t_{1/2}$ = no decline	$t_{1/2}$ = >1000 days	$t_{1/2}$ = 288 days	$t_{1/2}$ = 212 days	$t_{1/2}$ = 215 days
Pre-pandemic :	$t_{1/2}$ = no decline	$t_{1/2}$ = no decline	$t_{1/2}$ = no decline	$t_{1/2}$ = no decline	

COVID-19 — **Pre-pandemic** — **Exponential**

A

n=183

Exponential: $t_{1/2}$ = 150 days
Power Law: $t_{1/2}$ = 254 days

FRNT-mNG50

Days since symptom onset

B Spike

IgG AU/ml

Days since symptom onset

C RBD

IgG AU/ml

Days since symptom onset

D

FRNT-mNG50

r = 0.53
p<0.0001

Spike IgG AU/ml

E

FRNT-mNG50

r = 0.50
p<0.0001

RBD IgG AU/ml

A

	Response	P	No. of Participants
Ab	IgG N	**0.003**	210
	IgG NTD	0.103	208
	IgG RBD	0.105	208
	IgG Spike	0.059	208
	nAb	**<0.001**	122
B cells	S+ IgG+ MBC	**0.012**	99
	RBD+ IgG+ MBC	**0.019**	135
T cells	Total CD4+	**0.016**	93
	CD4+ S1	**0.020**	70
	CD4+ S2	0.316	82
	CD4+ E, M	0.180	68
	CD4+ N	0.125	68
	Total CD8+	0.202	69
	CD8+ S1	0.285	24
	CD8+ N	**0.035**	53

<– Lower responses   Higher responses –>

0.6 0.8 1 1.2 1.4 1.6

Estimated fold-change in immune response
per age decade (adjusted for disease severity)

B

	Response	P	No. of Participants
Ab	IgG N	**0.014**	210
	IgG NTD	**0.001**	208
	IgG RBD	**0.002**	208
	IgG Spike	**0.006**	208
	nAb	**0.020**	122
B cells	S+ IgG+ MBC	0.435	99
	RBD+ IgG+ MBC	0.160	135
T cells	Total CD4+	0.317	93
	CD4+ S1	**0.026**	70
	CD4+ S2	0.400	82
	CD4+ E, M	**0.038**	68
	CD4+ N	**0.013**	68
	Total CD8+	0.062	69
	CD8+ S1	0.518	24
	CD8+ N	0.288	53

<– Lower responses   Higher responses –>

0.1 0.2 0.5 1 2 5

Estimated fold-change in immune response
for severe disease (WHO > 4) (adjusted for age)

C

p=0.00015

Spike IgG AU/ml (ln)

Mild   Moderate   Severe

D

p=0.002

Neutralization titers (ln)

Mild   Moderate   Severe

Disease severity

E

T cells   B cells   Antibodies

Correlation

Negative   Positive