Multifactorial SARS-CoV-2 seroprofiling dissects interdependencies with human coronaviruses and

predicts neutralization activity

Irene A. Abela[1,2*], Chloé Pasin[1,2*], Magdalena Schwarzmüller[1*], Selina Epp[1], Michèle E. Sickmann[1],

Merle M. Schanz[1], Peter Rusert[1], Jacqueline Weber[1], Stefan Schmutz[1], Annette Audigé[1], Liridona

Maliqi[1], Annika Hunziker[1], Maria C. Hesselman[1], Cyrille R. Niklaus[1], Jochen Gottschalk[3], Eméry

Schindler[3], Alexander Wepf[4], Urs Karrer[5], Aline Wolfensberger[2], Silvana K. Rampini[6], Patrick M. Meyer

Sauteur[7], Christoph Berger[7], Michael Huber[1], Jürg Böni[1], Dominique L. Braun[1,2], Maddalena

Marconato[8], Markus G. Manz[8], Beat M. Frey[3], Huldrych F. Günthard[1,2$], Roger D. Kouyos[1,2$], Alexandra

Trkola[1$]

* shared authorship

[$] shared corresponding authors

Affiliations

[1] Institute of Medical Virology, University of Zurich, Switzerland

[2] Division of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, Switzerland

[3] Blood Transfusion Service Zurich, Switzerland

[4] Institute of Laboratory Medicine, Cantonal Hospital Winterthur, Switzerland

[5] Department of Medicine, Cantonal Hospital Winterthur, Switzerland

[6] Department of Internal Medicine, University Hospital Zurich, Switzerland

[7] Division of Infectious Diseases and Hospital Epidemiology, University Children's Hospital Zurich,

Switzerland

[8] Department of Medical Oncology and Hematology, University Hospital and University of Zurich,

Switzerland

Abstract

Definition of SARS-CoV-2 antibody responses is essential to verify protective immunity following

infection and vaccination. Here, we devised a versatile serological test, named ABCORA, that is based

on a multifactorial analysis of SARS-CoV-2 and circulating human coronavirus (HCoV) antibody

responses. Utilizing empirical tailored cut-offs and computational approaches based on training and

validation cohorts comprising pre-pandemic (N=825) and SARS-CoV-2 infected plasma donors (N=389),

we defined several analysis strategies that allow a highly accurate definition of SARS-CoV-2

seroconversion and prediction of neutralization activity. Intriguingly, HCoV reactivity was significantly

higher in SARS-CoV-2 negative donors. Amongst SARS-CoV-2 infected individuals, elevated SARS-CoV-

2 responses were linked to higher HCoV activity suggesting that pre-existing HCoV immunity may

confer protection against SARS-CoV-2 acquisition and promote development of SARS-CoV-2 specific

antibody responses. Deciphering interdependencies between SARS-CoV-2 and HCoV immunity should

be enforced as understanding their impact on infection may allow soliciting cross-protective activities

for broader coronavirus prevention.

Introduction

Monitoring the antibody response to SARS-CoV-2 is critical to define correlates of vaccine protection, differences in susceptibility to infection and in disease severity. The picture of the antibody landscape to SARS-CoV-2 that has thus far evolved is complex. The antibody response to SARS-CoV-2 is rapid, and triggers strong IgM, IgA and IgG responses [1,2]. Both binding and neutralizing responses increase with disease severity and show in part dependence on demographic parameters such as age and gender [3-5]. It remains, however, unclear which factors are independent drivers of antibody responses, reflect severe disease courses or are confounded by other factors including infection length and co-morbidities. Waning IgG binding and neutralizing antibody titers have been described in some but not all convalescent cohorts and may be particularly pronounced in individuals with asymptomatic or mild infection [6-9]. IgG responses to spike (S) glycoprotein may persist longer than to nucleocapsid protein (N)[7,10,11] and can in part undergo affinity maturation post virus clearance [5]. Current serological analyses predominantly focus on measuring reactivity to N, the spike glycoprotein S1 subunit and the ACE2 receptor-binding domain (RBD) in S1 [2,5,12-16]. Antibodies to RBD and the receptor-binding motif within the RBD constitute the main group of neutralizing antibodies, followed by S1 trimer specific, spike N-terminal domain, and spike S2 neutralizing antibodies [16-22]. S1 and RBD binding correlate with neutralizing activity in both natural and vaccine-induced immune responses providing means to estimate the potential for neutralization where neutralization capacity cannot be assessed directly [6,8]. Considering the complex antibody response patterns, possibilities to capture the dynamics of the SARS-CoV-2 response across diverse Immunoglobulin (Ig) classes and SARS-CoV-2 antigens are needed to ascertain sensitive detection of seroconversion and sero-reversion and to establish links to protective, neutralizing activity.

Infections with circulating human coronaviruses (HCoV), alphacoronavirus (HCoV-229E, HCoV-NL63) and betacoronavirus (HCoV-HKU1, HCoV-OC43), are common and contribute considerably to the seasonal respiratory disease burden in humans [23,24]. Despite an overall modest sequence homology between SARS-CoV-2 and circulating HCoVs, several conserved regions exist and antibody cross-

reactivity may occur [25-27]. While dismissed in the diagnostic setting as false-positives [28], cross-reactive

antibodies may bear biological relevance as suggested for S2 cross-neutralizing antibodies [29].

Uncertainty remains, however, whether cross-reactive HCoV antibody responses influence the

evolution of SARS-CoV-2 specific immunity. Positive impact by providing early low affinity memory

responses to build on and mature as well as negative influences following the antigenic sin principle [30]

by boosting non-protective cross-reactive antibodies on the expense of de-novo responses can be

envisaged. In view of this, the definition of pre-existing immunity due to prior infection with HCoVs

will remain complex in clinical diagnosis and strategies to record HCoV and SARS-CoV-2 responses side

by side are needed to fill this knowledge gap.

Here we report on the development of a serological assay that allows multifactorial seroprofiling of

SARS-CoV-2 and HCoV responses at high diagnostic accuracy. Seroprofiling of a large cohort of SARS-

CoV-2 infected and uninfected individuals provided key insights into the interdependencies of HCoV

and SARS-CoV-2 antibody responses. The results highlight a potential protective role of HCoV-specific

antibodies in SARS-CoV-2 acquisition as well as in shaping the SARS-CoV-2 response upon infection.

Results

Multifactorial seroprofiling defines SARS-CoV-2 specific responses

Recognizing the need for comprehensive SARS-CoV-2 serological profiling to elucidate central

questions in SARS-CoV-2 immunity and its interdependencies with HCoV responses, we created a bead-

based multiplex immunoassay using the Luminex technology© to measure specific IgG, IgA and IgM

responses to SARS-CoV-2 RBD, S1, S2 and N (Supplementary Fig. 1). The assay records in total 12 SARS-

CoV-2 specific antibody parameters (4 antigens across 3 Ig classes) with high diagnostic accuracy (see

methods, Supplementary Fig. 1-4 and Supplementary Tables 1, 2) and further includes the S1 protein

of HCoV-HKU1 to screen cross-reactive antibodies alongside SARS-CoV-2 responses. According to the

test's design to monitor antibodies to two coronaviruses, we termed the assay **A**nti**b**ody **COR**onavirus

Assay (ABCORA) 2.0.

Measurements in ABCORA are expressed as median fluorescence intensity (MFI) corrected for

background binding (fold over empty beads, FOE). To distinguish SARS-CoV-2-specific from cross-

reactive antibodies, we defined MFI-FOE thresholds for each of the 12 SARS-CoV-2 antigen and Ig class

combinations based on plasma antibody reactivity in training cohorts of pre-pandemic healthy donor

plasma (Training I, N=573) and in donors with confirmed SARS-CoV-2 infection (Training III, N=175)

(Fig. 1a, Supplementary Table 3). Thresholds were set to minimize false-positives while ensuring

sensitivity for SARS-CoV-2 antibody detection (Fig. 1a, c, Supplementary Table 3). Signal over cut-off

(SOC) values were calculated for each of the 12 SARS-CoV-2 antibody parameters based on the

established threshold values. Positive call criteria were defined to ascertain that in at least two of the

12 antigen and Ig combinations the threshold is reached (Supplementary Table 4). The final threshold

and positive call criteria allowed for a differentiation of partial (only IgM and IgA responses) to full

seroconversion (including IgG responses). In addition, the criteria denote samples with weak reactivity

and/or indeterminate reactivity (Supplementary Table 5).

We further evaluated the ABCORA 2.0 test in clinical samples of pre-pandemic patients with documented, recent HCoV infection (Training II, N= 75, OC43 (N=27), HKU1 (N=17), NL63 (N=22), 229E (N=9)). This group comprised individuals with different underlying severe diseases including immune compromised patients that underwent diagnostic screening for HCoV. We noted an overall lower reactivity with SARS-CoV-2 in this syndromic group but importantly observed no indication of cross-reactivity that affects the ABCORA readout. (Fig. 1a, b). Considering data of all training cohorts (I-III), ABCORA 2.0 exhibited a high sensitivity and specificity, reaching 94.29% sensitivity and 99.07% specificity (Fig. 1c, Supplementary Table 3).

To obtain insight into potential cross-reactivity with HCoV antibodies, we recorded reactivity to the S1 unit of HCoV-HKU1 in addition to the SARS-COV-2 antigens (Fig. 1a). Owing to the high prevalence of HCoV antibodies and the ensuing lack of true-negative controls, we set no thresholds to rate HKU1 reactivity as positive/negative. Overall, SARS-CoV-2 cross-reactivity was low in pre-pandemic samples despite notable HKU1 activity (Fig. 1a). Correlation analysis of SARS-CoV-2 and HKU1 plasma antibody reactivity in SARS-CoV-2 positive donors indicated no correlation in IgG but a substantial cross-reactivity between HKU1 S1 IgM and to lesser extent also IgA (Supplementary Fig. 5a). In pre-pandemic individuals, IgM binding to SARS-CoV-2 and HKU1 was highly correlated whereas IgG and IgA reactivities were less affected (Supplementary Fig. 5b). Of note, individuals with recent HCoV infection showed the highest correlation in IgM between HKU1 and SARS-CoV-2 (Supplementary Fig. 5d). These data underline that cross-reactivity needs to be considered but mostly affects low level responses as evidenced by the measurements in pre-pandemic individuals (Fig. 1a, b, Supplementary Fig. 5).

We verified the performance of ABCORA 2.0 on separate validation cohorts of pre-pandemic healthy adults (N=252), pre-pandemic children (N=169) and individuals with documented SARS-CoV-2 infection (N= 214) (Fig. 1 a, b, Supplementary Table 3). The outcome of this assessment confirmed the validity of the chosen assay criteria. Sensitivity and specificity of the combined negative cohorts were even slightly higher than in the training cohort (Fig. 1c). Of note, when analyzing children and adults in the validation cohort separately, we observed a slightly lower specificity amongst children (98.82%)

compared to adults (99.60%), raising the possibility that cross-reactive activity in children may be more

prevalent than in adults (Fig. 1a). Indeed, pre-pandemic children showed a higher correlation of IgM

HKU1 and SARS-CoV-2 (Supplementary Fig. 5c), highlighting that interpretation of IgM SARS-CoV-2

activity can be complex. Importantly, combining training and validation cohorts of SARS-CoV-2 positive

individuals (N=389) and negative controls (N= 825), ABCORA 2.0 achieved a sensitivity of 94.60% and

a specificity of 99.16%, underlining the high capacity of the test in accurately assessing SARS-CoV-2

seropositivity (Fig. 1c).

Computational analyses maximize specificity and sensitivity of SARS-CoV-2 seroprofiling

We next sought to devise computational analysis extensions to further increase specificity. We

compared two analysis approaches, the first was based on a logistic regression model (ABCORA 2.1),

the second on a random forest model (ABCORA 2.2). Instead of obtaining 12 individual thresholds (one

per antigen and Ig class) as in ABCORA 2.0, both computational models solely estimate the probability

of a sample to be positive by providing a composite result across all 12 measurements and ranking sera

positive or negative (1, 0 classification). Both analysis strategies were established on the training

dataset used for the setup of ABCORA 2.0 (Training I–III, Fig. 1). For the logistic regression ABCORA 2.1,

we grouped SARS-CoV-2 binding activities displaying high correlation (Supplementary Fig. 6a) and

included the mean value of their MFI-FOEs in the model. The analysis on the validation data sets

recorded 93.46% sensitivity and 99.60% specificity in adults and 99.41% specificity in children. When

assessed with the combined training and validation cohort data sets, 93.57% sensitivity and 99.35%

specificity were detected (Fig. 1c, Supplementary Table 3).

The random forest model ABCORA 2.2 included all 12 SARS-CoV-2 responses measured and aggregated

the result of 1000 classification trees. ABCORA 2.2 succeeded in tailoring thresholds tightly to the

training data sets, obtaining classification settings with 100% specificity and 100% sensitivity. This high

specificity was confirmed by analysis of the validation data sets (IV and V) comprising pre-pandemic

samples from healthy adults (specificity 100%) and children (specificity 99.41%). ABCORA 2.2 reached

a sensitivity of 95.33% in the validation set, outperforming both ABCORA 2.0 and 2.1. Of note, positive

calling by ABCORA 2.2 was dominated by IgG responses (Supplementary Fig. 6b). On the combined

training and validation cohorts of SARS-CoV-2 positive individuals (N=389) and negative controls (N=

825), ABCORA 2.2 achieved a striking sensitivity of 97.43% and a specificity of 99.91%. Prompted by

the observed correlation between SARS-CoV-2 and HKU1 IgM and to a lesser extent IgA responses, we

next explored whether incorporation of HKU1 reactivity into the random forest model may further

improve the calling specificity and sensitivity. Indeed, a model that included HKU1 S1 as additional

variable (ABCORA 2.3) increased sensitivity from 97.43% in ABCORA 2.2 to 98.20% (Fig. 1c,

Supplementary Table 3) without reduction of the specificity. Owing to its combined high sensitivity and

specificity, we therefore selected ABCORA 2.3 as the analysis strategy for rating global SARS-CoV-2

seroconversion.

We next verified the accuracy of ABCORA 2.0 and 2.3 in defining positive and negative SARS-CoV-2

immune status utilizing the National Institute for Biological Standards and Control (NIBSC) Anti SARS-

CoV-2 Verification Panel (20/B770)[31]. This verification panel for serology assays includes 23 positive

and 14 negative serum samples and allows direct comparison with other test systems [31]. Both ABCORA

versions showed 100% sensitivity and 100% specificity on the verification plasma panel and compared

favorably to commercial assay systems (Fig. 1d, Supplementary Table 6). To cross-reference these

external verification results, we next compared the sensitivity of the ABCORA tests and three

commercial serology test systems on a subset of the SARS-CoV-2 positive training cohort (cohort III,

N=171) for which sufficient material remained. Assays targeting the N protein (Elecsys® Anti-SARS-

CoV-2 (Roche Diagnostics GmbH)), the RBD region of the S protein (Elecsys® Anti-SARS-CoV-2 S assay

(Roche Diagnostics GmbH)), and the S1 subunit (EUROIMMUN Anti-SARS-CoV-2 ELISA (IgG)) were

included. The results confirmed the analysis on the international NIBSC 20/B770 plasma panel, with

ABCORA 2.0 and ABCORA 2.3 showing the highest sensitivity amongst the tested assays

(Supplementary Table 7).

We thus conclude that ABCORA 2.0 seroprofiling in combination with ABCORA 2.3 defines positivity

with the highest specificity and sensitivity. The individual antigen response evaluation by ABCORA 2.0

defines the stage of seroconversion status based on individual IgM, IgA and IgG cut-off values and

thereby complements and maximizes the information that can be obtained by ABCORA 2 seroprofiling.

Predicting SARS-CoV-2 neutralization based on ABCORA seroprofiling

Determining neutralization activity is critical to gauge protective immunity. While neutralization can

be directly measured with a range of authentic virus or pseudovirus SARS-CoV-2 neutralization tests

[5,22,32], applying direct binding or competition tests as surrogate for neutralization activity remains of

high interest as these test systems are faster, less labor-intensive, widely applicable and scalable [32,33].

In particular, S1 and RBD binding has been shown to correlate well with neutralization activity [5,8,19,32-

36]. To explore neutralization predictors based on ABCORA 2.0, we probed in a first step the capacity of

ABCORA to derive quantitative S1 and RBD readouts in a subset of SARS-CoV-2 positive patients (N=72).

ABCORA 2.0 measurements of serially diluted plasma were conducted to derive 50% effective

concentrations (EC50, expressed as reciprocal plasma dilution) and area under the curve values (AUC

expressed as MFI) for all 12 SARS-CoV-2 parameters (Fig. 2a, b). In addition, we quantified SARS-CoV-

2 RBD and S1 responses via the RBD specific mAb CR3022 [37] (Fig. 2c) and the WHO International

Standard Anti-SARS-CoV-2 Immunoglobulin NIBSC 20/136[31] (Supplementary Fig. 7b, Supplementary

Table 8). For this we quantified the respective antibody content of a positive control SARS-CoV-2 donor

pool included in all ABCORA measurements and expressed the antibody content of individual plasma

samples in relation to it (Fig. 2c, Supplementary Fig. 7). We then probed which of the ABCORA

quantitative readouts correlated best with each other, the basic readout of ABCORA 2.0 (MFI-FOE at

plasma dilution 1/100), and the quantitative Roche Elecsys S test (U/ml) (Fig. 2c, Supplementary Fig.

7). In addition to the individual antigen parameters, we also considered cumulative response values.

These were total spike reactivity (sum of RBD, S1, S2 across all Ig classes), Ig class spike reactivity (sum

of S1, RBD, S2 for one isotype) and antigen specific reactivity (sum of all Ig classes for one antigen). We

observed a genuinely good correlation across the diverse spike parameters tested (Fig. 2d). The

notable exception were classical EC50 values, which showed no to weak correlation across all

parameters including the commercial test. Interestingly, AUC values, which in contrast to EC50 are a

composite measure of concentration and signal strength, performed well. Of note, we observed highly

variable SARS-CoV-2 antibody dose response curves, reaching in our cohort individual plateaus over a

4-log range (Supplementary Fig. 7). These plateaus are respected in the AUC readout, and are also

recorded by the basic MFI-FOE ABCORA readout at 1/100 plasma dilution, but are not considered in

EC50 determinations. Indeed the basic MFI-FOE showed a high correlation with the quantitative

readouts across the tested variables including the quantitative commercial Roche Elecsys S test.

Based on these results, we concluded that the MFI-FOE readout at 1/100 provides an excellent

estimate for the S1 and RBD antibody content in plasma and that this straightforward and easily

obtainable readout, which only requires a single plasma dilution to be tested, can be used as a proxy

for quantification. We therefore employed the basic MFI-FOE in a next step to define neutralization

predictors.

Neutralization activity to Wuhan-Hu-1 in SARS-CoV-2 positive individuals (N=467) using an established

SARS-CoV-2 pseudovirus neutralization test [5,22,32] revealed a broad range of 50% neutralizing titers

(NT50) (post positive RT-PCR Fig. 3a (N=369), post onset of symptoms, Supplementary Fig. 8a (N=333)),

in line with previous findings [5,8]. Early in infection (within 30 days of positive RT-PCR) neutralization

titers were significantly higher (p<0.001) and correlated better with binding parameters. As expected,

IgG responses to spike antigens showed the highest correlation with neutralization activity (Fig. 3b,

Supplementary Fig. 8-9). We next grouped patients based on the population into high (NT50 >250,

N=135) and no or low neutralizers (NT50 <250, N=332)[38] (Fig. 4a) and compared the prediction ability

of six different classification models to assign individuals based on their ABCORA 2.0 binding patterns

to these groups. Univariable logistic regression (ULR) models included only one variable: either the

mean of MFI-FOE S1 reactivities (ULR-S1), or the mean of MFI-FOE RBD reactivities (ULR-RBD). A

multivariable logistic regression (MLR) included both S1 and RBD mean reactivities. The additional

models included all 12 antigen reactivities measured in ABCORA and comprised a random forest approach and two MLR strategies based on principle component analysis (PCA, 2 and 4 first axis). Models were compared based on AUC and the BIC (Bayesian information criterion[39]) by cross validation (Fig. 4b, c). All models performed similarly, with the univariable model based on the mean of S1 reactivities (ULR-S1) yielding the best BIC value. Receiver operating curve (ROC) analysis based on ULR-S1 showed a good capacity in predicting neutralization status yielding AUC 0.90 (N= 467, Fig. 4d). Exploring different cutoffs to balance sensitivity and specificity keeping both above 80%, we chose to assign samples to the high neutralizers group if its predicted probability was above 70%. This corresponds to an 83% specificity in correctly assigning non–neutralizers and 80% sensitivity in assigning neutralizers (Fig. 4d, e). To increase the utility of the ULR-S1 prediction model for clinical diagnostics, we devised a modified neutralization prediction model ULR-S1-SOC based on the SOC values reported for ABCORA 2.0. At 70% predicted probability, ULR-S1-SOC delivers neutralization prediction at similar sensitivity (81%) and specificity (81%) by examining if the composite S1 SOC value (sum of S1 SOC values for IgG, IgA and IgM) is below or above 9.7 (Fig. 4f). Thus, based on ULR-S1-SOC neutralization activity can be ranked directly from the basic SOC readout. We therefore conclude that the basic readout in ABCORA 2.0 can deliver a reliable prediction of high neutralization activity.

Resolution of temporal antibody dynamics by ABCORA seroprofiling

Cross-sectional analysis of antibody reactivity post SARS-CoV-2 diagnosis by RT-PCR (N=369) and post onset of symptoms (N=333) underlined the capacity of ABCORA seroprofiling to dissect onset, peak and waning of SARS-CoV-2 antibody responses (Fig. 5a, b, Supplementary Fig. 10). In individuals with known date of first SARS-CoV-2 RT-PCR diagnosis or onset of symptoms, ABCORA 2.3 detected early seroconversion in 98% (48 of 49) and 100% (9 of 9) of individuals within 7 days post RT-PCR and onset of symptoms, respectively. Besides IgM and IgA reactivity, IgG responses were readily detectable in ABCORA 2.0 after a few days of infection (Supplementary Fig. 10).

Longitudinal assessment of a cohort of convalescent patients up to 11 months post infection (251

measurements on 120 patients) highlighted the temporal dynamics of SARS-CoV-2 binding antibodies.

We estimated the decay of binding reactivity employing a linear mixed model and identified a

significant reduction in RBD, S1, and N in all Ig subtypes (Fig. 5b) with half-lives ranging from 116 to

836 days, with IgG N titers decaying the fastest, in line with previous reports [10]. Half-lives of the

neutralization relevant IgG responses to RBD and S1 where 209 and 323 days, respectively. Intriguingly,

the kinetics of neutralizing antibodies did not mirror the decay rates observed for binding antibodies.

Neutralization activity decreased overall at a slower rate, with a half-life of 410 days (Fig 5c). This was

in part due to a mixed reactivity pattern with some individuals showing an increase in neutralization

activity post positive SARS-CoV-2 RT-PCR [5], while neutralization activity in others rapidly decayed

(Supplementary Fig. 10c).

Interdependencies of SARS-CoV-2 and HCoV immunity

To enable a more detailed analysis of HCoV antibody reactivity alongside SARS-CoV-2, we expanded

the bead antigen array to include S1 proteins of all four circulating HCoVs (HCoV-NL63, HCoV-229E,

HCoV-HKU1, HCoV-OC43) (Supplementary Fig. 11a). According to its capacity to monitor antibodies to

five coronaviruses we termed the assay ABCORA 5.0 and trained and validated it on the same cohorts

as ABCORA 2.0 (Fig. 1). To allow direct comparison with ABCORA 2.0 and use of the neutralization

prediction models, we used the threshold-/SOC-based analysis settings of ABCORA 2.0 also for ABCORA

5.0. Based on ABCORA 5.0 measurements of training cohorts I-III, we devised two random forest-based

analysis models. ABCORA 5.4 included solely the 12 SARS-CoV-2 parameters, ABCORA 5.5 included in

addition the S1 HCoV measurements adding up to 24 parameters in total. In analogy to ABCORA 2.3,

incorporation of HCoV reactivity into the model was advantageous. ABCORA 5.5 provided the highest

sensitivity and specificity amongst the analysis algorithms probed ABCORA 5.0 (Supplementary Fig.

11b, Supplementary Table 10).

Interdependencies between antibody reactivity to the four HCoVs and SARS-CoV-2 mirrored what we previously observed for HKU1 with a particular high correlation of IgM reactivity of SARS-CoV-2 and HCoVs in pre-pandemic individuals, particularly in those with recent HCoV infection (Supplementary Fig. 12). HCoV infections are frequent but subject to seasonality and prevalence of individual HCoV infections fluctuates [40,41]. In line with this, the prevalence of HCoV responses measured by ABCORA 5.0 in local blood donors in January 2019 (N=285), May 2019 (N=288), and January 2020 (N=252) varied considerably (Fig. 6). To enable a time-controlled comparison of HCoV reactivity between SARS-CoV-2-infected and healthy donors, we screened blood donors from May 2020 (N=672), when SARS-CoV-2 prevalence was estimated below 2% in Zurich, Switzerland [42], by ABCORA 2.0/5.0 and excluded all samples with SARS-CoV-2 reactivity. The residual May 2020 cohort (N= 653) formed a pandemic, healthy donor control group. Interestingly, HCoV reactivity patterns in 2019 and 2020 differed substantially as assessed by one-way ANOVA, with January 2020 showing the comparatively lowest and May 2020 the highest IgA and IgG reactivity, which may indicate a later onset of an HCoV epidemic in 2020 compared to 2019 (t-tests of May 2020 versus other groups shown in Fig. 6).

Most intriguingly, a time-matched analysis comparing May 2020 healthy donors with SARS-CoV-2-positive patients sampled in April, May and June 2020 (N=65) revealed significantly lower HCoV reactivity in SARS-CoV-2 positive patients (Fig. 7a). This pattern was also evident when we extended the analysis to include the full cohort of SARS-CoV-2 infected individuals measured with ABCORA 5.0 (N=389, sampled from March 2020 to February 2021, Supplementary Fig. 13). Overall, this raised the possibility that pre-existing immune responses to HCoVs may to a certain degree protect against SARS-CoV-2 infection. We therefore investigated whether HCoV responses are linked to the evolution of SARS-CoV-2 antibodies in patients with known date of first positive RT-PCR (SARS-CoV-2 patients measured in ABCORA 5.0 with time since positive RT-PCR<60 days, N=204). Associations were investigated with a linear regression model adjusted for age, gender, time since positive RT-PCR and HCoV reactivity. To stratify HCoV reactivity into high and low HCoV activity, median logMFI-FOE were defined for each HCoV and antibody class. LogMFI-FOE higher than the corresponding median for at

least three HCoVs (HKU1, OC43, NL63 or 229E) in a specific Ig class were ranked as having high HCoV

activity within this class. First, only reactivities among the same antibody class were explored in the

model (i.e. HCoV IgG high on SARS-CoV-2 IgGs). We observed exceptionally strong interdependencies

for IgA and IgM responses to SARS-CoV-2, which all were significantly higher in individuals with high

HCoV reactivity (Fig. 7b). This strongly suggests that pre-existing HCoV immunity may provide an

advantage in mounting SARS-CoV-2 responses. Interdependencies between HCoV IgG and SARS-CoV-

2 specific IgG were only observed for the S2 response. To explore if SARS-CoV-2 IgG may build on recent

HCoV IgA and IgM responses we next probed whether HCoV IgM and IgA are linked to elevated SARS-

CoV-2 specific IgG levels. While no effect was evident for IgM, we observed a significant association of

high HCoV IgA activity on all four measured SARS-CoV-2 responses (Fig. 7c, d). This strongly suggests

that recent HCoV infection has a beneficial effect on mounting SARS-CoV-2 antibody responses.

Discussion

Definition of SARS-CoV-2 immunity post vaccination and infection is of immediate importance [43-45].

Deciphering antibody correlates of SARS-CoV-2 protection and monitoring vaccine responsiveness are

challenging tasks ahead. The magnitude and longevity of protective antibody responses to natural

infection and of different vaccines need to be examined to understand parameters that shape

protective responses and guide decisions on re-vaccination in non-responders and immunization

against novel arising SARS-CoV-2 variants [46]. Likewise, creating means to serologically distinguish

between de novo infection, re-infection, and vaccine responses, their durability and failures will be

critical for clinical diagnosis.

Here we demonstrate the high utility of multi-parameter seroprofiling in addressing key issues in

defining SARS-CoV-2 immunity. We designed SARS-CoV-2 seroprofiling tests, termed ABCORA, to

address research and clinical diagnostics needs. Simultaneous recording of antibody responses to a

range of SARS-CoV-2 antigens and different Ig classes with ABCORA seroprofiling yields an extensive

picture on the SARS-CoV-2 serology status in a single assessment. High accuracy in defining SARS-CoV-

2 seropositivity and staging of seroconversion was achieved by a computational analysis. Validation

and verification of the performance of ABCORA against internal and external standards proved its

exceptional performance enabling application in clinical diagnostics and research.

Measuring neutralization will remain the ultimate goal to determine protective immunity to infection

and immunization. Protective activity against novel arising SARS-CoV-2 variants needs to be

determined [46-51]. Collecting information on neutralizing activity in large population groups is however

challenging as cell-based neutralization assays are not in wide diagnostic use and have limited turnover

and scalability. Means to derive estimates of neutralization activity based on comparatively simpler

binding assays would therefore provide an enormous advantage as previously suggested [52-54]. As we

demonstrate, computational modeling allows predicting plasma neutralization capacity from ABCORA

seroprofiling results, obliterating the need to conduct labor-intensive and costly neutralization tests in

clinical diagnostics or in large-scale research-based screens. The same principle can be applied in future

assay extensions to predict neutralization activity against novel SARS-CoV-2 variants.

The successful development of a sensitive seroprofiling test enabled a precise analysis of SARS-CoV-2

antibody dynamics and their interplay with HCoV responses. We utilized two ABCORA assay versions

that both measured HCoV reactivity alongside the 12 SARS-CoV-2 parameters. ABCORA 2 included the

S1 antigen of HKU1. ABCORA 5 included S1 of all four circulating HCoVs. Notably, computational

models that included the HCoV measurements allowed a higher precision in determining SARS-CoV-2

seropositivity, highlighting interdependencies between HCoV and SARS-CoV-2 responses that need to

be resolved. Recording reactivity against all four HCoVs in SARS-CoV-2 uninfected and infected

individuals we observed intriguing associations. Uninfected individuals displayed higher HCoV

reactivity compared to infected individuals. However, amongst persons with documented SARS-CoV-2

infection, those with high HCoV reactivity developed higher SARS-CoV-2 antibody levels. While

association studies as ours cannot formally define causality, the implications of our findings are

evident: Prior immunity to HCoV may protect to some extent against SARS-CoV-2 acquisition and may

provide a boost to the development of SARS-CoV-2 specific immunity. While we solely measured

antibody responses, a potential protective effect of HCoV immunity against SARS-CoV-2 acquisition

should not be viewed restricted to antibody activity. Antibodies and cellular immunity may both be

relevant and act in concert [55-57]. Alternatively, antibody responses measured in the present study may

solely document recent HCoV infection and deliver a surrogate measurement of other protective HCoV

responses. The link between higher HCoV and SARS-CoV-2 reactivity in infected individuals is

particularly intriguing. Strongest effects were seen for IgM and IgA HCoV responses, suggesting that

recent HCoV immunity provides an early boost to SARS-CoV-2 antibody development. Whether this is

due to cross-reactive B cell responses on which the SARS-CoV-2 immunity builds on and matures or

whether cross-reactive T helper activities play a role will be important to resolve in forth-coming

studies. Likewise the precise role and timing of the HCoV responses that shape SARS-CoV-2 antibody

responses needs to be defined. Early, low affinity HCoV responses may have a positive impact for the

evolution of SARS-CoV-2 immunity by providing an immune memory to build on, while the boosting of

non-protective cross-reactive HCoV antibodies may have a negative impact following the antigenic sin

principle [30].

Our findings have multiple implications and warrant further investigation of the interplay between

HCoV and SARS-CoV-2 immunity. A modest protective effect by HCoV immunity would be a plausible

explanation for the high proportion of asymptomatic and mild SARS-CoV-2 infections [58,59]. Even more

intriguing are future perspectives. As others and we have shown, SARS-CoV-2 and HCoV immunity to

infection is often not long-lasting (Fig. 5, Supplementary Fig. 10) [5,60], a limitation that SARS-CoV-2

vaccines hope to overcome. Should SARS-CoV-2 responses in turn provide a degree of defense against

HCoV infection, broad protection against coronaviruses may be in reach.

Methods

Human specimen

Serum and plasma samples collected pre and post emergence of SARS-CoV-2 in Switzerland (pre and post February 2020, respectively) were included. No patient enrollment was conducted for the present study. All experiments involving samples from human donors were conducted with the approval of the responsible local ethics committee Zurich, Switzerland (BASEC Nrs 2020-01327, 2020-00363; Req-2021-00437; 2020-00787), in accordance with the provisions of the Declaration of Helsinki and the Good Clinical Practice guidelines of the International Conference on Harmonisation. Samples were obtained from the following sources: i) Zurich blood donation services (ZHBDS): Anonymized healthy adult plasma from pre-pandemic time points (January 2019, May 2019 and January 2020) and from the first wave of the pandemic in Zurich, Switzerland (May 2020) were provided by the ZHBDS internal serum repository (BASEC 2020-00363). ii) Anonymized leftover specimens from routine diagnostics at the Institute of Medical Virology, University of Zurich, the University Children Hospital Zurich and the Cantonal Hospital Winterthur (BASEC Nrs 2020-01327, Req-2021-00437). iii) Healthcare workers with RT-PCR confirmed SARS-CoV-2 infection participating in a study at the University Hospital Zurich (BASEC 2020-00363). iv) Male plasma donors participating in a SARS-CoV-2 plasma therapy study conducted at the University Hospital Zurich (CPT-ZHP, Swissmedic 2020TpP1004; BASEC 2020-00787). Pre-pandemic (SARS-CoV-2 negative, N=825) and confirmed SARS-CoV-2 positive samples (N=389) were divided into training and validation cohorts (Supplementary Table 3). The SARS-CoV-2 training cohort (N=175) included plasma collected during infection (N=114) and convalescence (N=61). Per donor only one sampling time point was included, longitudinal samples of donors included in the training cohort were not included in the validation cohort to ascertain independence when assessing the sensitivity and specificity of the different diagnostic methods. The SARS-CoV-2 validation cohort (N=214), comprised plasma collected during infection (N=90, one sampling time point per donor) and convalescence (N=124, 79 convalescent patients with 1-4 longitudinal samples). Multiple time points of convalescent patients were included in the validation data set to capture a wide spectrum of waning

antibody titers. Cross-sectional analysis was based on samples with known time since positive RT-PCR (N=369) or known time since symptom onset (N=333), both including the longitudinal analysis observations. Both, time since positive RT-PCR and time since onset were known for 330 samples, with a median time of 3 days between symptom onset and RT-PCR (1st-3rd quartile: 1-7 days). Longitudinal analysis of antibody reactivity was based on 251 observations from 120 convalescent patient with known time since positive RT-PCR and time since symptom onset. Neutralization was measured on 467 SARS-CoV-2 RT-PCR positive samples (N=369 with known time since positive RT-PCR, N=333 with known time since symptom onset).

We further evaluated cross-reactivity ABCORA 2.0 and 5.0 in left-over plasma from routine diagnostics in a pre-pandemic control group with documented, recent HCoV infection (Training II, N= 75, OC43 (N=27), HKU1 (N=17), NL63 (N=22), 229E (N=9)). Circulating HCoV are commonly only screened for in in hospitalized, severe respiratory infections and immune compromised individuals who routinely undergo a broad screening for respiratory infections. Hence, in this patient group both, reduced antibody reactivity due to immune compromising or elevated HCoV antibody reactivity due to recent or recurring HCoV infection may occur. As this group is diagnostically relevant we considered it prudent to include this cohort as Training II data set to verify if cross-reactivity with SARS-CoV-2 in ABCORA occurs in this setting. Training II data were not included in the threshold definition to not over-represent individuals with severe illness. This HCoV infected group displayed overall lower reactivity with SARS-CoV-2 than plasma from healthy adults but importantly showed no indication of cross-reactivity (Fig. 1a, b). Pandemic samples from anonymous blood donors with unknown SARS-CoV-2 status collected in May 2020 (N=672) were not included in training and validation cohorts.

Reagents and cell lines

His-tagged SARS-CoV-2-derived antigens (receptor binding domain (RBD), subunit S1 (S1), subunit S2 (S2), nucleocapsid protein (N)) and S1 of the four circulating HCoVs (HKU1, OC43, NL63, 229E) were

purchased from Sino Biological Europe GmbH, Eschborn, Germany (Supplementary Table 12). Sources,

specifics and concentration of detection and control antibodies and sera used for ABCORA and

neutralization tests are listed in Supplementary Table 13. 293-T cells were obtained from the American

Type Culture Collection (ATCC CRL-11268) [61]. HT1080/ACE2cl.14 cells [32] were kindly provided by P.

Bieniasz, Rockefeller University, NY. Both cell lines were cultured in DMEM containing 10% FCS.

Design of multiplex bead assay ABCORA 2.0

We established two bead-based multiplexed SARS-CoV-2 immunoassays (ABCORA 2.0 and ABCORA

5.0) that included a range of SARS-CoV-2 and HCoV antigens (Sino Biological Europe GmbH, Eschborn,

Germany, Supplementary Table 12). Four SARS-CoV-2 antigens - RBD, S1, S2 and N - were included in

both ABCORA 2.0 and ABCORA 5.0. ABCORA 2.0 included in addition S1 of HCoV-HKU1, ABCORA 5.0

included S1 of all circulating HCoVs (HCoV-NL63, HCoV-229E, HCoV-HKU1, HCoV-OC43). In brief,

individual MagPlex beads (Luminex Corporation, Austin, TX) with unique fluorescent bead regions were

chosen for each antigen, beads were coupled and mixtures of antigen-coupled beads incubated with

patient plasma in a 96-well plate set-up. Median Fluorescent Intensity (MFI) of bead-bound plasma

antibodies were measured utilizing a FlexMap 3D reader (Luminex Corporation, Austin, TX). We

designed the assay to fulfil the following criteria: i) high specificity, sensitivity and reproducibility, ii)

flexible multiplex design that allows straightforward addition and/or alteration of antigens; iii) wide

dynamic range; iv) optional quantification of antibody responses; v) optional recording of antibody

responses to HCoVs and vi) use in routine diagnostics and research.

We chose a sterically orientation capture via anti-His antibodies to ensure a homogenous antigen

display. Therefore, carboxylated MagPlex beads (Luminex Corporation, Austin, TX) were coupled with

anti-His antibody (Sino Biological Europe GmbH, Eschborn, Germany, Supplementary Table 10) and

then coupled with His-tagged antigens using Bio-Plex Amine coupling (Bio-Rad Laboratories AG,

Cressier, Switzerland) according to the manufacturer's instructions and as described [62].

Serum/plasma titration is in general considered the most accurate strategy to retrieve quantitative

information on antibody reactivity. However, in diagnostic use tests ideally should deliver (semi)-

quantitative information from a single serum dilution to permit a sufficient throughput. The finalized

assay conditions covered a 2-log MFI range across all probed antigen-Ig combinations (Supplementary

Fig. 2). Ratifying the validity of using a single plasma dilution, we confirmed that plasma from SARS-

CoV-2 positive patients and pre-pandemic SARS-CoV-2 negative plasma samples show optimal dose

response curves over a wide plasma dilution range (Supplementary Fig. 2d). Importantly, a 1/100

dilution of plasma was in all cases close to the maximum signal, underlining that increasing plasma

concentration would not increase signal intensity but rather endanger decreasing signals due to

prozone effects (Supplementary Fig. 2d).

Maximal anti-His antibody loading was achieved at 5 µg antibody per million beads (Supplementary

Fig. 2) and used as standard coupling condition. In the final protocol, five million anti-His antibody

coupled magnetic beads were incubated with His-tagged antigens diluted in PBS at a concentration of

320 nM. Phycoerythrin (PE)-labeled secondary antibodies specific to IgG, IgA or IgM were used as

detector antibodies (Supplementary Table 13). Quality control of the antigen loading was performed

by incubating the beads with monoclonal antibodies targeting the corresponding CoV-derived antigen

as detailed in Supplementary Table 12. Analysis was performed with the FlexMap 3D reader (Luminex

Corporation, Austin, TX) with the acquisition of at least of 50 beads per bead region. Results are

recorded as MFI per bead region.

Several control measures were installed to ascertain inter- and intra-assay performance. To ascertain

a low assay-to-assay variability, large batches of individual antigen-loaded beads were prepared and

frozen in aliquots until use at -20°C to circumvent decay of the antigen-coupled beads (Supplementary

Fig. 1). Individual coupled beads were mixed on the test day to yield the required antigen bead cocktail.

Cocktails contained 60 beads per bead region per µl. In addition to the SARS-CoV-2 and HCoV bead

regions, each cocktail included an empty bead region (no antigen coupled) to control for unspecific

binding. Quality control and validation procedures for the FlexMap 3D instrument were done on each

day of experiment according to manufacturer's instructions. The variability of the assay was analyzed

as described previously [62]. In brief, plasma samples from 20 RT-PCR confirmed SARS-CoV-2 infected

patients were pooled and tested over a range of seven dilutions in 31 different titrations performed

on 10 different days (Supplementary Fig. 3). Across all antigens and Ig classes, signals were retained

over the test period of 25 days post bead coupling. Coefficients of variation (CV) of the binding signal

across titrations of the 12 antigen-Ig class combinations proved low (range: 0.010-0.128, median 0.059,

Supplementary Fig. 4c, d, Supplementary Tables 1, 2). Same-day and day-to day variability proved low

and comparable (Supplementary Fig. 4e, f). Below a 1/100 plasma concentration, CV increased

markedly (Supplementary Fig. 4d), defining 1/100 as highest concentration (lowest plasma dilution) to

be tested in the assay. A 1/100 plasma dilution was thus defined as the basic dilution for screening

plasma in ABCORA 2.0 when a qualitative (i.e. presence or absence of SARS-CoV-2 specific antibodies)

or semi-quantitative (i.e. MFI signal intensity) readout is required.

All ABCORA measurements were derived from single measurements unless stated otherwise. To

measure SARS-CoV-2 specific antibodies in patient plasma, heat inactivated plasma (1 h at 56°C) was

diluted 1/100 in PBS-BSA 1% unless otherwise stated. 50 µl diluted plasma were incubated with 50 µl

of the ABCORA antigen bead cocktail for 30 minutes at room temperature in 96-well plates, washed

three times with PBS-BSA 1% and incubated in separate reactions with phycoerythrin (PE)-labeled

detector antibodies for IgG, IgA or IgM at a final concentration of 1/500 in PBS-BSA 1%. This dilution

was previously defined by titration of the detector antibodies to yield optimal MFI signals. After 45

minutes of incubation at room temperature, beads were washed three times with PBS-BSA 1% and

analyzed in 96-well plates on the FlexMap 3D reader (Luminex Corporation, Austin, TX). A minimum of

50 bead reads per antigen was acquired.

To control for genuine cross-reactive antibodies, each plasma sample was assessed with beads without

antigen (empty bead control) in combination which each detector antibody. For analysis, raw MFI

values were transformed to MFI-FOE to correct for background binding. We established mean empty

bead MFI-FOE for IgG, IgA and IgM of pre-pandemic healthy donors (N=1016) and set the mean MFI-

FOE + 4x standard deviation as threshold for the empty bead control. In absolute levels, these

thresholds amounted to MFI-FOE 41.58 (IgG), 55.91 (IgA) and 269.47 (IgM). Measurements for which

the empty bead control recorded values above this threshold were considered invalid and repeated.

Each Luminex analysis 96-well plate was set up to contain the same set of control samples

(Supplementary Fig. 1c), namely 7 serial 4-fold dilutions of a SARS-CoV-2 positive control donor pool

(N=20 donors, starting dilution 1/100, Supplementary Fig. 2d) and a pre-pandemic healthy donor pool

(dilution 1/100, N=20 donors, Supplementary Fig. 2d). These positive and negative controls allow to

control assay performance across independent measurements and in addition enable retrospective

standardization against external controls if needed.

Definition of SARS-CoV-2 seropositivity in ABCORA 2.0 and ABCORA 5.0

Individual MFI-FOE cut-off values for SARS-CoV-2 specific responses for each of the 12

antigen/immunoglobulin combinations were set for ABCORA 2.0 based on the training cohorts

(Supplementary Table 4). Threshold levels were set to reach an overall specificity above 99% and

included levels for border-line reactivity for IgG RBD, IgG S1 and IgG N to allow also modest antibody

reactivity to these antigens to be examined. MFI-FOE reads of individual samples were transformed

into signal-over cut-off (SOC) values (MFI-FOE/threshold). SOC values are used for assessing positive

reactivity for each individual antigen-antibody class combination, with SOC>1 denoting positive

reactivity, SOC<1 denoting negative reactivity. When setting individual thresholds, it must be

considered that for each of the 12 probed activities cross-reactivities may occur. With 12 individual

SOC parameters recorded, overall specificity will decrease if any positive SOC independently suffices

to rate a sample overall as SARS-CoV-2 antibody positive. To exemplify: Assuming independent

responses, even a high 99% specificity for each antigen will add up to an overall low 88% specificity

across the entire assay. We thus required for SARS-CoV-2 positive calling in ABCORA 2.0 a minimum of

two specificities to reach activity above threshold. The combined SOC values used to define the overall

serostatus of a given sample are detailed in Supplementary Table 5. For IgG RBD, S1 and N, for which

we also recorded border-line SOC activity, we allowed for a combination of 1 antigen reactivity SOC>1,

the second reactivity SOC> border line. The final threshold and positive call criteria allowed for a

differentiation of partial (partial (early seroconversion with only IgM and IgA responses) to full

seroconversion (including IgG responses) (Supplementary Table 5). In addition, the criteria denote

samples with weak reactivity and/or indeterminate reactivity (Supplementary Table 5).

To ease comparison between ABCORA 2.0 and ABCORA 5.0 the same threshold cut-offs were used for

ABCORA 5.0. We chose not to create specific cut-off thresholds for HCoV antibody reactivity as an

accurate definition of a negative response is complex due to the wide-spread exposure to HCoVs and

considerable antibody cross-reactivity between them. HCoV responses were however included in the

statistical analyses as MFI-FOE values.

Definition of SARS-CoV-2 seropositivity using logistic regression classification

Classification of seropositive versus seronegative samples in ABCORA 2.1 was realized using logistic

regression. The identical training and validation data used for the establishment for ABCORA 2.0 were

used. As the ABCORA 2.0 binding reactivities were highly correlated, we included the following

variables in the model (Supplementary Fig. 6a): the mean value of all IgG MFI-FOE responses (RBD, S1,

S2, N), the mean value of the IgA MFI-FOE responses against RBD, S1 and S2, and the mean value of

the IgM MFI FOE responses against RBD, S1 and S2. IgA and IgM responses to N were excluded as they

were not clustering with the other responses of the same Ig class (Supplementary Fig. 6a). The logistic

regression was used to estimate and predict the probability of a given sample to be positive (p) as

follows:

$$p = \frac{\exp(\beta0 + \beta G * mean(IgG:RBD,S1,S2,N) + \beta A * mean(IgA:RBD,S1,S2) + \beta M * mean(IgM:RBD,S1,S2))}{1 + \exp(\beta0 + \beta G * mean(IgG:RBD,S1,S2,N) + \beta A * mean(IgA:RBD,S1,S2) + \beta M * mean(IgM:RBD,S1,S2))}$$

Parameters $\beta0$, βG, βA and βM were estimed on the training dataset. A sample was then defined as

positive if its predicted probability of being positive was above a threshold c'. This threshold was

defined as to obtain a specificity of at least 0.99 and maximal sensitivity on the training dataset

(similarly to c for the random forest). In summary, in ABCORA 2.1, any new sample is defined as

seropositive if its probability of being seropositive as estimated by the logistic regression is above c'.

Analyses were performed in R version 3.6.3.

Definition of SARS-CoV-2 seropositivity using random forest classification

Classification of seropositive versus seronegative samples in context of ABCORA 2.0 and ABCORA 5.0

was realized using a random forest approach following the basic setup of random forests as described

in[63]. The random forest itself was built of an ensemble of 1000 classification trees using MFI-FOE

responses (IgA, IgG and IgM against RBD, S1, S2, N). The probability of a sample being positive as

predicted by the random forest is the average of the probabilities over all 1000 trees. Finally, a sample

is defined as positive if its probability of being positive is above a threshold c, which is defined as to

obtain a specificity of at least 0.99 and a maximal sensitivity on the training dataset. In summary, any

new sample is defined as seropositive if its probability of being seropositive as estimated by the

random forest is above the threshold c. We conducted a series of random forest analyses that

considered either only SARS-CoV-2 responses or SARS-CoV-2 and HCoV responses in ABCORA 2.0 and

ABCORA 5.0: ABCORA 2.2 and ABCORA 2.3 were trained and used for prediction on ABCORA 2.0 data

and included only SARS-CoV-2 responses or SARS-CoV-2 and HKU1 responses, respectively. ABCORA

5.4 (SARS-CoV-2 responses only) and ABCORA 5.5 (SARS-CoV-2 and HCoV responses) were trained on

ABCORA 5.0 data. Details on the data inclusion for the respective models are listed in Supplementary

Table 3 and Supplementary Table 10. Analyses were performed in R version 3.6.3 using packages

random Forest and ranger [64-67].

Validation and verification using external controls

We used the Anti-SARS-CoV-2 Verification Panel for Serology Assays (NIBSC code: 20/B770, NIBSC) to verify the performance of the ABCORA 2.0 and ABCORA 2.3 test. Serum samples of the verification panel measured by ABCORA 2.0/2.3 as described and results compared with the results of commercially available assays reported by the NIBSC ([31] and Supplementary Table 6). We further verified the sensitivity of the ABCORA 2 test in detecting SARS-CoV-2 infection in a direct comparison with commercial tests. Antibody status of plasma from SARS-CoV-2 positive individuals (N=171) were analyzed with the following test systems: Included test systems targeted the N protein (Elecsys® Anti-SARS-CoV-2 (Roche Diagnostics GmbH)), the RBD region of the S protein (Elecsys® Anti-SARS-CoV-2 S assay (Roche Diagnostics GmbH)), and the S1 subunit (EUROIMMUN Anti-SARS-CoV-2 ELISA (IgG)) (Supplementary Table 7). All assays were performed according to the manufacturer's instructions in the diagnostics unit of the Institute of Medical Virology, University of Zurich, Switzerland.

SARS-CoV-2 binding antibody titers

To define binding antibody titers, 8 serial 4-fold dilutions starting with a 1/25 dilution of plasma were prepared and measured in ABCORA 2.0. To derive quantitative information, MFI values were corrected for background activity (MFI-empty bead control) and we defined the area under the MFI curve (AUC) across the dilution series for each antigen-Ig combination. As a second quantitative readout, we calculated 50% effective titer concentrations (EC50) using a four-parameter logistic curve (y=Bottom+(Top-Bottom)/(1+10^((logEC50-X)*HillSlope)).

Quantification of SARS-CoV-2 S1 and RBD activity

We used two approaches to standardize SARS-CoV-2 S1 and RBD activity. The first was based on the S1/RBD specific antibody CR3022 ([37] and Supplementary Table 13). Serial dilutions of IgG, IgA and IgM

versions of CR3022 were used to create standard curves on RBD and S1 coated beads. The linear range

of the standard and sample dilution curve was used for quantitation. We fitted a four-parameter

logistic curve (y=Bottom+(Top-Bottom)/(1+10^((logEC50-X)*HillSlope)) (Supplementary Fig. 7b)

through which MFI values of measured samples are interpolated into a corresponding concentration

of antibody (µg/ml). We used this approach to quantify the concentration of RBD and S1 antibody

reactivity in the positive donor control, and used titrations of the donor pool included on each ABCORA

plate to calculate the S1 and RBD content of plasma samples in relation to it. We used the same

strategy in combination with the WHO International Standard Anti-SARS-CoV-2 Immunoglobulin

(NIBSC 20/136 [31]) to defer IU/ml content of the internal ABCORA positive donor pool and the individual

specimen tested (Supplementary Fig. 7). The WHO International Standard consists of a pool of plasma

from individuals with confirmed SARS-CoV-2 infection. RBD and S1 content of the ABCORA positive

donor pool quantified via the polyclonal WHO standard was highly similar within each Ig class

(Supplementary Fig. 7b). In contrast, RBD values estimated by the mAb CR3022 were a factor 2.4 – 3.9

lower than the corresponding S1 values, suggesting an affinity difference of CR3022 for the two

antigens (Supplementary Fig. 7b).

Temporal evolution of SARS-CoV-2 binding antibody response

Antibody binding of 140 convalescent patients was measured longitudinally in 274 measurements with

ABCORA 2.0, including 251 measurements from 120 patients with known time since positive RT-PCR.

We assumed the dynamics of antibodies (analyzed as logMFI-FOE) were following a linear trend with

time and estimated the slope using a linear mixed model with random effect on the intercept. As time

measures days post first positive RT-PCR result (Fig. 5b) or days post onset of symptoms

(Supplementary Fig. 9b) were employed. Half-lives (t1/2, in days) of significant response with negative

slopes were calculated based on the respective estimated slopes. Analyses were performed in R

version 3.6.3 using packages lme4 [68] and lmerTest [69].

SARS-CoV-2 pseudo-neutralization assay

SARS-CoV-2 plasma neutralization activity was defined using an HIV-based pseudovirus system as described [32]. The env-inactivated HIV-1 reporter construct pHIV-1NL4-3 ΔEnv-NanoLuc (pHIV-1Nanoluc) and HT1080/ACE2cl.14 cells were kindly provided by P. Bieniasz, Rockefeller University, NY, USA. To create a SARS-CoV-2 spike expression plasmid (P_CoV2_Wuhan), a codon-optimized C terminal truncated (AA 1255-1273) spike encoding gene of strain Wuhan-Hu-1 (GenBank accession no. MN908947) was synthesized (GeneArt, Thermo Fisher Scientific, Waltham, MA) and cloned into pcDNA.3.1. Pseudotyped SARS-CoV-2 spike expressing viruses were generated by co-transfecting 293-T cells with a mixture of pHIV-1Nanoluc, P_CoV2_Wuhan and PEI Max (Polysciences Europe GmbH, Hirschberg, Germany). After 48h virus supernatants were filtered (0.2 µm) and stored in aliquots at -80°C until use. Infectivity of virus stocks was measured by infection of HT1080/ACE2cl.14 cells. For this 384-well culture plate pre-treated with poly-L-Lysin were seeded with HT1080/ACE2cl.14 (2200 cells/well) one day before the assay. Cells were infected with titrated virus stocks and NanoLuc luciferase activity in cell lysates measured 48 h post infection using the Nano-Glo Luciferase Assay System (Promega, Fitchburg, WI). For this, cells were washed once with PBS, supernatant was removed and cells were lysed with 20 µl/well of Luciferase Cell Lysis reagent (Promega, Fitchburg, WI) for 15 min under continuous shaking at room temperature. 20 µl of 1/ 50 diluted NanoGlo buffer were added and NanoLuc luciferase activity (relative light units, RLU) was measured after 5 min incubation at room temperature on a Perkin Elmer EnVision reader. Input of SARS-CoV-2 pseudoviruses for neutralization assays was adjusted to yield virus infectivity corresponding to 5-10 x10^6 RLU (corresponding to 100-250-fold over background RLU values) in the absence of inhibitors. To measure plasma neutralization activity six serial 4-fold dilutions of plasma starting at a 1/25 dilution were prepared. 20 µl of the diluted plasma and 20 µl of virus were pre-incubated for 1 h at 37°C and then 30 µl of the virus/plasma mix were transferred to 384-well plates seeded with HT1080/ACE2cl.14 cells in a volume of 30 µl. This resulted in a final concentration of the plasma starting dilution of 1/100. Plasma neutralization titers causing 50%, 80% and 90% reduction in viral infectivity (NT50, NT80 and NT90, respectively) compared

to controls without plasma were calculated by fitting a sigmoid dose–response curve (variable slope)

to the RLU data, using GraphPad Prism with constraints (bottom=0, top=100). If 50% inhibition was not

achieved at the lowest plasma dilution of 1/100, a 'less than' value was recorded. All measurements

were conducted in duplicates.

Predicting neutralization based on ABCORA binding activity

To compare the ability of SARS-CoV-2 binding activity measured in ABCORA 2.0 to predict the

neutralization status, we measured neutralization activity to Wuhan-Hu-1 in SARS-CoV-2 positive

individuals (N=467) and classified individuals as high neutralizers (NT50 >250, N=332) and low

neutralizers (NT50 <250, N=135). Six different classification models were designed to assign individuals

to the high or low neutralizers category, based on their ABCORA2.0 binding patterns we established

two univariable logistic regression (ULR) models that included mean MFI-FOE spike antigen S1

reactivities and mean MFI-FOE spike antigen RBD reactivities, respectively. S1 and RBD were chosen

due to their highest correlation with NT50 (r= 0.82 and r=0.80 for the total spike reactivities of S1 and

RBD respectively, Supplementary Fig. 9). In addition to these two ULR, we established a multivariate

logistic regression model including both mean S1 and RBD reactivities (MFI). We further devised three

models that considered all 12 SARS-CoV-2 binding parameters recorded by ABCORA 2.0. Two

multivariable logistic regression models were based on a principal component analysis on all binding

activities and included the first two (respectively four) principal components, which explained 60%

(respectively 75%) of the variance in the data. We also included in our model comparison a

classification based on a random forest analysis that incorporated all 12 SARS-CoV-2 binding activity

variables.

For all 6 models, performance was assessed in 100 cross-validation sets: each set was built by randomly

sampling without replacement among the 467 measurements available. 80% of the data set was used

to train the model (N=374). Prediction of neutralization status was realized on the other 20% (N=93)

and compared to the true NT50 value and neutralization status, using a roc curve. The area under the

curve (AUC) was computed for all 6 models in each cross-validation run. The Bayesian information

criterion (BIC) was computed for all 5 logistic regressions in each cross-validation run.

To increase the utility of the ULR-S1 prediction model for clinical diagnostics we devised a modified

neutralization prediction model ULR-S1-SOC based on the signal over cut-off (SOC) values reported for

ABCORA 2.0. The ULR-S1-SOC estimates the probability of NT50>250 based on the sum of S1 SOC

values for IgG, IgA and IgM as follows:

702
$$P(NT50 > 250) = \frac{\exp(a + b * log10(sum\ S1\ SOC))}{1 + \exp(a + b * log10(sum\ S1\ SOC))}$$

With estimated values: a= -2.6447 and b= 3.5353.

Association between HCoVs and SARS-CoV-2 reactivities

To explore the association between HCoVs and SARS-CoV-2 reactivities, we defined a new HCoV

response variable (HCoV high/low) for each antibody class (IgG, IgA, IgM) as follows: a patient had high

HCoV Ig reactivity for a given antibody class if its measurements were higher than the population

median in at least three out of the four HCoV measurements (HKU1, OC43, NL63, 229E). To assess

interdependencies between HCoV and SARS-CoV-2 responses, we then included the HCoV response

variable in a linear regression model of SARS-CoV-2 reactivities in the same antibody class. The linear

regression models were estimated on a subset of SARS-CoV-2 positive patients (N=204), measured on

ABCORA 5.0 less than 60 days since positive RT-PCR. The restriction to 60 days was chosen to allow

modeling the effect of time with splines. This time period restriction further guaranteed a gender

balance, as convalescent donors with longer follow up were all males recruited for a plasma therapy

study (CPT-ZHP, Swissmedic 2020TpP1004). Regression analyses were adjusted on time (days post

positive RT-PCR or onset of symptoms; as a spline with 3 degrees of freedom), age (as a spline with 3

degrees of freedom) and gender.

Statistical analysis

Statistical analyses were performed in R (Version 3.6.3). Figures were made using the ggplot2

package[70]. When included, boxplots represent the following: median with the middle line, upper and

lower quartiles with the box limits, 1.5x interquartile ranges with the whiskers and outliers with points.

Significance of Spearman rank correlations were assessed through asymptotic t approximation.

Differences in means between two groups with independent measures were tested using two-tailed t-

tests. When applicable, multiple testing was adjusted using Bonferroni correction for multiple

comparisons. A one-way ANOVA with 3 degrees of freedom was used in addition to two-tailed t-tests

in Fig. 6 to provide insights on overall versus group comparison. When analyzing datasets including

repeated measurements of the same individuals (Fig. 3a, Fig. 5b, Fig. 5c, Supplementary Fig. 8a,

Supplementary Fig. 10b), we used linear mixed models with time since positive RT-PCR or time since

symptom onset (continuous or binary variable) as fixed effect and individual as random effect. In the

case of Fig. 5c and Supplementary Fig. 10c, the decreasing slope of neutralization titers was estimated

by considering only individuals with neutralization titers above the detection levels (NT50>100) at their

first measurement. For all linear mixed models, a Wald test with two-sided hypothesis was used to

determine if the estimated slope was significantly different from 0. In addition, half-lives were

obtained from the decreasing slope estimated on the log of either MFI-FOEs (Fig. 5b, Supplementary

Fig. 10b) or NT50s (Fig. 5c, Supplementary Fig. 10c) as follows: $t_{1/2}=\log(2)/(\text{slope}*\log(10))$. In Fig. 7b

and 7c, linear regressions were used to estimate the association between hCoV and SARS-CoV-2

reactivities: a Student t-test with two-sided hypothesis was used to assess if this association was

significantly different from 0.

Data availability

Serological measurements from all patients are made available in Supplementary Table 14.

Code availability

Codes to assess serostatus based on the ABCORA 2.3 method are available at:

https://github.com/chlpasin/SARS-CoV-2-serology

References

1   Rydyznski Moderbacher, C. *et al.* Antigen-Specific Adaptive Immunity to SARS-CoV-2 in Acute
COVID-19 and Associations with Age and Disease Severity. *Cell* **183**, 996-1012 e1019,
doi:10.1016/j.cell.2020.09.038 (2020).
2   Wheatley, A. K. *et al.* Evolution of immune responses to SARS-CoV-2 in mild-moderate COVID-
19. *Nature Communications* **12**, 1162, doi:10.1038/s41467-021-21444-5 (2021).
3   Wang, Z. *et al.* Enhanced SARS-CoV-2 neutralization by dimeric IgA. *Sci Transl Med* **13**,
doi:10.1126/scitranslmed.abf1555 (2021).
4   Robbiani, D. F. *et al.* Convergent antibody responses to SARS-CoV-2 in convalescent
individuals. *Nature* **584**, 437-442, doi:10.1038/s41586-020-2456-9 (2020).
5   Gaebler, C. *et al.* Evolution of antibody immunity to SARS-CoV-2. *Nature*, doi:10.1038/s41586-
021-03207-w (2021).
6   Muecksch, F. *et al.* Longitudinal analysis of serology and neutralizing antibody levels in
COVID19 convalescents. *J Infect Dis*, doi:10.1093/infdis/jiaa659 (2020).
7   Long, Q.-X. *et al.* Clinical and immunological assessment of asymptomatic SARS-CoV-2
infections. *Nature Medicine* **26**, 1200-1204, doi:10.1038/s41591-020-0965-6 (2020).
8   Robbiani, D. F. *et al.* Convergent antibody responses to SARS-CoV-2 in convalescent
individuals. *Nature* **584**, 437-442, doi:10.1038/s41586-020-2456-9 (2020).
9   Aziz, N. A. *et al.* Seroprevalence and correlates of SARS-CoV-2 neutralizing antibodies from a
population-based study in Bonn, Germany. *Nat Commun* **12**, 2117, doi:10.1038/s41467-021-
22351-5 (2021).
10   Fenwick, C. *et al.* Changes in SARS-CoV-2 Spike versus Nucleoprotein Antibody Responses
Impact the Estimates of Infections in Population-Based Seroprevalence Studies. *J Virol* **95**,
doi:10.1128/JVI.01828-20 (2021).
11   Buss, L. F. *et al.* Three-quarters attack rate of SARS-CoV-2 in the Brazilian Amazon during a
largely unmitigated epidemic. *Science* **371**, 288-292, doi:10.1126/science.abe9728 (2021).
12   Diagnostics, E. R. <https://diagnostics.roche.com/global/en/products/params/elecsys-anti-
sars-cov-2.html> (
13   Abbott.   <https://www.corelaboratory.abbott/us/en/offerings/segments/infectious-
disease/sars-cov-2> (
14   Euroimmun.   <https://www.euroimmun.com/products/infection-diagnostics/id/sars-cov-2-
infection-covid-19/> (
15   Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody.
Nature **583**, 290-295, doi:10.1038/s41586-020-2349-y (2020).
16   Piccoli, L. *et al.* Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike
Receptor-Binding Domain by Structure-Guided High-Resolution Serology. *Cell* **183**, 1024-
1042.e1021, doi:https://doi.org/10.1016/j.cell.2020.09.037 (2020).
17   Thomson, E. C. *et al.* Circulating SARS-CoV-2 spike N439K variants maintain fitness while
evading   antibody-mediated   immunity.   *Cell*   **184**,   1171-1187   e1120,
doi:10.1016/j.cell.2021.01.037 (2021).
18   Müller, L. *et al.* Sensitivity of anti-SARS-CoV-2 serological assays in a high-prevalence setting.
European Journal of Clinical Microbiology & Infectious Diseases, doi:10.1007/s10096-021-
04169-7 (2021).
19   Rogers, T. F. *et al.* Isolation of potent SARS-CoV-2 neutralizing antibodies and protection from
disease in a small animal model. *Science* **369**, 956-963, doi:10.1126/science.abc7520 (2020).
20   Barnes, C. O. *et al.* Structures of Human Antibodies Bound to SARS-CoV-2 Spike Reveal
Common Epitopes and Recurrent Features of Antibodies. *Cell* **182**, 828-842 e816,
doi:10.1016/j.cell.2020.06.025 (2020).

21   Starr, T. N. *et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain Reveals
Constraints   on   Folding   and   ACE2   Binding.   *Cell*   **182**,   1295-1310   e1220,
doi:10.1016/j.cell.2020.08.012 (2020).
22   Weisblum, Y. *et al.* Escape from neutralizing antibodies by SARS-CoV-2 spike protein variants.
Elife **9**, doi:10.7554/eLife.61312 (2020).
23   Zhang, S. F. *et al.* Epidemiology characteristics of human coronaviruses in patients with
respiratory infection symptoms and phylogenetic analysis of HCoV-OC43 during 2010-2015 in
Guangzhou. *PLoS One* **13**, e0191789, doi:10.1371/journal.pone.0191789 (2018).
24   van der Hoek, L. *et al.* Burden of disease due to human coronavirus NL63 infections and
periodicity   of   infection.   *Journal   of   Clinical   Virology*   **48**,   104-108,
doi:https://doi.org/10.1016/j.jcv.2010.02.023 (2010).
25   Huang, A. T. *et al.* A systematic review of antibody mediated immunity to coronaviruses:
kinetics, correlates of protection, and association with severity. *Nat Commun* **11**, 4704,
doi:10.1038/s41467-020-18450-4 (2020).
26   Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nat Med*
26, 1033-1036, doi:10.1038/s41591-020-0913-5 (2020).
27   Ladner, J. T. *et al.* Epitope-resolved profiling of the SARS-CoV-2 antibody response identifies
cross-reactivity   with   endemic   human   coronaviruses.   *Cell   Rep   Med*   **2**,   100189,
doi:10.1016/j.xcrm.2020.100189 (2021).
28   *Diagnostic   testing   for   SARS-CoV-2-   Interim   guidance*,
<https://www.who.int/publications/i/item/diagnostic-testing-for-sars-cov-2> (
29   Ng, K. W. *et al.* Preexisting and de novo humoral immunity to SARS-CoV-2 in humans. *Science*
370, 1339-1343, doi:10.1126/science.abe1107 (2020).
30   Francis, T. On the Doctrine of Original Antigenic Sin. *Proceedings of the American Philosophical*
Society **104**, 572-578 (1960).
31   NIBSC.org.
<https://www.nibsc.org/products/brm_product_catalogue/detail_page.aspx?catid=20/136>
(
32   Schmidt, F. *et al.* Measuring SARS-CoV-2 neutralizing antibody activity using pseudotyped and
chimeric viruses. *bioRxiv*, doi:10.1101/2020.06.08.140871 (2020).
33   Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody.
Nature **583**, 290-295, doi:10.1038/s41586-020-2349-y (2020).
34   Brouwer, P. J. M. *et al.* Potent neutralizing antibodies from COVID-19 patients define multiple
targets of vulnerability. *Science* **369**, 643-650, doi:10.1126/science.abc5902 (2020).
35   Wu, Y. *et al.* A noncompeting pair of human neutralizing antibodies block COVID-19 virus
binding to its receptor ACE2. *Science* **368**, 1274-1278, doi:10.1126/science.abc2241 (2020).
36   Chi, X. *et al.* A neutralizing human antibody binds to the N-terminal domain of the Spike protein
of SARS-CoV-2. *Science* **369**, 650-655, doi:10.1126/science.abc6952 (2020).
37   Yuan, M. *et al.* A highly conserved cryptic epitope in the receptor binding domains of SARS-
CoV-2 and SARS-CoV. *Science* **368**, 630-633, doi:10.1126/science.abb7269 (2020).
38   FDA. Decisional Memorandum - Neutralization titer. (2020).
39   Schwarz, G. Estimating the Dimension of a Model. *The Annals of Statistics* **6**, 461-464, 464
(1978).
40   Nickbakhsh, S. *et al.* Epidemiology of Seasonal Coronaviruses: Establishing the Context for the
Emergence of Coronavirus Disease 2019. *J Infect Dis* **222**, 17-25, doi:10.1093/infdis/jiaa185
(2020).
41   Lepiller, Q. *et al.* High incidence but low burden of coronaviruses and preferential associations
between respiratory viruses. *J Clin Microbiol* **51**, 3039-3046, doi:10.1128/jcm.01078-13 (2013).
42   Zurich, C. o. *Numbers and facts on COVID-19 [Kanton Zürich. Zahlen & Fakten zu COVID-19]*, <
https://www.zh.ch/de/gesundheit/coronavirus/zahlen-fakten-covid-
19.html?keyword=covid19#/home (accessed mar 30, 2021)> (

43 Bartsch, Y. C. *et al.* Discrete SARS-CoV-2 antibody titers track with functional humoral stability.
Nat Commun **12**, 1018, doi:10.1038/s41467-021-21336-8 (2021).
44 Stephenson, K. E. *et al.* Immunogenicity of the Ad26.COV2.S Vaccine for COVID-19. *JAMA*,
doi:10.1001/jama.2021.3645 (2021).
45 Alter, G. *et al.* Collaboration between the Fab and Fc contribute to maximal protection against
SARS-CoV-2 following NVX-CoV2373 subunit vaccine with Matrix-M vaccination. *Res Sq*,
doi:10.21203/rs.3.rs-200342/v1 (2021).
46 Garcia-Beltran, W. F. *et al.* Multiple SARS-CoV-2 variants escape neutralization by vaccine-
induced humoral immunity. *Cell*, doi:10.1016/j.cell.2021.03.013 (2021).
47 Wibmer, C. K. *et al.* SARS-CoV-2 501Y.V2 escapes neutralization by South African COVID-19
donor plasma. *Nat Med*, doi:10.1038/s41591-021-01285-x (2021).
48 Stamatatos, L. *et al.* mRNA vaccination boosts cross-variant neutralizing antibodies elicited by
SARS-CoV-2 infection. *Science*, doi:10.1126/science.abg9175 (2021).
49 Planas, D. *et al.* Sensitivity of infectious SARS-CoV-2 B.1.1.7 and B.1.351 variants to neutralizing
antibodies. *Nat Med*, doi:10.1038/s41591-021-01318-5 (2021).
50 Moyo-Gwete, T. *et al.* SARS-CoV-2 501Y.V2 (B.1.351) elicits cross-reactive neutralizing
antibodies. *bioRxiv*, doi:10.1101/2021.03.06.434193 (2021).
51 Li, Q. *et al.* SARS-CoV-2 501Y.V2 variants lack higher infectivity but do have immune escape.
Cell, doi:10.1016/j.cell.2021.02.042 (2021).
52 Wu, J. *et al.* SARS-CoV-2 infection induces sustained humoral immune responses in
convalescent patients following symptomatic COVID-19. *Nat Commun* **12**, 1813,
doi:10.1038/s41467-021-22034-1 (2021).
53 Ding, S. *et al.* Antibody Binding to SARS-CoV-2 S Glycoprotein Correlates with but Does Not
Predict Neutralization. *Viruses* **12**, doi:10.3390/v12111214 (2020).
54 Iyer, A. S. *et al.* Persistence and decay of human antibody responses to the receptor binding
domain of SARS-CoV-2 spike protein in COVID-19 patients. *Sci Immunol* **5**,
doi:10.1126/sciimmunol.abe0367 (2020).
55 Jiang, X. L. *et al.* Lasting antibody and T cell responses to SARS-CoV-2 in COVID-19 patients
three months after infection. *Nat Commun* **12**, 897, doi:10.1038/s41467-021-21155-x (2021).
56 Breton, G. *et al.* Persistent cellular immunity to SARS-CoV-2 infection. *J Exp Med* **218**,
doi:10.1084/jem.20202515 (2021).
57 Dan, J. M. *et al.* Immunological memory to SARS-CoV-2 assessed for up to 8 months after
infection. *Science* **371**, doi:10.1126/science.abf4063 (2021).
58 Song, G. *et al.* Cross-reactive serum and memory B cell responses to spike protein in SARS-CoV-
2 and endemic coronavirus infection. *bioRxiv*, doi:10.1101/2020.09.22.308965 (2020).
59 Anderson, E. M. *et al.* Seasonal human coronavirus antibodies are boosted upon SARS-CoV-2
infection but not associated with protection. *Cell*, doi:10.1016/j.cell.2021.02.010 (2021).
60 Marot, S. *et al.* Rapid decline of neutralizing antibodies against SARS-CoV-2 among infected
healthcare workers. *Nature Communications* **12**, 844, doi:10.1038/s41467-021-21111-9
(2021).
61 Wei, X. *et al.* Emergence of Resistant Human Immunodeficiency Virus Type 1 in Patients
Receiving Fusion Inhibitor (T-20) Monotherapy. *Antimicrobial Agents and Chemotherapy* **46**,
1896-1905, doi:10.1128/aac.46.6.1896-1905.2002 (2002).
62 Liechti, T. *et al.* Development of a high-throughput bead based assay system to measure HIV-
1 specific immune signatures in clinical samples. *J Immunol Methods* **454**, 48-58,
doi:10.1016/j.jim.2017.12.003 (2018).
63 Breiman, L. Random Forests. *Machine Learning* **45**, 5-32, doi:10.1023/A:1010933404324
(2001).
64 Liaw, A. a. W., Matthew. Classification and Regression by randomForest. *R News* (2002).
65 Team, R. C.   (R Foundation for Statistical Computing, Vienna, Austria, 2020).
66 Wright, M. N. & Ziegler, A. ranger: A Fast Implementation of Random Forests for High
Dimensional Data in C++ and R. *2017* **77**, 17, doi:10.18637/jss.v077.i01 (2017).

67   <https://github.com/chlpasin/SARS-CoV-2-serology> (

68   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4.
Journal of Statistical Software **067** (2015).

69   Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear Mixed
Effects Models. *2017* **82**, 26, doi:10.18637/jss.v082.i13 (2017).

70   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (2016).

Funding

This work was supported by a grant of the Pandemiefonds of the University of Zurich Foundation to

A.T., a grant of the Swiss Red Cross to B.F. and A.T., a grant of the University Hospital Zurich Innovation

Grant to M.G.M., the Swiss National Science Foundation grant 31CA30_196906 to H.F.G., A.T., R.K. and

the Gilead COVID-19 RFP Research Program COMMIT Grant #: IN-SW-983-6078 (to H.F.G., A.T., R.K.).

I.A.A. is supported by a research grant of the Promedica Foundation. Roche Diagnostics supported the

study with providing test material for a proportion of the Elecsys S tests.

Acknowledgments

We thank the staff of the Institute of Medical Virology diagnostics unit, sample triage and

administration for their support and the staff of the participating clinics for coordinating the sample

collection in the frame of the included clinical trials.

Author Contributions

I.A.A., C.P., M.S., A.T., H.F.G. and R.D.K. conceived and designed the study and analyzed data. I.A.A.,

M.S., M.M.S., S.E., M.C.H., L.M., M.E.S., A.H., A.A., C.R.N., J.B., M.H. designed and performed binding

antibody experiments. P.R., J.W. and S.E. conducted neutralization experiments and analyzed data. S.S.

developed an analysis app. C.P. and R.D.K. analyzed data. D.L.B., M.M., A.W., S.K.R., B.M.F., E.S., J.G.,

C.B., P.M.M.S., M.G.M., H.F.G., A.W., U.K., J.B. and M.H. were involved in patient recruitment, provided

samples from study and diagnostic repositories and analyzed patient data. A.T., I.A.A., C.P. and M.S.

wrote the manuscript, which all co-authors commented on.

Declaration of interests

The authors declare no competing financial interests.

Figure legends

Fig. 1. Seroprofiling SARS-CoV-2 responses. (a) Assessment of the multiplex SARS-CoV-2 ABCORA 2.0

on the indicated training (N= 823) and validation (N= 635) cohorts (Supplementary Table 3). Depicted

are MFI signals normalized to empty bead controls (MFI-FOE). Grey boxes indicate values above the

individually set MFI-FOE cut-offs for SARS-CoV-2 specific responses for each antigen (see

Supplementary Table 4). (b) Heatmap representing the measured MFI-FOE values and the outcomes

predicted with ABCORA 2.0 - 2.3 of training and validation cohort measurements shown in (a).

Negative, Positive, and Positive, partial refer to ranking according to ABCORA 2.0 as specified in

Supplementary Table 5. (c) Sensitivity and specificity of ABCORA 2 assay versions based on the

combined training and validation cohort data depicted in (a). Proportion of false negative samples

(sensitivity; green) and proportion of false positive samples (specificity; blue) are represented by the

reduction from 100% (outer circle) per segment. (d) Assessment of ABCORA 2.0 with the National

Institute for Biological Standards and Control (NIBSC) Anti SARS-CoV-2 Verification Panel (20/B770)

comprising SARS-CoV-2 positive (red) and negative (blue) panel serum samples. Grey boxes indicate

values above the ABCORA 2.0 MFI-FOE cut-offs for SARS-CoV-2 specific responses for individual

antigen-Ig combinations.

Fig. 2. Quantification of SARS-CoV-2 specific antibody responses. (a-c) Titrated plasma from SARS-

CoV-2 positive adults (N=72) were measured with ABCORA 2.0 and (a) 50% effective concentrations

(EC50; expressed as reciprocal plasma dilution) and (b) area under the curve values (AUC; expressed

as MFI) were calculated. (c) Titrated SARS-CoV-2 RBD and S1 responses were quantified using the RBD

specific monoclonal antibody CR3022 (produced as IgG, IgA and IgM; expressed as ng/ml) as external

standard. See Supplementary Fig.7 for additional quantification with the WHO International Standard

Anti-SARS-CoV-2 Immunoglobulin. (d) Spearman correlation matrix assessing agreement between

ABCORA 2.0 based quantification readouts (EC50, AUC, RBD Ab standardized), the basic MFI-FOE

measured at 1/100 plasma dilution (log), indicated summed logMFI-FOE values (1/100 dilution), and

Roche Elecsys Anti-SARS-CoV-2 (S) assay results (U/ml). Non-significant correlations are left blank.

Levels of significance are assessed by asymptotic t approximation of Spearman's rank correlation, and

corrected by the Bonferroni method for multiple testing ($p < 0.05/780$). Color shading denotes

correlation coefficient.

Fig. 3. Binding and neutralization activity are closely linked in early and late infection (a) 50%

Neutralization titers (NT50) titers against Wuhan-Hu-1 pseudotype post SARS-CoV-2 infection in

patients with known positive SARS-CoV-2 RT-PCR date (N= 369). Patients were stratified according to

time since first diagnosis to investigate early (less than 30 days post RT-PCR, lavender) and late (more

than 30 days post RT-PCR, turquoise) neutralization responses. Difference between these two groups

was assessed with a linear mixed model with time since RT-PCR (binary variable early/late) as fixed

effect and individual as random effect and using a Wald test on the parameter associated with time

since RT-PCR (***: $p < 0.001$). (b) Linear regression analysis to define association between neutralization

(reciprocal NT50) and antibody binding (MFI-FOE). Black lines indicate linear regression predictions

and grey shaded areas correspond to the 95% confidence intervals. Results depict early (lavender), late

(turquoise) and full cohort (black). n.s. denotes non-significant results. Levels of significance are

assessed by asymptotic t approximation of Spearman's rank correlation, and corrected by the

Bonferroni method for multiple testing ($p < 0.05/1200$, see Supplementary Figures 8b and 9).

Fig. 4. Predicting neutralization capacity as a function of binding activity. (a) SARS-CoV-2 positive

donors (N=467) were stratified into high neutralizers (NT50 >250, N=332; blue) and no/low neutralizers

(NT50 <250, N=135; grey), based on their neutralization activity against Wuhan-Hu-1. (b) and (c)

Comparison of the prediction ability of six different classification models using 100 cross-validation

sets (divided as 80% for training and 20% for validation. (b) Comparison of models by area under the

curve (AUC). Each dot corresponds to one cross-validation set. (c) Bayesian information criterion (BIC)

of the five models based on logistic regression. The different models are: Univariable logistic

regressions (ULR). ULR-RBD: mean of MFI-FOE RBD. ULR-S1: mean of MFI-FOE S1. Multivariable logistic

regression (MLR). MLR-S1, RBD: mean of S1 reactivity and mean of RBD reactivity. MLR-PCA2 and MLR-

PCA4: MLR of 2 and 4 first axis of PCA analysis, respectively. PCA was based on all 12 SARS-CoV-2

antibody reactivities measured by ABCORA 2.0. Random forest (RF) including all antibody reactivities

measured by ABCORA 2.0. (d) ULR-S1 estimated ROC curve based on full data set (N=467). (e)

Measured NT50 value versus probability of NT50 >250 as predicted by ULR-S1 in five randomly chosen

validation sets (each symbol corresponds to a validation set). Purple colored symbols indicate a higher

than 0.70 probability of the respective sample to be neutralizing at NT50 >250 and are therefore

denoted as high neutralizers. Grey indicates samples with predicted neutralization NT50 <250,

therefore classified as no/low neutralizers. (f) Neutralization prediction based on a modified ULR-S1

model utilizing the diagnostic readout SOC instead of MFI-FOE values as input. Measured NT50 value

versus sum of S1 SOC values (IgG, IgA, IgM) are depicted. Dashed lines correspond to a NT50=250

horizontally and the sum S1 SOCs=9.7 vertically. The sum S1 SOCs=9.7 corresponds to the thresholds

depicted for ULR-S1 in (d) and (e). The grey shaded area corresponds to true positives (individuals with

NT50 >250 predicted as high neutralizers).

Fig. 5. Monitoring temporal evolution of antibody responses. (a) ABCORA 2.3 definition of

seropositivity in donors with positive RT-PCR confirmed SARS-CoV-2 infection and known RT-PCR date

(N=369). Seropositivity rating in relation to plasma sampling time point post diagnosis is depicted. Grey

shaded area highlights the first seven days since positive RT-PCR detection. (b) Linear mixed model,

with time since RT-PCR as fixed effect and individual as random effect, estimating the decay of antibody

binding activity based on ABCORA 2.0 measurements at 1 - 4 longitudinal time points in 120 individuals

totaling in 251 measurements. Purple lines correspond to the models estimation and purple shaded

areas to the 95% confidence intervals. Antibody half-lives (t1/2 in days) from significant models are

depicted. Significance was assessed using Wald tests on the slope parameters. (c) Linear mixed model,

with time since RT-PCR as fixed effect and individual as random effect, estimating the decay of

neutralizing capacity on 251 measurements from 120 individuals. Only individuals with NT50>100 at

their first measurement were used to estimate the half-life. The purple line corresponds to the model

estimation and the purple shaded area to the 95% confidence intervals. Significance was assessed

using Wald tests on the slope parameters.

Fig. 6. Seasonal and annual fluctuation in HCoV reactivity. Reactivity to human coronaviruses (HCoV-

NL63, HCoV-229E, HCoV-HKU1, HCoV-OC43) was compared by ABCORA 5.0. Reactivity in healthy blood

donors from 2019 and 2020 was compared. Pre-pandemic samples included: January 2019 (N=285),

May 2019 (N=288), January 2020 (N=252). Samples from May 2020 (N=672) were collected during the

pandemic in Switzerland. Only samples without SARS-CoV-2 specific reactivity as defined by ABCORA

were included (N=653). Stars correspond to levels of significance of two-tailed t-tests comparing the

indicated groups. Levels of significance are corrected by the Bonferroni method for multiple testing

and indicated as follows: $*p<0.05/36$, $**p<0.01/36$, $***p<0.001/36$.

Fig. 7. Association between SARS-CoV-2 and HCoV antibody responses. (a-b) Time-matched

comparison of ABCORA 5.0 reactivity for SARS-CoV-2 (a) and HCoVs (b) in healthy and SARS-CoV-2

infected individuals. Healthy donors were sampled in May 2020 (N=653; blue). Plasma from SARS-CoV-

2 infected individuals were collected between April - June 2020 (N=65; red). See Supplementary Fig.

13 for analysis on the full SARS-CoV-2 positive cohort (N=389). Grey boxes indicate values above the

individual MFI-FOE cut-offs for SARS-CoV-2 specific responses for each antigen. Stars correspond to

levels of significance of t-tests comparing negative versus positive patients. Levels of significance are

corrected by the Bonferroni method for multiple testing and indicated as follows: $*p<0.05/12$,

$**p<0.01/12$, $***p<0.001/12$. (b) Linear regression models showing the association between SARS-

CoV-2 and HCoV signals in 204 SARS-CoV-2 positive patients with known dates of first positive RT-PCR

detection. Influences within the same antibody class are investigated. The models were adjusted on

age (spline with 3 degrees of freedom), gender, time since positive RT-PCR (spline with 3 degrees of

freedom) and level of HCoV reactivity. Samples are defined to harbor high HCoV reactivity if they show

ABCORA 5.0 HCoV logMFI-FOE values higher than the corresponding median in at least 3 HCoV

measurements (HKU1, OC43, NL63 or 229E). Curves correspond to the models estimation and shaded

areas to the 95% confidence intervals. p-values were obtained by running a Student t-test on the

parameter associated to HCoV reactivity in the linear regression. (c) Linear regression model showing

the association between SARS-CoV-2 IgG and HCoV IgA signals. Curves correspond to the models

estimation and shaded areas to the 95% confidence intervals. (d) Linear regression model showing the

association between SARS-CoV-2 IgG and HCoV IgM signals. Curves correspond to the models

estimation and shaded areas to the 95% confidence intervals.

1039

Supplementary Figures

Supplementary Fig.1. Schematic overview of ABCORA seroprofiling. (a) Directed coupling of His-

tagged antigens to magnetic beads covalently coupled with anti-His antibody. (b) Binding of patient

plasma antibodies to antigen-coupled beads and detection by PE-labeled secondary antibodies (IgG,

IgA or IgM) with the FlexMap 3D reader (Luminex). Median fluorescence intensity (MFI) proportional

to bound secondary antibody is recorded. Figure created with BioRender.

Supplementary Fig. 2. Determination of optimal concentration of anti-His antibodies and SARS-CoV-

2 derived antigens for coupling and loading of magnetic beads. (a) Titration of anti-His capture

antibody on magnetic beads. One of two independent experiments is depicted. (b) and (c) Optimization

of antigen loading. (b) Reactivity of beads loaded with increasing doses of SARS-CoV-2 nucleoprotein

(N) or SARS-CoV-2 spike protein subunit S1 (S1) with titered anti-N and anti-S1 mAbs and a SARS-CoV-

2 positive patient plasma pool. One of three independent experiments is depicted. (c) Median

fluorescence intensity (MFI) at 1/100 dilution and the 50 % effective concentration (EC50) values for

anti-N and anti-S1 mAbs, the SARS-CoV-2 positive patient pool (+PP), two individual SARS-CoV-2

positive patient plasma (P1 and P2) and a plasma pool of pre-pandemic healthy donors (-PP). One of

three independent experiments is depicted. (d) Final assessment of assay setup (5 µg anti-his Ab per

million beads, 320 nM His-tagged antigens, Phycoerythrin (PE)-labeled secondary antibodies at 17500).

Reactivity of the indicated SARS-CoV-2 antigens with serial dilutions of anti-N and anti-S mAbs, positive and negative donor plasma pools was probed. Three independent experiments are depicted.

Supplementary Fig. 3. Assessment of assay variability. Titration of the positive control plasma donor pool composed of 20 SARS-CoV-2 RT-PCR positive patients. Median fluorescence intensity (MFI) for IgG, IgA and IgM reactivities to SARS-CoV-2 proteins (RBD, S1, S2, N) and empty bead reactivity of 31 independent titrations are shown.

Supplementary Fig. 4. Temporal stability and variability analysis. (a) Assessment of the temporal stability of antigen coupled beads. The SARS-CoV-2 positive plasma pool was titrated on 25 days and the distribution of all signal intensities (pooled over plasma dilutions and antigens) for each day depicted. (b) Histogram of the overall assay variability (coefficient of variation) for all tested Ig classes based on the variability of mean log MFI values from 31 independent titrations (7 dilution steps) of the positive control plasma pool depicted in Supplementary Figure 3. (c-d) Boxplots depicting the overall assay variability stratified by the four different antigens (c) and plasma dilutions (d) based on 31 independent titrations (7 dilution steps) of a positive plasma pool. (e) Boxplots showing the intraday variability stratified by the four different antigens based on six independent titrations of a positive plasma pool performed on the same day. (f) Boxplots showing the intra-day variability stratified by the four different antigens based on a titration of a positive plasma pool performed on 10 different days.

Supplementary Fig. 5. Interdependency of SARS-CoV-2 and HCoV HKU1 antibody reactivity. Spearman correlation matrix of SARS-CoV-2 (RBD, S1, S2, N) and HKU1 S1 antigen reactivity (based on logMFI-FOE) in SARS-CoV-2 positive adults (N= 389), healthy, pre-pandemic adults (N=825), pre-pandemic samples from patients recently infected with a circulating HCoV strain (N=75) and pre-pandemic children (N=169). Non-significant correlations are left blank. Levels of significance are assessed by asymptotic t approximation of Spearman's rank correlation, and corrected by the Bonferroni method for multiple testing ($p<0.05/420$).

Supplementary Fig. 6. Variable importance for computational models. (a) Correlation matrix of all immunoglobulin variables in SARS-CoV-2 positive patients from the training dataset (N=175). Defining five clusters based on hierarchical clustering showed that IgA N and IgM N clustered separately from other IgA and IgM variables. Other variables (indicated by stars: all IgGs, IgAs without N and IgMs without N) were highly correlated. We therefore used the mean of these three clusters in the logistic regression. (b) Variable importance (measured as the mean decrease of node impurity with Gini index). Each of the 100 dots corresponds to a random forest performed on a bootstrap sample of the training dataset (N=823).

Supplementary Fig. 7. Quantification of SARS-CoV-2 specific antibodies. (a) Titration of SARS-CoV-2 positive plasma pool used as plate standard curve for quantification in Figure 3 (N=72). Uncorrected MFI values are depicted. (b) Quantification of SARS-CoV-2 specific antibodies in the positive plasma pool by interpolating RBD and S1 content based on standard curves of the RBD-specific mAB CR3022 (upper panel) or the WHO International Standard (lower panel). Data from three independent experiments are depicted. (c) Spearman correlation matrix assessing agreement between diverse ABCORA measurements and quantifications (individual and summed logMFI-FOE values at 1/100 dilution of plasma, EC50, AUC, IU/ml content based on WHO Standard NIBSC 20/136) and Roche Elecsys Anti-SARS-CoV-2 (S) (U/ml). Non-significant correlations are left blank. Levels of significance are assessed by asymptotic t approximation of Spearman's rank correlation, and corrected by the Bonferroni method for multiple testing ($p < 0.05/780$). Color shading denotes correlation coefficient.

Supplementary Fig. 8. Binding and neutralization activity are closely linked in early and late SARS-CoV-2 infection. (a) 50% Neutralization titers (NT50) titers against Wuhan-Hu-1 pseudotype in patients with known date of symptoms onset (N= 333). Patients were stratified according to time since first diagnosis to investigate early (less than 30 days post symptoms onset, lavender) and late (more than 30 days symptoms onset, turquoise) neutralization responses. Difference between these two groups was assessed with a linear mixed model with time since symptom onset (binary variable early/late) as fixed effect and individual as random effect and using a Wald test on the parameter associated with

time since symptom onset (***p<0.001). (b) Linear regression analysis to define association between

neutralization (reciprocal NT50) and antibody binding (MFI-FOE). Black lines indicate linear regression

predictions. Levels of significance are assessed by asymptotic t approximation of Spearman's rank

correlation, and corrected by the Bonferroni method for multiple testing (p<0.05/1200, see

Supplementary Figure 9).

Supplementary Fig. 9. Correlation of antibody binding and neutralization activity in early and late

infection. Spearman correlation matrix assessing agreement between SARS-CoV-2 antigen reactivity

(RBD, S1, S2, N) based on logMFI-FOE values and neutralization (NT50, NT80, NT90) in SARS-CoV-2

positive adults (N= 389) divided in in early and late time points corresponding to time since positive

RT-PCR diagnosis (a, N=118 – b, N=251) or symptom onset (c, N=66 – d, N=267). Non-significant

correlations are left blank. Levels of significance are assessed by asymptotic t approximation of

Spearman's rank correlation, and corrected by the Bonferroni method for multiple testing

(p<0.05/1200).

Supplementary Fig. 10. Monitoring temporal evolution of antibody responses (a) Heatmaps

representing the measured MFI-FOE values and the outcome predicted with ABCORA 2.0 - 2.3 of

measurements of SARS-CoV-2 positive patients with known dates of positive RT-PCR diagnosis (N=369)

(upper panel) or with known dates of onset of symptoms (N=333) (lower panel). Purple and orange

scales indicate days post positive RT-PCR or days post onset of symptoms, respectively, white-to-black

scale indicates seroconversion predicted with the different ABCORA approaches. (b) Linear mixed

model, with time since symptom onset as fixed effect and individual as random effect, estimating the

decay of antibody binding activity based on ABCORA 2.0 measurements at 1-4 longitudinal time points

in 120 individuals totaling in 251 measurements. Orange lines correspond to the models estimation

and orange shaded areas to the 95% confidence intervals. Antibody half-lives ($t_{1/2}$ in days) from

significant models are depicted. Significance was assessed using Wald tests on the slope parameters.

(c) Linear mixed model estimating the decay of neutralizing capacity in patients separated by their

neutralizing activity. Only individuals with NT50>100 at their first measurement were used to estimate

the half-life. The black line corresponds to the model estimation and the grey shaded area to the 95%

confidence interval.

Supplementary Fig. 11. ABCORA 5 seroprofiling records antibody reactivity to SARS-CoV-2 and four

HCoVs. (a) Assessment of the multiplex SARS-CoV-2 ABCORA 5.0 on the indicated training (N= 825) and

validation (N= 635) cohorts (Supplementary Table 3). Depicted are MFI signals normalized to empty

bead controls (MFI-FOE). Grey boxes indicate values above the individually set MFI-FOE cut-offs for

SARS-CoV-2 specific responses for each antigen (see Supplementary Table 4). (b) Sensitivity and

specificity of ABCORA 5 assay versions 5.0, 5.4 and 5.5 based on the combined training and validation

cohort data depicted in (a) (see also Supplementary Table 10). False negative proportion (sensitivity;

green) and false positive proportion (specificity; blue) samples are represented by the reduction from

100% (outer circle) per segment.

Supplementary Fig. 12. Interdependencies between antibody reactivity to SARS-CoV-2 and the four

HCoVs. Spearman correlation matrix assessing agreement between SARS-CoV-2 antigens (RBD, S1, S2,

N) and HCoVs (229E, NL63, OC43, HKU1) (based on logMFI-FOE) in SARS-CoV-2 positive adults (N= 389),

healthy, pre-pandemic adults (N= 825), pre-pandemic samples from patients recently infected with a

circulating HCoV strain (N=75) and pre-pandemic children (N=169). Non-significant correlations are left

blank. Levels of significance are assessed by asymptotic t approximation of Spearman's rank

correlation, and corrected by the Bonferroni method for multiple testing (p<0.05/1104).

Supplementary Fig. 13. Association between SARS-CoV-2 and HCoV antibody responses. Comparison

of ABCORA 5.0 reactivity for SARS-CoV-2 and HCoVs in healthy, SARS-CoV-2 negative and SARS-CoV-2

infected individuals. Healthy donors were sampled in May 2020 (N=653; blue). Plasma from SARS-CoV-

2 infected individuals were collected between April 2020 and February 2021 (N=389; red, Training III

and Validation VI). Grey boxes indicate values above the individual MFI-FOE cut-offs for SARS-CoV-2

specific responses for each antigen. Stars correspond to levels of significance of two-tailed t-tests

comparing negative versus positive patients. Levels of significance are corrected by the Bonferroni

method for multiple testing and indicated as follows: *p<0.05/12, **p<0.01/12, ***p< 0.001/12.

1159

1160

Figure 1

Fig 1. Seroprofiling SARS-CoV-2 responses. (a) Assessment of the multiplex SARS-CoV-2 ABCORA 2.0 on the indicated training (N= 823) and validation (N= 635) cohorts (Supplementary Table 3). Depicted are MFI signals normalized to empty bead controls (MFI-FOE). Grey boxes indicate values above the individually set MFI-FOE cut-offs for SARS-CoV-2 specific responses for each antigen (see Supplementary Table 4). (b) Heatmap representing the measured MFI-FOE values and the outcomes predicted with ABCORA 2.0 - 2.3 of training and validation cohort measurements shown in (a). Negative, Positive, Positive, partial refer to ranking according to ABCORA 2.0 as specified in Supplementary Table 5. (c) Sensitivity and specificity of ABCORA 2 assay versions based on the combined training and validation cohort data depicted in (a). Proportion of false negative samples (sensitivity; green) and proportion of false positive samples (specificity; blue) are represented by the reduction from 100% (outer circle) per segment. (d) Assessment of ABCORA 2 with the National Institute for Biological Standards and Control (NIBSC) Anti SARS-CoV-2 Verification Panel (20/B770) comprising SARS-CoV-2 positive (red) and negative (blue) panel serum samples. Grey boxes indicate values above the ABCORA 2.0 MFI-FOE cut-offs for SARS-CoV-2 specific responses for individual antigen-Ig combinations.

Figure 2

Fig 2. Quantification of SARS-CoV-2 specific antibody responses. (a-c) Titrated plasma from SARS-CoV-2 positive adults (N=72) were measured with ABCORA 2.0 and (a) 50% effective concentrations (EC50; expressed as reciprocal plasma dilution) and (b) area under the curve values (AUC; expressed as MFI) were calculated. (c) Titrated SARS-CoV-2 RBD and S1 responses were quantified using the RBD specific monoclonal antibody CR3022 (produced as IgG, IgA and IgM; expressed as ng/ml) as external standard. See Supplementary Fig.7 for additional quantification with the WHO International Standard Anti-SARS-CoV-2 Immunoglobulin. (d) Spearman correlation matrix assessing agreement between ABCORA 2.0 based quantification readouts (EC50, AUC, RBD Ab standardized), the basic MFI-FOE measured at 1/100 plasma dilution (log), indicated summed LogMFI-FOE values (1/100 dilution), and Roche Elecsys Anti-SARS-CoV-2 (S) assay results (U/ml). Non-significant correlations are left blank. Levels of significance are assessed by asymptotic t approximation of Spearman's rank correlation, and corrected by the Bonferroni method for multiple testing (p<0.05/780). Color shading denotes correlation coefficient.

Figure 3

Fig 3. Binding and neutralization activity are closely linked in early and late infection (a) 50% Neutralization titers (NT50) titers against Wuhan-Hu-1 pseudotype post SARS-CoV-2 infection in patients with known positive SARS-CoV-2 RT-PCR date (N= 369). Patients were stratified according to time since first diagnosis to investigate early (less than 30 days post RT-PCR, lavender) and late (more than 30 days post RT-PCR, turquoise) neutralization responses. Difference between these two groups was assessed with a linear mixed model with time since RT-PCR (binary variable early/late) as fixed effect and individual as random effect and using a Wald test on the parameter associated with time since RT-PCR (***: $p<0.001$). (b) Linear regression analysis to define association between neutralization (reciprocal NT50) and antibody binding (MFI-FOE). Black lines indicate linear regression predictions and grey shaded areas correspond to the 95% confidence intervals. Results depict early (lavender), late (turquoise) and full cohort (black). n.s. denotes non-significant results. Levels of significance are assessed by asymptotic t approximation of Spearman's rank correlation, and corrected by the Bonferroni method for multiple testing ($p<0.05/1200$, see Supplementary Figures 8b and 9).

Figure 4

Fig 4. Predicting neutralization capacity as a function of binding activity. (a) SARS-CoV-2 positive donors (N=467) were stratified into high neutralizers (NT50>250, N=332; blue) and no/low neutralizers (NT50<250, N=135; grey), based on their neutralization activity against Wuhan-Hu-1. (b) and (c) Comparison of the prediction ability of six different classification models using 100 cross-validation sets (divided as 80% for training and 20% for validation. (b) Comparison of models by area under the curve (AUC). Each dot corresponds to one cross-validation set. (c) Bayesian information criterion (BIC) of the five models based on logistic regression. The different models are: Univariable logistic regressions (ULR). ULR-RBD: mean of MFI-FOE RBD. ULR-S1: mean of MFI-FOE S1. Multivariable logistic regression (MLR). MLR-S1, RBD: mean of S1 reactivity and mean of RBD reactivity. MLR-PCA2 and MLR-PCA4: MLR of 2 and 4 first axis of PCA analysis, respectively. PCA was based on all 12 SARS-CoV-2 antibody reactivities measured by ABCORA 2.0. Random forest (RF) including all antibody reactivities measured by ABCORA 2.0. (d) ULR-S1 estimated ROC curve based on full data set (N=467). (e) Measured NT50 value versus probability of NT50>250 as predicted by ULR-S1 in five randomly chosen validation sets (each symbol corresponds to a validation set). Purple colored symbols indicate a higher than 0.70 probability of the respective sample to be neutralizing at NT50>250 and are therefore denoted as high neutralizers. Grey indicates samples with predicted neutralization NT50<250, therefore classified as no/low neutralizers. (f) Neutralization prediction based on a modified ULR-S1 model utilizing the diagnostic readout SOC instead of MFI-FOE values as input. Measured NT50 value versus sum of S1 SOC values (IgG, IgA, IgM) are depicted. Dashed lines correspond to a NT50=250 horizontally and the sum S1 SOCs=9.7 vertically. The sum S1 SOCs=9.7 corresponds to the thresholds depicted for ULR-S1 in (d) and (e). The grey shaded area corresponds to true positives (individuals with NT50 >250 predicted as high neutralizers).

Figure 5

Fig 5. Monitoring temporal evolution of antibody responses (a) ABCORA 2.3 definition of seropositivity in donors with positive RT-PCR confirmed SARS-CoV-2 infection and known RT-PCR date (N=369). Seropositivity rating in relation to plasma sampling time point post diagnosis is depicted. Grey shaded area highlights the first seven days since positive RT-PCR detection. (b) Linear mixed model, with time since RT-PCR as fixed effect and individual as random effect, estimating the decay of antibody binding activity based on ABCORA 2.0 measurements at 1 - 4 longitudinal time points in 120 individuals totaling in 251 measurements. Purple lines correspond to the models estimation and purple shaded areas to the 95% confidence intervals. Antibody half-lives (t1/2 in days) from significant models are depicted. Significance was assessed using Wald tests on the slope parameters. (c) Linear mixed model, with time since RT-PCR as fixed effect and individual as random effect, estimating the decay of neutralizing capacity on 251 measurements from 120 individuals. Only individuals with NT50>100 at their first measurement were used to estimate the half-life. The purple line corresponds to the model estimation and the purple shaded area to the 95% confidence intervals. Significance was assessed using Wald tests on the slope parameters.

Figure 6

Fig 6. Seasonal and annual fluctuation in HCoV reactivity Reactivity to human coronaviruses (HCoV-NL63, HCoV-229E, HCoV-HKU1, HCoV-OC43) was compared by ABCORA 5.0. Reactivity in healthy blood donors from 2019 and 2020 was compared. Pre-pandemic samples included: January 2019 (N=285), May 2019 (N=288), January 2020 (N=252). Samples from May 2020 (N=672) were collected during the pandemic in Switzerland. Only samples without SARS-CoV-2 specific reactivity as defined by ABCORA were included (N=653). Stars correspond to levels of significance of two-tailed t-tests comparing the indicated groups. Levels of significance are corrected by the Bonferroni method for multiple testing and indicated as follows: *p<0.05/36, **p<0.01/36, ***p<0.001/36.

Figure 7

Fig 7. Association between SARS-CoV-2 and HCoV antibody responses. (a-b) Time-matched comparison of ABCORA 5.0 reactivity for SARS-CoV-2 (a) and HCoVs (b) in healthy and SARS-CoV-2 infected individuals. Healthy donors were sampled in May 2020 (N=653; blue). Plasma from SARS-CoV-2 infected individuals were collected between April - June 2020 (N=65; red). See Supplementary Figure 13 for analysis on the full SARS-CoV-2 positive cohort (N=389). Grey boxes indicate values above the individual MFI-FOE cut-offs for SARS-CoV-2 specific responses for each antigen. Stars correspond to levels of significance of t-tests comparing negative versus positive patients. Levels of significance are corrected by the Bonferroni method for multiple testing and indicated as follows: *p<0.05/12, **p<0.01/12, ***p< 0.001/12. (b) Linear regression models showing the association between SARS-CoV-2 and HCoV signals in 204 SARS-CoV-2 positive patients with known dates of first positive RT-PCR detection. Influences within the same antibody class are investigated. The models were adjusted on age (spline with 3 degrees of freedom), gender, time since positive RT-PCR (spline with 3 degrees of freedom) and level of HCoV reactivity. Samples are defined to harbor high HCoV reactivity if they show ABCORA 5.0 HCoV LogMFI-FOE values higher than the corresponding median in at least 3 HCoV measurements (HKU1, OC43, NL63 or 229E). Curves correspond to the models estimation and shaded areas to the 95% confidence intervals. p-values were obtained by running a Student t-test on the parameter associated to HCoV reactivity in the linear regression. (c) Linear regression model showing the association between SARS-CoV-2 IgG and HCoV IgA signals. Curves correspond to the models estimation and shaded areas to the 95% confidence intervals. (d) Linear regression model showing the association between SARS-CoV-2 IgG and HCoV IgM signals. Curves correspond to the models estimation and shaded areas to the 95% confidence intervals.