Sales of over-the-counter products containing codeine in 31 countries,

2013-2019: a retrospective observational study

Georgia C Richards (0000-0003-0244-5620)[1,2];*

Jeffrey K Aronson (0000-0003-1139-655X)[2];

Brian MacKenna (0000-0002-3786-9063)[3];

Ben Goldacre (0000-0002-5127-4728)[3];

FD Richard Hobbs (0000-0001-7976-7172)[4];

Carl Heneghan (0000-0002-1009-1992)[1,2]

[1]Global Centre on Healthcare and Urbanisation, Kellogg College, University of Oxford,

60-62 Banbury Road, Oxford, OX2 6PN, UK.

[2]Centre for Evidence-Based Medicine, Nuffield Department of Primary Care Health

Sciences, University of Oxford, Radcliffe Observatory Quarter, Woodstock Road,

Oxford, OX2 6GG, UK.

[3]EBM Datalab, Nuffield Department of Primary Care Health Sciences, University of

Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG, UK.

[4]Nuffield Department of Primary Care Health Sciences, University of Oxford, Radcliffe

Observatory Quarter, Woodstock Road, Oxford, OX2 6GG, UK.

*corresponding author: GC Richards, georgia.richards@kellogg.ox.ac.uk

Word count: 2384

Running heading: Trends in sales of over-the-counter codeine in 31 countries

Acknowledgements: We thank the Primary Care Research Trust of Birmingham and

Midlands Research Practices Consortium for providing the funding to purchase the

sales data from IQVIA.

ABSTRACT

Introduction

Opioid prescribing trends have been investigated in many countries. However, the

patterns of over-the-counter purchases of opioids without a prescription, such as

codeine combinations, are mostly unknown.

Objective

We aimed to assess national sales and expenditure of over-the-counter codeine-

containing products purchased in 31 countries over six years.

Methods

We conducted a retrospective observational study using electronic point-of-sales data

from the human data science company, IQVIA, for Argentina, Belgium, Brazil,

Bulgaria, Canada, Croatia, Estonia, Finland, France, Germany, Greece, Ireland, Italy,

Japan, Latvia, Lithuania, Mexico, The Netherlands, Poland, Portugal, Romania, Russia,

Serbia, Slovakia, Slovenia, South Africa, Spain, Switzerland, Thailand, the UK, and the

USA. We calculated annual mean sales (units per 1000 of the population) and public

expenditure (GBP, £ per 1000) for each country between April 2013 and March 2019.

We quantified changes over time and the types of products sold.

Results

31.5 billion units of codeine, costing £2.55 billion, were sold over-the-counter in 31

countries between April 2013 and March 2019. Total sales increased by 3% (3025

units/1000 in 2013 to 3111 in 2019) and public expenditure doubled (£196/1000 in

2013 to £301/1000 in 2019). Sales were not equally distributed; South Africa sold the

most (31 units/person), followed by Ireland (24 units/person), France (16 units/person),

Latvia (15 units/person), and the UK (11 units/person). The types of products (n=569)

and formulations (n=12) sold varied.

Conclusion and Relevance

In many parts of the world, substantial numbers of people may be purchasing and

consuming codeine from over-the-counter products. Clinicians should ask patients

about their use of over-the-counter products and public health measures are required to

improve the collection of sales data and safety of such products.

Study protocol pre-registration: https://osf.io/ay4mc

Key points

- Total sales and public expenditure of over-the-counter products containing

codeine increased from April 2013 to March 2019 in line with increased trends

of global opioid use.

- There was substantial variation in mean sales of over-the-counter codeine

between countries.

- In countries with access to over-the-counter codeine, these sales data should be

reviewed to informed regulatory decisions and requirements for public health

measures to ensure safety.

1 Introduction

Prescribing patterns of opioids are documented in many countries [1–7]. However, opioids such as analgesic combinations containing codeine can be purchased over-the-counter (OTC) without a prescription or consultation with a doctor or prescriber in most countries. As access to granular data on OTC sales has been limited, previous research on the use of non-prescribed codeine has relied on case reports [8–10], self-reported questionnaires [11–18], qualitative studies [19–22], and data from poisons centres, hospital admissions, or coronial systems [23–27]. It is therefore unknown whether OTC sales of codeine have followed trends similar to the use of prescribed opioids.

Codeine (3-methylmorphine) is used for its analgesic, antidiarrheal, and antitussive effects [28–30]. It is often combined with other analgesics, such as paracetamol, and non-steroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen. These combinations have greater efficacy than codeine alone [28,31,32]. But most clinical trials testing the efficacy of codeine have used high doses (25–90 mg), which are not available OTC [31–33]. A Cochrane overview of systematic reviews on oral OTC analgesics for acute pain found no studies or data that could be extracted on combinations of analgesics containing low doses of codeine [33]. A systematic review of the efficacy and safety of low-dose (≤30 mg) codeine included ten RCTs [34]. It reported low- to moderate-quality evidence that combination products of low-dose codeine provided little to moderate pain relief for acute and chronic pain conditions in the short term [34]. In observational studies products containing codeine have been associated with dependence, misuse, death, and collateral toxicity from combinations with paracetamol and ibuprofen [9,25].

Access to codeine without a prescription can bolster the perception of safety, and in

some settings abuse of OTC codeine is normalised and encouraged. Mixing codeine

cough syrup with alcohol and or soft drinks has been popularised in rap music and by

American athletes, who make a drink called "purple drank" from codeine syrup and

soda [35]. OTC codeine has been used to manufacture illicit morphine and heroin and to

create a cheap heroin substitute called "krokodil" [36,37]. The use of "Krok" has been

reported in Russia, Europe, the UK, and North America, and its adverse effects from

intravenous use include damage to blood vessels, skin, muscles, and bones, multiorgan

failure, and death [36,38]. Thus, the growing opioid problem is incomplete without

evaluation of OTC sales of codeine.

Regulation of codeine-containing products varies across the world, making it difficult to

estimate how much they are used [39]. Under the 1961 Single Convention on Narcotic

Drugs, codeine is a Schedule III drug [40]. Drugs in this Schedule reportedly "are not

liable to abuse and cannot produce ill effects", and thus it is not mandatory to report

data on their consumption to the International Narcotics Control Board (INCB). In a

report presented at the WHO's Expert Committee on Drug Dependence in October

2019, reviewing codeine formulations listed in Schedule III, the INCB reported a 64%

increase in demand for codeine in the last decade [39].

However, governments can also mandate regulation of codeine. Minnesota (July 2013),

Manitoba (February 2016), France (July 2017), and Australia (February 2018) have

reclassified codeine to prescription-only [41–44]. A review of OTC codeine regulations

in the European Union showed that more than half of member countries did not permit

OTC sale of codeine as of March–August 2014 [45]. Despite variation in codeine's

regulatory status, studies have analysed consumption of OTC cough syrup containing

codeine in Taiwan [46] and the impact of rescheduling codeine to prescription-only in

Australia [47,48]. We aimed to assess national sales and public expenditure on OTC

codeine-containing products purchased in countries with available data.

2 Methods

2.1 Design and data source

We conducted a retrospective observational study using consumer health sales data

from *IQVIA* [49], which has previously been used in observational research on a range

of medications [46–48,50–53]. The data included products containing codeine in the

adult pain relief category, which were collected using scan track barcodes from

electronic point-of-sale (EPoS) store data in all countries that provided data, including

Argentina, Belgium, Brazil, Bulgaria, Canada, Croatia, Estonia, Finland, France,

Germany, Greece, Ireland, Italy, Japan, Latvia, Lithuania, Mexico, The Netherlands,

Poland, Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, South Africa, Spain,

Switzerland, Thailand, the UK, and the USA. *IQVIA* extracted the data on 16 September

2019 and provided quarterly sales from 1 April 2013 to 31 March 2019. *IQVIA's* sample

of data is based on audits and covers a median of 73% (IQR: 58–86%) of

pharmaceutical markets (Table S1 in Supplement 1). Annual population statistics in

calendar years (2013 to 2018) were sourced from the World Bank [54].

2.2 Data analysis

We extracted details from the pack information and used descriptive statistics to

determine the numbers and types of products sold across the 31 countries. Data on

dosages were missing from the pack information for most countries, so we used

IQVIA's standard units to account for liquid and solid dosage forms. We calculated the

total units sold over the study period and the totals for each year (e.g. from quarter two

in 2013 to quarter one in 2014). We also calculated the mean number of units sold over

six years, adjusted for population. We created an annual rate of units sold per 1000 of

each year's population for each country to examine trends over time.

For expenditure, *IQVIA* converted sales to pounds sterling (GBP, £) for each country on

the date of data extraction (16 September 2019). We calculated annual totals, mean

public expenditure for each country, adjusted for population, and a rate of GBP per

1000 to assess changes over time.

2.3 Software and data sharing

We used Stata v16 and Python v3 in Jupyter Notebooks with pandas [55], seaborn [56],

and matplotlib [57] libraries for analysis and figures. The commercial information we

used requires a fee to access. Thus, we cannot openly share the data, owing to

contractual agreements with *IQVIA*. However, we have openly shared our statistical

code at GitHub [58], preregistered [59] and published [60] our study protocol, and

shared all our study materials via the Open Science Framework (OSF) [61].

3 Results

31.5 billion units containing codeine were sold across 31 countries over the study period

(April 2013 to March 2019). Total sales increased by 2.8%, from 3025 units/1000 in

2013–14 to 3111 units/1000 in 2018–19. However, the distribution of sales between

countries was not uniform. Five countries represented 90% of all OTC codeine sales.

South Africa accounted for the greatest volume of sales (34%), followed by France

(20%), Japan (16.5%), the UK (14.5%), and Poland (5%). South Africa sold the most

(mean of 31 units/person, see Figure 1), followed by Ireland (24 units/person), France

(16 units/person), Latvia (15 units/person), and the UK (12 units/person).

[Figure 1 near here]

In the most recent year (April 2018–March 2019), South Africa sold the most (38

units/person), followed by Ireland (23 units/person), Latvia (16 units/person), the UK

(11 units/person), and Japan (10.7 units/person) (Table S2 in Supplement 1).

Over time, 48% of countries (15/31) had increased OTC codeine sales. For countries in

the top quartile of sales (Figure 2A), trends increased in South Africa (30%), Ireland

(8%), Latvia (11%), and Poland (19%), and sales fell in France (99%), the UK (7%),

Serbia (4%), and Croatia (17%). For countries in the second-largest quartile of sales

(Figure 2B), trends increased in Japan (1219%), Estonia (65%), Romania (10%), and

Bulgaria (18%), and sales fell in Switzerland (50%), Finland (22%), The Netherlands

(16%), and Slovenia (16%). Trends for countries in the bottom two quartiles are in

Figure S1 and S2 in Supplement 1.

[Figure 2 (Figure 2A and Figure 2B) near here]

The public spent £2.55 billion on OTC codeine-containing products in 31 countries over

six years; expenditure increased by 54%, from £196/1000 in 2013–14 to £301/1000 in

2018–19. Ireland had the largest mean public expenditure of £5.70 per person, followed

by the UK (mean of £1.60/person), South Africa (£1.26/person), Croatia (£1.25/person),

and Estonia (£1/person) (Figure 3). In April 2018 to March 2019, Ireland continued to

have the largest public expenditure (£6.60/person), followed by South Africa

(£1.64/person), the UK (£1.64/person), Japan (£1.47/person), and Estonia (£1.41)

(Table S2 in Supplement 1).

[Figure 3 near here]

Most countries (58%, 18/31) had increased public expenditure over time. There were

simultaneous increases (45%, 14/31) and decreases (39%, 12/31) in both sales and

expenditure in most countries, while other countries (16%, 5/31) had a discordance in

the direction of their sales and expenditure (Figure 4).

[Figure 4 near here]

There were 569 products and 12 formulations sold across 31 countries. Tablets were the

most common formulations sold, followed by syrups, soluble tablets, and coated tablets

(Figure S3 in Supplement 1). Products contained a median of three substances per

combination (IQR: 2–4, range: 1–16). Limited details were available in the pack

information: the dosages of codeine were available in 17% of products (98 of 569) in 15

countries.

4 Discussion

Many people are purchasing non-prescribed codeine in several parts of the world. Total

sales and public expenditure of OTC codeine products increased over time in line with

previous trends of global opioid use. However, sales were not equally distributed across

the 31 countries.

According to *IQVIA*'s data, South Africa consistently sold the greatest volume of OTC

codeine each year. In a study of opioid dependence in South Africa 5–8% of people in

addiction treatment facilities reported problems with OTC codeine medications and

cough mixtures as their primary or secondary drug of choice [62]. The availability of

non-prescribed codeine may also have ramifications for neighbouring countries that

restrict access. For example, there were reports in Zimbabwe that codeine-containing

cough syrup was being illegally smuggled in from South Africa and sold on the streets

after being outlawed in 2015 [63]. However, there are limited data in many countries on

the prevalence of such activities and the extent of codeine use and misuse.

Abuse of OTC codeine products has been encouraged and normalised on social media

and in hip-hop music [64–66], which is a growing public health concern. A scoping

review of the non-medical use of pharmaceuticals identified several ways in which

codeine was reportedly being abused, including mixtures with alcohol or soft drinks

(e.g. "purple drank") and in the production of home-made opiates (e.g. "krokodil") [67].

Public health interventions are needed to prevent abuse, misuse, and pharmacy shopping

of OTC codeine products, particularly in young adults. The hip-hop song titled "1-800-

273-8255", from the US suicide prevention lifeline, significantly increased public

awareness of and calls to the hotline [68]. Thus, hip-hop music can also be used as a

tool to tailor public health messages to young adults and counteract the glamourisation

of codeine misuse.

The growing recognition of codeine abuse and misuse has led governments in

Minnesota, Manitoba, France, and Australia to reclassify codeine to prescription-only

[41–44]. In South Africa, Canada, Switzerland, Ireland, and the UK, governments are

proposing or considering plans to reclassify codeine-containing products to

prescription-only [69–72]. Studies assessing the effect of rescheduling codeine to

prescription-only in Australia showed a reduction in all codeine-related poisonings and

no change in calls to poisons centres or sales of high-strength (>15 mg) prescribed

codeine after reclassification [47,73]. The success of Australia's rescheduling questions

whether governments worldwide should make codeine prescription-only. But since

many low- and middle-income countries experience barriers to accessing opioids [74–

76], the WHO recommends that codeine should not be upscheduled and for codeine to

be included in essential medicines lists [77]. If a consensus on the status of OTC

codeine products cannot be reached, data should be collected globally to monitor its use

and harms.

Changes to regulations of OTC codeine and differences in trade exemptions and

disclosures of commercial interests at the country level may explain some of the

variation in sales [39]. For example, in France, sales fell considerably after codeine

became prescription-only. In countries such as Canada, Germany, and the USA, which

had high rates of prescribed opioids [76], mean sales of OTC codeine products were

lower than countries without similar access to prescribed opioids. However, our figures

depended on coverage of data from *IQVIA* during this time. Thus, it is hard to determine

whether the variation in sales represents real differences between countries.

Strengths and limitations

We used *IQVIA's* standard units to measure sales, which allows liquid and solid dosage

forms to be combined. The figures represent population-level sales and expenditure of

OTC codeine in 31 countries, providing the best available proxy for actual use. Sales

represented adult pain relief, although we calculated rates using population statistics for

all age groups, including children. Codeine-containing products may also be purchased

in large quantities from online pharmacies or the black market [78], not captured in

these data. *IQVIA*'s coverage and the completeness of data may have also affected sales

trends; they provided percentages on data coverage and converted expenditure to GBP

at single time points, which may not accurately represent changes to such metrics over

time.

Better access to OTC sales data is required. Amendments to medicines legislation in the

UK show how such data could be collected. The Misuse of Drugs Regulation 2001, the

Medicines for Human Use (Administration and Sale or Supply) (Miscellaneous

Amendments) Order 2007, and the Medicines (Sale or Supply) (Miscellaneous

Provisions Amendment Regulations 2007) were updated to require pharmacies to

submit counts of private prescriptions for Schedule 2 and Schedule 3 controlled drugs to

the National Health Service (NHS) Prescription Services for analysis, audit, and

monitoring [79,80]. A similar system could be enforced through a public health

organisation such as the WHO or the International Narcotics Control Board (INCB),

which already collects governments' annual drug statistics on opioids [81]. Such data

could then be used by governments and researchers to monitor sales of OTC codeine

and measure the impact of regulatory changes.

Conclusions

Codeine is one of the most widely accessible and used opioids worldwide. However,

monitoring its use and preventing its misuse as an OTC product is a public health

challenge. Healthcare professionals should ask their patients about their use of OTC

products. Public health measures are needed to identify and prevent codeine misuse and

increase awareness and education of the harms of codeine, particularly in young adults.

Governments should review policies to improve the collection of sales data and safety

of products sold OTC containing codeine.

Supplementary material

Supplement 1: Supplementary tables and figures

Supplement 2: STROBE reporting checklist

Declarations

Funding

This research was supported by the Primary Care Research Trust of Birmingham and

Midlands Research Practices Consortium who provided the funding to purchase the

sales data from IQVIA.

Competing interests

GCR was financially supported by the National Institute for Health Research (NIHR)

School for Primary Care Research (SPCR), the Naji Foundation, and the Rotary

Foundation to study for a Doctor of Philosophy (2017-2020), but no longer has any

financial COIs. GCR is an Associate Editor of BMJ Evidence Based Medicine. JKA has

published articles and edited textbooks on adverse drug reactions and interactions and

has often given medicolegal advice, including appearances as an expert witness in

coroners' courts, often dealing with the adverse effects of opioids. BM works for NHS

England as a pharmacist adviser. BG has received research funding from the Laura and

John Arnold Foundation, the NIHR, the NIHR SPCR, the NIHR Oxford Biomedical

Research Centre, the Mohn-Westlake Foundation, NIHR Applied Research

Collaboration Oxford and Thames Valley, the Wellcome Trust, the Good Thinking

Foundation, Health Data Research UK (HDRUK), the Health Foundation, and the

World Health Organisation (WHO); he also receives personal income from speaking

and writing for lay audiences on the misuse of science. FDRH acknowledges part

support from the NIHR SPCR, the NIHR Applied Research Collaboration (ARC)

Oxford Thames Valley, and the NIHR Oxford OUH BRC. CH is an NIHR Senior

Investigator and has received expenses and fees for his media work, received expenses

from the WHO, FDA, and holds grant funding from the NIHR SPCR and the NIHR

SPCR Evidence Synthesis Working Group [Project 380], the NIHR BRC Oxford and

the WHO. On occasion, CH receives expenses for teaching EBM and is also paid for his

GP work in NHS out of hours (contract with Oxford Health NHS Foundation Trust).

CH is the Director of the CEBM. The views expressed are those of the authors and not

necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

349   Availability of data and material

350   Study materials are available on an open repository [61] (https://osf.io/yt6bf/). We

351   cannot openly share the data owing to contractual agreements with *IQVIA,* but the data

352   can be accessed directly from *IQVIA*, which will require a fee.

354   Code availability

355   Our statistical code is openly available at GitHub [58]

356   (https://github.com/georgiarichards/otc_codeine).

358   Authors' contributions

GCR devised the research question, designed the methods, wrote the protocol,

conducted a literature search, sourced the data, cleaned, managed, and analysed the

data, created the figures, and wrote the first draft of the manuscript. JKA and CH

reviewed the protocol and preliminary findings and provided supervisory support.

FDRH reviewed the protocol and facilitated the grant application. BM reviewed

preliminary findings and contributed to the interpretation of data. BG provided

supervisory support.

Ethics approval

Not applicable

Consent to participate

Not applicable

Consent to publication

All authors read and approved the final manuscript and consent to submit the

manuscript for publication.

Figure 1: Mean sales of over-the-counter (OTC) codeine-containing units sold over six

years in 31 countries with data in descending order

Figure 2: Sales of over-the-counter (OTC) products containing codeine per 1000 of the

population starting in April 2013 to March 2014, and ending in April 2018 to March

2019, for countries in the top quartile (A) and second quartile (B) of sales

387   **Figure 3:** Public expenditure on over-the-counter (OTC) codeine-containing units sold

388   over six years in 31 countries with data in descending order

391
392 **Figure 4:** Percentage changes in sales (units sold per 1000 of the population) and public

393   expenditure (£ per 1000) on products containing codeine sold over-the-counter (OTC)

394   in 31 countries between April 2013 and March 2019

References

1. Curtis HJ, Croker R, Walker AJ, Richards GC, Quinlan J, Goldacre B. Opioid prescribing trends and geographical variation in England, 1998-2018: a retrospective database study. Lancet Psych [Internet]. The DataLab, Nuffield Department of Primary Care Health Sciences, University of Oxford, Radcliffe Observatory Quarter, Oxford, UK. Centre for Evidence Based Medicine, Nuffield Department of Primary Care Health Sciences, University of Oxford, Radcliffe Obs; 2019;6:140–50. Available from: http://dx.doi.org/10.1016/S2215-0366(18)30471-1.

2. Davies E, Sewell B, Jones M, Phillips CJ, Rance JY. Examining opioid prescribing trends for non-cancer pain using an estimated oral morphine equivalence measure: a retrospective cohort study between 2005 and 2015. BJGP Open [Internet]. Royal College of General Practitioners; 2020 [cited 2021 Jan 7];bjgpopen20X101122. Available from: https://bjgpopen.org/content/early/2020/11/30/bjgpopen20X101122.

3. Gomes T, Mamdani MM, Paterson JM, Dhalla IA, Juurlink DN. Trends in high-dose opioid prescribing in Canada. Can Fam Physician [Internet]. Scientist at the Institute for Clinical Evaluative Sciences; Assistant Professor at the Institute for Health Policy, Management and Evaluation and the Leslie Dan Faculty of Pharmacy at the University of Toronto; and Scientist at the Li Ka Shing Knowledge ; 2014;60:826–32. Available from: https://www.ncbi.nlm.nih.gov/pubmed/25217680.

4. Wagemaakers FN, Hollingworth SA, Kreijkamp-Kaspers S, Tee EHL, Leendertse AJ, van Driel ML. Opioid analgesic use in Australia and The Netherlands: a cross-country comparison. Int J Clin Pharm. 2017;39.

5. Goodman-Meza D, Friedman J, Kalmin MM, Aguilar-Posada E, Seamans MJ, Velazquez-Moreno S, et al. Geographical and socioeconomic disparities in opioid access in Mexico, 2015–19: a retrospective analysis of surveillance data. Lancet Public Heal [Internet]. Elsevier; 2021 [cited 2021 Jan 29];6:e88–96. Available from: https://linkinghub.elsevier.com/retrieve/pii/S2468266720302607.

6. Kalkman GA, Kramers C, Van Dongen RT, Van Den Brink W, Schellekens A. Trends in use and misuse of opioids in the Netherlands: a retrospective, multi-source database study. Lancet Public Heal [Internet]. 2019 [cited 2019 Oct

29];4:498–505. Available from: www.thelancet.com.

7. Krawczyk N, Greene MC, Zorzanelli R, Bastos FI. Rising trends of prescription
opioid sales in contemporary Brazil, 2009-2015. Am J Public Health 2018.

8. Dutch MJ. Nurofen Plus misuse: An emerging cause of perforated gastric ulcer.
Med J Aust [Internet]. Australasian Medical Publishing Co. Ltd; 2008 [cited 2020
Sep 7];188:56–7. Available from:
https://www.mja.com.au/journal/2008/188/1/nurofen-plus-misuse-emerging-cause-
perforated-gastric-ulcer.

9. Frei MY, Nielsen S, Dobbin MDH, Tobin CL. Serious morbidity associated with
misuse of over-the-counter codeine-ibuprofen analgesics: A series of 27 cases. Med
J Aust. Australasian Medical Publishing Co. Ltd; 2010;193:294–6.

10. Lambert AP. Life-threatening hypokalaemia from abuse of Nurofen Plus. J R Soc
Med [Internet]. SAGE Publications; 2005 [cited 2020 Sep 7];98:21–21. Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1079233.

11. Roussin A, Bouyssi A, Pouché L, Pourcel L, Lapeyre-Mestre M. Misuse and
Dependence on Non-Prescription Codeine Analgesics or Sedative H1
Antihistamines by Adults: A Cross-Sectional Investigation in France. PLoS One
[Internet]. Public Library of Science; 2013 [cited 2020 Sep 24];8. Available from:
/pmc/articles/PMC3789666/?report=abstract.

12. Carney T, Wells J, Parry CDH, McGuinness P, Harris R, Van Hout MC. A
comparative analysis of pharmacists' perspectives on codeine use and misuse - A
three country survey. Subst Abus Treat Prev Policy. BioMed Central Ltd.; 2018;13.

13. Sproule BA, Busto UE, Somer G, Romach MK, Sellers EM. Characteristics of
dependent and nondependent regular users of codeine. J Clin Psychopharmacol
[Internet]. J Clin Psychopharmacol; 1999 [cited 2020 Sep 7];19:367–72. Available
from: https://pubmed.ncbi.nlm.nih.gov/10440466.

14. Wu Q, Yu J, Yang C, Chen J, Yang L, Zhang H, et al. Nonmedical use of cough
syrup among secondary vocational school students: A national survey in China.
Med (United States) [Internet]. Lippincott Williams and Wilkins; 2016 [cited 2021
Jan 18];95. Available from: https://pubmed.ncbi.nlm.nih.gov/26962800.

15. Kimergård A, Foley M, Davey Z, Dunne J, Drummond C, Deluca P. Codeine use,
dependence and help-seeking behaviour in the UK and Ireland: an online cross-

sectional survey. QJM An Int J Med [Internet]. Oxford University Press; 2017

[cited 2021 Jan 18];110:559–64. Available from:

https://academic.oup.com/qjmed/article/110/9/559/3098674.

16.   Agaba E, Agaba P, Wigwe C. Use and abuse of analgesics in Nigeria: a community

survey. Niger J Med [Internet]. 2004 [cited 2021 Jan 29];13:379–82. Available

from: https://pubmed.ncbi.nlm.nih.gov/15523865.

17.   Orriols L, Gaillard J, Lapeyre-Mestre M, Roussin A. Evaluation of abuse and

dependence on drugs used for self-medication: A pharmacoepidemiological pilot

study based on community pharmacies in France. Drug Saf [Internet]. Drug Saf;

2009 [cited 2020 Sep 11];32:859–73. Available from:

https://pubmed.ncbi.nlm.nih.gov/19722729.

18.   Kontogiorgis C, Nena E, Berberoglou E, Moschoni K, Polyzois S, Tselemponis A,

et al. Estimating Consumers' Knowledge and Attitudes Towards Over-The-Counter

Analgesic Medication in Greece in the Years of Financial Crisis: The Case of

Paracetamol. Pain Ther. 2016;5.

19.   Cooper RJ. "I can't be an addict. I am." Over-the-counter medicine abuse: a

qualitative study. BMJ Open [Internet]. British Medical Journal Publishing Group;

2013 [cited 2020 Oct 6];3:2913. Available from: http://bmjopen.bmj.com.

20.   Lee E, Cooper RJ. Codeine addiction and internet forum use and support:

Qualitative netnographic study. J Med Internet Res [Internet]. Journal of Medical

Internet Research; 2019 [cited 2020 Oct 6];21. Available from:

/pmc/articles/PMC6658256/?report=abstract.

21.   Van Hout MC, Rich E, Dada S, Bergin M. "codeine Is My Helper": Misuse of and

Dependence on Codeine-Containing Medicines in South Africa. Qual Health Res.

2017;27.

22.   Kinnaird E, Kimergård A, Jennings S, Drummond C, Deluca P. From pain

treatment to opioid dependence: A qualitative study of the environmental influence

on codeine use in UK adults. BMJ Open [Internet]. BMJ Publishing Group; 2019

[cited 2021 Jan 12];9:e025331. Available from: http://bmjopen.bmj.com.

23.   Roxburgh A, Hall WD, Burns L, Pilgrim J, Saar E, Nielsen S, et al. Trends and

characteristics of accidental and intentional codeine overdose deaths in Australia.

Med J Aust. Australasian Medical Publishing Co. Ltd; 2015;203:299.e1-299.e7.

24. Hopkins RE, Dobbin M, Pilgrim JL. Unintentional mortality associated with
paracetamol and codeine preparations, with and without doxylamine, in Australia.
Forensic Sci Int [Internet]. Elsevier Ireland Ltd; 2018 [cited 2020 Jul 2];282:122–6.
Available from: https://pubmed.ncbi.nlm.nih.gov/29182956.

25. Schifano F, Zamparutti G, Zambello F, Oyefeso A, Deluca P, Balestrieri M, et al.
Review of deaths related to analgesic- and cough suppressant-opioids; England and
Wales 1996-2002. Pharmacopsychiatry. Pharmacopsychiatry; 2006. p. 185–91.

26. Cairns R, Brown JA, Buckley NA. The impact of codeine re-scheduling on misuse:
a retrospective review of calls to Australia's largest poisons centre. Addiction
[Internet]. Blackwell Publishing Ltd; 2016 [cited 2020 Sep 7];111:1848–53.
Available from: https://pubmed.ncbi.nlm.nih.gov/27177599.

27. Mill D, Johnson JL, Cock V, Monaghan E, Hotham ED. Counting the cost of over-
the-counter codeine containing analgesic misuse: A retrospective review of hospital
admissions over a 5 year period. Drug Alcohol Rev [Internet]. Blackwell
Publishing; 2018 [cited 2020 Sep 7];37:247–56. Available from:
https://pubmed.ncbi.nlm.nih.gov/28925091.

28. Derry S, Moore RA, McQuay HJ. Single dose oral codeine, as a single agent, for
acute postoperative pain in adults. Cochrane Database Syst Rev [Internet]. Wiley;
2010 [cited 2021 Jan 18]; Available from:
https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD008099.pub2/full.

29. Omar MI, Alexander CE. Drug treatment for faecal incontinence in adults
[Internet]. Cochrane Database Syst. Rev. John Wiley and Sons Ltd; 2013 [cited
2020 Sep 3]. Available from:
http://doi.wiley.com/10.1002/14651858.CD002116.pub2.

30. Molassiotis A, Bailey C, Caress A, Tan JY. Interventions for cough in cancer
[Internet]. Cochrane Database Syst. Rev. John Wiley and Sons Ltd; 2015 [cited
2020 Sep 4]. Available from:
http://doi.wiley.com/10.1002/14651858.CD007881.pub3.

31. Derry S, Karlin SM, Moore RA. Single dose oral ibuprofen plus codeine for acute
postoperative pain in adults [Internet]. Cochrane Database Syst. Rev. John Wiley
and Sons Ltd; 2015 [cited 2020 Sep 4]. Available from:
http://doi.wiley.com/10.1002/14651858.CD010107.pub3.

32. Toms L, Derry S, Moore RA, McQuay HJ. Single dose oral paracetamol
(acetaminophen) with codeine for postoperative pain in adults [Internet]. Cochrane
Database Syst. Rev. John Wiley and Sons Ltd; 2009 [cited 2020 Sep 4]. Available
from: http://doi.wiley.com/10.1002/14651858.CD001547.pub2.
33. Moore RA, Wiffen PJ, Derry S, Maguire T, Roy YM, Tyrrell L. Non-prescription
(OTC) oral analgesics for acute pain - an overview of Cochrane reviews [Internet].
Cochrane Database Syst. Rev. John Wiley and Sons Ltd; 2015 [cited 2020 Sep 3].
Available from: http://doi.wiley.com/10.1002/14651858.CD010794.pub2.
34. Abdel Shaheed C, Maher CG, McLachlan AJ. Efficacy and Safety of Low-dose
Codeine-containing Combination Analgesics for Pain. Clin J Pain [Internet]. 2019
[cited 2019 Dec 14];35:836–43. Available from:
http://insights.ovid.com/crossref?an=00002508-201910000-00005.
35. Agnich LE, Stogner JM, Miller BL, Marcum CD. Purple drank prevalence and
characteristics of misusers of codeine cough syrup mixtures. Addict Behav
[Internet]. Addict Behav; 2013 [cited 2021 Jan 18];38:2445–9. Available from:
https://pubmed.ncbi.nlm.nih.gov/23688907.
36. Soares JX, Alves EA, Silva AMN, De Figueiredo NG, Neves JF, Cravo SM, et al.
Street-Like Synthesis of Krokodil Results in the Formation of an Enlarged Cluster
of Known and New Morphinans. Chem Res Toxicol [Internet]. American Chemical
Society; 2017 [cited 2021 Jan 18];30:1609–21. Available from:
https://pubs.acs.org/doi/abs/10.1021/acs.chemrestox.7b00126.
37. Bedford KR, Nolan SL, Onrust R, Siegers JD. The illicit preparation of morphine
and heroin from pharmaceutical products containing codeine: "homebake"
laboratories in New Zealand. Forensic Sci Int [Internet]. Forensic Sci Int; 1987
[cited 2021 Jan 18];34:197–204. Available from:
https://pubmed.ncbi.nlm.nih.gov/3666623.
38. Oliver T, Gheevarghese SJ, Gandhi U, Bhat ZY, Pillai U. "Krokodil"-A menace
slowly spreading across the Atlantic [Internet]. Am. J. Ther. Lippincott Williams
and Wilkins; 2015 [cited 2021 Jan 18]. p. 231–3. Available from:
https://pubmed.ncbi.nlm.nih.gov/25756471.
39. WHO. Pre-Review Report: Preparations of codeine listed in Schedule lll of the
1961 Single Convention on Narcotic Drugs [Internet]. Geneva; 2019 Oct. Available

from: https://www.who.int/medicines/access/controlled-
substances/Final_codeine.pdf.

40. United Nations. United Nations Single Convention on Narcotic Drugs, 1961
[Internet]. 1961 [cited 2021 Jan 20]. Available from:
https://www.emcdda.europa.eu/drugs-library/single-convention-narcotic-drugs-
1961_en.

41. Office of the Revisor of Statutes. 2020 Minnesota Statutes [Internet]. Minnesota
Legis. 2020 [cited 2020 Jan 24]. Available from:
https://www.revisor.mn.gov/statutes/cite/152.02.

42. College of Pharmacists of Manitoba. Exempted codeine changes [Internet]. 2016
[cited 2021 Jan 24]. Available from:
https://mpha.in1touch.org/site/exemptedcodeine?nav=public.

43. Wilkes D. France restricts dextromethorphan and codeine [Internet]. OTC Toolbox.
2017 [cited 2021 Jan 24]. Available from: https://www.otctoolbox.com/industry-
news/france-restricts-dextromethorphan-and-codeine.

44. Kirby T. The end of over-the-counter codeine in Australia. Lancet [Internet]. 2018
[cited 2019 Dec 13]; Available from: www.thelancet.com/psychiatry.

45. Foley M, Harris R, Rich E, Rapca A, Bergin M, Norman I, et al. The availability of
over-the-counter codeine medicines across the European Union. Public Health.
Elsevier; 2015;129:1465–70.

46. Lo MY, Ong MW, Lin JG, Sun WZ. Codeine consumption from over-the-counter
anti-cough syrup in Taiwan: A useful indicator for opioid abuse. Acta Anaesthesiol
Taiwanica [Internet]. Elsevier Taiwan LLC; 2015 [cited 2021 Jan 20];53:135–8.
Available from: https://pubmed.ncbi.nlm.nih.gov/26646544.

47. Cairns R, Schaffer AL, Brown JA, Pearson S, Buckley NA. Codeine use and harms
in Australia: evaluating the effects of re☐scheduling. Addiction [Internet].
Blackwell Publishing Ltd; 2020 [cited 2020 May 17];115:451–9. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/add.14798.

48. Gisev N, Nielsen S, Cama E, Larance B, Bruno R, Degenhardt L. An ecological
study of the extent and factors associated with the use of prescription and over-the-
counter codeine in Australia. Eur J Clin Pharmacol [Internet]. Springer Verlag;
2016 [cited 2020 Jul 7];72:469–94. Available from:

https://link.springer.com/article/10.1007/s00228-015-1995-8.

49. Consumer Health - IQVIA [Internet]. [cited 2020 Jun 12]. Available from:

https://www.iqvia.com/solutions/industry-segments/consumer-health.

50. Hsia Y, Sharland M, Jackson C, Wong ICK, Magrini N, Bielicki JA. Consumption

of oral antibiotic formulations for young children according to the WHO Access,

Watch, Reserve (AWaRe) antibiotic groups: an analysis of sales data from 70

middle-income and high-income countries. Lancet Infect Dis. Lancet Publishing

Group; 2019;19:67–75.

51. Suchard MA, Schuemie MJ, Krumholz HM, You SC, Chen RJ, Pratt N, et al.

Comprehensive comparative effectiveness and safety of first-line antihypertensive

drug classes: a systematic, multinational, large-scale analysis. Lancet. Lancet

Publishing Group; 2019;394:1816–26.

52. Guy GP, Haegerich TM, Evans ME, Losby JL, Young R, Jones CM. Vital Signs:

Pharmacy-Based Naloxone Dispensing — United States, 2012–2018. MMWR

Morb Mortal Wkly Rep [Internet]. NLM (Medline); 2019 [cited 2020 Jun

12];68:679–86. Available from:

http://www.cdc.gov/mmwr/volumes/68/wr/mm6831e1.htm?s_cid=mm6831e1_w.

53. Smith B, Culligan C, McFarlane A, Rickwood S, Clark C. The changing dynamic

of the UK retail pharmaceutical market [Internet]. 2018. Available from:

https://www.iqvia.com/-/media/iqvia/pdfs/uk/the-changing-dynamic-of-the-uk-

retail-pharmaceutical-market.pdf.

54. The World Bank. Population data [Internet]. 2019 [cited 2020 May 28]. Available

from:

https://data.worldbank.org/indicator/SP.POP.TOTL?end=2018&start=1960&view=

chart.

55. PANDAS Development Team. pandas-dev/pandas: PANDAS. Zenodo; 2020.

56. Waskom M. mwaskom/seaborn. Zenodo; 2020.

57. Hunter J. Matplotlib: a 2D graphics environment. Comput Sci Eng. 2009;9:90–5.

58. Richards G. georgiarichards/otc_codeine [Internet]. GitHub. 2020 [cited 2021 Apr

3]. Available from: https://github.com/georgiarichards/otc_codeine.

59. Richards G. Sales of over-the-counter (OTC) codeine-containing products in 31

countries, 2013-2019 [Internet]. OSF Regist. 2020 [cited 2021 Apr 3]. Available

from: https://osf.io/ay4mc.

60. Richards GC, Mahtani KR, Goldacre B, Heneghan C. 135 Trends and variation in

the sales of over-the-counter analgesics: a protocol for a retrospective database

study and policy review. BMJ Evidence-Based Med [Internet]. BMJ; 2018 [cited

2021 Jan 5]. p. A63.2-A64. Available from:

https://ebm.bmj.com/content/23/Suppl_2/A63.2.

61. Richards GC, Hobbs R, Aronson JK, Heneghan C. Sales of over-the-counter (OTC)

codeine-containing products in 31 countries, 2013-2019 [Internet]. OSF. OSF; 2021

[cited 2021 Jan 4]. Available from: https://osf.io/yt6bf.

62. Weich L, Perkel C, van Zyl N, Rataemane S, Naidoo L. Review Article: Medical

management of opioid dependence in South Africa. South African Med J [Internet].

2008 [cited 2021 Jan 31];98. Available from:

https://www.ajol.info/index.php/samj/article/view/13959.

63. VICE Video. Zimbabwe's Codeine Cough Syrup Epidemic [Internet]. 2018 [cited

2021 Feb 1]. Available from: https://video.vice.com/en_uk/video/vice-zimbabwes-

codeine-cough-syrup-epidemic-1-104/5c373790be407707904345c1.

64. Cherian R, Westbrook M, Ramo D, Sarkar U. Representations of codeine misuse

on instagram: Content analysis. J Med Internet Res [Internet]. JMIR Publications

Inc.; 2018 [cited 2021 Jan 19];20. Available from:

https://pubmed.ncbi.nlm.nih.gov/29559422.

65. Tettey N-S, Siddiqui K, Llamoca H, Nagamine S, Ahn S. Purple Drank, Sizurp, and

Lean: Hip-Hop Music and Codeine Use, A Call to Action for Public Health

Educators. Int J Psychol Stud [Internet]. Canadian Center of Science and

Education; 2020 [cited 2021 Jan 19];12:42. Available from:

http://www.ccsenet.org/journal/index.php/ijps/article/view/0/42126.

66. Peteet B, Roundtree C, Dixon S, Mosley C, Miller-Roenigk B, White J, et al.

'Codeine crazy:' a content analysis of prescription drug references in popular

music. J Youth Stud [Internet]. Routledge; 2020 [cited 2021 Jan 19]; Available

from: https://www.tandfonline.com/doi/abs/10.1080/13676261.2020.1801992.

67. Van Hout MC. Kitchen chemistry: A scoping review of the diversionary use of

pharmaceuticals for non-medicinal use and home production of drug solutions

[Internet]. Drug Test. Anal. John Wiley and Sons Ltd; 2014 [cited 2021 Jan 28]. p.

778–87. Available from: https://pubmed.ncbi.nlm.nih.gov/24619569.

68. Torgerson T, Swayze C, Sanghera S, Cooper C, Beaman J, Hartwell M, et al.

Public awareness of the National Suicide Prevention Lifeline following the release

of a hip-hop song. BMJ Evidence-Based Med [Internet]. BMJ; 2021 [cited 2021

Feb 12];bmjebm-2020-111509. Available from:

https://ebm.bmj.com/content/early/2021/01/28/bmjebm-2020-111509.

69. de Villiers J. Here are the over-the-counter medicines you may need a prescription

for in future - including Adcodol and Nurofen Plus [Internet]. Bus. Insid. SA. 2020

[cited 2021 Jan 25]. Available from: https://www.businessinsider.co.za/codeine-

containing-medicines-in-south-africa-to-be-affected-by-south-african-health-

products-regulatory-authoritys-up-scheduling-move-2020-1.

70. Ridley D. Swiss experts advise codeine reverse-switch [Internet]. LinkedIn. 2018

[cited 2021 Jan 25]. Available from: https://www.linkedin.com/pulse/swiss-experts-

advise-codeine-reverse-switch-david-ridley.

71. MHRA. UK regulator strengthens opioid warnings [Internet]. Vigil. Saf. alerts

Guid. 2020 [cited 2021 Jan 20]. Available from:

https://www.gov.uk/government/news/uk-regulator-strengthens-opioid-warnings.

72. Government of Canada. Forward Regulatory Plan 2019-2021: Regulations

Amending the Narcotic Control Regulations under the Controlled Drugs and

Substances Act to make all products containing codeine available by prescription

only while ensuring access to these medications [Internet]. 2019 [cited 2021 Jan

28]. Available from: https://www.canada.ca/en/health-canada/corporate/about-

health-canada/legislation-guidelines/acts-regulations/forward-regulatory-

plan/plan/codeine-prescription.html.

73. Schaffer AL, Cairns R, Brown JA, Gisev N, Buckley NA, Pearson SA. Changes in

sales of analgesics to pharmacies after codeine was rescheduled as a prescription

only medicine. Med J Aust. John Wiley and Sons Inc.; 2020;212:321–7.

74. Fadhil I, Lyons G, Payne S. Barriers to, and opportunities for, palliative care

development in the Eastern Mediterranean Region. Lancet Oncol. Lancet

Publishing Group; 2017. p. e176–84.

75. Berterame S, Erthal J, Thomas J, Fellner S, Vosse B, Clare P, et al. Use of and

barriers to access to opioid analgesics: A worldwide, regional, and national study.

Lancet. 2016;387.

76. Richards GC, Aronson JK, Mahtani KR, Heneghan C. Global, regional, and

national consumption of controlled opioids: a cross-sectional study of 214 countries

and non-metropolitan territories. Br J Pain [Internet]. SAGE Publications Ltd; 2021

[cited 2021 Jun 11]; Available from: https://doi.org/10.1177/20494637211013052.

77. Richards GC, Aronson JK, Heneghan C, Mahtani KR, Koshiaris C, Persaud N.

Relation between opioid consumption and inclusion of opioids in 137 national

essential medicines lists. BMJ Glob Heal. 2020;5:1–8.

78. Marsh S. "I could get 500 tablets a month": the ease of buying opiates online

[Internet]. Guard. 2019 [cited 2021 Jan 24]. Available from:

https://www.theguardian.com/society/2019/apr/26/i-could-get-500-tablets-a-month-

ease-of-buying-opiates-online.

79. NHS England. The Controlled Drugs (Supervision of Management and Use)

Regulations 2013 [Internet]. Single Oper. Model. 2013 [cited 2021 Apr 19].

Available from: https://www.england.nhs.uk/wp-content/uploads/2013/11/som-

cont-drugs.pdf.

80. PSNC. Controlled Drug prescription forms and validity [Internet]. Pharm. Serv.

Negot. Comm. 2020 [cited 2021 Apr 19]. Available from:

https://psnc.org.uk/dispensing-supply/dispensing-controlled-drugs/controlled-drug-

prescription-forms-and-validity.

81. INCB. Narcotic Drugs — Estimated World Requirements for 2020 — Statistics for

2018 [Internet]. Vienna; 2019. Available from:

https://apps.who.int/iris/bitstream/handle/10665/325771/WHO-MVP-EMP-IAU-

2019.06-eng.pdf.