Estimating local outbreak risks and the effects of non-pharmaceutical interventions in age-structured populations: SARS-CoV-2 as a case study

Authors

Francesca A. Lovell-Read[1*], Silvia Shen[1,2], Robin N. Thompson[3,4]

Affiliations

[1]Mathematical Institute, University of Oxford, Oxford, UK

[2]Pembroke College, University of Oxford, Oxford, UK

[3]Mathematics Institute, University of Warwick, Coventry, UK

[4]The Zeeman Institute for Systems Biology and Infectious Disease Epidemiology Research, University of Warwick, Coventry, UK

* Corresponding author

Address: Merton College, Merton Street, Oxford, OX1 4JD

E-mail: francesca.lovell-read@merton.ox.ac.uk. Tel: +44 (0)1865 273525.

Abstract

During the COVID-19 pandemic, non-pharmaceutical interventions (NPIs) including school closures, workplace closures and social distancing policies have been employed worldwide to reduce transmission and prevent local outbreaks. However, transmission and the effectiveness of

NPIs depend strongly on age-related factors including heterogeneities in contact patterns and

pathophysiology. Here, using SARS-CoV-2 as a case study, we develop a branching process

model for assessing the risk that an infectious case arriving in a new location will initiate a local

outbreak, accounting for the age distribution of the host population. We show that the risk of a

local outbreak depends on the age of the index case, and we explore the effects of NPIs targeting

individuals of different ages. Social distancing policies that reduce contacts outside of schools

and workplaces and target individuals of all ages are predicted to reduce local outbreak risks

substantially, whereas school closures have a more limited impact. In the scenarios considered

here, when different NPIs are used in combination the risk of local outbreaks can be eliminated.

We also show that heightened surveillance of infectious individuals reduces the level of NPIs

required to prevent local outbreaks, particularly if enhanced surveillance of symptomatic cases is

combined with efforts to find and isolate nonsymptomatic infected individuals. Our results

reflect real-world experience of the COVID-19 pandemic, during which combinations of intense

NPIs have reduced transmission and the risk of local outbreaks. The general modelling

framework that we present can be used to estimate local outbreak risks during future epidemics

of a range of pathogens, accounting fully for age-related factors.

Keywords

mathematical modelling; infectious disease epidemiology; outbreak probability; SARS-CoV-2;

COVID-19; age-structured models; non-pharmaceutical interventions.

1. Introduction

Throughout the COVID-19 pandemic, policy makers worldwide have relied on non-pharmaceutical interventions (NPIs) to limit the spread of SARS-CoV-2. Commonly introduced NPIs have included school closures, workplace closures and population-wide social distancing policies, all of which aim to reduce the numbers of contacts between individuals and disrupt potential chains of transmission [1-4]. Similar measures have previously been adopted for countering other infectious diseases such as Ebola and pandemic influenza [5-7], and are likely to remain a key line of defence against emerging pathogens that are directly transmitted between hosts. NPIs are particularly important when no effective treatment or vaccine is available, and they are also beneficial when vaccination programmes are being rolled out [8-10]. If vaccines do not prevent transmission completely, then NPIs may be important even when vaccination is widespread [11]. However, the negative economic, social and non-disease health consequences of NPIs have been widely discussed, with the impact of school closures on the academic progress and wellbeing of school-aged individuals a particular concern [7, 12-16]. Therefore, assessing the effectiveness of different NPIs at reducing transmission is critical for determining whether or not they should be used.

Since NPIs such as school and workplace closures affect distinct age groups within the population, when evaluating their effectiveness it is important to account for age-dependent factors that influence transmission. Multiple studies have documented marked heterogeneities in the patterns of contacts between individuals in different age groups, with school-aged individuals tending to have more contacts each day than older individuals [17-23]. Since close contact between individuals is a key driver of transmission for respiratory pathogens such as influenza

viruses and SARS-CoV-2, these contact patterns influence transmission dynamics and

consequently the effects of interventions that target different age groups [18, 19, 24-27].

Additionally, many diseases are characterised by significant age-related variations in

pathophysiology. For example, for SARS-CoV-2, children may be less susceptible to infection

than adults [27-31], and more likely to experience asymptomatic or subclinical courses of

infection [28, 31-36]. Since the secondary attack rate (the proportion of close contacts that lead

to new infections) from asymptomatic or subclinical hosts is lower than from hosts with clinical

symptoms [37-42], children are likely to be less infectious on average than older individuals who

are at increased risk of developing symptoms [43-45].

Previous studies have used age-stratified deterministic transmission models to investigate the

effects of NPIs on COVID-19 epidemic peak incidence and timing. Prem *et al.* [46] projected the

outbreak in Wuhan, China, over a one year period under different control scenarios, and

demonstrated that a period of intense control measures including school closures, a 90%

reduction in the workforce and a significant reduction in other social mixing could delay the

epidemic peak by several months. Zhang *et al.* [27] predicted that eliminating all school contacts

during the outbreak period would lead to a noticeable decrease in the peak incidence and a later

peak; however, they did not take differences between symptomatic and asymptomatic cases into

account explicitly. In contrast, Davies *et al.* [31] used estimates of age-dependent susceptibility

and clinical fraction fitted to the observed age distribution of cases in six countries to

demonstrate that school closures alone were unlikely to reduce SARS-CoV-2 transmission

substantially. Davies *et al.* [47] subsequently concluded that a combination of several strongly

enforced NPIs would be necessary to avoid COVID-19 cases exceeding available healthcare

capacity in the UK.

Rather than considering the entire epidemic curve, here we focus on estimating the probability

that cases introduced to a new location trigger a local outbreak as opposed to fading out with few

cases. Localised clusters of transmission have been a feature of the COVID-19 pandemic [48-

50], and assessing the risk that such local outbreaks occur requires a stochastic model in which

the pathogen can either invade or fade out. Stochastic branching process models have been

applied previously to assess outbreak risks for many pathogens without considering different age

groups explicitly [51-55], and extended to consider adults and children as two distinct groups

[56]. However, the significant heterogeneities in contact patterns and pathophysiology between

individuals across the full range of ages have never previously been considered in estimates of

local outbreak risks. Here, we develop an age-structured branching process model that can be

used to estimate the probability of a local outbreak occurring for index cases of different ages,

and demonstrate how the age-dependent risk profile changes when susceptibility to infection and

clinical fraction vary with age.

We use the model to investigate the effects on the local outbreak probability of NPIs that reduce

the numbers of contacts between individuals. Specifically, we use location-specific contact data

for the UK detailing the average numbers of daily contacts occurring in school, in the workplace

and elsewhere [17] to model the impacts of school closures, workplace closures and broader

social distancing policies. We demonstrate that, for SARS-CoV-2, contacts occurring outside

schools and the workplace are a key driver of sustained transmission. Thus, population-wide

social distancing policies that affect individuals of all ages lead to a substantial reduction in the

risk of local outbreaks. In contrast, since school-aged individuals only make up around one

quarter of the UK population and tend to have large numbers of contacts outside school, school

closures are predicted to have only a limited effect when applied as the sole NPI.

We then go on to consider the impacts of mixed strategies made up of multiple NPIs, as well as

additional NPIs that do not only reduce numbers of contacts. Specifically, we show that rigorous

surveillance and effective isolation of infected hosts can reduce the level of contact-reducing

NPIs required to achieve substantial reductions in the risk of local outbreaks. Although we use

SARS-CoV-2 as a case study, our approach can be applied more generally to explore the effects

of NPIs on the risk of outbreaks of any pathogen for which age-related heterogeneities play a

significant role in transmission dynamics.

2. Methods

2.1 Mathematical model

We considered a branching process model in which the population was divided into 16 age

groups, denoted $G_1, G_2, \dots, G_{16}$. The first 15 groups represent individuals aged 0-74, divided into

five-year intervals (0-4, 5-9, 10-14 etc.). The final group represents individuals aged 75 and over.

The total number of individuals in age group G_k is denoted N_k. Infected individuals in each age

group G_k are classified into compartments representing asymptomatic (A_k), presymptomatic (P_k)

or symptomatic (S_k) hosts, where an individual in the A_k compartment does not develop

symptoms at any time during their course of infection.

An infected individual of any type in group G_k may generate new infections in any age group. In

our model, the rate at which a single infected symptomatic individual in group G_k generates

infections in group G_j is given by

138
$$\beta_{kj} = B\tau_k\omega_j C_{kj}.$$

Here, τ_k represents the infectivity of individuals in group G_k, ω_j represents the susceptibility to

infection of individuals in group G_j, C_{kj} represents the daily number of unique contacts a single

individual in group G_k has with individuals in group G_j, and B is a scaling factor that can be used

to set the reproduction number of the pathogen being considered (see Section 2.2). Since the

initial phase of potential local outbreaks are the focus of this study, we did not account for

depletion of susceptible hosts explicitly. The relative transmission rates from presymptomatic

and asymptomatic individuals compared to symptomatic individuals are given by the scaled

quantities $\eta\beta_{kj}$ and $\theta\beta_{kj}$, respectively, where η and θ were chosen so that the proportions of

transmissions generated by presymptomatic and asymptomatic hosts were in line with literature

estimates [57]. The parameter ξ_k represents the proportion of asymptomatic infections in group

G_k, so that a new infection in group G_k either increases A_k by one (with probability ξ_k) or

increases P_k by one (with probability $1 - \xi_k$).

A presymptomatic individual in group G_k may go on to develop symptoms (transition from P_k

to S_k) or be detected and isolated (so that P_k decreases by one). A symptomatic individual in

group G_k may be detected and isolated as a result of successful surveillance, or may be removed

due to self-isolation, recovery or death (so that S_k decreases by one in either case). Similarly, an

asymptomatic individual in group G_k may be detected and isolated or recover (so that A_k

decreases by one). A schematic of the different possible events in the model is shown in Fig 1.

The parameter λ represents the rate at which presymptomatic individuals develop symptoms, so

that the expected duration of the presymptomatic infectious period is $1/\lambda$ days in the absence of

surveillance of nonsymptomatic infected individuals. Similarly, the expected duration of the

asymptomatic infectious period in the absence of surveillance is $1/\nu$ days. The parameter μ

represents the rate at which symptomatic individuals are removed as a result of self-isolation,

recovery or death, so that the duration of time for which they are able to infect others is $1/\mu$

days.

For each group G_k, the rate at which symptomatic individuals are detected and isolated as a result

of enhanced surveillance is determined by the parameter ρ_k. Analogously, the parameter σ_k

governs the rate at which presymptomatic and asymptomatic individuals in G_k are detected and

isolated. We assumed that surveillance measures targeting nonsymptomatic hosts are equally

effective for those who are presymptomatic and those who are asymptomatic, and therefore used

the same rate of isolation due to surveillance for both of these groups.

Fig 1. The branching process model used in our analyses. Schematic showing the different possible

events in the branching process model and the rates at which they occur. The parameters of the model

are described in the text and in Tables 1, 2 and 3.

2.2 Reproduction number

The effective reproduction number, R, represents the expected number of secondary infections

generated by a single infected individual during their entire course of infection, accounting for

interventions that are in place. Here, we take a heuristic approach to derive the following

expression for R:

$$R = \sum_{k=1}^{16} \frac{N_k}{N} \sum_{l=1}^{16} \left[\frac{\xi_k \theta \beta_{kl}}{\nu + \sigma_k} + (1 - \xi_k) \left(\frac{\eta \beta_{kl}}{\lambda + \sigma_k} + \frac{\lambda}{\lambda + \sigma_k} \frac{\beta_{kl}}{\mu + \rho_k} \right) \right], \quad (1)$$

where $N = N_1 + \cdots + N_{16}$ is the total population size. To obtain this expression, we first consider

the expected number of secondary infections an infected individual in age group G_k will generate

in age group G_l. If an individual in age group G_k experiences a fully asymptomatic course of

infection, which occurs with probability ξ_k, they will generate new infections in age group G_l at

rate $\theta \beta_{kl}$ and recover or be isolated at rate $\nu + \sigma_k$. Therefore, the total number of infections they

are expected to cause in age group G_l is $\theta \beta_{kl}/(\nu + \sigma_k)$. If instead the individual in age group G_k

experiences a symptomatic course of infection, which occurs with probability $1 - \xi_k$, whilst

presymptomatic they will generate new infections in age group G_l at rate $\eta \beta_{kl}$ and be isolated or

develop symptoms at rate $\lambda + \sigma_k$. Thus, the total number of infections they are expected to cause

in age group G_l whilst presymptomatic is $\eta \beta_{kl}/(\lambda + \sigma_k)$. If they go on to develop symptoms

before being isolated, which occurs with probability $\lambda/(\lambda + \sigma_k)$, applying similar reasoning they

are expected to cause $\beta_{kl}/(\mu + \rho_k)$ new infections in age group G_l whilst symptomatic.

Combining these possibilities leads to the term in square brackets in expression (1), which is then

summed over all possible age groups G_l of the infectee. Finally, to obtain the full expression (1)

we take a weighted average across all possible age groups G_k of the infector, where the weights

N_k/N represent the proportions of the population belonging to each age group. This corresponds

to the assumption that the initial infected host is more likely to belong to an age group containing

more individuals than an age group with fewer individuals.

In the absence of interventions, i.e. when $\sigma_k = \rho_k = 0$ (representing no enhanced isolation as a

result of surveillance) and β_{kl} is calculated using contact patterns that are characteristic of

normal behaviour, the effective reproduction number, R, is equal to the basic reproduction

number, R_0.

2.3 Model parameterisation

The numbers of individuals in each age group (values of N_k) were chosen according to United

Nations age demographic data for the UK [58] (Fig 2A). The daily numbers of contacts between

individuals in each age group (values of C_{kj}) were set according to the 16x16 'contact matrix'

for the UK, in which the (k, j)th entry represents the expected daily number of unique contacts

an individual in age group G_k has with individuals in age group G_j [17]. In addition to matrices

representing 'all' contacts (Figure 2B), we also considered matrices detailing a breakdown into

'school', 'work', 'home' and 'other' contacts (Figures 2C-F), allowing us to investigate the

effects of control interventions that reduce contacts in each of these settings.

Since we considered SARS-CoV-2 as a case study, we used studies conducted during the

COVID-19 pandemic to inform the epidemiological parameters of our model. Despite previous

research assessing the relationships between age and factors such as susceptibility to SARS-

CoV-2 infection or the propensity to develop symptoms, there is some variation in estimated

parameters between different studies. To test the robustness of our results to this uncertainty, we

conducted our analyses under three different scenarios (A, B and C). In scenario A, we assumed

that susceptibility to infection (values of ω_k) and the proportion of hosts who experience a fully

asymptomatic course of infection (values of ξ_k) are independent of age. In scenario B,

susceptibility was assumed to vary with age but the proportion of asymptomatic infections is

independent of age. In scenario C, we allowed both susceptibility and the asymptomatic

proportion to vary with age. The values used for the parameters ω_k and ξ_k in each of these three

scenarios are shown in Table 1 (see also [31]).

In all scenarios considered, the inherent infectivity was not assumed to be age-dependent (i.e.

$\tau_k = 1$ for all values of k). In other words, the expected infectiousness of infected hosts in

different age groups was governed solely by the proportion of asymptomatic infections in that

age group. We chose the scaling factors η and θ for the relative transmission rates from

presymptomatic and asymptomatic individuals compared to symptomatic individuals so that the

proportions of infections arising from each of these groups were in line with literature estimates

(see Table 3 and [57]).

In the absence of enhanced isolation, we set the expected duration of the presymptomatic

infectious period and the time for which symptomatic individuals are able to infect others to be

$1/\lambda = 2$ days and $1/\mu = 8$ days, respectively [59-62]. The asymptomatic infectious period was

then chosen so that all infected individuals are expected to be infectious for the same period (i.e.

$1/\nu = 10$ days). In our initial analysis, we set the isolation rates ρ_k and σ_k equal to 0 for all

k; later, we considered the effects of increasing these rates.

Initially, we fixed $R_0 = 3$ (in line with initial estimates of SARS-CoV-2 transmissibility [63-66],

before the emergence of more transmissible variants) and used expression (1) to determine the

appropriate corresponding value of the scaling factor B. Later, when considering the impact of

NPIs on the probability of a local outbreak, we retained this value of B and used expression (1)

to determine how the reproduction number changes as a result of the control implemented.

Table 1. Baseline values of age-dependent parameters. Values used for the age-dependent relative

susceptibility to infection (ω_k) and the proportion of infections that are asymptomatic (ξ_k) for each of the

scenarios A, B and C [31].

Age group (G_k)	Relative susceptibility (ω_k)		Asymptomatic proportion (ξ_k)	
	Scenario A	Scenarios B & C	Scenarios A & B	Scenario C
G_1, G_2 (0-9)		0.4	0.584	0.71
G_3, G_4 (10-19)	1.0	0.38	(weighted average	0.79
G_5, G_6 (20-29)		0.79	of age-dependent	0.73

G_7, G_8 (30-39)	0.86	values of ξ_k in scenario C)	0.67
G_9, G_{10} (40-49)	0.8		0.6
G_{11}, G_{12} (50-59)	0.82		0.51
G_{13}, G_{14} (60-69)	0.88		0.37
G_{15}, G_{16} (70+)	0.74		0.31

257 **Table 2. Baseline values of scenario-independent parameters.** Values used for the parameters that were

258 assumed not to vary between scenarios A, B and C. We also considered strategies involving enhanced

259 surveillance ($\rho_k, \sigma_k > 0$) – see Fig 6.

Parameter	Meaning	Baseline value	Justification
R_0	Expected number of secondary infections generated by a single infectious host in the absence of interventions	$R_0 = 3$	Within estimated range for original SARS-CoV-2 virus [63-66]
λ	Rate at which presymptomatic hosts develop symptoms	$\lambda = 1/2$ days^{-1}	[59]
μ	Rate at which symptomatic hosts are removed due to self-isolation, recovery or death	$\mu = 1/8$ days^{-1}	[60-62]
ν	Rate at which asymptomatic hosts are removed due to recovery or death	$\nu = 1/10$ days^{-1}	Chosen so that, in the absence of interventions, the expected duration of infection is identical for all infected hosts $\left(\frac{1}{\nu} = \frac{1}{\lambda} + \frac{1}{\mu}\right)$
τ_k	(Relative) infectivity parameter	$\tau_k = 1$ for $k = 1, ..., 16$	Assumed
ρ_k	Isolation rate due to surveillance of symptomatic individuals	$\rho_k = 0$ for $k = 1, ..., 16$	N/A
σ_k	Isolation rate due to surveillance of nonsymptomatic individuals	$\sigma_k = 0$ for $k = 1, ..., 16$	N/A

261 **Table 3. Baseline values of scenario-dependent scaling parameters.** Values used for the scaling parameters

262 B, η and θ for each of the scenarios A, B and C.

Parameter	Meaning	Value			Justification
		Scenario A	Scenario B	Scenario C	
B	Transmission rate scaling factor	0.0386	0.0559	0.0607	Chosen so that $R_0 = 3$ [63-66]
η	Relative transmission rate of presymptomatic hosts compared to symptomatic hosts	4.83	4.83	4.83	Chosen so that the proportion of all infections arising from presymptomatic hosts is 0.489 [57]

	Relative transmission rate of asymptomatic hosts compared to symptomatic hosts	0.149	0.149	0.130	Chosen so that the proportion of all infections arising from entirely asymptomatic hosts is 0.106 [57]
θ					

2.4 Probability of a local outbreak

The probability that an infected individual in a particular age group initiates a local outbreak

when they are introduced into the population was calculated using the branching process model.

One possible approach for approximating the age-dependent local outbreak probability using a

branching process model is to run a large number of stochastic simulations of the model starting

from a single infected individual in a particular age group, and record the proportion of

simulations in which the pathogen does not fade out after only a small number of infections [67].

This would then need to be repeated for index cases of different ages. Here, we instead take an

analytic approach, and derive a nonlinear system of simultaneous equations that determine the

age-dependent outbreak probabilities, as described below. The local outbreak probabilities are

then obtained by solving these equations numerically, and are analogous to the probabilities that

would be derived from the simulation approach in the limit of infinitely many simulations. The

benefit of our analytic approach is that it does not require a large number of stochastic

simulations to be run.

The probability of a local outbreak not occurring (i.e. pathogen fadeout occurs), starting from a

single symptomatic (or presymptomatic, asymptomatic respectively) infectious individual in age

group G_k, was denoted by x_k (y_k, z_k). Beginning with a single symptomatic individual in G_k,

the possibilities for the next event are as follows:

1. The infected individual in G_k infects a susceptible individual in G_j, so that either A_j

increases by one (with probability ξ_j) or P_j increases by one (with probability $(1 - \xi_j)$).

This occurs with probability

286
$$\alpha_{kj} = \frac{\beta_{kj}}{\mu + \rho_k + \sum_{l=1}^{16} \beta_{kl}}.$$

287   2. The infected individual in G_k recovers, dies or is isolated before infecting anyone else, so

288   that S_k decreases to zero (and there are no infected individuals left in the population).

289   This occurs with probability

290
$$\gamma_k = \frac{\mu + \rho_k}{\mu + \rho_k + \sum_{l=1}^{16} \beta_{kl}}.$$

291   If there are no infectious hosts present in the population, then a local outbreak will not occur.

292   Therefore, assuming that chains of transmission arising from infectious individuals are

293   independent, the probability that no local outbreak occurs beginning from a single symptomatic

294   individual in G_k is

295
$$x_k = x_k \left[\sum_{j=1}^{16} \alpha_{kj} \left(\xi_j z_j + (1 - \xi_j) y_j \right) \right] + \gamma_k. \quad (2)$$

296   Similarly, beginning instead with a single presymptomatic individual in G_k, the possibilities for

297   the next event are:

298   1. The presymptomatic infected individual in G_k infects a susceptible individual in G_j, so

299   that (as before) either A_j increases by one (with probability ξ_j) or P_j increases by one

300   (with probability $(1 - \xi_j)$). This occurs with probability

301
$$\tilde{\alpha}_{kj} = \frac{\eta \beta_{kj}}{\lambda + \sigma_k + \eta \sum_{l=1}^{16} \beta_{kl}}.$$

302   2. The infected individual in G_k develops symptoms (transitions from P_k to S_k). This occurs

303   with probability

304
$$\delta_k = \frac{\lambda}{\lambda + \sigma_k + \eta \sum_{l=1}^{16} \beta_{kl}}.$$

305   3. The infected individual in G_k is isolated before infecting anyone else, so that S_k decreases

306   by one. This occurs with probability

307
$$\phi_k = \frac{\sigma_k}{\lambda + \sigma_k + \eta \sum_{l=1}^{16} \beta_{kl}}.$$

308 Therefore, the probability that no local outbreak occurs beginning from a single presymptomatic

309 individual in G_k is

310
$$y_k = y_k \left[\sum_{j=1}^{16} \tilde{\alpha}_{kj} \big(\xi_j z_j + (1 - \xi_j) y_j \big) \right] + \delta_k x_k + \phi_k. \quad (3)$$

311 Similarly, the probability z_k that a local outbreak does not occur starting from a single

312 asymptomatic individual in G_k satisfies the equation

313
$$z_k = z_k \left[\sum_{j=1}^{16} \hat{a}_{kj} \big(\xi_j z_j + (1 - \xi_j) y_j \big) \right] + \epsilon_k, \quad (4)$$

314 where

315
$$\hat{\alpha}_{kj} = \frac{\theta \beta_{kj}}{\nu + \sigma_k + \theta \sum_{l=1}^{16} \beta_{kl}} \quad \text{and} \quad \epsilon_k = \frac{\nu + \sigma_k}{\nu + \sigma_k + \theta \sum_{l=1}^{16} \beta_{kl}}.$$

The system of simultaneous equations $(2) - (4)$ can be solved numerically to obtain x_k, y_k and

z_k (here, we did this using the MATLAB nonlinear system solver *'fsolve'*). Specifically, we take

the minimal non-negative solution, as is standard when calculating extinction probabilities using

branching process models [55, 68]. Then, for each k, the probability of a local outbreak

occurring beginning from a single symptomatic (or presymptomatic, asymptomatic respectively)

individual in group G_k is given by $1 - x_k \, (1 - y_k, 1 - z_k)$.

Throughout, we consider the probability p_k of a local outbreak occurring beginning from a single

nonsymptomatic individual in group G_k arriving in the population at the beginning of their

infection:

$$p_k = \xi_k(1 - z_k) + (1 - \xi_k)(1 - y_k).$$

The average local outbreak probability, P, which is defined as the probability of a local outbreak

when the index case is chosen randomly from the population, is also considered. The value of P

is therefore a weighted average of the p_k values, where the weights correspond to the proportion

of the population represented by each group:

$$P = \frac{1}{N} \sum_{k=1}^{16} N_k p_k.$$

This reflects an assumption that the index case is more likely to come from an age group with

more individuals than an age group with fewer individuals.

All computing code used to implement the above methods was written in MATLAB version

R2019a, and is available at https://github.com/francescalovellread/age-dependent-outbreak-risks.

Fig 2. Age demographic and age-structured contact patterns for the United Kingdom. A.

United Nations age demographic data for the UK in 2020, split into five-year age groups (where the

final age group contains all ages 75+) [58]. B. Heat map of UK 'all contacts' matrix, representing the

expected daily number of unique contacts that an individual in each age group G_k has with

individuals in each other age group G_j [17]. C. The analogous figure to B, but showing only the

subset of 'all' contacts that occur in schools ('school' contacts). D. The analogous figure to C, but

347   showing only 'workplace' contacts. E. The analogous figure to C, but showing only 'home' contacts.

348   F. The analogous figure to C, but showing only 'other' contacts (i.e. all contacts outside schools,

349   workplaces or homes).

351 3. Results

352 3.1 Effect of the age of the index case on the risk of a local outbreak

We first considered the probability that a single infected individual in a particular age group G_k

initiates a local outbreak when introduced into a new host population. This quantity was

calculated for each of the three scenarios A, B and C (Fig 3).

In scenario A, the variation in the local outbreak risk for introduced cases of different ages is

driven solely by the numbers of contacts between individuals. As a result, due to their higher

numbers of daily contacts, school- and working-age individuals are more likely to trigger a local

outbreak than children under five or adults over 60, with index cases aged 15-19 posing the

highest risk (Fig 3A). These findings do not change significantly when susceptibility is allowed

to vary with age in scenario B (Fig 3B). However, in scenario C, assuming that the clinical

fraction also varies between age groups alters the age-dependent risk profile substantially. This is

because asymptomatic individuals are assumed to be less infectious than symptomatic

individuals, and therefore an index case in an age group with a high proportion of asymptomatic

infections is less likely to initiate a local outbreak. In this scenario, index cases aged 40 or over

had a disproportionately high probability of generating a local outbreak, with individuals aged

70-74 presenting the highest risk (Fig 3C). These individuals are more likely to develop

symptoms than younger individuals (Table 1), leading to a higher expected infectiousness. In

contrast, individuals under the age of 40 had a below average probability of generating a local

outbreak, with individuals aged 10-14 presenting the lowest risk. Noticeably, individuals aged 5-

19 presented relatively low risks, despite the high numbers of contacts occurring among these

age groups (Fig 2B). In this scenario, the large number of contacts was offset by the fact that

individuals in these age groups are more likely to be asymptomatic and consequently less

infectious than older individuals (Table 1). Therefore, an index case in one of these age groups is

likely to lead to fewer secondary transmissions. Furthermore, the contact patterns between

individuals in these age groups are highly assortative with respect to age (Fig 2B). Therefore, in

addition to the index case being less infectious, a high proportion of the contacts they make are

with individuals who are also likely to be less infectious, as well as being less susceptible to

infection in the first place.

We performed our subsequent analyses for each of the three scenarios A, B and C, with

qualitatively similar results. The figures shown in the main text are for scenario C, since we

deem this scenario to be the most realistic for SARS-CoV-2 transmission, but the analogous

results for scenarios A and B are presented in Supplementary Figs S1-6.

Fig 3. The probability of a local outbreak depends on the age of the index case. A. The

probability that a single infected individual in any given age group triggers a local outbreak (grey

bars) for scenario A, in which susceptibility and clinical fraction are assumed constant across all age

groups. The weighted average local outbreak probability P is shown by the black horizontal line. B.

The analogous figure to A but for scenario B, in which clinical fraction is assumed constant across

all age groups but susceptibility varies with age (Table 1). C. The analogous figure to A but for

scenario C, in which both susceptibility and clinical fraction vary with age (Table 1).

3.2 Effect of the target age group on NPI effectiveness

We next considered the effects of NPIs that reduce the number of contacts between individuals

on the probability that an introduced case will lead to a local outbreak. To approximate the

relative effects of school closures, workplace closures and population-wide social distancing

policies, we calculated the age-dependent risk profiles when each of these types of contact were

excluded from the overall contact matrix.

First, we removed all 'school' contacts from the total contact matrix (Fig 4A). For scenario C,

removing 'school' contacts led to a 4.2% reduction in the average probability of a local outbreak

(from 0.449 to 0.430). This small reduction is unsurprising for scenario C, since in that scenario

school-aged infected individuals are assumed to be more likely to be asymptomatic than other

infected individuals, and therefore their expected infectiousness is lower. However, even for

scenarios A and B, in which school-aged individuals present the greatest risk of triggering a local

outbreak, the effectiveness of removing 'school' contacts alone at reducing the local outbreak

probability was limited (reductions of 7.2% and 4.75% respectively; see Supplementary Figs

S1A, S4A). In each scenario, the reduction in risk was predominantly for school-aged index

cases, with the risk from index cases of other ages only slightly reduced. Second, we considered

the effects of removing 'work' contacts from the total contact matrix (Fig 4B). This led to a more

substantial 25.4% reduction in the average probability of a local outbreak for scenario C (with

corresponding reductions of 19.0% and 24.0% for scenarios A and B respectively; see

Supplementary Figs S1B, S4B). As well as reducing the risk of a local outbreak from an index

case of working age, removing 'work' contacts also reduced the probability of a local outbreak

occurring starting from a school-aged individual. This is because closing workplaces helps to

block chains of transmission that begin with an infected child. For example, a transmission chain

involving a child transmitting to an adult at home, followed by subsequent spread around the

adult's workplace, will be less likely to occur. Third, we investigated the effect of removing all

'other' contacts, reflecting perfect social distancing being observed outside of the home, school

or workplace (Fig 4C). This had the most significant effect of the three types of contact-reducing

intervention considered, reducing the probability of a local outbreak by 41.7% for scenario C

(and 30.7% or 33.2% for scenarios A and B, respectively).

In the three cases described above, we considered complete reductions in 'school', 'work' and

'other' contacts, respectively. In practice, such complete elimination of contacts is unlikely. We

therefore also considered partial reductions in 'school', 'work' and 'other' contacts, and

compared the resulting reductions in the local outbreak probability (Fig 4D). For any given

percentage reduction in contacts, reducing 'other' contacts always led to the largest reduction in

the local outbreak probability (see also Supplementary Figs S1D, S4D). This suggests that

reducing social contacts outside schools and workplaces can be an important component of

strategies to reduce the risk of local outbreaks of SARS-CoV-2. However, this alone is not

enough to eliminate the risk of local outbreaks entirely. For greater risk reductions using contact-

reducing NPIs, a mixed approach involving combinations of reductions in 'school', 'workplace'

and 'other' contacts is needed.

Fig 4. The effects of interventions that reduce contacts between individuals on the probability of a

local outbreak. A. The effect of removing all 'school' contacts on the probability of a local outbreak.

Pale grey bars and black dash-dotted line represent the local outbreak probabilities without any contacts

removed (as in Fig 3C). Red bars and the solid red line represent the local outbreak probabilities and

their weighted average when 'school' contacts are removed. B. The analogous figure to A, but with all

'work' contacts removed. C. The analogous figure to A, but with all 'other' contacts removed. D. Partial

reductions in 'school', 'work' and 'other' contacts, and the resulting reductions in the average local

outbreak probability (solid red, dashed blue and dotted green lines respectively).

3.3 Mixed strategies for reducing the local outbreak risk

Next, we considered the effects of combining reductions in 'school', 'work' and 'other' contacts

on the local outbreak probability (Fig 5; analogous results for scenarios A and B are shown in

Supplementary Figs S2 and S5). We allowed reductions in 'school' and 'work' contacts to vary

between 0% and 100% whilst 'other' contacts were reduced by 25%, 50% or 75% (Fig 5A,B,C,

respectively).

Since NPIs have negative economic, social and non-disease health consequences, policy makers

may choose to implement public health measures in which the risk of local outbreaks is not

eliminated completely. These results provide contact reduction targets for mixed strategies in

which the local outbreak probability is reduced to a pre-specified 'acceptable' level. For

example, to reduce the local outbreak probability to 0.25, 'other' contacts could be reduced by

25% from the baseline level, and 'school' and 'home' contacts reduced as indicated by the red

dotted contour marked '0.25' in Fig 5A. Alternatively, to achieve the same local outbreak risk,

'other' contacts can instead be reduced by 50% or by 75% with the degree of 'school' and

'work' reductions chosen according to the contours marked '0.25' in Figs 5B,C respectively.

If a policy maker wishes to eliminate the local outbreak risk entirely using contact-reducing

NPIs, for the model parameterisation considered very significant reductions in multiple types of

contacts are needed in combination. For example, even if all 'school' and 'work' contacts are

removed, 'other' contacts must be reduced by 66% for the overall average local outbreak

probability to fall below 0.01 (Fig 5D). Since such substantial reductions in multiple types of

contacts are unlikely to be possible, this suggests that contact-reducing NPIs must be combined

with other interventions, such as effective surveillance and isolation strategies, to eliminate local

outbreaks.

Fig 5. The effects of intervention strategies that combine reductions in 'school', 'work' and 'other'

contacts. A. The effect of reducing 'school' and 'work' contacts on the weighted average probability of

a local outbreak (*P*), when 'other' contacts are reduced by 25% across all age groups. Red dotted lines

indicate contours along which the local outbreak probability is constant. B. The analogous figure to A,

but with a 50% reduction in 'other' contacts. C. The analogous figure to A, but with a 75% reduction in

'other' contacts. D. The effect of reducing 'other' contacts on the average local outbreak probability

when 'school' and 'work' contacts are not reduced at all (dotted line) and when 'school' and 'work'

contacts are reduced by 100% (solid line).

3.4 Effect of surveillance on the contact-reducing NPIs required for local outbreak control

We considered whether or not low local outbreak probabilities can be achieved using limited

contact-reducing NPIs in combination with other interventions. Specifically, the effects of

surveillance and isolation of infected individuals (through e.g. contact tracing) as well as

reducing contacts in schools, workplaces and other locations, were assessed. While results are

shown for scenario C in Fig 6, analogous results for scenarios A and B are presented in

Supplementary Figs S3 and S6.

Initially, we considered the effect of increasing the rate at which symptomatic and/or

nonsymptomatic infected individuals are detected and isolated as a result of surveillance, in the

absence of contact-reducing NPIs (i.e. with no reduction in the number of contacts between

individuals compared to the baseline case in Fig 2B) (Fig 6A,B). For symptomatic hosts, this

represents an enhanced rate of isolation compared to the baseline rate of self-isolation already

present in the model. Isolation of nonsymptomatic hosts was more effective at reducing the local

outbreak probability than isolation of symptomatic hosts (Figs 6A,B), although of course this is

more challenging to achieve [55]. However, if fast isolation of nonsymptomatic hosts could be

achieved through efficient large-scale testing (potentially in combination with contact tracing

[69]), the probability of local outbreaks could be reduced substantially through this measure

alone.

We then demonstrated the effects of combining contact-reducing NPIs with enhanced isolation

of infected hosts due to infection surveillance. First, we increased the enhanced isolation rate of

symptomatic individuals to $\rho_k = 1/2$ days^{-1}. In the absence of other interventions, this reduced

the local outbreak probability by 22.0% (Figure 6C). With this level of surveillance, the local

outbreak risk could be reduced below 0.01 with a reduction in 'work' and 'other' contacts of

around 73% each, for example.

Finally, keeping the enhanced isolation rate of symptomatic individuals equal to $\rho_k =$

$1/2 \text{ days}^{-1}$, we increased the isolation rate of nonsymptomatic individuals to $\sigma_k = 1/7 \text{ days}^{-1}$.

In this case, the local outbreak probability without contact-reducing NPIs fell by 59.4%

compared to a situation without enhanced surveillance (Fig 6E), and the reductions in 'work' and

'other' contacts needed to bring the local outbreak probability below 0.01 were significantly

smaller (Figure 6F). For example, if 'work' contacts can be reduced by 50%, then 'other'

contacts only need to be reduced by 43%. This indicates that effective surveillance of both

symptomatic and nonsymptomatic individuals can substantially lower the extent of contact-

reducing NPIs that are required to achieve substantial reductions in local outbreak risks.

Fig 6. Surveillance as part of a mixed strategy to reduce the local outbreak probability. A. The

effect of increasing the isolation rate of symptomatic (red line) or nonsymptomatic infected hosts (blue

line) on the average probability of a local outbreak (P), in the absence of contact-reducing NPIs. The

isolation rates ρ_k and σ_k are varied in turn between 0 days^{-1} and 1 days^{-1}. B. The effect of

simultaneously varying the isolation rate of symptomatic and nonsymptomatic hosts on the average

probability of a local outbreak (P), again without contact-reducing NPIs. C. The age-dependent

probability of a local outbreak when the isolation rate of symptomatic individuals is $\rho_k = 1/2$ days^{-1},

without contact-reducing NPIs or surveillance of nonsymptomatic infected individuals (purple bars and

solid line). Pale grey bars and black dash-dotted line represent the original local outbreak probabilities

without any contact-reducing NPIs or enhanced surveillance (as in Fig 3C). D. The effect of reducing

'work' and 'other' contacts when the isolation rate of symptomatic infected individuals is $\rho_k =$

$1/2$ days^{-1}, as in C, without surveillance of nonsymptomatic infected individuals. E,F. The analogous

figures to C,D, with enhanced surveillance of both symptomatic and nonsymptomatic infected hosts

($\rho_k = 1/2$ days^{-1} and $\sigma_k = 1/7$ days^{-1}).

4. Discussion

During the COVID-19 pandemic, public health measures that reduce the numbers of contacts

between individuals have been implemented in countries globally. These measures include

school closures, workplace closures and population-wide social distancing policies. Contact-

reducing NPIs have been shown to be effective at reducing SARS-CoV-2 transmission, and have

also been used previously during influenza pandemics [5, 70-72]. However, long-term

implementation of these measures has negative social, psychological and economic

consequences [7, 12-16]. It is therefore important to assess the effectiveness of different contact-

reducing NPIs at lowering transmission and preventing local outbreaks, in order to design

effective targeted control strategies that avoid unnecessarily strict measures.

Here, we constructed a branching process model to estimate the risk of local outbreaks under

different contact-reducing NPIs and different levels of surveillance for symptomatic and

nonsymptomatic infected individuals. Unlike previous approaches for estimating outbreak risks

using branching processes [51-56], we considered the effects of age-related heterogeneities

affecting transmission for infected individuals of a wide range of ages, including age-dependent

variations in social mixing patterns, susceptibility to infection and clinical fraction. Using SARS-

CoV-2 as a case study, we demonstrated that the risk that an introduced case initiates a local

outbreak depends on these age-related factors and on the age of the introduced case (Fig 3), as

well as the age-structure of the local population.

We used our model to assess the effects of reducing the numbers of contacts that occur in school,

in the workplace and elsewhere. Of the three contact-reducing NPIs considered, removing

'school' contacts had the smallest effect on the probability of observing a local outbreak, even

when age-dependent variations in susceptibility and clinical fraction were ignored (Figs 4A,D,

Supplementary Figs S1A,D, and S4,A,D). This can be attributed to the fact that school closures

predominantly reduce contacts between individuals aged 5-19, who only account for

approximately 23% of the total population [58]. Additionally, these individuals tend to have

large numbers of contacts outside of the school environment (Figs 2E,F). Therefore, compared to

other measures, interrupting within-school transmission may have only a limited effect on

transmission in the wider population, particularly when school-aged individuals are less

susceptible to infection and more likely to experience subclinical courses of infection. In

contrast, reducing contacts that occur outside schools or workplaces was the most effective

intervention, significantly lowering the local outbreak risk across all age groups, and for those

aged over 60 in particular (Figs 4C,D, and Supplementary Figs S1C,D and S4C,D). This could

explain the success of social distancing strategies worldwide for reducing observed COVID-19

cases and deaths.

Mixed strategies combining reductions in 'school', 'work' and 'other' contacts led to greater

reductions in the local outbreak probability than individual interventions (Figs 5A-C), but very

large reductions in all three types of contact were required to eliminate the risk of local outbreaks

entirely (Fig 5D). However, implementing effective surveillance to identify infected hosts

(followed by isolation) led to substantial reductions in the risk of local outbreaks even in the

absence of other control measures (Figs 6A,B). In the scenarios considered here, with an

efficient surveillance strategy in place, significantly smaller reductions in 'work' and 'other'

contacts were needed to render the local outbreak probability negligible, even when 'school'

contacts were not reduced at all (Figs 6C-F). This supports the use of surveillance that targets

both symptomatic and nonsymptomatic individuals, such as contact tracing and isolation

strategies or population-wide diagnostic testing, to help prevent local outbreaks [55].

Although here we used SARS-CoV-2 as a case study, our model provides a framework for

estimating the risk of local outbreaks in age-structured populations that can be adapted for other

pathogens, provided sufficient data are available to parametrise the model appropriately. The

effects of age-structure on local outbreak risks may vary for pathogens with different

epidemiological characteristics. For influenza-A viruses, for example, susceptibility to infection

tends to decrease with age, whilst the risk of an infection leading to severe symptoms is greater

both for the elderly and for the very young [31, 73, 74]. This is in contrast to SARS-CoV-2, for

which children are more likely to experience subclinical courses of infection. In this study, we

used age demographic and contact data for the UK, but equivalent data for other countries are

available and can easily be substituted into our model to estimate outbreak risks elsewhere [17,

58].

One caveat of the results for SARS-CoV-2 presented here is that, although the epidemiological

parameters of our model were chosen to be consistent with reported literature estimates, there is

considerable variation between studies. In particular, the precise age-dependent variation in

susceptibility and clinical fraction remains unclear, and the relative infectiousness of

asymptomatic, presymptomatic and symptomatic hosts has not been determined exactly.

Furthermore, the inherent transmissibility of SARS-CoV-2 is now higher than in the initial stage

of the pandemic, due to the appearance of novel variants. To explore ongoing local outbreak

risks due to SARS-CoV-2, it would be necessary to update the model to reflect the increased

transmissibility of the Delta variant [75]. Due to the uncertainty in model parameter values, we

conducted sensitivity analyses to explore the effects of varying the parameters of the model on

our results (Supplementary Figs S1-12). In each case that we considered, our main conclusions

were unchanged: the probability that an introduced case initiates a local outbreak depends on

age-dependent factors affecting pathogen transmission and control, with widespread

interventions and combinations of NPIs reducing the risk of local outbreaks most significantly.

An important limitation of our approach to modelling contact-reducing NPIs is that we made a

standard assumption in our main analyses that 'school', 'work' and 'other' contacts are

independent [27, 31, 47]. In other words, reducing the numbers of contacts in one location did

not affect the numbers of contacts occurring in another. In reality, this is unlikely to be the case.

For example, closing schools also affects workplace contacts, as adults may then work from

home in order to fulfil childcare requirements. Additionally, the contact data that we used

represent the number of unique contacts per day and do not reflect the numbers of repeated

contacts with the same person, which affect the risk of transmission between individuals. These

assumptions could in principle be removed, if relevant data become available – for example, data

describing the effects of school closures on numbers of contacts in other locations. To

demonstrate how changes in multiple types of contact due to NPIs could be implemented in the

model, we conducted a supplementary analysis in which we investigated the effects of removing

all 'school' contacts and allowing for concurrent changes in 'work', 'home' and 'other' contacts

(Supplementary Figs S13-14). These results support our conclusion that school closures are

unlikely to have a substantial impact on SARS-CoV-2 transmission when applied as the sole

NPI. The benefit of school closures could even potentially be outweighed by the possible

secondary effects on other types of contacts. An improved understanding of how NPIs affect

different types of contact is important for more accurate assessments of interventions in future.

Despite the simplifications made, our model provides a useful framework for estimating the risk

of local outbreaks and the effects of NPIs. Different measures can be considered in combination

in the model to develop strategies for lowering local outbreak risks. Our results emphasise the

importance of quantifying age-dependent factors that affect transmission dynamics, such as

susceptibility to infection and the proportion of hosts who develop clinical symptoms, for

individuals of different ages. As we have shown, it is crucial to take age-dependent factors into

account when assessing local outbreak risks and designing public health measures.

Declaration of interests

We declare no competing interests.

Authors' contributions

Conceptualisation: All authors. Methodology: FALR, RNT. Investigation: FALR, SS.

Writing - original draft: FALR, RNT. Writing - review and editing: All authors. Supervision:

RNT.

Data availability

All computer code used in this paper was written in MATLAB version R2019a, and is available

at the following GitHub repository: https://github.com/francescalovellread/age-dependent-

outbreak-risks.

Funding

FALR acknowledges funding from the Biotechnology and Biological Sciences Research Council

(BBSRC), grant number BB/M011224/1. SS was funded by a BBSRC Research Experience

Placement and a Rokos award for undergraduate research from Pembroke College Oxford. RNT

was supported by UK Research and Innovation (UKRI) via grant EP/V053507/1. The funders

had no role in the study design; the collection, analysis and interpretation of data; the writing of

the report; or the decision to submit the article for publication.

References

[1] Perra, N. 2021 Non-pharmaceutical interventions during the COVID-19 pandemic: A review.

Phys. Rep. **913**, 1-52. (doi:10.1016/j.physrep.2021.02.001).

[2] Regmi, K. & Lwin, C.M. 2020 Impact of non-pharmaceutical interventions for reducing

transmission of COVID-19: a systematic review and meta-analysis protocol. *BMJ Open* **10**,

e041383. (doi:10.1136/bmjopen-2020-041383).

[3] Imai, N., Gaythorpe, K.A.M., Abbott, S., Bhatia, S., van Elsland, S., Prem, K., Liu, Y. &

Ferguson, N.M. 2020 Adoption and impact of non-pharmaceutical interventions for COVID-19.

Wellcome Open Res **5**, 59-59. (doi:10.12688/wellcomeopenres.15808.1).

[4] Thompson, R.N., Hollingsworth, T.D., Isham, V., Arribas-Bel, D., Ashby, B., Britton, T.,

Challenor, P., Chappell, L.H.K., Clapham, H., Cunniffe, N.J., et al. 2020 Key questions for

modelling COVID-19 exit strategies. *Proc. R. Soc. Lond. B. Biol. Sci.* **287**, 20201405.

(doi:10.1098/rspb.2020.1405).

[5] Hatchett, R.J., Mecher, C.E. & Lipsitch, M. 2007 Public health interventions and epidemic

intensity during the 1918 influenza pandemic. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 7582.

(doi:10.1073/pnas.0610941104).

[6] Kirsch, T.D., Moseson, H., Massaquoi, M., Nyenswah, T.G., Goodermote, R., Rodriguez-

Barraquer, I., Lessler, J., Cumings, D.A.T. & Peters, D.H. 2017 Impact of interventions and the

incidence of ebola virus disease in Liberia—implications for future epidemics. *Health Policy*

Plan. **32**, 205-214. (doi:10.1093/heapol/czw113).

[7] Berkman, B.E. 2008 Mitigating Pandemic Influenza: The Ethics of Implementing a School

Closure Policy. *J. Public Health Manag. Pract.* **14**, 372-378.

(doi:10.1097/01.PHH.0000324566.72533.0b).

[8] Moore, S., Hill, E.M., Tildesley, M.J., Dyson, L. & Keeling, M.J. 2021 Vaccination and non-

pharmaceutical interventions for COVID-19: a mathematical modelling study. *Lancet Infect Dis*

21, 793-802. (doi:10.1016/S1473-3099(21)00143-2).

[9] Spinelli, M.A., Glidden, D.V., Gennatas, E.D., Bielecki, M., Beyrer, C., Rutherford, G.,

Chambers, H., Goosby, E. & Gandhi, M. 2021 Importance of non-pharmaceutical interventions

in lowering the viral inoculum to reduce susceptibility to infection by SARS-CoV-2 and

potentially disease severity. *Lancet Infect Dis* **21**, E296-E301. (doi:10.1016/S1473-

3099(20)30982-8).

[10] Thompson, R.N., Hill, E.M. & Gog, J.R. 2021 SARS-CoV-2 incidence and vaccine escape.

Lancet Infect Dis **21**, 913-914. (doi:10.1016/S1473-3099(21)00202-4).

[11] Sachak-Patwa, R., Byrne, H.M., Dyson, L. & Thompson, R.N. 2021 The risk of SARS-

CoV-2 outbreaks in low prevalence settings following the removal of travel restrictions.

Communications Medicine **1**, 39. (doi:10.1038/s43856-021-00038-8).

[12] Donohue, J.M. & Miller, E. 2020 COVID-19 and School Closures. *JAMA* **324**, 845-847.

(doi:10.1001/jama.2020.13092).

[13] Silverman, M., Sibbald, R. & Stranges, S. 2020 Ethics of COVID-19-related school

closures. *Can. J. Public Health* **111**, 462-465. (doi:10.17269/s41997-020-00396-1).

[14] Sadique, M.Z., Adams, E.J. & Edmunds, W.J. 2008 Estimating the costs of school closure

for mitigating an influenza pandemic. *BMC Public Health* **8**, 135. (doi:10.1186/1471-2458-8-

135).

[15] Brooks, S.K., Webster, R.K., Smith, L.E., Woodland, L., Wessely, S., Greenberg, N. &

Rubin, G.J. 2020 The psychological impact of quarantine and how to reduce it: rapid review of

the evidence. *The Lancet* **395**, 912-920. (doi:10.1016/S0140-6736(20)30460-8).

[16] Loades, M.E., Chatburn, E., Higson-Sweeney, N., Reynolds, S., Shafran, R., Brigden, A.,

Linney, C., McManus, M.N., Borwick, C. & Crawley, E. 2020 Rapid Systematic Review: The

Impact of Social Isolation and Loneliness on the Mental Health of Children and Adolescents in

the Context of COVID-19. *J. Am. Acad. Child Adolesc. Psychiatry* **59**, 1218-1239.e1213.

(doi:10.1016/j.jaac.2020.05.009).

[17] Prem, K., Cook, A.R. & Jit, M. 2017 Projecting social contact matrices in 152 countries

using contact surveys and demographic data. *PLoS Comp. Biol.* **13**, e1005697.

(doi:10.1371/journal.pcbi.1005697).

[18] Mossong, J., Hens, N., Jit, M., Beutels, P., Auranen, K., Mikolajczyk, R., Massari, M.,

Salmaso, S., Tomba, G.S., Wallinga, J., et al. 2008 Social Contacts and Mixing Patterns Relevant

to the Spread of Infectious Diseases. *PLoS Med.* **5**, e74. (doi:10.1371/journal.pmed.0050074).

[19] Danon, L., Read, J.M., House, T.A., Vernon, M.C. & Keeling, M.J. 2013 Social encounter

networks: characterizing Great Britain. *Proc. R. Soc. Lond. B. Biol. Sci.* **280**, 20131037.

(doi:10.1098/rspb.2013.1037).

[20] Leung, K., Jit, M., Lau, E.H.Y. & Wu, J.T. 2017 Social contact patterns relevant to the

spread of respiratory infectious diseases in Hong Kong. *Sci. Rep.* **7**. (doi:10.1038/s41598-017-

08241-1).

[21] Béraud, G., Kazmercziak, S., Beutels, P., Levy-Bruhl, D., Lenne, X., Mielcarek, N.,

Yazdanpanah, Y., Boëlle, P.-Y., Hens, N. & Dervaux, B. 2015 The French Connection: The First

Large Population-Based Contact Survey in France Relevant for the Spread of Infectious

Diseases. *PLoS One* **10**, e0133203. (doi:10.1371/journal.pone.0133203).

[22] Ibuka, Y., Ohkusa, Y., Sugawara, T., Chapman, G.B., Yamin, D., Atkins, K.E., Taniguchi,

K., Okabe, N. & Galvani, A.P. 2016 Social contacts, vaccination decisions and influenza in

Japan. *J. Epidemiol. Community Health* **70**, 162-167. (doi:10.1136/jech-2015-205777).

[23] Read, J.M., Lessler, J., Riley, S., Wang, S., Tan, L.J., Kwok, K.O., Guan, Y., Jiang, C.Q. &

Cummings, D.A.T. 2014 Social mixing patterns in rural and urban areas of southern China. *Proc.*

R. Soc. Lond. B. Biol. Sci. **281**, 20140268. (doi:10.1098/rspb.2014.0268).

[24] Read, J.M., Edmunds, W.J., Riley, S., Lessler, J. & Cummings, D.A.T. 2012 Close

encounters of the infectious kind: methods to measure social mixing behaviour. *Epidemiol.*

Infect. **140**, 2117-2130. (doi:10.1017/S0950268812000842).

[25] Wallinga, J., Teunis, P. & Kretzschmar, M. 2006 Using Data on Social Contacts to Estimate

Age-specific Transmission Parameters for Respiratory-spread Infectious Agents. *Am. J.*

Epidemiol. **164**, 936-944. (doi:10.1093/aje/kwj317).

[26] Kucharski, A.J., Kwok, K.O., Wei, V.W.I., Cowling, B.J., Read, J.M., Lessler, J.,

Cummings, D.A. & Riley, S. 2014 The Contribution of Social Behaviour to the Transmission of

Influenza A in a Human Population. *PLoS Pathog.* **10**, e1004206.

(doi:10.1371/journal.ppat.1004206).

[27] Zhang, J., Litvinova, M., Liang, Y., Wang, Y., Wang, W., Zhao, S., Wu, Q., Merler, S.,

Viboud, C., Vespignani, A., et al. 2020 Changes in contact patterns shape the dynamics of the

COVID-19 outbreak in China. *Science* **368**, 1481-1486. (doi:10.1126/science.abb8001).

[28] Munro, A.P. & Faust, S.N. 2020 COVID-19 in children: current evidence and key

questions. *Curr. Opin. Infect. Dis.* **33**, 540-547. (doi:10.1097/QCO.0000000000000690).

[29] Viner, R.M., Mytton, O.T., Bonell, C., Melendez-Torres, G.J., Ward, J., Hudson, L.,

Waddington, C., Thomas, J., Russell, S., van der Klis, F., et al. 2021 Susceptibility to SARS-

CoV-2 Infection Among Children and Adolescents Compared With Adults: A Systematic

Review and Meta-analysis. *JAMA Pediatrics* **175**, 143-156.

(doi:10.1001/jamapediatrics.2020.4573).

[30] Goldstein, E., Lipsitch, M. & Cevik, M. 2020 On the effect of age on the transmission of

SARS-CoV-2 in households, schools and the community. *J. Infect. Dis.* **223**, 362-369.

(doi:10.1093/infdis/jiaa691).

[31] Davies, N.G., Klepac, P., Liu, Y., Prem, K., Jit, M., Pearson, C.A.B., Quilty, B.J.,

Kucharski, A.J., Gibbs, H., Clifford, S., et al. 2020 Age-dependent effects in the transmission

and control of COVID-19 epidemics. *Nat. Med.* **26**, 1205-1211. (doi:10.1038/s41591-020-0962-

9).

[32] Ludvigsson, J.F. 2020 Systematic review of COVID-19 in children shows milder cases and

a better prognosis than adults. *Acta paediatrica (Oslo, Norway : 1992)* **109**, 1088-1095.

(doi:10.1111/apa.15270).

[33] Dong, Y., Mo, X., Hu, Y., Qi, X., Jiang, F., Jiang, Z. & Tong, S. 2020 Epidemiology of

COVID-19 Among Children in China. *Pediatrics* **145**, e20200702. (doi:10.1542/peds.2020-

0702).

[34] Siebach, M.K., Piedimonte, G. & Ley, S.H. 2021 COVID-19 in childhood: Transmission,

clinical presentation, complications and risk factors. *Pediatr. Pulmonol.* **56**, 1342-1356.

(doi:10.1002/ppul.25344).

[35] Mehta, N.S., Mytton, O.T., Mullins, E.W.S., Fowler, T.A., Falconer, C.L., Murphy, O.B.,

Langenberg, C., Jayatunga, W.J.P., Eddy, D.H. & Nguyen-Van-Tam, J.S. 2020 SARS-CoV-2

(COVID-19): What Do We Know About Children? A Systematic Review. *Clin. Infect. Dis.* **71**,

2469-2479. (doi:10.1093/cid/ciaa556).

[36] Han, M.S., Choi, E.H., Chang, S.H., Jin, B.-L., Lee, E.J., Kim, B.N., Kim, M.K., Doo, K.,

Seo, J.-H., Kim, Y.-J., et al. 2021 Clinical Characteristics and Viral RNA Detection in Children

With Coronavirus Disease 2019 in the Republic of Korea. *JAMA Pediatrics* **175**, 73-80.

(doi:10.1001/jamapediatrics.2020.3988).

[37] Pollock, A.M. & Lancaster, J. 2020 Asymptomatic transmission of covid-19. *BMJ* **371**,

m4851. (doi:10.1136/bmj.m4851).

[38] Buitrago-Garcia, D., Egli-Gany, D., Counotte, M.J., Hossmann, S., Imeri, H., Ipekci, A.M.,

Salanti, G. & Low, N. 2020 Occurrence and transmission potential of asymptomatic and

presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis. *PLoS*

Med. **17**, e1003346. (doi:10.1371/journal.pmed.1003346).

[39] Byambasuren, O., Cardona, M., Bell, K., Clark, J., McLaws, M.-L. & Glasziou, P. 2020

Estimating the extent of asymptomatic COVID-19 and its potential for community transmission:

Systematic review and meta-analysis. *JAMMI* **5**, 223-234. (doi:10.3138/jammi-2020-0030).

[40] Koh, W.C., Naing, L., Chaw, L., Rosledzana, M.A., Alikhan, M.F., Jamaludin, S.A., Amin,

F., Omar, A., Shazli, A., Griffith, M., et al. 2020 What do we know about SARS-CoV-2

transmission? A systematic review and meta-analysis of the secondary attack rate and associated

risk factors. *PLoS One* **15**, e0240205. (doi:10.1371/journal.pone.0240205).

[41] Madewell, Z.J., Yang, Y., Longini, I.M., Jr., Halloran, M.E. & Dean, N.E. 2020 Household

Transmission of SARS-CoV-2: A Systematic Review and Meta-analysis. *JAMA Network Open*

3, e2031756-e2031756. (doi:10.1001/jamanetworkopen.2020.31756).

[42] Qiu, X., Nergiz, A.I., Maraolo, A.E., Bogoch, I.I., Low, N. & Cevik, M. 2021 Defining the

role of asymptomatic and pre-symptomatic SARS-CoV-2 transmission - a living systematic

review. *Clin. Microbiol. Infect.* **27**, 511-519. (doi:10.1016/j.cmi.2021.01.011).

[43] Pijls, B.G., Jolani, S., Atherley, A., Derckx, R.T., Dijkstra, J.I.R., Franssen, G.H.L.,

Hendriks, S., Richters, A., Venemans-Jellema, A., Zalpuri, S., et al. 2021 Demographic risk

factors for COVID-19 infection, severity, ICU admission and death: a meta-analysis of 59

studies. *BMJ Open* **11**, e044640. (doi:10.1136/bmjopen-2020-044640).

[44] Verity, R., Okell, L.C., Dorigatti, I., Winskill, P., Whittaker, C., Imai, N., Cuomo-

Dannenburg, G., Thompson, H., Walker, P.G.T., Fu, H., et al. 2020 Estimates of the severity of

coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis* **20**, 669-677.

(doi:10.1016/S1473-3099(20)30243-7).

[45] Chen, T., Dai, Z., Mo, P., Li, X., Ma, Z., Song, S., Chen, X., Luo, M., Liang, K., Gao, S., et

al. 2020 Clinical Characteristics and Outcomes of Older Patients with Coronavirus Disease 2019

(COVID-19) in Wuhan, China: A Single-Centered, Retrospective Study. *J. Gerontol. A Biolo.*

Sci. Med. Sci. **75**, 1788-1795. (doi:10.1093/gerona/glaa089).

[46] Prem, K., Liu, Y., Russell, T.W., Kucharski, A.J., Eggo, R.M., Davies, N., Flasche, S.,

Clifford, S., Pearson, C.A.B., Munday, J.D., et al. 2020 The effect of control strategies to reduce

social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a modelling study.

Lancet Public Health **5**, e261-e270. (doi:10.1016/S2468-2667(20)30073-6).

[47] Davies, N.G., Kucharski, A.J., Eggo, R.M., Gimma, A., Edmunds, W.J. & Centre for the

Mathematical Modelling of Infectious Diseases COVID-19 working group. 2020 Effects of non-

pharmaceutical interventions on COVID-19 cases, deaths, and demand for hospital services in

the UK: a modelling study. *Lancet Public Health* **5**, e375-e385. (doi:10.1016/S2468-

2667(20)30133-X).

[48] Liu, T., Gong, D., Xiao, J., Hu, J., He, G., Rong, Z. & Ma, W. 2020 Cluster infections play

important roles in the rapid evolution of COVID-19 transmission: A systematic review. *Int. J.*

Infect. Dis. **99**, 374-380. (doi:10.1016/j.ijid.2020.07.073).

[49] Leclerc, Q., Fuller, N., Knight, L., CMMID COVID-19 Working Group, Funk, S. & Knight,

G. 2020 What settings have been linked to SARS-CoV-2 transmission clusters? [version 2; peer

review: 2 approved]. *Wellcome Open Res* **5**. (doi:10.12688/wellcomeopenres.15889.2).

[50] Rossman, H., Keshet, A., Shilo, S., Gavrieli, A., Bauman, T., Cohen, O., Shelly, E., Balicer,

R., Geiger, B., Dor, Y., et al. 2020 A framework for identifying regional outbreak and spread of

COVID-19 from one-minute population-wide surveys. *Nat. Med.* **26**, 634-638.

(doi:10.1038/s41591-020-0857-9).

[51] Althaus, C.L., Low, N., Musa, E.O., Shuaib, F. & Gsteiger, S. 2015 Ebola virus disease

outbreak in Nigeria: Transmission dynamics and rapid control. *Epidemics* **11**, 80-84.

(doi:10.1016/j.epidem.2015.03.001).

[52] Thompson, R.N., Jalava, K. & Obolski, U. 2019 Sustained transmission of Ebola in new

locations: more likely than previously thought. *Lancet Infect Dis* **19**, 1058-1059.

(doi:10.1016/S1473-3099(19)30483-9).

[53] Thompson, R.N. 2020 Novel Coronavirus Outbreak in Wuhan, China, 2020: Intense

Surveillance Is Vital for Preventing Sustained Transmission in New Locations. *J. Clin. Med.* **9**,

498. (doi:10.3390/jcm9020498).

[54] Hellewell, J., Abbott, S., Gimma, A., Bosse, N.I., Jarvis, C.I., Russell, T.W., Munday, J.D.,

Kucharski, A.J., Edmunds, W.J., Sun, F., et al. 2020 Feasibility of controlling COVID-19

outbreaks by isolation of cases and contacts. *Lancet Glob. Health* **8**, e488-e496.

(doi:10.1016/S2214-109X(20)30074-7).

[55] Lovell-Read, F.A., Funk, S., Obolski, U., Donnelly, C.A. & Thompson, R.N. 2021

Interventions targeting nonsymptomatic cases can be important to prevent local outbreaks:

SARS-CoV-2 as a case-study. *J. R. Soc. Interface* **18**, 20201014. (doi:10.1098/rsif.2020.1014).

[56] Nishiura, H., Cook, A.R. & Cowling, B.J. 2011 Assortativity and the Probability of

Epidemic Extinction: A Case Study of Pandemic Influenza A (H1N1-2009). *Interdiscip.*

Perspect. Infect. Dis. **2011**, 194507. (doi:10.1155/2011/194507).

[57] Ferretti, L., Wymant, C., Kendall, M., Zhao, L., Nurtay, A., Abeler-Dorner, L., Parker, M.,

Bonsall, D. & Fraser, C. 2020 Quantifying SARS-CoV-2 transmission suggests epidemic control

with digital contact tracing. *Science* **368**, eabb6936. (doi:10.1126/science.abb6936).

[58] United Nations, Department of Economic and Social Affairs, Population Division. 2019

World Population Prospects 2019, Online Edition, Rev. 1. *Available at:*

https://population.un.org/wpp/Download/Standard/Population/.

[59] Wei, W.E., Li, Z., Chiew, C.J., Yong, S.E., Toh, M.P. & Lee, V.J. 2020 Presymptomatic

Transmission of SARS-CoV-2 — Singapore, January 23–March 16, 2020. *Morb. Mortal. Weekly*

Rep. **69**, 411-415. (doi:10.15585/mmwr.mm6914e1external icon).

[60] Arons, M.M., Hatfield, K.M., Reddy, S.C., Kimball, A., James, A., Jacobs, J.R., Taylor, J.,

Spicer, K., Bardossy, A.C., Oakley, L.P., et al. 2020 Presymptomatic SARS-CoV-2 Infections

and Transmission in a Skilled Nursing Facility. *N. Engl. J. Med.* **382**, 2081-2090.

(doi:10.1056/NEJMoa2008457).

[61] Bullard, J., Dust, K., Funk, D., Strong, J.E., Alexander, D., Garnett, L., Boodman, C., Bello,

A., Hedley, A., Schiffman, Z., et al. 2020 Predicting Infectious Severe Acute Respiratory

Syndrome Coronavirus 2 From Diagnostic Samples. *Clin. Infect. Dis.* **71**, 2663-2666.

(doi:10.1093/cid/ciaa638).

[62] Wolfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Muller, M.A.,

Niemeyer, D., Jones, T.C., Vollmar, P., Rothe, C., et al. 2020 Virological assessment of

hospitalized patients with COVID-2019. *Nature* **581**, 465-469. (doi:10.1038/s41586-020-2196-

x).

[63] Lai, C.-C., Shih, T.-P., Ko, W.-C., Tang, H.-J. & Hsueh, P.-R. 2020 Severe acute respiratory

syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The

epidemic and the challenges. *Int. J. Antimicrob. Agents* **55**, 105924.

(doi:10.1016/j.ijantimicag.2020.105924).

[64] Liu, Y., Gayle, A.A., Wilder-Smith, A. & Rocklov, J. 2020 The reproductive number of

COVID-19 is higher compared to SARS coronavirus. *J. Travel Med.* **27**, taaa021.

(doi:10.1093/jtm/taaa021).

[65] Zhai, P., Ding, Y., Wu, X., Long, J., Zhong, Y. & Li, Y. 2020 The epidemiology, diagnosis

and treatment of COVID-19. *Int. J. Antimicrob. Agents* **55**, 105955-105955.

(doi:10.1016/j.ijantimicag.2020.105955).

[66] Zhao, S., Lin, Q., Ran, J., Musa, S.S., Yang, G., Wang, W., Lou, Y., Gao, D., Yang, L., He,

D., et al. 2020 Preliminary estimation of the basic reproduction number of novel coronavirus

(2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the

outbreak. *Int. J. Infect. Dis.* **92**, 214-217. (doi:10.1016/j.ijid.2020.01.050).

[67] Thompson, R.N., Gilligan, C.A. & Cunniffe, N.J. 2020 Will an outbreak exceed available

resources for control? Estimating the risk from invading pathogens using practical definitions of

a severe epidemic. *J. R. Soc. Interface* **17**, 20200690. (doi:10.1098/rsif.2020.0690).

[68] Norris, J.R. 1997 *Markov Chains (Cambridge Series in Statistical and Probabilistic*

Mathematics). Cambridge, Cambridge University Press.

[69] Hart, W.S., Maini, P.K. & Thompson, R.N. 2021 High infectiousness immediately before

COVID-19 symptom onset highlights the importance of continued contact tracing. *Elife* **10**.

(doi:10.7554/eLife.65534).

[70] Huang, Q.S., Wood, T., Jelley, L., Jennings, T., Jefferies, S., Daniells, K., Nesdale, A.,

Dowell, T., Turner, N., Campbell-Stokes, P., et al. 2021 Impact of the COVID-19

nonpharmaceutical interventions on influenza and other respiratory viral infections in New

Zealand. *Nat. Commun.* **12**, 1001. (doi:10.1038/s41467-021-21157-9).

[71] Flaxman, S., Mishra, S., Gandy, A., Unwin, H.J.T., Mellan, T.A., Coupland, H., Whittaker,

C., Zhu, H., Berah, T., Eaton, J.W., et al. 2020 Estimating the effects of non-pharmaceutical

interventions on COVID-19 in Europe. *Nature* **584**, 257-261. (doi:10.1038/s41586-020-2405-7).

[72] Borse, R.H., Behravesh, C.B., Dumanovsky, T., Zucker, J.R., Swerdlow, D., Edelson, P.,

Choe-Castillo, J. & Meltzer, M.I. 2011 Closing Schools in Response to the 2009 Pandemic

Influenza A H1N1 Virus in New York City: Economic Impact on Households. *Clin. Infect. Dis.*

52, S168-S172. (doi:10.1093/cid/ciq033).

[73] Greer, A.L., Tuite, A. & Fisman, D.N. 2010 Age, influenza pandemics and disease

dynamics. *Epidemiol. Infect.* **138**, 1542-1549. (doi:10.1017/S0950268810000579).

[74] Clohisey, S. & Baillie, J.K. 2019 Host susceptibility to severe influenza A virus infection.

Crit. Care **23**, 303. (doi:10.1186/s13054-019-2566-7).

[75] Campbell, F., Archer, B., Laurenson-Schafer, H., Jinnai, Y., Konings, F., Batra, N., Pavlin,

B., Vandemaele, K., Van Kerkhove, M.D., Jombart, T., et al. 2021 Increased transmissibility and

global spread of SARS-CoV-2 variants of concern as at June 2021. *Eurosurveillance* **26**,

2100509. (doi:10.2807/1560-7917.ES.2021.26.24.2100509).