Psychosocial family-level mediators in the intergenerational transmission of trauma: Protocol for a systematic review and meta-analysis

Author names and affiliations
Emma J. Mew[1]
Kate Nyhan[2,3]
Jessica L. Bonumwezi[4]
Vanessa Blas[5]
Hannah Gorman[6]
Rachel Hennein[7,8]
Kevin Quach[5]
Nicola L. Hawley[1]
Sarah R. Lowe[6]

1. Department of Chronic Disease Epidemiology, Yale School of Public Health, Yale University, New Haven, Connecticut, USA; 2. Harvey Cushing / John Hay Whitney Medical Library, Yale University, New Haven, Connecticut, USA. 3. Department of Environmental Health Sciences, Yale School of Public Health, Yale University, New Haven, Connecticut, USA; 4. Department of Psychology, Montclair State University, Montclair, New Jersey, USA; 5. Yale College, Yale University, New Haven, Connecticut, USA; 6. Department of Social and Behavioral Sciences, Yale School of Public Health, Yale University, New Haven, Connecticut, USA; 7. Department of Epidemiology of Microbial Diseases, Yale School of Public Health, Yale University, New Haven, Connecticut, USA; 8. Yale School of Medicine, Yale University, New Haven, Connecticut, USA.

Corresponding author
Emma J. Mew
Email: emma.mew@yale.edu

Title word count: 136 / 250 characters

Abstract word count: 224 / 300 words

Total manuscript word count: 3236 words

Keywords: intergenerational trauma; psychosocial mechanisms; family-level; causal diagram; meta-analysis

ABSTRACT

Introduction: Family-level psychosocial factors appear to play a critical role in mediating the intergenerational transmission of trauma; however, no review article has quantitatively synthesized causal mechanisms across a diversity of trauma types. This study aims to systematically consolidate the epidemiological research on family-level psychosocial mediators and moderators to ultimately produce causal diagram(s) of the intergenerational transmission of trauma.

Methods: We will identify epidemiological peer-reviewed publications, dissertations, and conference abstracts that measure the impact of at least one psychosocial family-level factor mediating or moderating the relationship between parental trauma exposure and a child mental health outcome. English, French, Kinyarwanda, and Spanish articles will be eligible. We searched MEDLINE, PsycINFO, PTSDpubs, Scopus and ProQuest Dissertations and Theses and will conduct forward citation chaining of included documents. Two reviewers will perform screening independently. We will extract reported mediators, moderators, and relevant study characteristics for included studies. Findings will be presented using narrative syntheses, descriptive analyses, mediation meta-analyses, moderating meta-analyses, and causal diagram(s), where possible. We will perform a risk of bias assessment and will assess for publication bias.

Discussion: The development of evidence-based causal diagram(s) would provide more detailed understanding of the paths by which unresolved trauma can be transmitted intergenerationally at the family-level. This review could provide evidence to better support interventions that interrupt the cycle of intergenerational trauma.

Systematic review registration: PROSPERO registration ID #CRD42021251053.

INTRODUCTION

Intergenerational trauma is the process by which the psychological impact of a traumatic event is transmitted from one generation to the next.(1-4) This results in subsequent generations experiencing the psychological effects of a traumatic event without exposure to it, such as increased risk of developing trauma- and stressor-related disorders.(1-4) Epidemiological studies provide some evidence for intergenerational trauma. Many studies to date have focused on the descendants of Holocaust survivors,(5) whereas smaller bodies of literature including other populations and traumatic events.(1-4, 6, 7), and on transmission of trauma-related symptoms from the first (G1) to the second (G2) generation.

Researchers have evaluated intergenerational trauma through two main paths: (1) the path from parental trauma exposure to child psychopathology; and (2) the path from parental trauma-related symptoms to child psychopathology. One recent meta-analysis showed that being a child of a Holocaust survivor was modestly associated with increased PTSD symptoms (8) and another meta-analysis of studies across any population or trauma-type identified a significant association between parental PTSD symptoms and child PTSD symptoms.(9) However, evidence is mixed, as additional reviews have been less conclusive (10, 11). Researchers also speculate transmission from G1 to the third (G3) generation which is supported by animal models.(12, 13) However, few epidemiological studies have evaluated transmission from G1 to G3 and the few that have obtained mixed results.(14)

In addition to empirically demonstrating intergenerational trauma, it would be helpful to identify the causal processes underpinning this phenomenon as such findings could inform interventions aimed to prevent or mitigate it. Mediation studies examine the extent to which an intermediate variable (mediator) explains the effect of exposure on an outcome in an attempt to tease apart causal mechanisms.(15) Mediation studies quantify both direct and indirect effects, meaning the

effect of exposure on the outcome and the effect of the exposure on the outcome through a

mediating variable.(15) Epidemiological studies can also identify potential moderators, or

variables that influence the degree of association between an exposure and an outcome. These

relationships are often demonstrated visually through causal diagrams, most formally directed

acyclic graphs (DAGs). DAGs can powerfully deduce causal relationships as they incorporate

statistical approaches into a visual diagram.(16) Figure 1 demonstrates simplified DAGs for the

intergenerational transmission of trauma including mediation (Figure 1A) and moderation (Figure

1B) paths.

Figure 1. Simplified directed acyclic graphs for the intergenerational transmission of trauma from

first (G1) to second (G2) generation through parental trauma exposure → child mental health

outcome. (A) Single mediator model showing indirect, direct, and total effects. (B) Simple

moderation model.

Evidence suggests several complex, multifaceted causal processes underpin the

intergenerational transmission of trauma. Although the most notable evidence is for genetic and

epigenetic mechanisms, these do not fully account for transmission of trauma, suggesting the

involvement of other psychosocial and interpersonal factors.(17, 18). Recent studies have

provided some evidence that psychosocial factors that operate within the household play some

role in mediating the intergenerational transmission of trauma,(1, 3, 4, 17) with the strongest

evidence supporting parenting behaviors, parental mental health symptoms, and attachment.(1,

3, 4)

Understanding the role of psychosocial factors within the family in mediating and moderating the

intergenerational transmission of trauma is critical to develop household-level psychosocial

interventions that prevent transmission among high-risk families. Some psychosocial

interventions have been explicitly developed to prevent the intergenerational transmission of

trauma or have been evaluated for this purpose;(19, 20) however, these interventions are

heterogeneous in their theoretical and therapeutic approaches and in evidence used to inform

development. Understanding mediating and moderating factors could also optimize existing

interventions, helping researchers understand what intervention component(s) are most effective

and in what context(s).

To assess the need for a systematic review on this topic,(21) we conducted a preliminary search

to identify relevant literature reviews (S1 eTable1). We located 21 reviews, most of which were

scoping reviews focused on narrow sub-populations and trauma-event types (S1 eTable 2). Only

two reviews conducted moderation meta-analyses: one restricted to mothers with childhood

maltreatment, and the other only including studies published until 2011. Based on these findings:

(1) no review has published a diagram to demonstrate and contextualize epidemiological causal

relationships at the family-level in the intergenerational transmission of trauma; (2) no review has

meta-synthesized family-level mediators; and (3) there is a need to perform an updated meta-

analysis of family-level moderators across diverse trauma-types.

We aim to fill these gaps by systematically consolidating the family-level psychosocial mediators,

their respective direct and indirect effects, and moderators of such effects, on the

intergenerational transmission of trauma. We will focus on a broad scope of populations and

trauma types to increase the sample size available for meta-synthesis and to understand general

mechanisms. We will also include impacts on the third generation (G3). By using the consolidation

and meta-synthesis of epidemiological data, we will generate the first causal diagram to describe

mechanisms through which trauma is transmitted intergenerationally.

REVIEW QUESTIONS

(1) What family-level psychosocial mediator(s) and family-level psychosocial and

demographic moderator(s) have epidemiological evidence supporting their involvement

in the intergenerational transmission of trauma?

(2) What are the combined quantitative effects across mediators and moderators in the

intergenerational transmission of trauma?

METHODS

This project will follow review guidelines from the Joanna Briggs Institute (JBI) Manual for

Evidence Synthesis (21) and the Cochrane Handbook for Systematic Reviews.(22) We followed

the PRISMA-P(23) (S1 eTable 3) reporting guideline to develop this protocol. This study has been

registered on PROSPERO (registration ID #251053). We will post project materials, statistical

code, and protocol amendments on our Open Science Framework webpage

(https://osf.io/k5ezm/).

Protocol registration and timeline

Preliminary searching and development of the search strategy occurred from July 2020 to March

2021. Reviewer training occurred from January to February 2021. Title/abstract screening is

ongoing (having begun in March 2021) and is anticipated to finish in Summer 2022. Full-text

screening will finish in Fall 2022. Data extraction, data charting, and meta-analyses will finish in

December 2022. We anticipate submitting the manuscript for publication in March 2023.

Inclusion criteria

We will use the systematic review PICOTS (Population, Intervention, Comparator, Outcome,

Timeframe, Study design) framework to define eligibility, adapted to accommodate our research

question focused on observational studies reporting meditator(s) and/or moderator(s) (Table 1).

Definitions/elaborations documents will be available on Open Science Framework

(https://osf.io/k5ezm/).

188 **Table 1**. Eligibility criteria for systematic review.

Inclusion concept	Eligibility criteria
Population	● Human population
Exposure	● At least one primary caregiver (G1) who experienced a traumatic event and their child (G2) and/or grandchild (G3) who did not experience this traumatic event ● The traumatic event in G1 must have occurred before or during the birth of the child under study
Mediator or moderator	● Psychosocial factor(s) operating at the family-level that influence the interpersonal relationship between the child and their immediate family and/or primary caregiver before the child has reached 18 years of age. Factors can be a mediator or moderator ● Comparators will be defined as the absence, or varying levels, of factor(s) measured in at least some sub-group of participants
Outcome	● Quantitative measurement of at least one total, direct, or indirect effect of trauma (psychological symptoms/functioning) in the child under observation at any point during the child's lifespan (including >18 years old)
Timeframe and language	● No date restrictions, but limited to English, French, Kinyarwanda, and Spanish languages (research team members are fluent in these languages)
Study design and document type	● Original research article of a quantitative or mixed methods study published in journals, dissertations (any level), or conference abstracts

Population and Exposure(s)

Included studies will have a population composed of at least one primary caregiver (G1) who

experienced a traumatic event and their child (G2) and/or grandchild (G3) who did not experience

this traumatic event firsthand. The G1 individual would not need to directly report their traumatic

exposure, as long as there is some method to assess this, such as indirect reporting by G2. The

traumatic event in G1 must have occurred before the birth of G2. G1 or G2 generation parents do

not need to be biologically related to the child of interest. We will include trauma-exposed parents

regardless of their trauma-related symptomatology. Animal studies will be excluded.

A traumatic event will be defined as exposure to threatened death, serious injury, or sexual

violence, as outlined in the Diagnostic and Statistical Manual of Mental Disorders.(24). We will

also consider parental trauma-related symptoms as the exposure, contingent that these

symptoms are a direct result of an eligible traumatic exposure.

Mediator(s) and moderator(s)

Included studies will have at least one psychosocial factor operating at the family-level (including

household- and parent-levels) that measured the interpersonal relationship between the child and

their immediate family and/or primary caregiver before the child has reached 18 years of age.

These could be framed or examined as mediators or moderators. Original articles would not need

to explicitly use these terms or conceptualize factors in this way to be eligible.

Psychosocial factors might include parental psychiatric symptoms, suggesting that parental

trauma-related mental health symptoms could be considered as the exposure or the mechanism,

depending on the context. Factors might also include parental psychological factors, interpersonal

parent-child relationships, or social factors within the household-level (which include demographic

factors) that would influence child-rearing behaviors.(25) Comparators will be defined the

measurement of the absence of the factor(s) or varying levels of continuously-measured factor(s)

of interest in at least some sub-group of participants in the study sample. We will also include

demographic factors as moderators.

Outcome(s)

Included studies will quantify either the presence or absence of direct or indirect effects of trauma

in the child (G2 or G3) under observation. Given the heterogeneity in clinical presentations of

intergenerationally-transmitted trauma, we will only include quantitatively measured markers of

psychological symptoms/functioning (i.e., child psychological development, psychopathology,

and/or indicators of wellbeing). To increase our yield, we will be inclusive in what outcomes would

qualify as child psychopathology and/or indicators of wellbeing for screening and may restrict this

criterion after we assess the final sample of included studies. Outcomes will be considered in

offspring. Psychiatric symptoms in the child would not need to be directly observed; for example,

it would be acceptable for G1 to report on G2 symptoms.

Date and language restrictions

This review will have no date restrictions, but will be limited to studies published in English,

French, Kinyarwanda, and Spanish languages, for feasibility, as research team members are

fluent in these languages.

Study design and document type

We will include any epidemiologic study design, such as cross-sectional studies, case-control

studies, cohort studies, randomized controlled trials, or the quantitative component of mixed-

methods studies. Systematic review papers, meta-analyses, case-studies, qualitative studies,

and study protocols will be excluded.

Due to feasibility, we will only include published literature (peer-reviewed literature and

dissertations). One exception is that we will include conference abstracts during the screening

phase contingent upon gaining access to the full text document (see Source of evidence

selection). We included conference abstracts and dissertations to reduce publication bias.

Searches

The search strategy was developed and refined by a medical research librarian (KN) with the

assistance of the first author. The final search strategy was developed using the following Boolean

logic: *[intergenerational concept] AND [family level concept] AND [transmission of trauma*

concept] using appropriate keywords and controlled vocabulary.

We searched MEDLINE, PsycINFO, and PTSDpubs (formerly PILOTS) in January 2021. We also

searched Scopus (conference papers only) and ProQuest Dissertations and Theses (doctoral

dissertations only) in March 2021. S1 eTables 4-8 shows the final search strategy for each

bibliographic database. Although we only retrieved doctoral dissertations from Scopus, our

PsycINFO search captured dissertations of all levels. The final search was peer-reviewed by a

second medical librarian using the Peer Review of Electronic Search Strategies (PRESS)

checklist before the search was finalized.(26)

We limited the search to five databases for feasibility given large search yields. However, to

compensate for this limitation, we will implement three methods to increase sensitivity: (1) robust

forward citation chaining of included studies; (2) reference list screening of relevant reviews from

the authors' personal libraries; and (3) screening additional papers from the authors' personal

libraries.

Source of evidence selection

Title/abstract and full-text screening will be performed independently by two reviewers using

Covidence software(27). Discrepancies will be resolved through screening from a third reviewer.

During full-text screening, we will record the primary reason for exclusion.

We will screen the full-text versions of conference abstracts during full-text review. For each

conference abstract, we will first attempt to locate full-text document online or by emailing the

corresponding author. If we cannot obtain the full text using this process, we will exclude the

record.

Data extraction

We will use Research Electronic Data Capture (REDCap) software to extract data from included

studies.(28) If feasible, data will be extracted independently by two reviewers. Discrepancies will

be resolved through consensus, and when needed, discussion with a third reviewer. We will de-

duplicate the results of individual studies reporting on the same underlying project and will use

this combined information for our analyses. S1 eTable 9 shows the drafted data extraction items,

which will likely be refined after piloting.

Training and piloting procedures

We held formal piloting and training procedures for reviewers before each screening stage. To

prepare for title/abstract screening, reviewers first reviewed training materials, which included a

draft definitions/elaborations document. We then underwent a series of training rounds to assess

inter-rater agreement and identify discrepancies. In each round, reviewers independently

screened a random sample of 25 records from our final search yield to calculate percent

agreement. Then, after each round, the team discussed discrepancies and modified the

definitions/elaborations document. We repeated this process until we achieved team agreement

of >75%, and then began title/abstract screening. We followed similar procedures for full-text

review training. Reviewers first reviewed training materials and draft definitions/elaborations

document for full-text screening. We then underwent a series of training rounds where each

reviewer independently screened a sample of approximately 10 records. Each round was followed

by discussion to modify screening procedures. We began full-text screening after we achieved

agreement of >75%.

Prior to beginning data extraction, the first author will prepare and disseminate training materials

for data extractors. Then, the team of data extractors will pilot data extraction and critical appraisal

of 1-2 papers, discuss discrepancies, and repeat the process until achieving an agreement of

>75% on each item before beginning extraction and appraisal.

Data charting

We will present summary information for all included studies in tabular format (S1 eTable 10). We

will then group the mediators presented in S1 eTable 10 together into similar thematic concepts,

through discussion between the first and senior author. We will present these findings in a second

table (S1 eTable 11).

Meta-analyses

Our statistical analysis plan is presented below with the understanding that our analytical

strategies may be refined based on the distribution of data collected and from input of statistical

experts at the Yale School of Public Health. Not all included studies will be eligible for meta-

analysis. The first author in consultation from the senior author will use information on the quality

of reporting of results to decide what included studies will be eligible for meta-analyses. We will

conduct meta-analyses using R statistical software and will consider a p-value < 0.05 as sufficient

evidence to reject a null hypothesis.

Mediation meta-analyses

We will use a mediation meta-analysis approach to pool data from individual mediation studies.

Our primary analyses will assess the following pathway: parental trauma exposure →

psychosocial mechanisms (including parental trauma-related symptoms) → child mental health

outcome. We will also assess the following pathway: parental trauma-related symptoms

→ psychosocial mechanisms → child mental health outcome. We will conduct several meta-

analyses according to each mediating construct and whether it mediated G1 to G2, G2 to G3, or

G1 to G3 effects. We will conduct sensitivity analyses to test whether we could consolidate

findings of individual studies with different study designs and consolidate individual studies that

measured the exposure as parental trauma versus parental post-traumatic symptoms. We plan

to merge outcomes together to assess the global effect on child psychopathology; if there is

enough data, we will sub-divide outcomes into different meta-analytic models. We will conduct

meta-analyses following guidelines reported in the literature(29) that would be most suitable to

for the methods reported in our sample of included studies.

Moderation meta-analyses

We will aim to use a moderation meta-analysis approach to pool moderator data from individual

studies, using similar methods as described above. Our primary analyses will assess moderating

effects on parental trauma exposure → child mental health outcome; our secondary analyses will

assess moderating effects on parental trauma-related symptoms → child mental health outcome.

We will perform moderation meta-analyses using standard statistical methods.(30)

Assessment of methodological quality

For studies included in any meta-analysis, we will assess the quality of all included articles using

the JBI critical appraisal tools, selected based on the study design of the included paper.(31, 32)

We will apply this framework at the study level. We will also assess the quality of mediation paths

using the quality assessment tool used in other mediation meta-analyses.(15) If performed in

duplicate, discrepancies will be resolved through discussion and consensus, and when needed,

consultation with a third reviewer. If we gather enough studies, we will conduct sensitivity analyses

to assess whether study methodological quality impacts pooled estimates.

Publication bias

We aim to assess for publication bias using visual funnel plot inspection.

Causal diagram

We will develop at least one causal diagram. These diagram(s) will visually demonstrate the

consolidated quantified relationships between variables and their relative magnitudes of

association. We will include paths that are explicitly supported in our data and theoretical paths

not explicitly supported (differentiated using solid versus dashed path lines). We will construct

DAGs where possible if we obtain sufficient information, which would include combined effect

estimates from our meta-analyses.

Assessing confidence in cumulative evidence

We will interpret our results using frameworks for causal inference(33) to provide a narrative

summary and recommendations for interventions.

DISCUSSON

Strengths and limitations

Our methods are rigorous and follow the Cochrane Handbook for Systematic Reviews and

Joanna Briggs Institute systematic review methods manual. We will take additional precautions

to reduce likelihood of publication bias through forward citation chaining and including conference

abstracts and dissertations. This review has several potential limitations. Although we consider

our search strategy comprehensive, we only searched five databases and did not include

Embase. We will not cover all languages, which might contribute language bias. We also might

not locate enough studies to perform meta-analyses, and would instead produce narrative

summaries of quantitative results. Our causal diagrams(s) will be restricted to psychosocial

household-level factors and will exclude other relevant mechanisms outside of this scope such

as biological factors. This review will also be unable to assess complex causal relationships

between mediators, moderators, and their respective interactions; however, we will include this

information in our narrative synthesis and causal diagram(s) if this information happens to be

reported in individual studies.

CONCLUSION

The development of evidence-based causal diagram(s) would provide more detailed understanding of the paths by which unresolved trauma can be transmitted intergenerationally at the family level, including the relative strength of each factor in mediating and moderating cause and effect. These results could then be applied to design and optimize evidence-based interventions that target mechanisms with the strongest mediating effect. Taken together, this review could provide evidence to better support interventions that interrupt the cycle of intergenerational trauma for future generations.

DECLARATIONS

Ethics approval and consent to participate

This systematic review does not require ethical approval.

Availability of data and materials

We will post detailed project materials, resources, statistical code, and protocol amendments on our Open Science Framework webpage (https://osf.io/k5ezm/).

Competing interests

The authors declare no conflicts of interest.

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. EM is supported by the Michael B. Bracken Doctoral Fellowship at the Yale School of Public Health and a Canadian Institutes of Health Research Doctoral Foreign Study

Award (#DFD-175791). SL was supported by an Early-Career Research Fellowship from the Gulf

Research Program of the National Academies of Sciences, Engineering, and Medicine; the

content is solely the responsibility of the authors and does not necessarily represent the official

views of the Gulf Research Program of the National Academies of Sciences, Engineering, and

Medicine.

Authors' contributions

EM and SL developed the study concept and question. EM provided project leadership and

management. EM developed the initial draft of this protocol, with primary feedback from SL, and

all authors reviewed and refined the protocol manuscript. KN and EM developed the search

strategy. KN executed the search strategy. EM developed and revised the protocol for

PROSPERO registration; all authors provided manuscript feedback. EM is the guarantor of the

review.

Acknowledgements

We thank Rhayna Poulin for administrative support.

REFERENCES

1.   Flanagan N, Travers A, Vallières F, Hansen M, Halpin R, Sheaf G, et al. Crossing borders: a systematic review identifying potential mechanisms of intergenerational trauma transmission in asylum-seeking and refugee families. European journal of psychotraumatology. 2020;11(1):1790283-83.

2.   Sangalang CC, Vang C. Intergenerational trauma in refugee families: a systematic review. Journal of immigrant and minority health. 2017;19(3):745-54.

3.   Cerdeña JP, Rivera LM, Spak JM. Intergenerational trauma in Latinxs: A scoping review. Social Science & Medicine. 2021;270:113662.

4.   Isobel S, Goodyear M, Furness T, Foster K. Preventing intergenerational trauma transmission: A critical interpretive synthesis. Journal of clinical nursing. 2019;28(7-8):1100-13.

5.   Kellermann NP. Past, present and future perspectives of Holocaust trauma transmission. ResearchGate. 2019.

6.   O'Neill L, Fraser T, Kitchenham A, McDonald V. Hidden Burdens: a Review of Intergenerational, Historical and Complex Trauma, Implications for Indigenous Families. J Child Adolesc Trauma. 2018;11(2):173-86.

7.   Dekel R, Goldblatt H. Is there intergenerational transmission of trauma? The case of combat veterans' children. Am J Orthopsychiatry. 2008;78(3):281-9.

8.   Payne EA, Berle D. Posttraumatic stress disorder symptoms among offspring of Holocaust survivors: A systematic review and meta-analysis. Traumatology. 2020.

9.   Morris A, Gabert-Quillen C, Delahanty D. The Association Between Parent PTSD/Depression Symptoms and Child PTSD Symptoms: A Meta-Analysis. Journal of Pediatric Psychology. 2012;37(10):1076-88.

10.   van IMH, Bakermans-Kranenburg MJ, Sagi-Schwartz A. Are children of Holocaust survivors less well-adapted? A meta-analytic investigation of secondary traumatization. J Trauma Stress. 2003;16(5):459-69.

11.   Lindert J, Knobler HY, Kawachi I, Bain PA, Abramowitz MZ, McKee C, et al. Psychopathology of children of genocide survivors: a systematic review on the impact of genocide on their children`s psychopathology from five countries. Int J Epidemiol. 2017;46(1):246-57.

12.   Palacios JF, Portillo CJ. Understanding Native women's health: historical legacies. J Transcult Nurs. 2009;20(1):15-27.

13.   van Steenwyk G, Roszkowski M, Manuella F, Franklin TB, Mansuy IM. Transgenerational inheritance of behavioral and metabolic effects of paternal exposure to traumatic stress in early postnatal life: evidence in the 4th generation. Environmental epigenetics. 2018;4(2):dvy023-dvy23.

14.   Sagi-Schwartz A, van Ijzendoorn M, Bakermans-Kranenburg M. Does intergenerational transmission of trauma skip a generation? No meta-analytic evidence for tertiary traumatization with third generation of Holocaust survivors. Attachment & human development. 2008;10:105-21.

15.   Lee H, Hübscher M, Moseley GL, Kamper SJ, Traeger AC, Mansell G, et al. How does pain lead to disability? A systematic review and meta-analysis of mediation studies in people with back and neck pain. Pain. 2015;156(6):988-97.

16.   Glymour MM, Greenland S. Causal diagrams. Modern epidemiology. 2008;3:183-209.

17.   Dashorst P, Mooren TM, Kleber RJ, de Jong PJ, Huntjens RJC. Intergenerational consequences of the Holocaust on offspring mental health: a systematic review of associated factors and mechanisms. European journal of psychotraumatology. 2019;10(1):1654065-65.

18.   Yehuda R, Lehrner A. Intergenerational transmission of trauma effects: putative role of epigenetic mechanisms. World Psychiatry. 2018;17(3):243-57.

19.   Colegrove VM, Havighurst SS, Kehoe CE, Jacobsen SL. Pilot randomized controlled trial
of Tuning Relationships with Music: Intervention for parents with a trauma history and their
adolescent. Child Abuse and Neglect. 2018;79:259-68.
20.   Lawler J, Rosenblum K, Schuster M, Muzik M. Mom Power: A Parenting Group
Intervention for Mothers with Trauma Histories. In: Muzik M, Rosenblum K, editors. Motherhood
in the Face of Trauma. Integrating Psychiatry and Primary Care. Springer; 2018.p. 165-80.
https://doi.org/10.1007/978-3-319-65724-0_11.
21.   Aromataris E MZ. JBI Manual for Evidence Synthesis 2020 [Available from:
https://synthesismanual.jbi.global.
22.   Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. Cochrane
handbook for systematic reviews of interventions: John Wiley & Sons; 2019.
23.   Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred
reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement.
Systematic reviews. 2015;4(1):1-9.
24.   Association AP. Diagnostic and statistical manual of mental disorders (DSM-5®):
American Psychiatric Pub; 2013.
25.   Suzuki S-i, Takei Y. Psychosocial Factors and Traumatic Events. In: Gellman MD,
Turner JR, editors. Encyclopedia of Behavioral Medicine. New York, NY: Springer New York;
2013. p. 1582-3.
26.   McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer
Review of Electronic Search Strategies: 2015 Guideline Statement. Journal of Clinical
Epidemiology. 2016;75:40-6.
27.   Covidence software [Available from: https://www.covidence.org/.
28.   REDCap [Internet]. 2021. Available from: https://www.project-redcap.org/.
29.   Jak S, Li H, Kolbe L, de Jonge H, Cheung MW. Meta-analytic structural equation
modeling made easy: A tutorial and web application for one-stage MASEM. Res Synth Methods.
2021;12(5):590-606.
30.   Lipsey MW, Wilson DB. Practical meta-analysis: SAGE publications, Inc; 2001.
31.   Joanna Briggs Institute. The Joanna Briggs Institute Critical Appraisal tools for use in JBI
Systematic Reviews: Checklist for Analytical Cross Sectional Studies. Adelaide: The Joanna
Briggs Institute. 2017.
32.   Joanna Briggs Institute. The Joanna Briggs Institute Critical Appraisal tools for use in JBI
Systematic Reviews: Checklist for Cohort Studies. Adelaide: The Joanna Briggs Institute. 2017.
33.   Rothman KJ, Greenland S. Causation and causal inference in epidemiology. Am J
Public Health. 2005;95(S1):S144-S50.
520