TITLE PAGE

Article type: Original article

Characterization of the skin microbiota in bullous pemphigoid patients and controls reveals novel microbial indicators of disease

Meriem Belheouane, PhD[1,2,3,*], Britt M. Hermes, MS[1,2,*], Nina Van Beek, MD[4], Sandrine Benoit, MD[5], Philippe Bernard, MD, PhD[6], Kossara Drenovska, MD, PhD[7], Sascha Gerdes, MD[8], Regine Gläser, MD[8], Matthias Goebeler, MD[5], Claudia Günther, MD[9], Anabelle von Georg, MD[4], Christoph M. Hammers, MD, PhD[4,18], Maike M. Holtsche, MD[4], Bernhard Homey, MD[10], Orsolya N. Horváth, MD, PhD[11], Franziska Hübner, MD[4], Beke Linnemann, MD[4], Pascal Joly, MD[12], Dalma Márton, MD[13], Aikaterini Patsatsi, MD, PhD[14], Claudia Pföhler, MD[15], Miklós Sárdy, MD[11,13], Laura Huilaja, MD, PhD[16], Snejina Vassileva, MD, PhD[7], Detlef Zillikens, MD[4,17], Saleh Ibrahim, MD, PhD[17,18,19], Christian D. Sadik, MD[4,17], Enno Schmidt, MD, PhD[4,17,18+], John F. Baines, PhD[1,2+]

Author Affiliations

[1] Max Planck Institute for Evolutionary Biology, August-Thienemann-Str. 2, 24306 Plön, Germany

[2] Section of Evolutionary Medicine, Institute for Experimental Medicine, Kiel University, Arnold-Heller-Str. 3, 24105 Kiel, Germany

[3] Research Center Borstel, Leibniz Lung Center, Parkallee 1-40, 23845, Borstel, Germany.

[4] Department of Dermatology, Allergy, and Venereology, University of Lübeck, 23538 Lübeck, Germany

[5] Department of Dermatology, Venereology and Allergology, University Hospital Würzburg, Würzburg, Germany

[6] Department of Dermatology, Reims University Hospital, University of Reims Champagne-Ardenne, Reims, France

[7] Department of Dermatology and Venereology, Medical University - Sofia, 1, St. G. Sofiiski str., 1431 Sofia, Bulgaria

[8] Department of Dermatology, Venereology and Allergology, University of Kiel, 24105 Kiel, Germany

[9] Department of Dermatology, University Hospital, TU Dresden, 01307 Dresden, Germany

[10] Department of Dermatology, University Hospital Düsseldorf, Medical Faculty, Heinrich-Heine-University Düsseldorf, Moorenstr. 5, 40225 Düsseldorf

[11] Department of Dermatology and Allergy, University Hospital, LMU Munich, Munich, Germany

[12] Department of Dermatology, Rouen University Hospital, INSERM U1234, Normandie University, Rouen France

[13] Department of Dermatology, Venereology and Dermatooncology, Semmelweis University, Budapest, Hungary

[14] Autoimmune Bullous Diseases Unit, 2nd Dermatology Department, Aristotle University School of Medicine, Papageorgiou General Hospital, 56403 Thessaloniki, Greece

[15] Department of Dermatology, University of Saarland, Homburg/ Saar, Germany

[16] PEDEGO Research Unit, University of Oulu; Department of Dermatology and Medical Research Center Oulu, Oulu University Hospital, Oulu, Finland

[17] Center for Research on Inflammation of the Skin (CRIS), University of Lübeck, 23538

Lübeck, Germany

[18] Lübeck Institute of Experimental Dermatology (LIED), University of Lübeck, Lübeck,

Germany

[19] College of Medicine and Sharjah Institute for Medical Research, University of Sharjah, 27272

Sharjah, UAE.

*Authors contributed equally to this work.

+Authors contributed equally to this work.

Corresponding author:

John F. Baines

Max Planck Institute for Evolutionary Biology,

August-Thienemann-Str. 2, 24306 Plön, Germany

Email: baines@mpg.evolbio.de

ORCiDS

John F. Baines: https://orcid.org/0000-0002-8132-4909

Meriem Belheouane: https://orcid.org/0000-0002-2939-4862

Britt Marie Hermes: https://orcid.org/0000-0001-6655-6667

Saleh Ibrahim: https://orcid.org/0000-0001-7827-2290

Enno Schmidt: https://orcid.org/0000-0002-1206-8913

Detlef Zillikens: https://orcid.org/0000-0002-5668-307X

Christian D. Sadik: https://orcid.org/0000-0001-6701-048X

Nina Van Beek: https://orcid.org/0000-0002-3265-0011

Pascal Joly: https://orcid.org/0000-0002-5734-0255

Christoph Matthias Hammers: https://orcid.org/0000-0001-5631-2415

Aikaterini Patsatsi: https://orcid.org/0000-0001-9616-2001

Matthias Goebeler: https://orcid.org/ 0000-0001-7095-9848

Anabelle von Georg: https://orcid.org/0000-0003-2436-8937

Laura Huilaja: https://orcid.org/0000-0002-7597-4323

Claudia Günther: https://orcid.org/0000-0002-4330-1861

Maike Marleen Holtsche: https://orcid.org/0000-0002-5318-5796

Regine Gläser: https://orcid.org/0000-0002-5819-777X

Kossara Drenovska: https://orcid.org/0000-0002-8254-1630

Snejina Vassileva: https://orcid.org/0000-0003-1623-2482

Sascha Gerdes: https://orcid.org/ 0000-0002-6667-7757

Claudia Pföhler: https://orcid.org/0000-0003-0384-114X

Sandrine Benoit: https://orcid.org/0000-0002-2932-7539

Orsolya N. Horváth: https://orcid.org/0000-0002-2201-349X

Bernhard Homey: https://orcid.org/0000-0002-5784-4146

Dalma Márton: https://orcid.org/0000-0003-1686-1131

Philippe Bernard: https://orcid.org/0000-0002-9115-0690

Beke Linnemann: https://orcid.org/0000-0002-1921-8726

Franziska Hübner: https://orcid.org/0000-0003-2886-9499

Miklós Sárdy: https://orcid.org/0000-0003-4306-5093

Cities, States, and Countries in which the work was done:

Budapest, Central Hungary (Region) Hungary

Düsseldorf, North Rhine-Westphalia Germany

Dresden, Saxony Germany

Freiburg, Saxony Germany

Lübeck, Schleswig-Holstein Germany

Homburg, Saarland Germany

Kiel, Schleswig-Holstein Germany

München, Bavaria Germany

Würzburg Bavaria Germany

Oulu, North Ostrobothnia (Region) Finland

Reims, Grand Est (Region) France

Rouen, Normandy (Region) France

Sofia, Sofia City (Province) Bulgaria

Thessaloniki, Macedonia (Region) Greece

Short title: Skin microbiota in bullous pemphigoid

Abbreviation: AI: autoimmune; BP: bullous pemphigoid; AIBD: Autoimmune blistering disease; EBA: Epidermolysis bullosa acquisita; DF: degrees of freedom; IS: indicator species**,** ASV: amplicon sequence variant

ABSTRACT

Bullous pemphigoid (BP) is an autoimmune skin blistering disease afflicting mostly the elderly and is associated with significantly increased mortality. Here, we conducted the most extensive sampling effort of skin microbiota in BP to date to analyze whether intra-individual, body-site-specific, and/or geographical variation contributes to the emergence of BP. We find marked differences in the skin microbiota of BP patients compared to that of control subjects, and moreover that disease status rather than skin biogeography governs the skin microbiota composition in BP. Our data reveal a discernible transitional stage between normal and diseased skin in BP characterized by a loss of protective microbiota and an increase in sequences matching *Staphylococcus aureus,* a known inflammation-promoting species. Notably, *S. aureus* is ubiquitously associated with disease status, suggesting that this taxon is an important indicator of BP. Importantly, differences in a few key indicator taxa are able to reliably discriminate between BP patients and controls, characterized by their opposing abundance patterns. This may serve as valuable information for assessing disease risk and treatment outcomes. Future research will focus on the functional analysis of host-microbe and microbe-microbe interactions and the relevance of the host genome for microbiota abundances to identify novel BP treatment approaches.

INTRODUCTION

Bullous pemphigoid (BP) is the most common autoimmune skin blistering disease (AIBD) in

Europe, with an annual incidence of about 20 new cases per million in this region (Joly et al.,

2012; Schmidt and Zillikens, 2013; van Beek et al., 2021). It occurs when autoantibodies attack

two structural hemidesmosomal proteins of the epidermal basement membrane, i.e. BP180 (type

XVII collagen) and BP230, resulting in subepidermal blistering (Schmidt and Zillikens, 2013;

Stevens et al., 2019; Amber et al., 2018). The severity of this highly pruritic AIBD considerably

affects quality of life and is associated with significantly increased mortality (Amber et al.,

2018). The incidence of BP is increasing with the ageing European population (van Beek et al.,

2021; Schmidt and Zillikens, 2013). It is thus intriguing that a recent multinational study of

9,000 participants showed that the skin microbiota are a predictor of age, more so than oral or

gut microbiota (Huang et al., 2020).

The observation of age-dependent changes in the skin microbiota as well as the skin

microbiota's involvement in other inflammatory skin disorders raise the possibility that it may

play a role in the emergence of AIBD. Previous efforts by Srinivas et al. (2013) demonstrated

that genotype-dependent microbiota affect disease susceptibility in a mouse model of

epidermolysis bullosa acquisita (EBA), an AIBD similar to BP. Similarly, Miodovnik et al.

(2017) present pilot human data suggesting that skin microbiota contribute to the pathogenesis of

BP. However, identification of candidate bacterial taxa or important changes in microbial

diversity associated with BP remain uncharacterized.

Here, we conducted a large-scale investigation of BP patients and age- and sex-matched

control subjects within Europe to clarify relationships between microbiota and BP pathogenesis.

By examining perilesional sites and comparing them to unaffected but BP-susceptible body sites

in patients and in matched controls, we reveal clear microbial indicators of BP patients versus controls. Genetic detection of microbial signatures associated with AIBD could enable early intervention and thus, better clinical outcomes.

RESULTS

Sampling

Two-hundred twenty individuals with BP and 190 age- and sex-matched control subjects from fourteen study sites across Europe were included (Methods). We performed 16S rRNA gene sequencing on bacterial genomic DNA derived from swabbing four categories of body sites (Figure 1a). These include areas adjacent to a fresh blister or erosion ("perilesional"), unaffected skin from the contralateral site from the same patient ("contralateral"), and the same body site of an age- and sex-matched control subject ("corresponding"). Sites that are typically unaffected by BP (forehead, upper back, and antecubital fossa) were sampled to obtain a better picture of BP across skin biogeography (Figure 1a).

Figure 1. Sampling of various body sites for patients and controls; Box plots of Shannon (alpha) diversity. 1a. Grey figure represents age-and sex-matched control; orange figure represents BP patient. Sites typically unaffected by BP include the forehead (purple), upper back (turquoise), and antecubital fossa (dark blue) are represented on both figures. An example perilesional sampling site (red), unaffected contralateral site (yellow), and control matched corresponding site (green) are shown. **1b.** Shannon diversity at the ASV-level for sites typically unaffected by BP for controls and BP patients. **1c.** Shannon diversity at the ASV-level for patient perilesional, patient contralateral, and control corresponding sites. Significance represented by: * ≤ 0.05; ** ≤ 0.01; *** ≤ 0.001; **** ≤ 0.0001. Supplementary Table S1 reports summary statistics.

Reduced alpha diversity within lesional and BP-susceptible sites

At the ASV-level, we found that the Shannon (Figure 1b) and Chao1 (Supplementary Figure S1a) indices are similar at sites typically unaffected by BP in patients and controls. In contrast, control corresponding sites display higher bacterial diversity than patient contralateral sites, which in turn are more diverse than perilesional sites for Shannon (Figure 1c) and Chao1 indices (Supplementary Figure S1b).

Critically, study center, disease status, and sex significantly correlate with Shannon diversity for patient perilesional and contralateral sites as well as for control corresponding sites ($F_{37,1118} = 7.24$; R^2_{adj}: 0.17; $p < 0.001$), with disease status explaining 8.25% of the variance. Likewise, disease status and study center significantly correlate with Chao1 richness ($F_{37,1118} = 6.03$; $R^2_{adj} = 0.14$; $p < 0.001$), with study center explaining 7.86% and disease status explaining 1.41% of the Chao1 variance. Disease status associates with a decrease in Shannon diversity in patient perilesional and contralateral sites ($\beta = -0.72, -0.38$, respectively), and a decrease in Chao1 richness, whereby perilesional and contralateral sites display a similar reduction in diversity ($\beta = -39.38, -30.90$, respectively). Thus, disease status associates with a significant decrease in alpha diversity, which is still present after accounting for potential confounding variables.

Supplementary Material R1 reports results for sites typically unaffected by BP. Analysis of sum of squares shows that the effects of disease status on alpha diversity does not extend to sites typically unaffected by BP and that skin biogeography likely characterizes microbial diversity at these sites.

Beta diversity in relation to disease, individual, and sampling features

We conducted a non-parametric multivariate analysis of variance using distance matrices analysis (PERMANOVA) applied to the Bray-Curtis index as a measure of beta (between sample) diversity of the sites typically unaffected by BP to evaluate effects of potential confounding variables (Supplementary Material R2). Analysis of disease status per body site reveals a significant association with disease status (adonis: Bray-Curtis~disease status: body site; R^2=0.003; $p < 0.001$). Partial constrained principal coordinate analysis reveals that patients and controls cluster according to body site along the first and second axes and that forehead and upper back (typical sebaceous zones) are more similar to each other compared to the antecubital fossa (Figure 2b). These findings suggest that the microbial variation among sites typically unaffected by BP is likely linked to skin biogeography rather than disease or study center (Figure 2a, 2c).

   After analyzing beta diversity between perilesional, contralateral, and corresponding sites as described above (Supplementary Material R2), significant associations with disease and blistering status are observed (i.e., perilesional and contralateral sites within patients), which explain 2.12% and 0.29% of the variance, respectively (adonis: disease status R^2=0.02; blistering status $R^2 = 0.003$; $p < 0.001$). However, study center explains 3.41% of the variance (adonis: $R^2 = 0.03$; $p < 0.001$), suggesting environmental effects. Analysis of interaction between variables reveals that disease status accounts for significant differences between study centers (adonis: disease status: study center, $R^2 = 0.03$; $p < 0.001$). Additionally, partial constrained principal coordinate analysis reveals that on the first and second axes control corresponding sites are distinguishable from patient contralateral and perilesional sites, which largely cluster together

(Figure 2d, e). Figure 2f, on the other hand, shows comparatively little clustering according to study center.

Figure 2. Partial constrained principal coordinate analyses of Bray-Curtis

2a to 2c. Body sites typically unaffected by BP. (anova.cca, Full model: $p = 0.0009$; terms: disease status, body site (constrained inertia=5.04%, conditioned inertia=4.5%), study center: $p < 0.001$; axes: CAP1, CAP2: $p = 0.09$; 1,000 permutations). **2d to 2f**. Patient perilesional, contralateral sites and control corresponding sites. (anova.cca, Full model: $p < 0.001$; terms: disease status, blistering status, study center: $p < 0.001$; axes: CAP1, CAP2: $p = 0.009$; 1,000 permutations). "+" represents centroid. SD: standard deviation. Site abbreviations: Budapest, Hungary (BUD); Düsseldorf, (D), Dresden, (DD), Freiburg, (FR), Lübeck, (HL), Homburg, (HOM), Kiel, (KI), München, (M), Würzburg (WUE), all Germany; Oulu, Finland (OU); Reims, (RE), Rouen, (RO), both France, Sofia, Bulgaria (SO), Thessaloniki, Greece, (TH).

Indicator species of BP patients and controls

We conducted four indicator species analyses at the ASV-level (Methods). To refine taxonomic classification of indicator ASVs, we queried representative sequences using RDP SeqMatch

(Supplementary Table S2). ASVs strongly associated with BP patients or controls

(Supplementary Table S3) are shown in Figure 3.

Several indicator ASVs known to be human commensals associate with sites typically

unaffected by BP. Importantly, we identify a greater number of ASVs associating with control

sites, coherent with the loss of diversity in BP. ASV_1, which closely matches

Propionibacterium acnes (*P. acnes*) [now *Cutibacterium acnes;* (Scholz and Kilian, 2016)], is an

indicator for controls for all sampling categories. Among sites unaffected by BP, body site is

associated with 13% of the variance in *P. acnes* abundance; abundance is accordingly higher at

sebaceous sampling sites (forehead, upper back). Disease status accounts for 0.79% variance of

P. acnes abundance ($F_{39,1123}$ = 11.58; R^2_{adj} = 0.26; *p < 0.001*). Interestingly, disease status

accounts for 0.63% of the variance in *P. acnes* abundance among sites unaffected by BP

(interaction model, $F_{5,1157}$ = 39.06; R^2_{adj} = 0.14; *p < 0.001*) and is associated with a decrease in

P. acnes abundance at the upper back (β = –0.11).

Important patterns are also apparent among corresponding, patient perilesional, and

patient contralateral sites. *P. acnes* abundance is associated with study center, blistering status,

and sex ($F_{37,1118}$ = 5.40; R^2_{adj} = 0.12, *p < 0.001*), with sex explaining 7.17% of the variance.

Furthermore, its relative abundance is higher at corresponding sites and relatively lower at

perilesional sites (β = +0.03, –0.003, respectively). Additionally, ASV_4 [which closely matches

Corynebacterium tuberculostearicum (*C. tuberculostearicum*)] is an indicator for control

corresponding sites and for patient contralateral sites. Study center, blistering status, and sex are

significantly associated with *C. tuberculostearicum* abundance ($F_{37,1188}$ = 3.92; R^2_{adj} = 0.09; *p <*

0.001), with blistering status explaining 4.06% of the variance and correlating with an increase in

abundance in corresponding sites, but a decrease in perilesional sites (β = +0.05, –0.04, respectively).

Figure 3. Bar plots of mean relative abundance for ten most important indicator species. 3a. Bar plot showing relative abundance of important indicator species, at the ASV-level, for controls and BP patients, at sites typically unaffected by BP [antecubital fossa (AF), forehead (FH), and upper back (UB)]. **3b.** Bar plot showing the relative abundances of important indicator species at the ASV-level for patient perilesional, contralateral sites and control corresponding sites. RDP SeqMatch results for the representative ASV sequences are shown in the legend and provided in full in Supplementary Table S2. Supplementary tables S3, S4 provide statistical parameters for indicator species analyses and summary statistics of all indicator ASVs, respectively.

Contrasting patterns of *Staphylococcus* ASVs in BP patients and controls

Among the indicator ASVs, six belong to *Staphylococcus* and display contrasting patterns associated with disease status. *Staphylococcus* ASV_5 (which closely matches *S. hominis*) correlates with disease status at control corresponding and typically unaffected body sites. Both

disease status and body site are significantly associated with *Staphylococcus* ASV_5 abundance ($F_{39,1123}$ = 6.45; R^2_{adj} = 0.16; *p < 0.001*). However, body site explains a greater proportion of variance (11.17%) compared to disease status (1.24%), whereby the latter is associated with a decrease in abundance (β = –0.04).

In contrast, *Staphylococcus* ASV_2 (which closely matches *S. aureus*) is a strong indicator for disease even at the sites unaffected by BP. Here, disease status associates with the abundance of *Staphylococcus* ASV_2 (7.36%), but body site only slightly associates with this indicator (1.52%; $F_{39,1123}$ = 5.12; R^2_{adj} = 0.12; *p < 0.001*). Disease status associates with an increase in *Staphylococcus* ASV_2 abundance at sites unaffected by BP (β = 0.08). Among perilesional, contralateral and corresponding sites, body site associates with the greatest amount of variance for *Staphylococcus* ASV_2 abundances, followed by study center (6.45%) and sex (0.77%; $F_{37,1118}$ = 9.09; R^2_{adj} = 0.21, *p < 0.001*). This is characterized by a decrease of *Staphylococcus* ASV_2 abundance in corresponding sites (β = –0.12) compared to an increase at perilesional sites (β = +0.07).

Because individual members of *Staphylococcus* can display antagonistic interactions in the context of inflammatory skin disorders (Nakatsuji et al., 2017), we examined pairwise correlations among the top ten indicator ASVs (Supplementary Figures S3, S4; Supplementary Table S6). Importantly, *Staphylococcus* ASV_2 and ASV_5 display significant negative correlations among patient perilesional, contralateral, and antecubital fossa sites, whereas no significant correlation is present among any control sites. Furthermore, *Staphylococcus* ASV_2 is significantly negatively correlated with sequences matching *P. acnes* (ASV_1) at all categories of patient sites, which is absent at all control sites. This suggests a fundamental alteration in the nature of community interactions among members of *Staphylococcus* in the context of BP.

Staphylococcus ASVs predict disease status in random forest classification

Random forest classification analyses (Methods) reveal indicator ASVs to accurately classify samples (mtry = 15; 849/868 controls and 1,443/1,451 BP patients; mean classification accuracy 99.00%) when applied to all sampling sites. Prediction accuracy approaches 100% when applied to only corresponding, contralateral, and perilesional sites (mtry = 18 ; 324/334 controls and 822/822 BP patients; mean classification accuracy 99.15%; Supplementary Figures S2a, S2b). By inspecting the mean decrease accuracy components for ASVs, those belonging to the *Staphylococcus* genus are identified as being most important to both models (Supplementary Table S5).

To estimate the discriminatory power of *Staphylococcus* ASVs alone, we limited the random forest classification analyses to *Staphylococcus* ASVs with an abundance greater than 2% within each sample. We found that *Staphylococcus* ASVs accurately distinguish between controls and BP patients (mtry = 52; 790/868 controls and 1,446/1,451 BP patients; mean classification accuracy 96.40%) when applied to all sampling sites, and are similarly accurate when applied using only corresponding, perilesional, and contralateral sites (mtry = 62; controls 294/334 and 819/822 BP patients; 96.20%; Supplementary Figures S2c, S2d). Notably, inspection of mean decrease accuracy components indicates that *S. aureus* ASV_2 is the most important ASV for model accuracy.

DISCUSSION

This study reveals marked differences in the skin microbiota of BP patients compared to that of sex- and age-matched control subjects with non-inflammatory/ non-infectious dermatoses. This was accomplished by conducting a large-scale sampling- and bacterial 16S rRNA gene analysis,

utilizing a sampling scheme that accounts for both skin biogeography and disease status (Figure 1a). This study represents the most substantial sampling effort of skin microbiota in BP to date.

Among the first notable results is that we observe a significant reduction in alpha diversity at both perilesional sites and contralateral sites in BP patients compared to site-matched areas from control subjects. Furthermore, blistering status (i.e., patient perilesional versus unaffected, contralateral sites) is associated with a fewer number of indicator ASVs compared to matched sampling sites from control subjects. This reduction in alpha diversity in BP patients is consistent with findings from other studies of inflammatory skin diseases, including psoriasis (Quan et al., 2020; Yerushalmi et al., 2019), atopic dermatitis (Clausen et al., 2018) as well as a mouse model of the BP-like disease EBA (Srinivas et al., 2013).

BP blisters characteristically occur on the flexor surfaces of the extremities and the trunk (Schmidt and Zillikens, 2013). The clear biogeography of human skin microbiota, whereby distinct assemblages colonize different body sites depending upon numerous factors, suggest that conditions affecting the skin micro-environment, and thereby skin microbiota, may influence susceptibility to blistering (Kong and Segre, 2017; Oh et al., 2014). Our data show that in the control subjects' skin areas that correspond to blistering sites in BP patients, biogeography— rather than disease effect—is associated with a greater proportion of variance in *P. acnes* and *S. hominis* abundances, both of which can act as beneficial skin commensals (Nakatsuji et al., 2017; O'Neill and Gallo, 2018).

Our data reveal that BP might also lead to a loss of protective microbiota in sites typically unaffected by BP. At the upper back, an interaction model revealed that the effect of disease status significantly associates with a decrease in *P. acnes* relative abundance in BP patients. This is notable given that the upper back represents a sebaceous skin zone where we would expect

relatively high amounts of *P. acnes* (McLaughlin et al., 2019; O'Neill and Gallo, 2018).

Although *P. acnes* is commonly thought of as a potential pathogenic species responsible for

acne, it also acts as an important commensal that aids in preventing the colonization and invasion

of pathogens via the production of antimicrobials and hydrolysis of triglycerides (Gribbon et al.,

1993; O'Neill and Gallo, 2018), as well as the production of short-chain fatty acids (Nakamura et

al., 2020; Shu et al., 2013). For *S. hominis,* we found that disease status is associated with a

decrease in abundance at typically unaffected sites. In contrast, *S. aureus* relative abundance is

increased at these sites. It is thus possible that the nature of antagonistic effects between different

Staphylococcus species may be fundamentally altered in BP patients, e.g. based on strain-level

differences in the production of antimicrobials as in atopic dermatitis (Nakatsuji et al., 2021,

2017). Decreased commensal microbiota perhaps translate to fewer protective immune functions

in the skin, which in turn could allow for an increased colonization of inflammation-promoting

species like *S. aureus* (Nakatsuji et al., 2017), which can exacerbate or accelerate blister

formation. Thus, our observations might be capturing a transitional step between areas without

lesions and areas with blisters or erosions in BP patients.

Accordingly, *S. aureus* is known to dominate the skin microbiota of patients with atopic

dermatitis and exacerbates the disease through inflammation (Kong et al., 2012; Nakatsuji et al.,

2017). Our data suggest that *S. aureus* may similarly drive the development of BP lesions.

Notably, we found that *S. aureus* is associated with disease status regardless of sampling site,

suggesting that this taxon is an important indicator of BP. The mean relative abundance of *S.

aureus* is increased in sites typically unaffected by BP as well, as in the perilesional and

contralateral sites. The specific role of this microbe and its functional components, e.g., how it

might drive BP pathogenicity, will require further exploration.

In addition to cutaneous micro-environmental differences, geographic locations of BP patients should be considered, as there is significant global variation in microbial colonization, especially as it relates to disease susceptibility (Blaser et al., 2013; Chen and Tsao, 2013; Hospodsky et al., 2014; Rehman et al., 2016). Population differences observed in the gut microbiota in patients with inflammatory bowel disease, for example, suggest a complex interplay between geography and gut diseases that are in part driven by microbial factors (Rehman et al., 2016). Therefore, a broad-scale sampling of BP patients across regions with variable incidences could reveal population-specific characteristics that might affect disease predisposition. We found that the study center is significantly associated with the relative abundances of *S. aureus, C. tuberculostearicum*, and *P. acnes* within perilesional and contralateral sites in BP patients as well as control corresponding sites, suggesting an interplay between disease status, skin biogeography, and human geography. Further, an interactive analysis of beta diversity profiles reveals that disease status significantly differs between study centers. We recognize that geography represents an assemblage of factors including diet, culture, ancestry, and environmental features. Our results suggest the need for a large, global study in order to disentangle the relative importance of these features on the assembly of the skin microbiota, especially as it pertains to disease onset in AIBD.

In summary, our study provides evidence that the cutaneous microbiota may play an important role in the emergence of skin lesions in BP. Given the clear discriminatory power provided by differences in a few key indicator taxa, their relative proportions have the potential to provide critical information for assessing disease risk as well as treatment outcomes. Future research may focus on functional analysis of host-microbe and microbe-microbe interactions as a means to identify novel treatment approaches for BP.

MATERIALS AND METHODS

Study participants

Four-hundred twenty-seven volunteers were recruited from fourteen study centers across Europe (Supplementary Methods M1), approved by the University of Lübeck ethics committee (15-051, 18-046), as well as the respective committees of the study centers, following the Declaration of Helsinki. Written, informed consent was obtained from each participant. BP patients (n = 227) included 114 males, 112 females, and one sex "unspecified" participant, average age 80 years, with newly diagnosed or relapsed BP without previous systemic treatment of prednisolone, dapsone, doxycycline, immunosuppressants, and/or use of topical therapies for more than seven days prior to sampling or topical antiseptic use within the last two weeks. Inclusion and exclusion criteria are described in detail in Supplementary Methods M1. Age- and sex-matched subjects (n = 190) included 104 males and 86 females with non-inflammatory/non-infectious dermatoses (mostly basal cell or squamous cell carcinoma), with an average age of 80 years. Contemporaneous controls were sought for each BP subject in minimal time after sampling, which was largely successful (controls per case for males and females are 0.91 and 0.77, respectively). None of the study participants received systemic antibiotics within seven days of sampling.

Sampling, DNA extraction, and 16S rRNA sequencing

Briefly, cotton swabs were used to sample skin surfaces, placed in buffer solution, and stored at –80°C. Ambient air (n = 19) and negative extraction controls (n = 43) were processed alongside samples. ZymoBIOMICS Microbial Community Standard cells (Zymo Research) were used as

extraction and sequencing controls to assess contamination (Karstens et al., 2019). Hypervariable regions V1-V2 of the bacterial 16S rRNA gene were amplified and sequencing was performed using the dual-index sequencing strategy for amplicon sequencing on the MiSeq Illumina platform (Kozich et al., 2013). Final sample sizes included 2,319 skin swabs comprising 1,451 patient and 868 matched control subject swabs. Detailed methods are described in Supplementary Methods M2.

Data processing and taxonomic classification

Data processing and statistical analyses were performed using R (version 4.0.2). Briefly, sequences were processed using DADA2 (version 1.16.0), resulting in abundance tables of ASVs (Callahan et al., 2016). To normalize sequencing coverage, random sub-sampling to 5,000 sequences per sample was performed (Belheouane et al., 2020). Decontam (version 1.8.0; (Davis et al., 2018) was used within Phyloseq (version 1.32.0; (McMurdie and Holmes, 2013) to identify potential contaminant ASVs, according to the prevalence method (Karstens et al., 2019). ASVs classified to families Halomonadaceae (n = 1,040) and Shewanellaceae (n = 211) were removed, following recommendations of Weyrich et al. (2019). Taxonomic assignment of ASVs was completed in DADA2 with the Bayesian classifier using the NR Silva database training set, version 138 (Quast et al., 2013). Representative 16S rRNA gene sequences were queried via the Ribosomal Database Project (RDP; release 11.6; (Wang et al., 2007) SeqMatch (version 3; (Cole et al., 2005); Supplementary Table S2). Further details of 16S rRNA gene sequence processing are described in Supplementary Methods M3.

Ecological and statistical analyses

Alpha diversity was measured using Shannon and Chao1 indices with vegan (version 2.5-6) on absolute abundance data. Beta diversity was calculated using the Bray-Curtis dissimilarity index. We performed a non-parametric multivariable analysis of variance using distance matrices (PERMANOVA) using the "adonis" function with 1,000 permutations and a partial constrained principal coordinate analysis of beta diversity measures using the "capscale" function in vegan (Oksanen et al., 2005). The significance of models, axes, and terms were assessed using the "anova.cca" function with 1,000 permutations.

Indicator species analysis was applied using indicspecies (version 1.7.9) with the "r.g." function (Cáceres and Legendre, 2009) and 100,000 permutations. Random Forest classification and regression analyses were performed using randomForest (version 4-6-14; (Liaw and Wiener, 2002). Models were constructed with 100,000 trees, with "mtry" parameter set for each model and linear models constructed to evaluate potential disease effects. Adjusted R^2 values reported, beta coefficient values express directionality. Further details of ecological and statistical analyses are given in Supplementary Methods M4.

Funding sources: This work was supported by the German Research Foundation (DFG), through Clinical Research Unit 303, project number 269234613, subproject P2, jointly awarded to Prof. Dr. John F. Baines and Prof. Dr. Dr. Enno Schmidt.

DATA AVAILABILITY

Datasets related to this article can be found under BioProject accession number PRJNA715468 at https://www.ncbi.nlm.nih.gov/bioproject/, an open-source online data repository hosted at the NCBI SRA BioProject database.

CONFLICTS OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGEMENTS

We would like to thank Jan Schubert, Katja Cloppenborg-Schmidt, and Olga Eitel for excellent technical assistance. We are grateful to Sarah Gaugel and Stephanie Freyher, Lübeck, for technical assistance with sample storage. We are indebted to Ana Luiza Lima, Kaan Yilmaz, and Onur Dikmen, Lübeck, for assistance with sample storage and communication with study centers during various phases of the study.

AUTHOR CONTRIBUTIONS

Conceptualization: JFB, ES, SI, CDS; **Data curation:** MB, BMH; **Formal analysis:** MB, BMH; **Funding acquisition:** JFB, ES, CDS; **Investigation:** MB, BMH; **Methodology:** JFB, ES, CDS; **Project administration:** JFB, ES, NVB, MB, CMH, CDS, MMH; **Resources:** MB, NVB, SB, PB, KD, SG, RG, MG, CG, AVG, CMH, MMH, BH, FH, BL, PJ, AP, CP, MS, LH, SV, CDS, ES; **Software:** MB, BMH; **Supervision:** JFB, ES, MB; **Validation:** JFB, MB; **Visualization:** JFB, MB, BMH; **Writing–Original draft preparation:** JFB, BMH, MB; **Writing–Review and editing:** JFB, BMH, MB, NVB, SB, PB, KD, SG, RG, MG, CG, AVG, CMH, MMH, BH, FH, BL, PJ, AP, CP, MS, LH, SV, CDS, ES

REFERENCES:

Amber KT, Murrell DF, Schmidt E, Joly P, Borradori L. Autoimmune Subepidermal Bullous
Diseases of the Skin and Mucosae: Clinical Features, Diagnosis, and Management. Clin Rev
Allergy Immunol 2018;54:26–51. https://doi.org/10.1007/s12016-017-8633-4.

van Beek N, Weidinger A, Schneider SW, Kleinheinz A, Gläser R, Holtsche MM, et al.
Incidence of pemphigoid diseases in Northern Germany in 2016 – first data from the
Schleswig-Holstein Registry of Autoimmune Bullous Diseases. Journal of the European
Academy of Dermatology and Venereology 2021;n/a. https://doi.org/10.1111/jdv.17107.

Belheouane M, Vallier M, Čepić A, Chung CJ, Ibrahim S, Baines JF. Assessing similarities and
disparities in the skin microbiota between wild and laboratory populations of house mice.
ISME J 2020. https://doi.org/10.1038/s41396-020-0690-7.

Blaser MJ, Dominguez-Bello MG, Contreras M, Magris M, Hidalgo G, Estrada I, et al. Distinct
cutaneous bacterial assemblages in a sampling of South American Amerindians and US
residents. ISME J 2013;7:85–95. https://doi.org/10.1038/ismej.2012.81.

Cáceres MD, Legendre P. Associations between species and groups of sites: indices and
statistical inference. Ecology 2009;90:3566–74. https://doi.org/10.1890/08-1823.1.

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods 2016;13:581–3.
https://doi.org/10.1038/nmeth.3869.

Chen YE, Tsao H. The skin microbiome: current perspectives and future challenges. J Am Acad
Dermatol 2013;69:143-155.e3. https://doi.org/10.1016/j.jaad.2013.01.016.

Clausen M-L, Agner T, Lilje B, Edslev SM, Johannesen TB, Andersen PS. Association of
Disease Severity With Skin Microbiome and Filaggrin Gene Mutations in Adult Atopic

Dermatitis. JAMA Dermatol 2018;154:293–300.

https://doi.org/10.1001/jamadermatol.2017.5440.

Cole JR, Chai B, Farris RJ, Wang Q, Kulam SA, McGarrell DM, et al. The Ribosomal Database

Project (RDP-II): sequences and tools for high-throughput rRNA analysis. Nucleic Acids Res

2005;33:D294-296. https://doi.org/10.1093/nar/gki038.

Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical identification

and removal of contaminant sequences in marker-gene and metagenomics data. Microbiome

2018;6:226. https://doi.org/10.1186/s40168-018-0605-2.

Gribbon EM, Cunliffe WJ, Holland KT. Interaction of Propionibacterium acnes with skin lipids

in vitro. Microbiology, 1993;139:1745–51. https://doi.org/10.1099/00221287-139-8-1745.

Hospodsky D, Pickering AJ, Julian TR, Miller D, Gorthala S, Boehm AB, et al. Hand bacterial

communities vary across two different human populations. Microbiology 2014;160:1144–52.

https://doi.org/10.1099/mic.0.075390-0.

Huang S, Haiminen N, Carrieri A-P, Hu R, Jiang L, Parida L, et al. Human Skin, Oral, and Gut

Microbiomes Predict Chronological Age. MSystems 2020;5:e00630-19,

/msystems/5/1/msys.00630-19.atom. https://doi.org/10.1128/mSystems.00630-19.

Joly P, Baricault S, Sparsa A, Bernard P, Bédane C, Duvert-Lehembre S, et al. Incidence and

Mortality of Bullous Pemphigoid in France. Journal of Investigative Dermatology

2012;132:1998–2004. https://doi.org/10.1038/jid.2012.35.

Karstens L, Asquith M, Davin S, Fair D, Gregory WT, Wolfe AJ, et al. Controlling for

Contaminants in Low-Biomass 16S rRNA Gene Sequencing Experiments. MSystems

2019;4:e00290-19, /msystems/4/4/mSys.00290-19.atom.

https://doi.org/10.1128/mSystems.00290-19.

Kong HH, Oh J, Deming C, Conlan S, Grice EA, Beatson MA, et al. Temporal shifts in the skin microbiome associated with disease flares and treatment in children with atopic dermatitis. Genome Res 2012;22:850–9. https://doi.org/10.1101/gr.131029.111.

Kong HH, Segre JA. The Molecular Revolution in Cutaneous Biology: Investigating the Skin Microbiome. Journal of Investigative Dermatology 2017;137:e119–22. https://doi.org/10.1016/j.jid.2016.07.045.

Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. Appl Environ Microbiol 2013;79:5112–20. https://doi.org/10.1128/AEM.01043-13.

Liaw A, Wiener M. Classification and Regression by randomForest. Newsl R Project 2002;2:5.

McLaughlin J, Watterson S, Layton AM, Bjourson AJ, Barnard E, McDowell A. Propionibacterium acnes and Acne Vulgaris: New Insights from the Integration of Population Genetic, Multi-Omic, Biochemical and Host-Microbe Studies. Microorganisms 2019;7. https://doi.org/10.3390/microorganisms7050128.

McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. PLOS ONE 2013;8:e61217. https://doi.org/10.1371/journal.pone.0061217.

Miodovnik M, Künstner A, Langan EA, Zillikens D, Gläser R, Sprecher E, et al. A distinct cutaneous microbiota profile in autoimmune bullous disease patients. Exp Dermatol 2017. https://doi.org/10.1111/exd.13357.

Nakamura K, O'Neill AM, Williams MR, Cau L, Nakatsuji T, Horswill AR, et al. Short chain fatty acids produced by Cutibacterium acnes inhibit biofilm formation by Staphylococcus epidermidis. Scientific Reports 2020;10:21237. https://doi.org/10.1038/s41598-020-77790-9.

Nakatsuji T, Chen TH, Narala S, Chun KA, Two AM, Yun T, et al. Antimicrobials from human skin commensal bacteria protect against Staphylococcus aureus and are deficient in atopic dermatitis. Science Translational Medicine 2017;9. https://doi.org/10.1126/scitranslmed.aah4680.

Nakatsuji T, Hata TR, Tong Y, Cheng JY, Shafiq F, Butcher AM, et al. Development of a human skin commensal microbe for bacteriotherapy of atopic dermatitis and use in a phase 1 randomized clinical trial. Nature Medicine 2021:1–10. https://doi.org/10.1038/s41591-021-01256-2.

Oh J, Byrd AL, Deming C, Conlan S, Kong HH, Segre JA. Biogeography and individuality shape function in the human skin metagenome. Nature 2014;514:59–64. https://doi.org/10.1038/nature13786.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. Community Ecology Package. 2005.

O'Neill AM, Gallo RL. Host-microbiome interactions and recent progress into understanding the biology of acne vulgaris. Microbiome 2018;6:177. https://doi.org/10.1186/s40168-018-0558-5.

Quan C, Chen X-Y, Li X, Xue F, Chen L-H, Liu N, et al. Psoriatic lesions are characterized by higher bacterial load and imbalance between Cutibacterium and Corynebacterium. Journal of the American Academy of Dermatology 2020;82:955–61. https://doi.org/10.1016/j.jaad.2019.06.024.

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res 2013;41:D590-596. https://doi.org/10.1093/nar/gks1219.

Rehman A, Rausch P, Wang J, Skieceviciene J, Kiudelis G, Bhagalia K, et al. Geographical patterns of the standing and active human gut microbiome in health and IBD. Gut 2016;65:238–48. https://doi.org/10.1136/gutjnl-2014-308341.

Schmidt E, Zillikens D. Pemphigoid diseases. The Lancet 2013;381:320–32. https://doi.org/10.1016/S0140-6736(12)61140-4.

Scholz CFP, Kilian M. The natural history of cutaneous propionibacteria, and reclassification of selected species within the genus Propionibacterium to the proposed novel genera Acidipropionibacterium gen. nov., Cutibacterium gen. nov. and Pseudopropionibacterium gen. nov. Int J Syst Evol Microbiol 2016;66:4422–32. https://doi.org/10.1099/ijsem.0.001367.

Shu M, Wang Y, Yu J, Kuo S, Coda A, Jiang Y, et al. Fermentation of Propionibacterium acnes, a Commensal Bacterium in the Human Skin Microbiome, as Skin Probiotics against Methicillin-Resistant Staphylococcus aureus. PLOS ONE 2013;8:e55380. https://doi.org/10.1371/journal.pone.0055380.

Srinivas G, Möller S, Wang J, Künzel S, Zillikens D, Baines JF, et al. Genome-wide mapping of gene–microbiota interactions in susceptibility to autoimmune skin blistering. Nat Commun 2013;4:1–7. https://doi.org/10.1038/ncomms3462.

Stevens NE, Cowin AJ, Kopecki Z. Skin Barrier and Autoimmunity—Mechanisms and Novel Therapeutic Approaches for Autoimmune Blistering Diseases of the Skin. Front Immunol 2019;10. https://doi.org/10.3389/fimmu.2019.01089.

Wang Q, Garrity GM, Tiedje JM, Cole JR. Naïve Bayesian Classifier for Rapid Assignment of

rRNA Sequences into the New Bacterial Taxonomy. Appl Environ Microbiol 2007;73:5261–

7. https://doi.org/10.1128/AEM.00062-07.

Weyrich LS, Farrer AG, Eisenhofer R, Arriola LA, Young J, Selway CA, et al. Laboratory

contamination over time during low-biomass sample analysis. Molecular Ecology Resources

2019;19:982–96. https://doi.org/10.1111/1755-0998.13011.

Yerushalmi M, Elalouf O, Anderson M, Chandran V. The skin microbiome in psoriatic disease:

A systematic review and critical appraisal | Elsevier Enhanced Reader. Journal of

Translational Autoimmunity 2019;2. https://doi.org/10.1016/j.jtauto.2019.100009.