Illness duration and symptom profile in a large cohort of

symptomatic UK school-aged children tested for SARS-

CoV-2

Erika Molteni PhD[1]*, Carole H. Sudre PhD[1,2,3]*, Liane S. Canas PhD[1], Sunil S. Bhopal PhD[4],

Robert C. Hughes MPH MB ChB[5], Michela Antonelli PhD[1], Benjamin Murray MSc[1], Kerstin

Kläser MSc[1], Eric Kerfoot PhD[1], Liyuan Chen MSc[1], Jie Deng PhD[1], Christina Hu BA[6],

Somesh Selvachandran MEng[6], Kenneth Read BSc[6], Joan Capdevila Pujol PhD[6], Alexander

Hammers PhD[1,7], Tim D. Spector PhD[8], Sebastien Ourselin PhD[1], Claire J. Steves PhD[8],

Marc Modat PhD[1], Michael Absoud PhD[9,10], Emma L. Duncan PhD[8,10].

1. School of Biomedical Engineering & Imaging Sciences, King's College London,
London, UK.
2. MRC Unit for Lifelong Health and Ageing, Department of Population Health Sciences,
University College London, London, UK.
3. Centre for Medical Image Computing, Department of Computer Science, University
College London, London, UK.
4. Population Health Sciences Institute, Faculty of Medical Sciences, Newcastle
University, Newcastle upon Tyne, Tyne and Wear, UK.
5. Department of Population Health, Faculty of Epidemiology & Population Health,
London School of Hygiene & Tropical Medicine, Keppel Street, London, UK.
6. Zoe Global, London, UK.
7. King's College London & Guy's and St Thomas' PET Centre, London, UK.
8. Department of Twin Research and Genetic Epidemiology, King's College London,
London, UK.
9. Children's Neurosciences, Evelina London Children' Hospital, St Thomas' Hospital,
King's Health Partners, Academic Health Science Centre, London, UK.
10. Department of Women and Children's Health, Faculty of Life Sciences and Medicine,
School of Life Course Sciences, King's College London, London, UK.
* Equal contribution
31
32
Corresponding Author:
Emma L. Duncan, MBBS, FRACP, FRCP, PhD

Department of Twin Research and Genetic Epidemiology, and Department of Women and

Children's Health

King's College London,

WC2R 2LS, Strand, London, UK.

emma.duncan@kcl.ac.uk

Key words

Long COVID; ongoing symptomatic COVID-19; post-COVID-19 syndrome; SARS-CoV-2

prolonged infection; school-children; paediatric COVID-19.

Word count: 5975

Number of Tables: 2

Number of Figures: 4

Number of Supplementary Tables: 4

Number of Supplementary Figures: 6

Research in context

Evidence before this study: SARS-CoV-2 in children is usually asymptomatic or manifests as a mild illness of short duration. However, anecdotal cases have raised concerns of prolonged illness in children, with no clear resolution of symptoms several weeks after onset, as is observed in some adults. How common this might be in children, the clinical features of such prolonged illness in children, and how it might compare with illnesses from other respiratory viruses (and with general population prevalence of these symptoms) is unclear.

Added value of this study: We provide systematic description of COVID-19 in school-aged children. Our data, collected in a digital surveillance platform through one of the largest citizen science initiatives, show that long illness duration after SARS-CoV-2 infection in school-aged children does occur, but is uncommon. Only a small proportion of children have illness beyond four weeks; and the symptom burden in these children usually decreases over time. Most children resolve all symptoms by eight weeks, providing reassurance about long term outcomes. Importantly, symptom burden does not outnumber that observed in long illness due to causes other than SARS-CoV-2 infection.

Implications of all the available evidence:
Our results confirm that COVID-19 illness in children and adolescents is resolved in the community, after short duration, low symptom burden and with no need of hospitalization in most cases. Our findings also highlight that allocation of appropriate resources will be necessary for any child with prolonged illness, whether due to SARS-CoV-2 infection or other illness. Our study provides timely and critical data to inform discussions around the impact and implications of the pandemic on UK paediatric healthcare resource allocation.

Abstract

Background

In children, SARS-CoV-2 is usually asymptomatic or causes a mild illness of short duration. Persistent illness has been reported; however, its prevalence and characteristics are unclear. We aimed to determine illness duration and characteristics in symptomatic UK school-aged children tested for SARS-CoV-2 using data from the COVID Symptom Study, the largest citizen participatory epidemiological study to date.

Methods

Data from 258,790 children aged 5-17 years were reported by an adult proxy between 24 March 2020 and 22 February 2021. Illness duration and symptom profiles were analysed for all children testing positive for SARS-CoV-2 for whom illness duration could be determined, considered overall and within younger (5-11 years) and older (12-17 years) age groups. Data from symptomatic children testing negative for SARS-CoV-2, matched 1:1 for age, gender, and week of testing, were also assessed.

Findings

1,734 children (588 younger children, 1,146 older children) had a positive SARS-CoV-2 test result and calculable duration of illness within the study time frame. The commonest symptoms were headache (62.2%) and fatigue (55.0%). Median illness duration was six days (vs. three days in children testing negative), and was positively associated with age (rs 0.19, p<1.e-4) with median duration of seven days in older vs. five days in younger children.

Seventy-seven (4.4%) children had illness duration ≥28 days (LC28), more commonly experienced by older vs. younger children (59 (5.1%) vs. 18 (3.1%), p=0.046). The commonest symptoms experienced by these children were fatigue (84%), headache (80%) and anosmia (80%); however, by day 28 the median symptom burden was two. Only 25 (1.8%) of 1,379 children experienced symptoms for ≥56 days. Few children (15 children, 0.9%) in the negatively-tested cohort experienced prolonged symptom duration; however, these children experienced greater symptom burden (both throughout their illness and at day 28) than children positive for SARS-CoV-2.

Interpretation

Some children with COVID-19 experience prolonged illness duration. Reassuringly, symptom burden in these children did not increase with time, and most recovered by day 56. Some children who tested negative for SARS-CoV-2 also had persistent and burdensome

114   illness. A holistic approach for all children with persistent illness during the pandemic is

115   required.

118   **Funding**

Introduction

To date, the COVID-19 pandemic has resulted in >120 million cases of infection and 2.5 million deaths globally,[1] with widespread health, economic, and social chaos. In adults, severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2) causes a predominantly respiratory illness[2] of median duration 11 days.[3] In contrast, children with SARS-CoV-2 infection are often asymptomatic (43%-68%[4]) or have relatively mild symptoms, most commonly cough and fever;[4–9] and life-threatening illness or death is rare. 1,575 children (aged 0-17 years) hospitalised in England from 19 March 2020 to 3 March 2021 tested positive for SARS-CoV-2, noting that COVID-19 was not necessarily the reason for hospitalisation,[10] and 29 deaths due to COVID-19 were reported in children and young people across the UK from 1 March 2020 to 29 January 2021 (0-9 years: seven deaths; 10-19 years: 22 deaths, 0.19% and 1.98% respectively of estimated all-cause deaths during this time period[11]). The pandemic has also seen emergence of a new rare condition, multisystem inflammatory syndrome in children (MIS-C), typically presenting 2-4 weeks after acute SARS-CoV-2 infection.[12]

Some adults with SARS-CoV-2 infection experience prolonged duration of illness, known variably as Long COVID (LC),[3,13] ongoing symptomatic COVID-19, post-COVID-19 syndrome or post-COVID condition.[14,15] Data from the King's College London (KCL) COVID Symptom Study (CSS),[16] currently the largest citizen participatory epidemiological study in the UK with over >4.5 million UK participants, showed that 13.3% adults with a positive SARS-CoV-2 test had symptoms for ≥4 weeks (LC28) and 4.5% for ≥8 weeks (LC56).[3] Predictors of LC28 included older age, female gender, and higher symptom burden in the first week of illness.[3] Whether some children may also experience prolonged illness duration after infection with SARS-CoV-2[15] and, if so, how this compares with illness after infection with other respiratory viruses, is currently unclear.

In September 2020, coinciding with full re-opening of primary and secondary schools in the UK, governance for CSS data usage was extended to allow analysis of data from children (i.e., individuals aged <18 years). The UK subsequently experienced further waves of the pandemic (up to 30,000 new cases per day in November 2020 and 60,000 new cases per day in January 2021),[1,17] with 29 December 2020 the peak date for new specimens testing positive over the entire pandemic to date.[17] During this time, there was widespread testing availability for individuals with key symptoms of fever, cough and anosmia, in contrast to the

157 very limited access during the first wave, when testing was mostly restricted to individuals
158 presenting to hospital.[18,19]
159
Stay-at-home directives and school closures during these later waves resulted in unusually
low incidences of commonly circulating viruses such as influenza, adenovirus, and
respiratory syncytial virus over the 2020-21 UK winter, both generally and in school-aged
children specifically.[20] However, overlap in symptomatology meant many individuals (both
adults and children) with respiratory illnesses other than COVID-19 were tested for SARS-
CoV-2.
166
167 Here we report overall illness duration, individual symptom prevalence and duration, and
168 symptom burden in UK school-aged children testing positive for SARS-CoV-2, whose data
169 were logged through the KCL CSS. We provide similar data for symptomatic children tested
170 for SARS-CoV-2 but who were negative, during the same period. Lastly, we present the
171 prevalence and characteristics of long COVID in children.
172

Methods

Data were acquired within the COVID Symptom Study (CSS), through a mobile application
launched jointly by Zoe Global Ltd. and KCL on 24 March 2020.[16] Briefly, individuals are
prompted to report through a smartphone application and provide daily updates of their
health status, symptomatology, any SARS-CoV-2 testing, vaccination, and health care
access. Symptom assessment includes both direct questions for a list of symptoms (yes/no
or other defined options, listed in Supplementary Table 1), and free-text entry.[16] Adult
contributors can also report as a proxy for other persons (children, spouses, elderly relatives,
etc.). The relationship between the contributor and proxy-reported individual(s) is not
solicited and there is no data linkage between the contributor and proxy-reported
individual(s). Children aged 16-17 years can use the app directly as independent
contributors or be proxy-reported by an adult.
185
UK data from school-aged children (aged 5-17 years) were available from the launch of the
app to 22 February 2021, which latter date corresponds to eight weeks after the peak SARS-
CoV-2 positive specimen date in the UK.[17] Data were considered from launch of the app,
and from 1 September 2020 (i.e., return-to-school). The cohort was analysed overall and
within two age groups: younger children, aged 5-11 years, corresponding to primary school-
aged children in the UK; and older children, aged 12-17 years, corresponding to secondary

school-aged children in the UK (acknowledging that not all older teenagers attend school).

Preschool-aged children (≤4 years) were not included in the current study.

Aligning with our previous publications in adults,[3,16] children were considered symptomatic of

COVID-19 if they were reported by proxy as having any of a specific list of symptoms

associated with COVID-19[16] (Supplementary Table 1), mainly informed by adult

epidemiology,[21] with first symptom(s) presenting within a timeframe of 1 week before and 2

weeks after confirmation of infection (either a SARS-CoV-2 polymerase chain reaction or a

lateral flow test). Illness duration was calculated as time from first symptom(s) reported

within the requisite time frame (having been previously asymptomatic) until time of recovery

(defined as return to asymptomatic state or, if proxy-reporting ceased prior to logging of an

asymptomatic report, time of final report). Individuals who had returned to asymptomatic

state but for whom another symptomatic report was logged within one week of the last

symptomatic report were considered as still being unwell from their initial presentation (i.e.,

allowing for waxing/waning of illness); calculation of illness duration incorporated these short

periods of asymptomatic status. Individuals with reporting gaps longer than one week

between symptomatic reports were excluded. Individual symptom prevalence and duration

were assessed, with individual symptom duration calculated as time between the first and

last report for that symptom. Symptom burden was calculated as the number of different

symptoms reported at least once over a defined timeframe (during the first week, the first 28

days, at ≥28 days until end of illness, and over the entire duration of illness). Consistent with

our previous study in adults,[3] we termed illness with symptoms lasting longer than 28 days

as LC28; and longer than 56 days, LC56. Thus, by virtue of data census dates LC28 could

only be determined in proxy-reported children whose symptoms commenced on or before 24

January 2021, and for LC56 on or before 29 Dec 2020 (noting the peak positive specimen

date in the UK was 29 December 2020[17]).

Symptom profiles were also assessed in children with COVID-19 who presented for hospital-

based care (presenting to the accident and emergency department or admitted to hospital),

where presentation to hospital followed development of symptoms (i.e., when the positive

SARS-CoV-2 test result was not a secondary finding in a hospitalised child).

Several additional direct symptom questions were added to the app on 4 November 2020

(Supplementary Table 2), some of which overlapped with existing questions (e.g., for

cutaneous manifestations, questions on "rash" and "sensitive skin" were added to existing

questions on "red welts", "blisters", and "sores"). These additional questions and resultant

data are presented in Supplementary Table 2 and Supplementary Figure 1 respectively but
were not included in illness duration or symptom burden calculations for the main analysis.
230
Free text reporting was also possible, across the entire period. Free text data were divided
into themes using frequency of descriptive words; each item within the themes were
subsequently independently scrutinised by two clinicians (MA, ELD) to ensure appropriate
categorisation, and individuals reporting free text symptoms within those themes were then
counted. Free text data are reported here as descriptive statistics (Supplementary Table 3)
and were not included in calculating illness duration or symptom burden. Free text was also
searched for specific neurological terms of interest (e.g., weakness, difficulties with balance,
paralysis, seizures, fits, convulsions, paroxysms, tics) and symptoms potentially affecting
attention, behaviour, learning, and school performance (e.g., anxiety, irritability).[22] Symptoms
already assessed by direct question (Supplementary Table 1) were excluded from the free
text search, to avoid duplication.
242
Symptom profile and duration were also assessed in children who were tested but had a
negative result for SARS-CoV-2. The same parameters regarding test timing relative to
symptom onset were applied as for children with a positive test result. Illness duration and
symptom profile were determined in a randomly selected control cohort of these children
(matched 1:1 for age, gender, and week of testing), and compared to children with a positive
test.
249
250 Proxy-reporting density (defined as the number of episodes of proxy-reporting over the
251 duration of illness) and persistence (defined as proxy-reporting until return to healthy state)
252 were also determined.
253
254 Prevalence data for common winter circulating viruses were obtained from the Public Health
255 England weekly national influenza and COVID-19 surveillance report.[20]
256
Data are presented using descriptive statistics. Results are presented as median with
interquartile ranges. Due to rarity (with most percentages less than 5%) confidence intervals
were calculated using Poisson distribution. Comparisons of data between groups used
Wilcoxon signed-rank test or Chi-squared test/Fisher's exact test, as appropriate. Spearman
correlation testing (which does not require Gaussian distribution of the data) was used to
assess correlation of illness duration with age.
263

Ethics approval for this study was granted by the KCL Ethics Committee REMAS ID 18210,

review reference LRS-19/20-18210 and all participants (here, the proxy-reporting adult)

provided consent.

Results

Across the UK, 258,790 children aged 5-17 years were reported by proxy between 24 March

2020 and 22 February 2021, with a positive SARS-CoV-2 test result reported in 6,975

children. Among those, 1,912 (666 younger and 1,246 older) children had a calculable

duration of illness with requisite logging frequency (at least once weekly). As only 36 children

had illness onset prior to 1st September 2020, and as there was very limited community

access to testing early in the UK pandemic experience,[19] analyses of illness duration and

symptom profiles were restricted to children with illness onset after 1 September 2020.

Further, illness onset had to commence on or before 24 January 2021 to allow sufficient time

for symptom duration ≥4 weeks to be evident, as per LC28 definition.[3] Thus, overall, 1,734

children (588 younger children, 1,146 older children) had a calculable illness duration within

the requisite time frame. Similarly, 1,379 individuals (445 younger; 934 older children) had

symptoms commencing on or before 29 December 2020, allowing sufficient time for

symptom duration ≥8 weeks to be evident, as per LC56 definition.[3] A flow chart showing

inclusion/exclusion steps is shown in Figure 1.

Among 16- and 17-year-old individuals, 29,047 contributed self-logged data (447 reporting

testing positive) compared to 32,271 reported by proxy (1,197 reported as testing positive).

Illness duration could only be calculated in 10 self-logged 16- and 17-year-olds, compared

with 381 proxy-logged 16- and 17-year-olds; and self-reporting and proxy-reporting of the

same young person could not be excluded. Thus, only proxy-reported data are presented

here.

Illness in children who tested positive for SARS-CoV-2.

The median illness duration in children with COVID-19 was six days [IQR 3;11] (Table 1).

Illness duration was significantly shorter in younger compared with older children (five [IQR

2;9] vs. seven days [IQR 3;12]) (Mann-Whitney U test p<1.e-5); and age correlated strongly

with illness duration (r_s 0.19, p<1.e-4).

Individual symptom prevalence and duration are shown in Table 2, Figure 2 and Figure 3.

Overall, the most reported symptoms were headache (62.2% overall: 55.1% younger

children, 65.9% older children) and fatigue (55.0% overall: 43.9% younger children and

60.7% older children). Subsequent symptoms ranked by frequency were fever (43.7%), sore

throat (36.2%), abdominal pain (27.7%), and persistent cough (24.7%) in younger children;

and in older children sore throat (51.0%), anosmia (48.3%), fever (34.6%), and persistent

cough (26.0%). During the first week of illness, median symptom burden was three [IQR 2;6]

overall (three [IQR 2;5] symptoms in younger children, four [IQR 2;6] in older children).

Sixteen younger children and 21 older children who tested positive for SARS-CoV-2

subsequently attended for hospital care. The symptom profiles of children attending hospital

compared with children managed in the community are shown in Supplementary Figure 2.

No formal statistical comparisons were undertaken between hospital and community cases,

given the low numbers of hospital attendees.

Long illness duration in children who tested positive for SARS-CoV-2.

Overall, 77 (4.4% [95% CI 3.5-5.5]) of 1,734 children (18 younger children, 59 older children)

had symptoms for ≥28 days, meeting the definition of LC28. The median symptom burden in

these children was six symptoms [IQR 4;8] reported at least once during the first week of

illness, and eight symptoms [IQR 6;9] reported at least once over the duration of their illness.

However, by day 28 median symptom burden was low, at two [IQR 1;4] (younger children:

three [IQR 1;4]; in older children, one [IQR 1;3]). The commonest symptoms experienced by

children with LC28 over the duration of their illness were fatigue (experienced by 84.4% of

children), headache (77.9%), anosmia (77.9%) and sore throat (74.0%). Figure 4 is a heat

map of symptom profile and progression over the first 28 days in children with LC28.

Twenty-five (1.8% [95% CI 1.2-2.7]) of 1,379 children had symptoms for ≥56 days, meeting

the definition of LC56.[3] The median burden of symptoms in these children was six [IQR 4;8]

symptoms reported at least once during the first week, and eight [IQR 6;10] symptoms

reported at least once over the duration of their illness. The commonest symptoms

experienced by children with LC56 over the duration of their illness were anosmia (84.0%),

headache (80.0%), sore throat (80.0%) and fatigue (76.0%).

Consistent with correlation of illness duration with age across the cohort overall, older

children were more likely to manifest symptoms ≥28 days, compared with younger children

(59 (5.1%) of 1146 older children vs. 18 (3.1%) of 588 younger children, Chi-squared two-tail

test p=0.046). However, this difference was no longer significant in the smaller number of

children with illness duration ≥ 56 days (6 (1.3%) of 445 younger children vs. 19 (2.0%) of
934 older children, Fisher's exact test p=0.52).
337
338   ***Symptom Reporting from Additional Questions (added 4 November 2020) and Free***
339   ***Text Analysis***
340
341   Considering the additional questions added to the CSS app on 4 November 2020,
342   rhinorrhoea was the most reported symptom (45.8% younger children; 53.5% older children)
343   followed by sneezing (36.3% younger children; 36.6% older children) (Supplementary Figure
344   1), with similar prevalence when considered over the entire illness duration in children with
345   LC28 overall (51.4% rhinorrhoea, 48.9% sneezing).
346
347   ***Potential neurological symptoms***
348
As mentioned, headache and fatigue were the commonest symptoms in children testing
positive for SARS-CoV-2, overall and in each age group. Dizziness (without differentiation
between light-headedness and vertigo) was reported in 84 (14.3%) younger and 300
(26.2%) older children, with median duration of two [IQR 1;4] and two [IQR 1;5] days
respectively. Symptoms consistent with confusion (encompassing confusion, disorientation,
and drowsiness) were reported in 15 (2.6%) younger and 81 (7.1%) older children (median
duration 2 day [IQR 1;2] and one day [IQR 1;3] respectively). No formal statistical
comparisons of these symptoms in children with shorter or longer illness duration were made
given the low prevalence of these symptoms in children with LC28 (Figure 4).
358
With respect to questions added to the app on 4 November 2020 (Supplementary Table 2),
and with the sample size reduced because of the reduced timeframe, 'brain fog' was
reported for 44 (8.7%) of 507 younger children (median duration one day [IQR 1;4]) and 188
(20.2%) of 932 older children (median duration two days [IQR 1;5]); and low mood was
reported for 40 (8.0%) younger children (median duration two days [IQR 1;3.5]), and 145
(15.6%) older children (median duration two days [IQR 1;4.25]) (Supplementary Figure 1).
365
Free text searching for specific neurologic symptoms disclosed very few reports of weakness
(two children) or tics (one child). No severe neurological manifestations (paralysis, ataxia,
epileptic seizures, fits, convulsions, paroxysms) were reported (Supplementary Table 3).
Irritability (three children), emotional difficulties (two children), and behavioural difficulties
(one child) were reported rarely; impaired attention, impaired concentration, and anxiety
were not reported.

Illness in symptomatic children who tested negative for SARS-CoV-2.

15,597 symptomatic children (8,761 younger children, 6,836 older children), tested for

SARS-CoV-2 infection but with a negative result, with logging at least once weekly and with

calculable illness duration within the requisite time frame, were proxy-reported.

Demographic details of the randomly selected matched control sample are shown in Table 1.

The median illness duration in these children was three days [IQR 2;7] (younger children

three days [IQR 2;7]; older children four days [IQR 2;7]), significantly shorter than for

children with a positive SARS-CoV-2 test (Wilcoxon signed rank test p<1.e-5). The individual

symptom profile is shown in Supplementary Figure 4. The most frequently reported

symptoms over the entire illness duration were sore throat (46.6%), headache (38.8%), fever

(30.4%), fatigue (26.9%), and abdominal pain (24.7%) in younger children; and sore throat

(60.6%), headache (48.8%), fatigue (37.2%), fever (20.4%) and persistent cough (20.6%) in

older children.

Long illness duration in children who tested negative for SARS-CoV-2.

Few children with a negative test for SARS-CoV-2 had illness duration for ≥28 days (15 of

1,734 children; 0.9%; 95% CI [0.5-1.4]). This was significantly fewer than children with a

positive test, considered overall (Chi-squared 2-tail test p<1.e-10) and by age group (in

younger children, three (0.5%) vs. 18 (3.1%); Fisher's exact test: p=0.001; in older children,

12 (1.1%) vs. 59 (5.1%); Fisher's exact test: p<1.e-8). The symptom profile of these children

over the first 28 days is shown in Supplementary Figure 4. However, older children with a

negative test for SARS-CoV-2 whose illness duration was ≥28 days had a greater symptom

burden than was experienced by children with LC28, both at ≥28 days (Supplementary

Figure 5), and over their entire illness duration (Supplementary Figure 6) (p=0.005 and

p=0.025 respectively, Mann-Whitney-U tests).

Proxy-reporting density

Overall, proxy-reporting density (number of logging episodes/illness duration in days) was

assiduous for all children. Reporting density was significantly higher in children with a

negative test for SARS-CoV-2, compared to children with a positive test (reporting density in

children negative vs positive: in younger children: 1 [IQR 0.68;1] vs. 0.89 [IQR 0.6;1]; and in

older children 1 [IQR 0.67;1] vs. 0.8 [IQR 0.57;1]), noting the shorter illness duration in

children with a negative test.

Proxy-reporting perseverance

Children were included in this study if their illness onset corresponded temporally to testing

for SARS-CoV-2, and either a healthy report was received (1,551 (89.4%) of children with a

positive test and 1,674 (96.5%) of children with a negative test) or proxy-reporting ceased.

Amongst children with a positive SARS-CoV-2 test, logging ceased in 183 children prior to a

healthy report (22 children with LC28, 161 with [assumed] shorter illness duration). In

children with LC28, a healthy report was received in 71.4% [55/77]. However, the remaining

22 children had already had symptoms logged for more than 28 days and thus fulfil the

definition of LC28. In children with [assumed] shorter illness duration, a healthy report was

received for 90.3% [1496/1657]. For the remaining 161 children, median symptom burden at

last report was two [IQR 1;3]; and proxy-reporting usually ceased early in illness (logging

cessation rates: 11.3% of children with illness duration <10 days; 5.5% ≥10 days). Thus, for

children for whom proxy-reporting ceased prior to logging of a healthy report, we have

assumed that proxy-reporting cessation coincided with illness resolution (i.e., that adults

stopped proxy-reporting because the child had recovered) and illness duration was

calculated assuming last day of logging corresponded to last day of illness.

We considered the impact of this assumption. Excluding all children with a positive SARS-

CoV-2 test for whom a healthy report was not logged, median duration of illness in the

remaining 1,551 was unchanged (6 days [IQR 3;11]) with prevalence of LC28 3.5%

[55/1551], within the confidence intervals for LC28 using data from the entire cohort.

Excluding only the 161 children with [assumed] short symptoms but including all children

who had fulfilled the definition of LC28 (regardless of receipt of healthy report), prevalence of

LC28 was 4.9% (77/1,573 children), again within the confidence intervals for LC28 using the

entire cohort.

In children with a negative SARS-CoV-2 test, logging ceased in 60 children prior to a healthy

report (four of 15 children with long illness duration, 56 of 1,719 children with [assumed]

short illness duration). Proxy-reporting until logging of a healthy report was higher in children

with a negative test, compared with children with a positive test (Chi-squared 2-tail test

p<1.e-15).

Data governance did not allow assessment of familial relationships between proxy-reported

individuals and the contributor; thus, we cannot comment whether concurrent illness of the

contributor or other family members affected proxy-reporting.

Discussion

In this large study of UK school-aged children recruited via the CSS study app, we have

shown that symptomatic infection with SARS-CoV-2 in children is usually short, with a

median duration of illness of six days compared to 11 days in adults,[3] and with low symptom

burden (Figure 3). Prolonged illness duration can occur but is infrequent (4.4% with

symptoms ≥28 days; 1.8% with symptoms ≥56 days), considerably lower than observed in

adults using the same disease definitions (prevalence in adults of LC28: 13.3% and of LC56:

4.5%).[3] Age is a risk factor for longer symptom duration (both overall, and specifically for

illness duration ≥28 days), consistent with our previous findings in adults.[3]

Similar to adults,[3] the commonest symptoms in children with COVID-19 were headache

(62.2%) and fatigue (55.0%) (Table 2, Figure 2). A previous meta-analysis of studies in

children with COVID-19, which included community-based and hospitalized children,

identified fever (47%) and cough (42%) as the commonest symptoms.[23] However, headache

and fatigue were only assessed in half of contributing studies. In our (almost entirely

community-managed) cohort, prevalence of fever was 37.7%, and persistent cough 25.5%.

Anosmia (here, both anosmia and dysosmia) was also common in our cohort (39.6% overall:

22.4% of younger children, 48.3% of older children). Certainly, for older children this

symptom was reported more often than was observed in a previous small study of

adolescents (aged 10-19 years) with 'mild to moderate' COVID-19, ascertained from a single

centre, in whom prevalence of anosmia was 24.1%,[24] noting that anosmia was one of the

core symptoms determining access to testing in the UK during our study period.

In children with symptoms ≥28 days, symptom burden was low by day 28 (median symptom

burden of two). However, fatigue was a symptom at some stage in almost all (84.4%) of

these children (Figure 4, Supplementary Figure 6). In adults, fatigue has been reported in

many studies as the commonest symptom of long COVID, although prevalence varies. In our

previous study in adults,[3] fatigue was almost universal in LC28 (97.7% of cases experienced

fatigue during their illness). Other studies have also reported persistence of fatigue after

acute infection in adults (e.g., 53.1% at 60 days[25]; and 52.3% at 10 weeks[26]).

Considering the prevalence and persistence of headache and fatigue within the context of
usual prevalence of these symptoms within the general paediatric population is difficult, as
few large-scale epidemiological studies have been undertaken in children to provide
normative population data. Considering headache, a systematic review of headache
comprising 38 unselected population-based studies in individuals under 20 years reported
that around 60% of children are "prone to headache".[27] A random sample of 2,165 school
children from Aberdeen aged 5-15 years found that 1,166 (66%) had headaches over the
previous year, which in 391 (22%) children were of sufficient severity to stop normal
activities.[28] Considering fatigue, a study of 2,936 children found that 129 (4.4%) had "more
than a few days of disabling fatigue".[29] Prevalence of chronic fatigue syndrome (defined here
as "disabling fatigue lasting >3 months with one additional symptom, where no other cause
for the fatigue could be established") was 1% in a study of 11-16-year-old children from three
secondary schools in England.[30] Prevalence of "chronic disabling fatigue" reported by the
Avon Longitudinal Study of Parents and Children (defined here as fatigue lasting >6 months
associated with absence from full-time school or that had prevented the child from taking
part in activities 'quite a lot' or 'a great deal', and not due to excess of physical activity) was
1.5% and 2.2% at 13- and 16- years respectively.[31] Considering fatigue after viral infection
specifically, and choosing one of the younger published cohorts, median symptom duration
with Epstein–Barr virus in symptomatic university students (i.e., older than the cohort
reported in our study) was 10 days (mean 17, range 3-66 days), with fatigue persisting for a
median of 15.5 days.[32] These studies indicate the wide range of prevalence of these
symptoms in paediatric populations generally.
502
The paucity of contemporaneous data comparing illness duration and symptom profiles from
different viruses also complicates contextualisation of illness due to SARS-CoV-2 infection in
children. A preprint paper comparing data from 55,270 children/adolescents with COVID-19
(3,693 of whom were hospitalized) with a non-contemporaneous cohort of 1,952,693
children with influenza during 2017-19 (hospitalised numbers unclear) suggests that
dyspnoea, anosmia, and gastrointestinal tract symptoms are more common in children with
COVID-19 than in children with influenza.[33] However, symptoms and overall illness duration
were only reported as present or absent at 30 days in both groups, preventing more granular
comparisons.
512
Here, a strength of our study was our ability to compare contemporaneous illness profiles of
symptomatic children testing positive vs. negative for SARS-CoV-2, matched for age,
gender, and week of testing. Children testing positive for SARS-CoV-2 had longer median
illness duration (six days vs. three days in negatives); and were more likely to have illness

duration ≥28 days (4.4% vs. 0.9%). However, some children testing negative for SARS-CoV-

2 also had symptoms that lasted >28 days, and these children had a higher symptom count

both over the duration of their illness and at day 28 (Supplementary Figures 4, 5),

acknowledging here that our sample size is small. We considered whether some children

with long illness duration who tested negative for SARS-CoV-2 might have false negative

results. However, there is no evidence that sensitivity and specificity of testing for SARS-

CoV-2 are different in adults compared with children, with sensitivity for PCR SARS-CoV-2

tests around 95%.[34]

Additionally, the symptom profiles of children negative for SARS-CoV-2 (Supplementary

Figures 4, 5, 6) suggest some differences in their illness compared with children with positive

results. Relevantly, the prevalence of non-SARS-CoV-2 respiratory viruses (influenza A,

influenza B, parainfluenza, adenovirus, rhinovirus, and respiratory syncytial virus) was

unusually and extremely low over the UK winter of 2020/2021[35–37], with the exception of the

commonly observed rhinovirus peak coinciding with the September return-to-school.[20] With

relaxation of personal protection and social distancing measures, it is likely that these

illnesses will return to more usual (i.e., higher) levels of circulation in future UK winters. Our

data highlight that other illnesses may also have a protracted and burdensome course in

children, which will also need consideration in post-pandemic service delivery planning.

Short and long-term effects of COVID-19 on school performance and learning have been a

recent matter of concern.[38] In our cohort, neither attentional problems nor memory

complaints nor anxiety were reported. Isolated cases of low mood and/or irritability were

consistent with, if not lower than, previously reported statistics in the general school-aged

healthy population.[39] Our data do not support anecdotal reports of weakness and seizures

as common features in children with COVID-19, whether of short or longer illness duration;

and no severe neurological symptoms were reported. However, persistence of symptoms

from any illness can be associated with low mood, with adverse long-term outcomes

including school refusal and separation anxiety.[40] Although self-reporting adults were able to

participate in specific questions regarding mental health, these data could not be proxy-

reported. This limits our ability to assess mental health issues comprehensively, and any

potential impact on school performance and learning, in children during the COVID-19

pandemic whether testing positive or negative for SARS-CoV-2.

In considering how our data relates to other sources, the UK Office for National Statistics

(ONS) conducted a round of testing for SARS-CoV-2 (irrespective of symptoms) from 2 to 10

December 2020, prior to the UK peak of infection, with deliberate oversampling of schools

which had had the highest infection rates at the start of the academic year. 121 schools (41
primary, 80 secondary) in 15 local authorities were examined, with a total of 7,089 pupils.[41]
Primary school pupils and staff had a slightly lower percentage testing positive (0.94% and
0.99% respectively) for SARS-CoV-2 than secondary pupils and staff (1.22% and 1.64%).
ONS warned that their conclusions might not have general validity; and these figures do not
capture time fluctuations as the pandemic progressed. In our dataset, with data captured
until 22 February 2021 (thus including the subsequent rise in numbers over Christmas and
New Year), 6,043 of 258,790 (2.3%) of proxy-reported children were reported with a positive
test (2.3% of younger children, 3.7% of older children), noting that reporting was voluntary,
by proxy, and through a specific COVID-19 platform, and that testing was only available for
symptomatic individuals. Additionally, the CSS app user base (considered as a whole) is not
fully representative of the wider UK population, due to over-representation of female gender,
white background, and above-average socioeconomic status.[18] Further, we cannot
characterise regional variability, as geographic information was not available for many
participants.
569
We considered what proportion of the total number of school-aged children in the UK testing
positive for SARS-CoV-2 were proxy-reported to the CSS. The different countries of the UK
report data for children within different age groups and over varying time periods. In England
there were 390,866 positive tests in young people aged 5-19 years from 1 September 2020
to 24 January 2021 (5-9 years: 69,641 children; 10-19 years: 321,225 individuals).[20] In
Scotland,15,869 children aged 0-14 years, and 14,597 young people aged 15-19 years have
tested positive from the start of the pandemic to 30 March 2021.[42] In Wales there were
~28,300 cases in children and young people from the start of the pandemic to 30 March
2021.[43] In Northern Ireland 13,268 individuals aged 0-19 years from the start of the
pandemic to 30 March 2021.[44] Thus, our study represents ~1-2% of all school-aged children
with positive tests across the UK during this time, noting that many of these national figures
include young adults and preschool-aged children.
582
The ONS has also provided estimates of long COVID in children aged 2-16 years, with the
most recent data release (April 2021) suggesting that 9.8% of children aged 2-11 years and
13.0% of those aged 12-16 years experience ongoing symptoms five weeks after testing
positive for SARS-CoV-2, and 7.4% and 8.2% respectively are still reporting symptoms at 12
weeks.[45] These latest figures include a control group (defined as those who were never
symptomatic, never tested, never-self isolated, and never a contact of anyone testing
positive for SARS-CoV-2) which suggested 'baseline' rates for the same symptoms of 2%
amongst 2- to 11-year-olds and 1.7% amongst 12- to 16-year-olds. Earlier, in January 2021,

ONS had reported long COVID symptoms by age; however, updated age-specific estimates

for specific symptoms were not included in April 2021 data release. Overall ONS long

COVID prevalence estimates were adjusted downwards between January and April 2021

(e.g., from 12.9% to 9.8% in 2- to 11-year-olds).[45,46]

There is some disparity between our prevalence data for LC28 in children and those of the

ONS. At least part of the reason may be that the ONS required two consecutive

asymptomatic visits to define the end of illness. Thus, individuals with relapsing/remitting

symptoms in whom asymptomatic periods lasted longer than one week would be captured

by ONS but not by our study. Consistent with this, the ONS' sensitivity analysis of the impact

of defining symptom discontinuity to a single asymptomatic visit radically lowered their

prevalence estimates, especially for ongoing symptoms at 12 weeks, with estimates falling

from 13.7% to 0.9% (ref[47] - Table 11), much more in keeping with our results. Additionally,

ONS estimates are based on both current and recalled data, collected in the first week of

each month. In contrast, our app-based data collection was conducted in real time, with high

density and persistence of proxy-reporting of children (whether the child had tested positive

or negative for SARS-CoV-2). A recent Australian study reported 151 (of an initial cohort of

171) young children (median age three years [IQR 1;8]) who tested positive for SARS-CoV-2

and were followed for three to six months. Twelve (8%) children (median age two years)

were reported to have symptoms 3-8 weeks after initial presentation (most commonly cough

and/or fatigue), all of whom resolved by the end of the study to their baseline health.[23] This

small study concords more closely with our results. None-the-less, and with the methodology

for the ONS experimental estimates not yet published, the ONS estimates have been used

in many service provisions and government discussions, and reported widely in social and

other media.[48,49]

Our study is part of one of the largest citizen science initiatives ever in the UK, with

generation of real-time epidemiological data from over 4.5 million users across the UK. We

leveraged previously published methodologies assessing illness duration and symptom

profiling in adults, including assessment of long COVID.[3] Our data census points allowed us

to capture all children with illness duration ≥8 weeks, if they had presented before the date of

peak specimen receipt in the UK; and our restriction to individuals whose symptom

presentation concorded with test timing allowed an accurate determination of symptom

onset. Moreover, by restricting analyses to start from 1 September, we avoided bias due to

limited test availability during the first wave of the pandemic. However, despite general

availability there were still some blocks to testing - in particular, that individuals were

required at least one of a defined list of symptoms (specifically, fever, cough and anosmia),[50]

which list was largely informed by adult symptomatology and might not be sufficiently broad

to capture some common paediatric manifestations of COVID-19, (e.g., abdominal pain,

reported in 4% of paediatric cases[4] and in 27.8% of our younger children). Related to this,

the set of questions asked through the app, although refined between 24 March and 1

September, and again on 4 November 2020, was largely informed by research in adults.

Although we cannot exclude the possibility that a paediatric version might capture other

manifestations of COVID-19, the free text data did not suggest common themes emerging

unique to paediatric populations. We did not undertake a formal qualitative analysis of the

free-text data given (a) its *ad hoc* collection rather than a formal qualitative study, and (b) the

potential bias arising from introduction of additional direct symptom questions from 4

November 2020 (i.e., once a symptom was asked about directly, it was unlikely that to be

reported as free text). None-the-less, we have provided the data from free text responses,

and from questions asked after 4 November 2020 in the Supplementary Material. We

acknowledge that symptoms were reported by proxy rather than directly ascertained;

however, this is common in clinical assessment of children, particularly younger children.

Additionally, children cannot directly consent to research participation. We also acknowledge

the possibility that older children may have reported for themselves using their relatives'

phone/log-in function and/or misused proxy-reporting from their own mobile phones;

however, we have no means of capturing such activity.

Conclusions

Our national cohort of children and young people provides the first systematic description of

COVID-19 in children. Our data show that long illness duration after SARS-CoV-2 infection

in school-aged children is not common; however, a small proportion of children do have

prolonged illness duration and persistent symptoms, validating these children's experiences.

Our LC56 data provide reassurance regarding the longer term outcome for these children.

The symptom burden in children who tested negative for SARS-CoV-2 but had long illness

duration highlights that allocation of appropriate resources will be necessary for any child

with prolonged illness, whether due to SARS-CoV-2 infection or other illness. Our study

provides timely and critical data to inform discussions around the impact and implications of

the pandemic on UK paediatric healthcare resource allocation.

Data sharing

Data collected in the COVID Symptom Study smartphone application are being shared with

other health researchers through the UK National Health Service-funded Health Data

Research UK (HDRUK) and Secure Anonymised Information Linkage consortium, housed in

the UK Secure Research Platform (Swansea, UK). Anonymised data are available to be

shared with researchers according to their protocols in the public interest

(https://web.www.healthdatagateway.org/dataset/fddcb382-3051-4394-8436-b92295f14259).

Acknowledgements

This work is supported by the Wellcome EPSRC Centre for Medical Engineering at King's

College London (WT 203148/Z/16/Z) and the UK Department of Health via the National

Institute for Health Research (NIHR) comprehensive Biomedical Research Centre award to

Guy's & St Thomas' NHS Foundation Trust in partnership with King's College London and

King's College Hospital NHS Foundation Trust. Investigators also received support from the

Medical Research Council (MRC) and British Heart Foundation, the UK Research and

Innovation London Medical Imaging & Artificial Intelligence Centre for Value Based

Healthcare, and the Wellcome Flagship Programme (WT213038/Z/18/Z). EM is funded by

an MRC Skills Development Fellowship Scheme at KCL. CHS is supported by an

Alzheimer's Society Junior Fellowship (AS-JF-17-011). ZOE Global supported all aspects of

building and running the app and service to all users worldwide.

Declaration of interests

CH, SS, KR, JCP are employees of Zoe Global Ltd.

All other authors have nothing to declare.

References

Worldometer. Reported Cases and Dearths by Country or Territory. 2020. https://www.worldometers.info/coronavirus/#countries.

Xie Y, Wang Z, Liao H, Marley G, Wu D, Tang W. Epidemiologic, clinical, and laboratory findings of the COVID-19 in the current pandemic: Systematic review and meta-analysis. BMC Infectious Diseases. 2020; **20**. DOI:10.1186/s12879-020-05371-2.

Sudre C, Murray B, Varsavsky T, *et al.* Attributes and predictors of long COVID. *Nature Medicine* 2021; **Mar 10**. DOI:10.1038/s41591-021-01292-y.

Li B, Zhang S, Zhang R, Chen X, Wang Y, Zhu C. Epidemiological and Clinical Characteristics of COVID-19 in Children: A Systematic Review and Meta-Analysis. Frontiers in Pediatrics. 2020; **8**. DOI:10.3389/fped.2020.591132.

Ma N, Li P, Wang X, *et al.* Ocular Manifestations and Clinical Characteristics of Children with Laboratory-Confirmed COVID-19 in Wuhan, China. *JAMA Ophthalmology* 2020; **138**. DOI:10.1001/jamaophthalmol.2020.3690.

Alsharrah D, Alhaddad F, Alyaseen M, *et al.* Clinical characteristics of pediatric SARS-CoV-2 infection and coronavirus disease 2019 (COVID-19) in Kuwait. *Journal of Medical Virology* 2020. DOI:10.1002/jmv.26684.

Cai J, Wang X, Zhao J, *et al.* Comparison of Clinical and Epidemiological Characteristics of Asymptomatic and Symptomatic SARS-CoV-2 Infection in Children. *Virologica Sinica* 2020; **35**. DOI:10.1007/s12250-020-00312-4.

Chen Z, Tong L, Zhou Y, *et al.* Childhood COVID-19: a multicentre retrospective study. *Clinical Microbiology and Infection* 2020; **26**. DOI:10.1016/j.cmi.2020.06.015.

Xia W, Guo Y, Tian Z, *et al.* Clinical Features and Temporal Changes of RT-PCR and Chest CT in COVID-19 Pediatric Patients. *Frontiers in Pediatrics* 2020; **8**. DOI:10.3389/fped.2020.579512.

Rubens JH, Akindele NP, Tschudy MM, Sick-Samuels AC. Acute covid-19 and multisystem inflammatory syndrome in children. *The BMJ* 2021; **372**. DOI:10.1136/bmj.n385.

Evans C, Davies P. SARS-CoV-2 paediatric inflammatory syndrome. Paediatrics and Child Health (United Kingdom). 2021; **31**. DOI:10.1016/j.paed.2020.12.003.

Godfred-Cato S, Bryant B, Leung J, *et al.* COVID-19-Associated Multisystem Inflammatory Syndrome in Children - United States, March-July 2020. *MMWR Morb Mortal Wkly Rep*; **69**: 1074–80.

Sykes DL, Holdsworth L, Jawad N, Gunasekera P, Morice AH, Crooks MG. Post-COVID-19 Symptom Burden: What is Long-COVID and How Should We Manage It? *Lung* 2021. DOI:10.1007/s00408-021-00423-z.

Nalbandian A, Sehgal K, Gupta A, *et al.* Post-acute COVID-19 syndrome. *Nat Med* 2021. DOI:10.1038/s41591-021-01283-z.

National Institute for Health and Care Excellence (NICE). COVID-19 rapid guideline: managing the long- term effects of COVID-19 (NG188). Evidence reviews 2 and 3: prevalence. 2020.

Menni C, Valdes AM, Freidin MB, *et al.* Real-time tracking of self-reported symptoms to predict potential COVID-19. *Nature Medicine* 2020. DOI:10.1038/s41591-020-0916-2.

17   Government of the United Kingdom. GOV.UK Coronavirus (COVID-19) in the UK -
Cases by specimen date. 2021.

18   Varsavsky T, Graham MS, Canas LS, *et al.* Detecting COVID-19 infection hotspots
in England using large-scale self-reported data from a mobile application: a
prospective, observational study. *The Lancet Public Health* 2021; **6**.
DOI:10.1016/S2468-2667(20)30269-3.

19   Government of the United Kingdom. Coronavirus (COVID-19) in the UK - Lab-based
testing and capacity, by test type. 2021.
https://coronavirus.data.gov.uk/details/testing.

20   Public Health England. Weekly national Influenza and COVID-19 surveillance report
- Week 11 report (up to week 10 data) 18 March 2021. 2021.
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachm
ent_data/file/971212/Weekly_Flu_and_COVID-19_report_w11_v2.pdf (accessed April
27, 2021).

21   Struyf T, Deeks JJ, Dinnes J, *et al.* Signs and symptoms to determine if a patient
presenting in primary care or hospital outpatient settings has COVID-19. Cochrane
Database of Systematic Reviews. 2021; **2021**.
DOI:10.1002/14651858.CD013665.pub2.

22   O. A-M, M. E, U. L, *et al.* Neurologic and Radiographic Findings Associated with
COVID-19 Infection in Children. *JAMA Neurology* 2020.

23   Say D, Crawford N, McNab S, Wurzel D, Steer A, Tosif S. Post-acute COVID-19
outcomes in children with mild and asymptomatic disease. *The Lancet Child &*
Adolescent Health 2021; published online April. DOI:10.1016/S2352-4642(21)00124-
3.

24   Kumar L, Kahlon N, Jain A, Kaur J, Singh M, Pandey AK. Loss of smell and taste in
COVID-19 infection in adolescents. *International Journal of Pediatric*
Otorhinolaryngology 2021; **142**. DOI:10.1016/j.ijporl.2021.110626.

25   Carfì A, Bernabei R, Landi F. Persistent symptoms in patients after acute COVID-19.
JAMA - Journal of the American Medical Association. 2020; **324**.
DOI:10.1001/jama.2020.12603.

26   Townsend L, Dyer AH, Jones K, *et al.* Persistent fatigue following SARS-CoV-2
infection is common and independent of severity of initial infection. *PLoS ONE* 2020;
15. DOI:10.1371/journal.pone.0240784.

27   Abu-Arafeh I, Razak S, Sivaraman B, Graham C. Prevalence of headache and
migraine in children and adolescents: A systematic review of population-based
studies. Developmental Medicine and Child Neurology. 2010; **52**. DOI:10.1111/j.1469-
8749.2010.03793.x.

28   Abu-Arefeh I, Russell G. Prevalence of headache and migraine in schoolchildren.
BMJ 1994; **309**. DOI:10.1136/bmj.309.6957.765.

29   Farmer A, Fowler T, Scourfield J, Thapar A. Prevalence of chronic disabling fatigue
in children and adolescents. British Journal of Psychiatry. 2004; **184**.
DOI:10.1192/bjp.184.6.477.

30   Crawley EM, Emond AM, Sterne JAC. Unidentified chronic fatigue
syndrome/myalgic encephalomyelitis (CFS/ME) is a major cause of school absence:
Surveillance outcomes from school-based clinics. *BMJ Open* 2011; **1**.
DOI:10.1136/bmjopen-2011-000252.

31   Norris T, Collin SM, Tilling K, *et al.* Natural course of chronic fatigue
syndrome/myalgic encephalomyelitis in adolescents. *Archives of Disease in*
Childhood 2017; **102**. DOI:10.1136/archdischild-2016-311198.
32   Balfour HH, Odumade OA, Schmeling DO, *et al.* Behavioral, virologic, and
immunologic factors associated with acquisition and severity of primary epstein-barr
virus infection in university students. *Journal of Infectious Diseases* 2013; **207**.
DOI:10.1093/infdis/jis646.
33   Duarte-Salles T, Vizcaya D, Pistillo A, *et al.* Baseline characteristics, management,
and outcomes of 55,270 children and adolescents diagnosed with COVID-19 and
1,952,693 with influenza in France, Germany, Spain, South Korea and the United
States: An international network cohort study. medRxiv. 2020.
DOI:10.1101/2020.10.29.20222083.
34   UK Office for National Statistics (ONS), Steel K, Davies B. 5. Test sensitivity and
Specificity. 2021.
https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditio
nsanddiseases/methodologies/covid19infectionsurveypilotmethodsandfurtherinformati
on#test-sensitivity-and-specificity.
35   National Health Service (NHS). COVID-19 Hospital Activity. 2021.
https://www.england.nhs.uk/statistics/statistical-work-areas/covid-19-hospital-activity/.
36   Public Health Scotland. Weekly national seasonal respiratory report - Week ending
28 March 2021 – week 12. https://beta.isdscotland.org/media/8300/week-12-
respiratory-report-full-report-final.pdf. 2021; published online March 31.
37   Public Health Wales - National Health Service. Weekly Influenza Activity in Wales
Report. http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=34338. 2021;
published online April 7.
38   Ludvigsson JF. Case report and systematic review suggest that children may
experience similar long-term effects to adults after clinical COVID-19. *Acta*
Paediatrica, International Journal of Paediatrics 2021; **110**. DOI:10.1111/apa.15673.
39   UK Office for National Statistics (ONS) ER. Young people's well-being in the UK:
2020.
https://www.ons.gov.uk/peoplepopulationandcommunity/wellbeing/bulletins/youngpeo
pleswellbeingintheuk/2020. 2020; published online Oct 2.
40   Hysing M, Elgen I, Gillberg C, Lie SA, Lundervold AJ. Chronic physical illness and
mental health in children. Results from a large-scale population study. *Journal of Child*
Psychology and Psychiatry and Allied Disciplines 2007; **48**. DOI:10.1111/j.1469-
7610.2007.01755.x.
41   UK Office for National Statistics (ONS). COVID-19 Schools Infection Survey Round
2, England: December 2020. 2021.
https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditio
nsanddiseases/bulletins/covid19schoolsinfectionsurveyround2england/december2020
.
42   Public Health Scotland. Total Cases by Age and Sex. 2021.
43   Public Health Wales TP. Rapid COVID-19 surveillance.
https://public.tableau.com/profile/public.health.wales.health.protection#!/vizhome/Rapi
dCOVID-19virology-Public/Headlinesummary. 2021; published online April 16.
44   Ulster University. Northern Ireland COVID-19 Tracker - Cumulative Number of
Individuals Tested Positive by Age. 2021.
https://www.ulster.ac.uk/coronavirus/research/impact/ni-covid-19-tracker.

45   UK Office for National Statistics (ONS) DA. Prevalence of long COVID symptoms
and COVID-19 complications.
https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthan
dlifeexpectancies/datasets/prevalenceoflongcovidsymptomsandcovid19complications.
2020; published online Dec 16.
46   UK Office for National Statistics (ONS). Updated estimates of the prevalence of long
COVID symptoms.
https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthan
dlifeexpectancies/adhocs/12788updatedestimatesoftheprevalenceoflongcovidsympto
ms. 2021; published online Jan 21.
47   UK Office for National Statistics (ONS). All data related to Prevalence of ongoing
symptoms following coronavirus (COVID-19) infection in the UK: 1 April 2021.
https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditio
nsanddiseases/bulletins/prevalenceofongoingsymptomsfollowingcoronaviruscovid19in
fectionintheuk/1april2021/relateddata. 2021; published online April 1.
48   Katie Gibbons. Forgotten children living with long Covid,
https://www.thetimes.co.uk/article/forgotten-children-living-long-covid-s0c6z0f20. The
Times. 2021; published online March 13.
49   Scally Gabriel. Why it's not too late for the UK to pursue a zero-Covid strategy.
NewStatesman, https://www.newstatesman.com/science-
tech/coronavirus/2021/02/why-it-s-not-too-late-uk-pursue-zero-covid-strategy. 2021;
published online Feb 23.
50   Graham MS, May A, Varsavsky T, *et al.* Knowledge barriers in the symptomatic-
COVID-19 testing programme in the UK: an observational study. *medRxiv* 2021.

Table 1. Characteristics of school-aged children who tested positive for SARS-CoV-2, and the control cohort of children (matched 1:1 for age, gender, and week of testing) who tested negative for SARS-CoV-2. The cohort of children with positive SARS-CoV-2 testing is presented here both as younger and older groups; and for usual (i.e., short) vs. extended illness duration.

	Children with positive SARS-CoV-2 test					Children with negative SARS-CoV-2 test (matched cohort)
	Younger group	Older group	Symptom duration <10 days	Symptom duration ≥ 28 days	Full cohort	
Number	588	1,146	1,183	77	1,734	1,734
Males (%)	287 (48.9)	577 (50.3)	618 (52.2)	35 (45.5)	864 (49.8)	865 (49.9)
Age, years (median, [IQR])	9 [7;10]	15 [13;16]	13 [10;15]	14 [12;16]	13 [10;15]	13 [10;15]
BMI (kg/m^2) (median, [IQR])	17.0 [15.1;19.7]	20.1 [17.8;22.3]	19.0 [16.5;21.8]	18.6 [16.3;21.8]	19.2 [16.6;21.8]	19.0 [16.5;21.5]
Asthma (%)	69 (11.7)	147 (12.8)	134 (11.7)	10 (13.0)	216 (12.8)	229 (13.2)
Visit to hospital (%)	16 (2.7)	21 (1.8)	20 (1.7)	1 (1.3)	37 (2.1)	26 (1.5)

Illness duration (median, [IQR])	5 [2;9]	7 [3;12]	4 [2;6]	46 [32;58]	6 [3;11]	3 [2;7]
Number of symptoms in the first week (median, [IQR])	3 [2;5]	4 [2;6]	3 [2;5]	6 [4;8]	3 [2;6]	2 [1;4]

BMI: Body Mass Index. IQR: Inter Quartile Range.

Table 2. Symptoms over the duration of illness in children testing positive for SARS-CoV-2.

	Sample tested positive for SARS-CoV-2		
	Younger children (aged 5-11 years, n=588)	Older children (aged 12-17 years, n=1,146)	Overall cohort (n=1,734)
Headache	324	755	1,079 (62.2%)
Fatigue	258	696	954 (55.0%)
Sore Throat	213	585	798 (46.0%)
Anosmia	132	554	686 (39.6%)
Fever	257	396	653 (37.7%)
Abdominal Pain	163	194	357 (20.6%)
Dizziness	84	300	384 (22.1%)
Persistent Cough	145	298	443 (25.5%)
Loss of Appetite	120	254	374 (21.6%)
Eye Soreness	89	248	337 (19.4%)
Myalgias	54	231	285 (16.4%)
Nausea	95	193	288 (16.6%)
Hoarse voice	63	166	229 (13.2%)
Chest Pain	37	143	180 (10.4%)
Dyspnoea	24	143	167 (9.6%)
Diarrhoea	48	79	127 (7.3%)
Confusion	15	81	96 (5.5%)
Red Welts	16	36	52 (3.0%)
Blisters	4	22	26 (1.5%)

Figure 1. Flowchart of Inclusion and Exclusion Criteria for the Study.

Legend: overall number for the entire cohort of children is given first; numbers within brackets separated by oblique refer to younger children and older children in that order. 'Not valid result' – PCR test result proxy-reported as "failed test" or "still waiting". 'Duration calculable' – illness onset within defined timeframe of testing for SARS-CoV-2, and with defined endpoint (for details, please see Methods). 'Irregular logging' – proxy-reporting with intervals of >7 days between proxy-reports during illness duration). 'Illness onset outside of study bounds' - symptom onset before 1 September 2020 or after 24 January 2021. 'Hosp' – presenting to hospital (either admitted to hospital or seen in Accident and Emergency ward).

[image at following page]

[caption at previous page]

Figure 2. Prevalence of symptoms reported over the course of illness in younger (5-11 years, n=588) and older (12-17 years, n=1,146) children testing positive for SARS-CoV-2.

Figure 3. Median duration of each symptom [IQR] in younger (5-11 years) and older (12-17 years) children.

Figure 4. Heat maps showing symptom duration in school-aged children (aged 5-17 years) in whom at least one symptom persisted for ≥28 days.

Children with a positive SARS-CoV-2 test (n = 77 children)

Legend:

X-axis, duration in days.

Y axis, symptoms.

Legend: BT, blisters; RW, red welts; ES, eye soreness; DZ, dizziness and light-headedness; AN, anosmia; PC, persistent cough; FV, fever; DI, diarrhoea; CO, confusion; HV, hoarse voice; HA, headache; UMP, myalgias [unusual muscle pains]; LA, loss of appetite; SOB, dyspnoea [shortness of breath]; ST, sore throat; CP, chest pain; AP, abdominal pain; FA, fatigue. Colour bar provides percentage comparison.