

1 **Hypomethylation of Monoamine Oxidase A promoter/exon 1 region** 2 **is associated with temper outbursts in Prader-Willi syndrome**

3
4 Maximilian Deest¹, Vanessa Buchholz¹, Kirsten Jahn¹, Christian Eberlein, Stefan Bleich¹, Helge
5 Frieling¹

6
7 Department of Psychiatry, Social Psychiatry and Psychotherapy, Medical School Hannover,
8 Hannover, Germany

9 Corresponding author:

10 Dr. med. Maximilian Deest, Department Psychiatry, Social Psychiatry and Psychotherapy,
11 Medical School Hannover, Carl-Neuberg-Straße 1, 30625 Hannover, Germany

12 Phone: +49 176 1532 2402

13 Mail: deest.maximilian@mh-hannover.de

14

15 Conflict of interest

16 All authors declare, that they do not have a conflict of interest

17

18 Source of funding

19 This project was supported by PRACTIS – Clinician Scientist Program of Hannover Medical
20 School, funded by the German Research Foundation (DFG, ME 3696/3-1).

21

22 Abstract

23

24 Background: Prader-Willi syndrome (PWS) is a rare neurodevelopmental disorder caused by
25 the absence of paternally expressed and maternally imprinted genes on chromosome 15q
26 11.2-13. It is associated with a certain behavioural phenotype with repetitive and ritualistic
27 behaviours, skin-picking and temper outbursts. Temper outbursts are characterized by verbal
28 and physical aggression with screaming, destroying property, and/or physical aggression
29 towards others. They drastically effect the quality of life of the individuals as well as the
30 relatives and caregivers. Recent studies show a promising therapeutic effect of serotonin
31 reuptake inhibitors like sertraline on frequency and intensity of outbursts. Monoamine
32 oxidase A (MAOA) (X p11.23) plays a crucial role in the metabolism of monoamines such as
33 serotonin, norepinephrine, and dopamine. Dysregulation in methylation of the CpG island

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

34 spanning the promoter region and exon 1 of MAOA is implicated in impulsive and aggressive
35 behaviour.

36 Methods: In the present study, methylation rates of CpG dinucleotides in the MAOA promoter
37 and exon 1 region were determined from DNA derived from whole blood samples of PWS
38 patients (n=32) and controls (n=14) matched for age, sex and BMI via bisulfite sequencing.
39 PWS patients were grouped into those showing temper outbursts, and those who do not.

40

41 Results: Overall, PWS patients show a significant lower methylation rate at the promoter/exon
42 1 region than healthy controls in both sexes. Furthermore, PWS patients, male as well female
43 with temper outbursts show a significant lower methylation rate than those without temper
44 outbursts ($p < 0.001$ and $p = 0.006$)

45

46 Conclusion: The MAOA promoter/exon 1 region methylation seems to be dysregulated in PWS
47 patients in sense of a hypomethylation, especially in those suffering from temper outbursts.
48 As MAOA is involved in the metabolism of serotonin, we conclude that this dysregulation plays
49 a crucial role in the pathophysiology of temper outbursts in PWS. Furthermore, our findings
50 suggest, that dysregulation of certain genes outside the PWS locus contribute to the
51 behavioural phenotype of PWS.

52

53 Keywords: Prader-Willi syndrome, Monoamine Oxidase A, temper outbursts, Methylation

54

55

56

57 1. Introduction

58 Prader-Willi syndrome (PWS) is a rare neurodevelopmental disorder associated with a certain
59 behavioural phenotype with repetitive and ritualistic behaviours, skin-picking, and temper
60 outbursts (Sinnema *et al.*, 2011; Whittington and Holland, 2018; Butler, Miller and Forster,
61 2019). These temper outbursts are characterized by verbal and physical aggression with
62 screaming, destroying property, and/or physical aggression towards others (punching, kicking
63 etc) (Rice, Woodcock and Einfeld, 2018a).

64 PWS, in general, is characterised by hypogonadism and growth hormone deficiency, mild to
65 moderate intellectual disability, and infantile hypotonia with an early failure to thrive.

66 Newborns with PWS typically display feeding difficulties that drastically changes during early
67 childhood with the development of excessive hyperphagia leading to severe obesity, if food
68 intake is not restricted (Dykens and Shah, 2003; Cassidy *et al.*, 2012; Angulo, Butler and
69 Cataletto, 2015). The cause of PWS is the lack of expression of paternally expressed but
70 maternally imprinted genes on the chromosome 15q11.2 – 13 region. This can either result
71 from a paternal deletion (delPWS; 70% of cases), maternal uniparental disomy (mUPD;25%),
72 or an imprinting centre defect (IC;3-5%) (Nicholls and Knepper, 2001; Horsthemke and
73 Wagstaff, 2008). Several mental disorders, such as autism spectrum disorders (ASD),
74 psychosis, and depression, are more frequent in PWS than in other populations. The
75 prevalence of mental disorders differs among genetic subtypes of PWS; with a higher rate of
76 ASD and psychosis in those with a mUPD than those with a delPWS (Holland *et al.*, 2003;
77 Vogels *et al.*, 2003; Soni *et al.*, 2007, 2008).

78

79 Temper outbursts are displayed by up to 88% of individuals with PWS (Dykens, Cassidy and
80 King, 1999), typically start in childhood and increase in intensity and frequency with age
81 (Dykens *et al.*, 1992). They drastically affect the quality of life of the individual as well as the
82 relatives and caregivers (Mazaheri *et al.*, 2013). Recent studies identified the provocation of
83 those temper outbursts as goal blockage, difficulty in dealing with change, and feeling of social
84 injustice (Rice, Woodcock and Einfeld, 2018a). Current treatment options include clinical
85 behavior interventions like token boards or behavior management plans. Several studies have
86 shown that the serotonergic system is disturbed in PWS, possibly contributing to the
87 behavioural phenotype. For instance, SNORD115 (HB-52 III small nuclear RNA), located at the
88 PWS region is involved in alternative splicing of the serotonergic 2c receptor (Kishore and
89 Stamm, 2006), and mice with a deletion of the IC show reduced functioning of the serotonin
90 2c receptor. Those mice showed increased impulsivity and compulsivity. The use of a
91 serotonin receptor agonist improved impulsivity in these mice (Davies *et al.*, 2019).
92 Accordingly, we recently showed that the selective serotonin reuptake inhibitor (SSRI)
93 sertraline is a promising treatment option for temper outbursts in PWS leading to a decrease
94 of frequency and intensity of outbursts (Deest *et al.*, 2020). All in all, these findings strongly
95 suggest that the serotonergic system may play a role in the pathophysiology of behaviour
96 problems in PWS.

97

98 The *monoamine oxidase A gene*, located on Xp11.4-p11.3, and its product MAOA play a crucial
99 role in the metabolism of monoamines such as serotonin, norepinephrine, and dopamine (Lan
100 *et al.*, 1989; Bortolato, Chen and Shih, 2008). MAOA consists of 14 exons and 14 introns with
101 two cytosine-guanine (CpG) islands. One CpG island is located at the promoter region and the
102 second one spans the promoter region as well as exon I and the beginning of intron I (Shih,
103 Wu and Chen, 2011). Previous studies showed significant disturbances in methylation of these
104 CpG islands in several mental disorders like anxiety disorders (Domschke *et al.*, 2012),
105 depression (Melas *et al.*, 2013; Melas and Forsell, 2015), posttraumatic stress disorder (PTSD)
106 (Ziegler *et al.*, 2018) and antisocial personality disorder/conduct disorder (Cecil *et al.*, 2018).
107 Individuals with Brunner syndrome, caused by a MAOA deficiency, show serotonergic
108 symptoms (flushing, diarrhea, headache, palpitations) and aggressive outbursts (Palmer *et al.*,
109 2016). In PWS, Akefeldt *et al.* showed, that MAO activity in platelets is significantly higher than
110 in controls (Åkefeldt and Månsson, 1998).

111

112 As disturbances of the serotonergic pathway most likely contribute to temper outbursts and
113 MAOA is crucially involved in the metabolism of serotonin, we here investigate the
114 methylation of MAOA in PWS. Therefore, we investigated the methylation of 47 CpG
115 nucleotides at the CpG island spanning exon 1 and part of the promoter to investigate
116 methylation between PWS and healthy individuals with a focus on temper outbursts.

117

118 2. Methods

119

120 2.1 Subjects

121 This study was approved by the local Ethics Committee of Hannover Medical School and
122 adhered to the Declaration of Helsinki. All participants in this study gave their written
123 informed consent for participation in this study and were recruited at the Outpatient
124 Department for Mental Health in Rare Genetic Disorders of the Department of Psychiatry,
125 Social Psychiatry, and Psychotherapy of Medical School Hannover, Hannover, Germany. All
126 participants were patients of the Outpatient Department. In patients, the PWS diagnosis was
127 genetically confirmed. Characteristics such as age, gender, and BMI were documented. The
128 controls were matched for gender, body mass index (BMI), and age. EDTA-blood was collected
129 and stored at Hannover Unified Biobank.

130

131 2.2 Assessment of temper outbursts

132 As currently no reliable psychometric test instrument in german language for temper
133 outbursts in PWS exist, the caregivers of the patients had to state if temper outbursts take
134 place and have a significant impact on the patients daily life. A criterion for this was that the
135 patient must exhibit physical aggression towards others and/or destroy property.

136

137 2.3 DNA isolation and bisulfite reaction

138 DNA for methylation analysis was extracted from blood directly after sampling by the
139 Hannover Unified Biobank using the Hamilton ChemagicStar (Hamilton Germany Robotics,
140 Graefelfing, Germany) and the chemagicStar DNA-Blood1k kit (PerkinElmer chemagen
141 Technology, Baesweiler, Germany).

142 Bisulfite conversion and purification was performed using the EpiTect® 96 Bisulfite Kit
143 (Qiagen, Hilden, Germany) following the manufacturer recommendations.

144

145 2.4 Amplification of target sequences

146 Amplification of *MAOA* promoter, exon 1/ intron 1 target sequences of the purified bisulfite-
147 converted DNA was conducted using touchdown PCRs. We used the forward primer
148 'GTTATAAAGGGGTTAGATTAAGTGG' and the reverse primer 'AAACCTAAACAATTATACCCCC'.
149 Cycling conditions apply as follows: 15 min 95 °C, 1 min 97 °C , 15 x (30s 95 °C, 45s 65 °C
150 with -1 °C per cycle, 1 min 72 °C), 30 x (30s 95°C, 45s 50 °C, 1 min 70 °C), 5 min 65 °C.
151 Primers were ordered from Integrated DNA Technologies (Coral Ville, IA, USA). PCRs were
152 performed on a C1000 Thermal Cycler (BIO-RAD, Hercules, CA, USA) using the HotStarTaq®
153 Master Mix Kit (QIAGEN, Hilden, Germany). The PCR products were purified automatically on
154 a Biomek® NxP (Beckman Coulter, Brea, CA, USA) using paramagnetic beads (Clean-NGS, GC
155 Biotech®).

156

157 2.5 Sequencing

158 The Big-Dye® Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA,
159 USA) was used for sequencing PCR of the target fragments. Oligonucleotides used for
160 sequencing PCR were 'GTTATAAAGGGGTTAGATTAAGTGG'. Cycling conditions apply as
161 follows: 1 min 96°C, 25 x (5s 96°C, 90s 60°C, 90s 50°C). Clean-up of the sequencing PCR was

162 also automatized on our Biomek Nxp by use of CleanDTR beads (CG Biotech®). Sequencing
163 was performed on a 3500XL genetic analyzer from ABI Life Technologies (Grand Island, NY,
164 USA) according to the manufacturer's instructions. The amplified fragments covered 47 CpG
165 sites within exon 1 and the promoter region of the *MAOA* gene (from -241 bp to +328 bp of
166 the first exon). Position of CpG sites is given in relation to the starting point of exon 1:
167 GRCh38:X:43654907 (from ENSEMBL#ENSG00000189221).

168

169 2.6 Determination of methylation Rates

170

171 Determination of methylation rates for each CpG site and sequence analysis were performed
172 using the Epigenetic Sequencing Methylation analysis software (ESME) (Lewin *et al.*, 2004).
173 The ratio between normalized peak values of cytosine and thymine was used to calculate the
174 methylation rate of each CpG site per subject.

175

176 2.7 Statistical analysis

177

178 Sequence quality was determined via Sequence Scanner v2.0 software (ABI Life Technologies).
179 All sequences showed a Quality Value > 20 for Trace Score in the Quality Control Report and
180 could therefore be included for further analyses. All statistical analysis were performed using
181 the Statistical Package for Social Sciences (SPSS, IBM, Armonk, NY). GraphPad Prism was used
182 for data illustration. Single CpGs with less than 95% sequencing success among the samples
183 were excluded which applied for three CpGs ahead of the first exon as well as twenty of the
184 first exon. All samples showed more than 95% valid measurements and thus had been
185 included. No CpG site showed less than 5% inter-individual variability. After applying these
186 criteria for data exclusion, the number of analyzed CpGs was 24. Mixed linear models were
187 calculated to detect significant fixed effects of different variables on methylation rate.
188 Bonferroni correction method was used to correct P values for multiple testing. In each
189 analysis, a p value of < 0.05 was considered significant.

190

191

192

193

194 3. Results

195

196 3.1 Demographic data

197

198 In this study we included 32 patients with genetically confirmed diagnosis of PWS (13 female,
199 19 male) with a mean age of 27.9 years (min: 12y, max:55y) and a mean BMI of 28.1 kg/m²
200 (min: 19.1 kg/m², max: 55.58 kg/m²). The control group consisted of 14 healthy individuals
201 without PWS (5 female, 9 male) with a mean age of 31.2 years (min: 18y, max:47y) and a mean
202 BMI of 26.2 kg/m² (min: 19.4 kg/m², max: 40.1 kg/m²) (see Table 1 for characteristics of study
203 population). In 15 cases PWS was caused by a deletion (46.9%), with four having a deletion
204 type I, 10 having a deletion type II. In one case no differentiation between deletion subtype
205 was performed during genetic counseling. In nine patients PWS was caused by a mUPD
206 (28.1%), in 4 cases by IC (9.4%). In four cases distinguishing between mUPD or IC did not take
207 place due to various reasons (e.g. no blood from the parents could be obtained).

208 At the timepoint of counseling, caregivers had to state (yes/no) if temper outbursts are
209 common and have a significant impact on the daily life. Of the 32 individuals, 17 (eleven male,
210 six female) showed temper outbursts and 15 (eight male, seven female) did not (see Table 1
211 for more characteristics).

212 Overall, 18 patients (56%) received an antipsychotic medication (e.g. risperidone, aripiprazole,
213 quetiapine) and four received an antidepressant medication (e.g. sertraline, fluoxetine) at
214 timepoint of consultation. Seven patients suffered from a psychosis (acute or chronic), two
215 from an obsessive-compulsive disorder and two from a ASD. The other patients received
216 medication because of behavioural problems. In the subgroup of those showing temper
217 outbursts eight (44%) received an antipsychotic medication and one (5.5%) an antidepressant.
218 In those showing no outbursts, eight (57%) received antipsychotic medication and three
219 (21.4%) received an antidepressant medication.

220

221

222

223

	control (n=14)	PWS		
		total (n=32)	with temper outbursts (n=17)	no temper outbursts (n=15)
Sex	9 male (64.3%) 5 female (35.7%)	19 male (59.4%) 13 female (40.6%)	11 male (64.7%) 6 female (35.3%)	8 male (53.3%) 7 female (46.7%)
Age	18-47 y (mean 31.21y)	12-55y (mean 27.9y)	12-41y (mean 24.8y)	15 - 55y (mean 29.3y)
BMI	19.4-40 kg/m ² (mean 26.2 kg/m ²)	19.1 - 55.1 kg/m ² (mean 28.1 kg/m ²)	19.1-55.6kg/m ² (mean 29kg/m ²)	19.9 - 36.5y (mean 26.96kg/m ²)
Subtyp	/	del: 15 (4 del I, 10 del II, 1 ns) UPD/IC : 17 (9 UPD, 4 IC, 4 ns)	del: 9 (2 del I, 6 del II, 1 ns) UPD/IC: 8 (5 UPD, 2 IC, 1 ns)	del: 6 (2 del I, 4 del II) UPD/IC: 9 (4 UPD, 2 IC, 3 ns)
Antipsychotic medication	/	n= 18 (56%)	n = 8 (44%)	n=8 (57%)
Antidepressant medication	/	n=4 (12.5%)	n=1 (5.5%)	n=3 (21.4%)
Psychiatric diagnosis	/	psychosis = 7 OCD = 2 ASD = 2	psychosis = 2 ASD = 2	psychosis=5 OCD=2

224


225 *Table 1 Demographics of study population. Study population is grouped by controls and PWS. PWS patients are then grouped*
 226 *by showing temper outbursts vs. not showing outbursts. Abbreviations: del, deletion; del I, deletion type I; del II, deletion type*
 227 *II, UPD maternal uniparental disomie; IC, Imprinting defect; OCD, obsessive-compulsive disorder; ASD, autism spectrum*
 228 *disease, ns, not specified because no differentiation of specific subtype could be obtained (del I vs del II or UPD vs IC) due to*
 229 *different reasons (e.g.) no blood from the parents could be obtained*

230

231 3.2 The difference in Methylation Rate between PWS and controls

232

233 Comparison of the mean methylation levels of the investigated CpGs within the *MAOA* exon
 234 1/promoter region showed significantly lower methylation rates in PWS than in healthy
 235 controls ($p < 0.001$) with a mean methylation of 17.8% \pm 1% for controls and 11.8% \pm 0.6%
 236 for PWS. (see Fig. 1A). Methylation levels are known to be gender-dependent in general
 237 (Singmann *et al.*, 2015) and for *MAOA* in particular (Chen *et al.*, 2012; Domschke *et al.*, 2012),
 238 so we performed a gender-specific analysis. Looking at the mean methylation rate in males,
 239 PWS subjects showed a mean methylation rate of 1.1% \pm 0.4%, and male controls had a value
 240 of 6.9% \pm 0.6%. The difference between those two groups was significant ($p < 0.001$) (see Fig.
 241 1B). In the female PWS group mean methylation rate was 26.3% \pm 0.9% and in the control
 242 group the value was 37.2% \pm 1.4%. The difference in methylation rate between female PWS
 243 subjects and female controls was also significant ($p < 0.001$) (see Fig. 1C). As mean methylation
 244 rates showed a strong differences between the sexes, we performed only gender specific
 245 analysis for the next steps.



246

247 *Figure 1 Overview of methylation rates for single CpGs. (A) PWS patients show a significant lower methylation than controls,*
248 *(B), (C) Methylation rates between PWS and controls in regard of sex show, that methylation in PWS is significantly lower than*
249 *in healthy controls in both sexes*


250

251 3.3 The difference in methylation rate between the genetic subtype of PWS

252

253 As there are certain differences, especially in the distribution of psychiatric diseases between
254 the genotypes of PWS, we were interested in differences in methylation between those
255 groups. As an IC and a mUPD lead to the same imprinting pattern in the PWS region
256 (Horsthemke and Wagstaff, 2008), we split our cohort into two groups: a delPWS subgroup
257 and a mUPD/IC subgroup. This resulted in groups of 15 individuals with a delPWS (ten male,
258 five female) and 17 with a mUPD/IC (nine male, eight female). When looking at the males,
259 those with the delPWS subtype showed a mean methylation rate of 2.1% +/- 0.6% compared
260 with those with an mUPD/IC with a mean methylation rate of 0% +/- 0.7%. Those with a
261 deletion subtype showed significantly lower methylation rates ($p < 0.001$) compared to healthy
262 individuals as well as those with an mUPD/IC ($p < 0.001$). The difference between the subtypes
263 was not significant ($p = 0.053$) (see Fig. 2A). In the female group, the mean methylation rate of
264 those with a delPWS was 28.4% +/- 1.4% and those with mUPD/IC was 24.9% +/- 1.1%. Both

265 delPWS as well mUPD/IC groups were significantly different ($p < 0.001$) from healthy controls.
266 However, there was no significant difference between the genetic subgroups ($p = 0.147$) (see
267 Fig. 2B).



268
269 *Figure 2 Comparison of methylation rates between genetic subtypes of PWS and healthy controls in male (A) and female (B).*
270 *No difference between the subtypes was observed.*

271

272 3.4 Temper outbursts and Methylation Rate

273


274 As temper outbursts are common among all genetic subtypes we next investigated the
275 methylation rate among those showing temper outbursts and those not showing temper
276 outbursts. The caregivers had been asked if temper outbursts are common and have a
277 significant impact on daily life. Of the 32 individuals with PWS in this study, 17 showed temper
278 outbursts (11 male, six female) and 14 did not (eight male, six female).

279

280 We again compared the mean methylation rate in a sex-specific manner. In healthy males the
281 mean methylation rate was 6.9% +/- 0.6%, in male PWS showing no temper outbursts the
282 value was 2.7% +/- 0.7% and in those with temper outbursts 0% +/- 0.6%. Thus male PWS
283 without temper outbursts shows significantly lower methylation than healthy male controls
284 ($p < 0.001$) but significantly higher methylation than those PWS individuals with temper
285 outbursts ($p = 0.006$) (see Fig. 3A).

286 In female healthy controls the mean methylation rate was 37.2% +/- 1.4%, in female PWS
287 without temper outbursts the value was 30.4% +/- 1.1% and in female PWS showing temper
288 outbursts the value was 21.4% +/- 1.2 resulting in no significant difference between healthy
289 controls and female PWS showing no temper outbursts ($p = 0.265$). However, females with
290 PWS showing temper outbursts had significantly lower methylation than healthy controls
291 ($p < 0.001$) and PWS individuals without temper outbursts ($p < 0.001$) (see Fig. 3B).

292



293

294 *Figure 3 Comparison of methylation rates between PWS individuals showing temper outbursts, PWS individuals not showing*
 295 *temper outbursts and healthy controls in male (A) and female (B). In both sexes, patients suffering from temper outbursts*
 296 *have a significant hypomethylation than healthy controls and PWS patients without outbursts.*

297

298

299 4. Discussion

300

301 In this study, we investigated the methylation pattern of the CpG island flanking exon 1 and
 302 part of the promoter region in *MAOA* in PWS in comparison with healthy controls matched for
 303 age, sex, and BMI. In our study group, male controls as well as male PWS individuals, show
 304 significantly lower methylation than females which is in line with previous findings (Domschke
 305 *et al.*, 2012; Singmann *et al.*, 2015). However, in both sexes, the methylation rate in PWS is

306 significantly lower than in healthy controls. There are several reported findings of differences
307 in the prevalence of certain psychiatric disorders, such as psychosis or ASD, between genetic
308 subtypes of PWS (Vogels *et al.*, 2003; Soni *et al.*, 2007, 2008; Dykens *et al.*, 2017). However,
309 we did not find a significant difference in methylation rates between delPWS or mUPD/IC.
310 Thus differences in *MAOA* methylation probably do not contribute to the described
311 differences between the subtypes. Temper outbursts, often associated with physical
312 aggressiveness and/or self-harm are a major problem across all genetic subtypes of PWS
313 (Manzardo *et al.*, 2018; Rice, Woodcock and Einfeld, 2018b). In our study cohort, 53% showed
314 frequent temper outbursts having a significant impact on daily life as rated by the caregivers.
315 As dysregulation of methylation in exon 1 of *MAOA* is known to be associated with
316 impulsiveness and aggression (Cecil *et al.*, 2018), we compared the methylation patterns of
317 PWS individuals with temper outbursts and those without. In both sexes, individuals with
318 temper outbursts showed a significantly lower mean methylation rate than those without
319 temper outbursts and healthy controls. As the rate of those receiving an antipsychotic or
320 antidepressant medication was similar between those groups, we think that the observed
321 difference is not caused by the medication.

322 Methylation can be tissue-specific and as we investigated the methylation rate using
323 peripheral blood, it could be argued that this might not predict central enzyme activity.
324 However previous results showed that peripheral methylation of *MAOA* correlates with the
325 central enzyme activity of *MAOA* using in vivo brain imaging via positron emission tomography
326 (Shumay *et al.*, 2012). Furthermore, it has been shown that methylation of the first exon
327 predicts protein expression in general (Brenet *et al.*, 2011).

328 Thus, we conclude that individuals with PWS, especially those with temper outbursts, have an
329 increased *MAOA* activity in the brain leading to increased metabolism of monoamines,
330 especially serotonin. It has been shown that platelet MAO activity, both very low and very
331 high, correlates with impulsivity in children but not with intellectual levels (Shekim *et al.*,
332 1984). This would be in line with previous findings from Akesfeldt *et al.* which showed
333 increased serotonin metabolites in cerebrospinal fluid in PWS indicating an increased
334 metabolism of serotonin (Åkefeldt *et al.*, 1998). We previously showed that PWS patients
335 having temper outbursts benefitted from medication with the SSRI sertraline (Deest *et al.*,
336 2020). This inhibits the re-uptake of serotonin and thus increases the intrasynaptic serotonin
337 concentration and thus counteracts the increased *MAOA* activity. Disturbances of the

338 serotonergic system have long been proposed to contribute to the behaviour problems of
339 PWS. One hypothesis is, that *SNORD 115* (HBII-52), a small nuclear RNA located at the PWS IC
340 regulates alternative splicing of the serotonin 2c receptor (Kishore and Stamm, 2006; Davies
341 *et al.*, 2019). However, Hebras *et al.* recently showed that in knockout *SNORD115* mice
342 alternative RNA splicing is unchanged and mice did not show significant behaviour
343 aberrations. They raised doubts about the involvement of alternative RNA splicing of
344 serotonin 2c receptor in the pathophysiology of behavioural problems (Jade *et al.*, 2020). Our
345 findings could lead to the hypothesis that it is not the dysregulation of serotonin receptors but
346 dysregulation of MAOA activity that contributes to the disturbance of the serotonergic system
347 in PWS. This could be in line with findings that mouse models of PWS showing aberrant
348 behaviour have decreased impulsivity when treated with a serotonin receptor agonist (Davies
349 *et al.*, 2019). Interestingly, MAOA inhibitors are treatment options for atypical depression that
350 is usually associated with symptoms like overeating and sleep increase, which are particular
351 key symptoms of PWS (Davidson *et al.*, 1982).

352 Our findings could also contribute to the understanding of temper outbursts in PWS in general.
353 Several mental disorders are associated with disruptions of methylation patterns in *MAOA*.
354 Two studies showed lower methylation at the CpG sites in *MAOA* promoter as well as the exon
355 I /intron I region in female patients with an anxiety disorder (Domschke *et al.*, 2012; Ziegler *et*
356 *al.*, 2016), as well as in Acrophobia (Schiele *et al.*, 2018). Furthermore, several studies showed
357 lower methylation rates in female patients with depression (Melas *et al.*, 2013) and mixed
358 anxiety depression and dysthymia (Melas and Forsell, 2015). It was also shown that lower
359 methylation was associated with a higher score in Beck's Depression Inventory (Peng *et al.*,
360 2018). On the other hand, it was shown that higher methylation rates are found in conduct
361 disorder (Cecil *et al.*, 2018), Borderline personality disorder (Dammann *et al.*, 2011), PTSD
362 (Ziegler *et al.*, 2018) and schizophrenia (Chen *et al.*, 2012). It has been shown that temper
363 outbursts in PWS occur more frequently during cognitive challenges or at an unexpected
364 change of routines resulting in a cognitive overload with a loss of control (Woodcock, Oliver
365 and Humphreys, 2009; Woodcock *et al.*, 2010). Manning *et al.* showed that transcutaneous
366 vagus nerve stimulation is a promising treatment option for temper outbursts (Manning *et al.*,
367 2019). They propose that the rigidity in thinking leads to a flight or fight response to feeling
368 under threat associated with increased anxiety. This rising anxiety consequently serves as the
369 motor for the outbursts. We think that our findings, which are in line with those in patients

370 with anxiety or depression, support this hypothesis. Furthermore, our data support the
371 hypothesis that impulsiveness in PWS should not be seen in a link with mental disorders such
372 as conduct disorder or personality disorders.

373

374 5. Conclusion

375 In this study we investigated the methylation pattern of exon 1 and part of the promoter
376 region of *MAOA* in PWS. Overall, individuals in PWS who show temper outbursts have a
377 significantly lower methylation rate at the investigated CpGs than those PWS individuals who
378 do not have temper outbursts and healthy controls. This leads us to the conclusion that *MAOA*
379 is critically involved in the disturbance of the serotonergic pathway in PWS and contributes to
380 the high frequency of temper outbursts.

381

382 Acknowledgement

383 We thank Prof. Anthony J. Holland, University of Cambridge, for his expertise and comments
384 that greatly improved the manuscript.

385

386

387 References

388

389 Åkefeldt, A. *et al.* (1998) 'Cerebrospinal fluid monoamines in Prader-Willi syndrome',
390 *Biological Psychiatry*, 44(12), pp. 1321–1328. doi: 10.1016/S0006-3223(97)00519-2.

391 Åkefeldt, A. and Månsson, J. E. (1998) 'Is monoamine oxidase activity elevated in Prader-Willi
392 syndrome?', *European Child and Adolescent Psychiatry*, 7(3), pp. 163–165. doi:

393 10.1007/s007870050062.

394 Angulo, M. A., Butler, M. G. and Cataletto, M. E. (2015) 'Prader-Willi syndrome: A review of
395 clinical, genetic, and endocrine findings', *Journal of Endocrinological Investigation*. Springer
396 International Publishing, 38(12), pp. 1249–1263. doi: 10.1007/s40618-015-0312-9.

397 Bortolato, M., Chen, K. and Shih, J. C. (2008) 'Monoamine oxidase inactivation: From
398 pathophysiology to therapeutics', *Advanced Drug Delivery Reviews*, 60(13–14), pp. 1527–
399 1533. doi: 10.1016/j.addr.2008.06.002.

400 Brenet, F. *et al.* (2011) 'DNA methylation of the first exon is tightly linked to transcriptional
401 silencing', *PLoS ONE*, 6(1). doi: 10.1371/journal.pone.0014524.

- 402 Butler, M. G., Miller, J. L. and Forster, J. L. (2019) 'Prader-Willi Syndrome - Clinical Genetics,
403 Diagnosis and Treatment Approaches: An Update', *Current Pediatric Reviews*, 15(4), pp. 207–
404 244. doi: 10.2174/1573396315666190716120925.
- 405 Cassidy, S. B. *et al.* (2012) 'Prader-Willi syndrome', *Genetics in Medicine*, 14(1), pp. 10–26.
406 doi: 10.1038/gim.0b013e31822bead0.
- 407 Cecil, C. A. M. *et al.* (2018) 'Neonatal DNA methylation and early-onset conduct problems: A
408 genome-wide, prospective study', *Development and Psychopathology*, 30(2), pp. 383–397.
409 doi: 10.1017/S095457941700092X.
- 410 Chen, Y. *et al.* (2012) 'Effects of MAOA promoter methylation on susceptibility to paranoid
411 schizophrenia', *Human Genetics*, 131(7), pp. 1081–1087. doi: 10.1007/s00439-011-1131-5.
- 412 Dammann, G. *et al.* (2011) 'Increased DNA methylation of neuropsychiatric genes occurs in
413 borderline personality disorder View supplementary material', *Epigenetics*, 6, pp. 1454–
414 1462. doi: 10.4161/epi.6.12.18363.
- 415 Davidson, J. R. T. *et al.* (1982) 'Atypical Depression', *Archives of General Psychiatry*, 39(5), pp.
416 527–534. doi: 10.1001/archpsyc.1982.04290050015005.
- 417 Davies, J. R. *et al.* (2019) 'Prader-Willi syndrome imprinting centre deletion mice have
418 impaired baseline and 5-HT₂CR-mediated response inhibition', *Human Molecular Genetics*,
419 28(18), pp. 3013–3023. doi: 10.1093/hmg/ddz100.
- 420 Deest, M. *et al.* (2020) 'Sertraline as a treatment option for temper outbursts in Prader-Willi
421 syndrome', *American Journal of Medical Genetics, Part A*, (November), pp. 1–8. doi:
422 10.1002/ajmg.a.62041.
- 423 Domschke, K. *et al.* (2012) 'Monoamine oxidase A gene DNA hypomethylation—a risk factor
424 for panic disorder?', *International Journal of Neuropsychopharmacology*, 15(9), pp. 1217–
425 1228. doi: 10.1017/S146114571200020X.
- 426 Dykens, E. M. *et al.* (1992) 'Adaptive and Maladaptive Behavior in Prader-Willi Syndrome',
427 *Journal of the American Academy of Child and Adolescent Psychiatry*. The American Academy
428 of Child and Adolescent Psychiatry, 31(6), pp. 1131–1136. doi: 10.1097/00004583-
429 199211000-00023.
- 430 Dykens, E. M. *et al.* (2017) 'Diagnoses and characteristics of autism spectrum disorders in
431 children with Prader-Willi syndrome', *Journal of Neurodevelopmental Disorders*. *Journal of*
432 *Neurodevelopmental Disorders*, 9(1), pp. 1–12. doi: 10.1186/s11689-017-9200-2.
- 433 Dykens, E. M., Cassidy, S. B. and King, B. H. (1999) 'Maladaptive behavior differences in

- 434 Prader-Willi syndrome due to paternal deletion versus maternal uniparental disomy',
435 *American Journal on Mental Retardation*, 104(1), pp. 67–77. doi: 10.1352/0895-
436 8017(1999)104<0067:MBDIPS>2.0.CO;2.
- 437 Dykens, E. and Shah, B. (2003) 'Psychiatric Disorders in Prader-Willi Syndrome', *CNS Drugs*,
438 17(3), pp. 167–178. doi: 10.2165/00023210-200317030-00003.
- 439 Holland, A. J. *et al.* (2003) 'Behavioral phenotypes associated with specific genetic disorders:
440 Evidence from a population-based study of people with Prader-Willi syndrome',
441 *Psychological Medicine*, 33(1), pp. 141–153. doi: 10.1017/S0033291702006736.
- 442 Horsthemke, B. and Wagstaff, J. (2008) 'Mechanisms of imprinting of the Prader-
443 Willi/Angelman region', *American Journal of Medical Genetics, Part A*, 146(16), pp. 2041–
444 2052. doi: 10.1002/ajmg.a.32364.
- 445 Jade, H. *et al.* (2020) 'Re-assessment of the involvement of snord115 in the serotonin 2c
446 receptor pathway in a genetically relevant mouse model', *eLife*, 9, pp. 1–34. doi:
447 10.7554/eLife.60862.
- 448 Kishore, S. and Stamm, S. (2006) 'Regulation of alternative splicing by snoRNAs', *Cold Spring*
449 *Harbor Symposia on Quantitative Biology*, 71, pp. 329–334. doi: 10.1101/sqb.2006.71.024.
- 450 Lan, N. C. *et al.* (1989) 'Human monoamine oxidase A and B genes map to xp11.23 and are
451 deleted in a patient with norrie disease', *Genomics*, 4(4), pp. 552–559. doi: 10.1016/0888-
452 7543(89)90279-6.
- 453 Lewin, J. *et al.* (2004) 'Quantitative DNA methylation analysis based on four-dye trace data
454 from direct sequencing of PCR amplicates', *Bioinformatics*. Oxford Academic, 20(17), pp.
455 3005–3012. doi: 10.1093/bioinformatics/bth346.
- 456 Manning, K. E. *et al.* (2019) 'Transcutaneous vagus nerve stimulation (t-VNS): A novel
457 effective treatment for temper outbursts in adults with Prader-Willi Syndrome indicated by
458 results from a non-blind study', *PLoS one*, 14(12), p. e0223750. doi:
459 10.1371/journal.pone.0223750.
- 460 Manzardo, A. M. *et al.* (2018) 'Prader-Willi syndrome genetic subtypes and clinical
461 neuropsychiatric diagnoses in residential care adults', *Clinical Genetics*, 93(3), pp. 622–631.
462 doi: 10.1111/cge.13142.
- 463 Mazaheri, M. M. *et al.* (2013) 'The impact of Prader-Willi syndrome on the family's quality of
464 life and caregiving, and the unaffected siblings' psychosocial adjustment', *Journal of*
465 *Intellectual Disability Research*, 57(9), pp. 861–873. doi: 10.1111/j.1365-2788.2012.01634.x.

- 466 Melas, P. A. *et al.* (2013) 'Genetic and epigenetic associations of MAOA and NR3C1 with
467 depression and childhood adversities', *International Journal of Neuropsychopharmacology*,
468 16(7), pp. 1513–1528. doi: 10.1017/S1461145713000102.
- 469 Melas, P. A. and Forsell, Y. (2015) 'Hypomethylation of MAOA's first exon region in
470 depression: A replication study', *Psychiatry Research*, 226(1), pp. 389–391. doi:
471 10.1016/j.psychres.2015.01.003.
- 472 Nicholls, R. D. and Knepper, J. L. (2001) 'Genome Organization, Function, and Imprinting in
473 Prader-Willi and Angelman Syndromes', *Annual Review of Genomics and Human Genetics*,
474 2(1), pp. 153–175. doi: 10.1146/annurev.genom.2.1.153.
- 475 Palmer, E. E. *et al.* (2016) 'New insights into Brunner syndrome and potential for targeted
476 therapy', *Clinical Genetics*, 89(1), pp. 120–127. doi: 10.1111/cge.12589.
- 477 Peng, H. *et al.* (2018) 'Childhood Trauma, DNA Methylation of Stress-Related Genes, and
478 Depression', *Psychosomatic Medicine*. Lippincott Williams and Wilkins, 80(7), pp. 599–608.
479 doi: 10.1097/PSY.0000000000000604.
- 480 Rice, L. J., Woodcock, K. and Einfeld, S. L. (2018a) 'The characteristics of temper outbursts in
481 Prader–Willi syndrome', *American Journal of Medical Genetics, Part A*, 176(11), pp. 2292–
482 2300. doi: 10.1002/ajmg.a.40480.
- 483 Rice, L. J., Woodcock, K. and Einfeld, S. L. (2018b) 'The characteristics of temper outbursts in
484 Prader–Willi syndrome', *American Journal of Medical Genetics, Part A*, 176(11), pp. 2292–
485 2300. doi: 10.1002/ajmg.a.40480.
- 486 Schiele, M. A. *et al.* (2018) 'Plasticity of functional MAOA gene methylation in Acrophobia',
487 *International Journal of Neuropsychopharmacology*, 21(9), pp. 822–827. doi:
488 10.1093/ijnp/pyy050.
- 489 Shekim, W. O. *et al.* (1984) 'Psychoeducational and impulsivity correlates of platelet MAO in
490 normal children', *Psychiatry Research*, 11(2), pp. 99–106. doi: 10.1016/0165-1781(84)90092-
491 1.
- 492 Shih, C. J., Wu, J. B. and Chen, K. (2011) 'Transcriptional regulation and multiple functions of
493 MAO genes', *Journal of Neural Transmission*, 118(7), pp. 979–986. doi: 10.1007/s00702-010-
494 0562-9.
- 495 Shumay, E. *et al.* (2012) 'Evidence that the methylation state of the monoamine oxidase A
496 (MAOA) gene predicts brain activity of MAOA enzyme in healthy men', *Epigenetics*, 7(10),
497 pp. 1151–1160. doi: 10.4161/epi.21976.

- 498 Singmann, P. *et al.* (2015) 'Characterization of whole-genome autosomal differences of DNA
499 methylation between men and women', *Epigenetics and Chromatin*. BioMed Central, 8(1),
500 pp. 1–13. doi: 10.1186/s13072-015-0035-3.
- 501 Sinnema, M. *et al.* (2011) 'Behavioral phenotype in adults with Prader-Willi syndrome',
502 *Research in Developmental Disabilities*, 32(2), pp. 604–612. doi: 10.1016/j.ridd.2010.12.014.
- 503 Soni, S. *et al.* (2007) 'The course and outcome of psychiatric illness in people with Prader-
504 Willi syndrome: Implications for management and treatment', *Journal of Intellectual*
505 *Disability Research*, 51(1), pp. 32–42. doi: 10.1111/j.1365-2788.2006.00895.x.
- 506 Soni, S. *et al.* (2008) 'The phenomenology and diagnosis of psychiatric illness in people with
507 Prader-Willi syndrome', *Psychological Medicine*, 38(10), pp. 1505–1514. doi:
508 10.1017/S0033291707002504.
- 509 Vogels, A. *et al.* (2003) 'Chromosome 15 maternal uniparental disomy and psychosis in
510 Prader-Willi syndrome [7]', *Journal of Medical Genetics*, 40(1), pp. 72–73. doi:
511 10.1136/jmg.40.1.72.
- 512 Whittington, J. and Holland, A. (2018) 'A review of psychiatric conceptions of mental and
513 behavioural disorders in Prader-Willi syndrome.', *Neuroscience and biobehavioral reviews*.
514 Elsevier, 95(October), pp. 396–405. doi: 10.1016/j.neubiorev.2018.10.006.
- 515 Woodcock, K. A. *et al.* (2010) 'Neural correlates of task switching in paternal 15q11-q13
516 deletion Prader-Willi syndrome', *Brain Research*. Elsevier B.V., 1363, pp. 128–142. doi:
517 10.1016/j.brainres.2010.09.093.
- 518 Woodcock, K. A., Oliver, C. and Humphreys, G. (2009) 'Task-switching deficits and repetitive
519 behaviour in genetic neurodevelopmental disorders: Data from children with Prader - Willi
520 syndrome chromosome 15 q11 - q13 deletion and boys with Fragile X syndrome', *Cognitive*
521 *Neuropsychology*, 26(2), pp. 172–194. doi: 10.1080/02643290802685921.
- 522 Ziegler, C. *et al.* (2016) 'MAOA gene hypomethylation in panic disorder - Reversibility of an
523 epigenetic risk pattern by psychotherapy', *Translational Psychiatry*. Nature Publishing Group,
524 6(November 2015), pp. 1–8. doi: 10.1038/tp.2016.41.
- 525 Ziegler, C. *et al.* (2018) 'Monoamine Oxidase A Gene Methylation and Its Role in
526 Posttraumatic Stress Disorder: First Evidence from the South Eastern Europe (SEE)-PTSD
527 Study', *International Journal of Neuropsychopharmacology*, 21(5), pp. 423–432. doi:
528 10.1093/ijnp/pyx111.
- 529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548