Characterization of direct and/or indirect genetic associations for multiple traits in
longitudinal studies of disease progression

Myriam Brossard[1*], Andrew D. Paterson[2,3], Osvaldo Espin-Garcia[3,4], Radu V. Craiu[5], Shelley B.
Bull[1,3,*]

[1]Lunenfeld-Tanenbaum Research Institute, Sinai Health System, Toronto, ON, Canada;

[2]Program in Genetics and Genome Biology, Hospital for Sick Children Research Institute,
Toronto, ON, Canada;

[3]Division of Biostatistics, Dalla Lana School of Public Health, University of Toronto, Toronto, ON,
Canada;

[4] Department of Biostatistics, Princess Margaret Cancer Centre, Toronto, ON, Canada;

[5]Department of Statistical Sciences, University of Toronto, Toronto, ON, Canada.

Running title (40/40 characters): Joint models for multiple trait genetics.

Abstract (239/250)

When quantitative longitudinal traits are risk factors for disease progression, endogenous, and/or
subject to random errors, joint model specification of multiple time-to-event and multiple
longitudinal traits can effectively identify direct and/or indirect genetic association of single
nucleotide polymorphisms (SNPs) with time-to-event traits. Here, we present a joint model that
integrates: *i)* a linear mixed model describing the trajectory of each longitudinal trait as a function
of time, SNP effects and subject-specific random effects, and *ii)* a frailty Cox survival model that
depends on SNPs, longitudinal trajectory effects, and a subject-specific frailty term accounting for
unexplained dependency between time-to-event traits. Inference is based on a two-stage
approach with bootstrap joint covariance estimation. We develop a hypothesis testing procedure
to identify direct and/or indirect SNP association with each time-to-event trait. Motivated by
complex genetic architecture of type 1 diabetes complications (T1DC) observed in the Diabetes
Control and Complications Trial (DCCT), we show by realistic simulation study that joint modelling
of two time-to-T1DC (retinopathy, nephropathy) and two longitudinal risk factors (HbA1c, systolic
blood pressure) reduces bias and improves identification of direct and/or indirect SNP
associations, compared to alternative methods ignoring measurement errors in intermediate risk
factors. Through analysis of DCCT, we identify two SNPs with indirect associations with multiple
time-to-T1DC traits and obtain similar conclusions using alternative formulations of time-
dependent HbA1c effects on T1DC. In total, joint analysis of multiple longitudinal and multiple
time-to-event traits provides insight into aetiology of complex traits.

Key words: joint models; longitudinal study; direct and/or indirect genetic association; pleiotropy;
complex genetic architecture; multiple-trait analysis; measurement error; patient's trajectory;
mixed model; frailty model.

*****Corresponding authors:** Lunenfeld-Tanenbaum Research Institute, 60 Murray Street, Box #18,
M5T 3L9, Toronto, ON, Canada. E-mails: bull@lunenfeld.ca, brossard@lunenfeld.ca.
38

Introduction

Despite their known ability to improve inference in clinical and epidemiological studies, particularly in the presence of informative censoring/dropout or when longitudinal traits are measured with error[1–3], joint models for longitudinal and time-to-event outcomes have received limited attention in genetic association studies. Genome-wide association studies (GWAS) of intermediate quantitative traits (QTs) typically require follow-up studies to identify whether SNP associations detected with each of the QT(s), analyzed separately, also affect related outcomes through direct and/or indirect effects induced by those traits. Such intermediate traits may include established risk factors for related clinical outcomes and be measured with errors (e.g. biological variation). By accounting for QT measurement error and dependencies among traits, joint models can improve accuracy and efficiency of effect estimation, as well as detection of SNP associations.

The so-called shared-random-effects joint model[4,5] consists of a sub-model for a single longitudinal trait and a sub-model for a single right-censored time-to-event trait. The longitudinal sub-model describes the QT as an underlying smoothed trajectory that depends on fixed effects of time and baseline covariates, as well as subject-specific random effects. The joint model association structure is induced via the functional dependence between the hazard of an event at any time t and the longitudinal trait trajectory[6,7]. This relationship can be based on prior biological knowledge of the link between that longitudinal and time-to-event traits. As elucidated by Ibrahim et al[1], this class of joint models lends itself to interpretations of direct/indirect effects because the relationship between a baseline covariate, such as a SNP genotype, and each of the longitudinal and time-to-event traits, as well as the relationship between the longitudinal and time-to-event traits can be specified via model parameters corresponding to direct, indirect and overall effects. Extensions of joint models for multiple longitudinal and/or multiple time-to-event traits can further improve inference by borrowing information among related traits. Extensions of joint models have been reviewed for multiple longitudinal traits[6,7] or multiple time-to-event traits[8]. A few extensions for both multiple longitudinal and multiple time-to-event traits have been developed ([9–11], among others), but these models are often formulated for a specific study question, and thus can lack generalizability. Such extensions raise computational challenges for maximisation of the likelihood under the multivariate random effect's distribution. Two-stage approaches[12–15] for joint model fitting appear more computationally efficient and lends itself to more flexible model formulation but inference can be mis-calibrated because parameter estimates and predictions from Stage 1 are obtained from the longitudinal model without consideration of the time-to-event outcome, and the uncertainty in Stage 1 estimates is ignored during Stage 2 estimation, a problem known as propagation of errors[16].

Motivated by the complex genetic architecture of long-term type 1 diabetes complications (T1DC), we propose joint-model extension to evaluate genetic associations with multiple time-to-event traits and multiple longitudinal risk factors. Risk of development of T1DC, including diabetic retinopathy (DR) and diabetic nephropathy (DN), is hypothesized to result from multiple genetic factors with potential direct and/or indirect effects via multiple - shared and/or specific - risk factors[17]. In addition to genetic factors, hyperglycemia (measured by Hemoglobin A1c, referred as HbA1c) represents the major QT risk factor for T1DC. Other measured longitudinal traits, that can also be influenced by genetic factors, have postulated associations with T1DC, for example,

association of systolic blood pressure (SBP) with DN. Because conventional analysis methods in
GWAS ignores dependency between related traits, SNPs can appear associated with multiple
traits (including SNPs with direct and/or indirect effects induced by intermediate QT(s) measured
or unmeasured). Accurately distinguishing between direct and/or indirect associations is crucial
to elucidate the biological pathways through which genetic factors operate into the etiology of
those complex traits, like in our motivating example of the T1DC genetic architecture. A naïve
approach to disentangle direct from indirect associations can be achieved by including the
intermediate QT as a time-dependent covariate in a Cox Proportional Hazard (PH) model.
However, measurement errors in intermediate QT(s), presence of informative censoring in
longitudinal trait, unmeasured shared risk factors between longitudinal/time-to-event traits can
challenge accurate classification of direct and/or indirect SNP associations.

The *major* contribution of our work is a general formulation for the joint analysis of multiple time-
to-event traits and multiple longitudinal QT risk factors in genetic association studies. We develop
inference methods for statistical genetic analysis based on joint model parameters estimation,
including a hypothesis testing procedure to classify direct and/or indirect SNP associations with
each time-to-event trait. A *second* contribution of the paper is the implementation of a data-
informed simulation algorithm to generate multiple causal SNPs with various direct effects on
simulated time-to-event traits and/or indirect effects via observed (*measured*) longitudinal QTs in
the Diabetes Control and Complications Trial (DCCT) study[18,19] and unobserved (*simulated*)
longitudinal QTs. This algorithm also provides a general approach to estimate achievable power
given sample size. Our simulation results show bias reduction and improved classification of direct
and/or indirect SNP associations in comparison with alternative approaches ignoring
measurement errors in longitudinal QT(s). *Thirdly*, we apply the joint model in DCCT data and
clarifies two SNPs as having indirect associations via the HbA1c longitudinal risk factor, with
consistent conclusions using alternative time-dependent association structures that account for
established cumulative and time-weighted effects of HbA1c on T1DC traits[20,21]. Example R codes
for data simulation and for application of the proposed joint model are available on GitHub.

Methods

Joint Modelling Approach

Model Formulation

We assume that a set of *M* SNPs are available with *K* (*1 ≤ k ≤ K)* unordered and non-competing
time-to-event traits, such as multiple disease complications, and *L* (*1 ≤ l ≤ L)* longitudinal QT traits
(*i.e.* intermediate risk factors) possibly measured with error (*i.e.* biological variation) in *N* unrelated
individuals indexed by *i* (*1 ≤ i ≤ N*). For each SNP *m* (*1 ≤ m ≤ M*), with minor allele frequency (MAF)
p_m, the genotypes are coded as the number of copies of the minor allele, under an additive genetic
model. For each individual *i*, $Y_{i(l)} = (y_{i(l)}(t_{i1}), \dots y_{i(l)}(t_{ij}), \dots, y_{i(l)}(t_{in_{i(l)}}))'$ denotes the vector of
quantitative measures collected over scheduled visits at t_{ij} for *j*=1,…, n_i for the *l*th longitudinal trait
(1≤l≤ L) with n_i represents the maximum number of visits recorded. Let $(T_{i(k)}, \delta_{i(k)})$ be the vector
of observed right-censored event time $T_{i(k)}$ and event indicator $\delta_{i(k)}$ for the *k*th time-to-event trait.

We assume $T_{i(k)} = \min(T^*_{i(k)}, C_i)$, where $T^*_{i(k)}$ is the latent (uncensored) time-to-event k and C_i is

the censoring time (e.g. administrative censoring). We define $\delta_{i(k)} = I(T^*_{i(k)} \leq C_i)$, with $\delta_{i(k)} = 1$ if

the event occurs during the observation period ($T^*_{i(k)} \leq C_i$) and $\delta_{i(k)} = 0$ otherwise.

To characterize the genetic architecture of a system of *multiple* time-to-events and *multiple*
longitudinal risk factors, we formulate a shared-random-effects joint model, as specified in **Figure**
1, combining: ***(i) longitudinal*** and ***(ii) time-to-event sub-models*** connected by ***(iii) specified***
time-dependent association structures.

[Insertion of Figure 1]

(i)   **Longitudinal sub-model**

The *longitudinal sub-model* is specified by a *multivariate mixed model* for the *L* longitudinal QTs
(or risk factors), based on the multivariate extension of the Laird and Ware linear mixed model[22].
It describes each vector of *observed* longitudinal QT measures $Y_{i(l)}(t_{ij})$ as noisy observations of
a *true* and *unobserved smoothed* subject-specific longitudinal trajectory, $Y^*_{i(l)}(t_{ij})$, with noise
terms $\varepsilon_{ij(l)} \sim N(0, \sigma^2_{(l)})$. Each *smooth* trajectory describes subject-specific evolution and depends
on time, SNP effect, individual-level random effects $b_{i(l)}$ and can also include effects of other
baseline covariates, for example confounding factors or ancestry-related principal components.
To simplify the presentation, we assume a linear trajectory, but the longitudinal sub-models can
be adapted for nonlinear trajectories using, for example, higher order polynomials or splines[23].

Multivariate mixed model

$Y_{i(l)}(t_{ij}) = Y^*_{i(l)}(t_{ij}) + \varepsilon_{ij(l)}$ (*observed* trait values, Equation 1)

$Y^*_{i(l)}(t_{ij}) = \beta_{0(l)} + b_{i0(l)} + (\beta_{1(l)} + b_{i1(l)})t_{ij} + \beta_{g(l)}SNP_i + \beta_{h(l)}H_{i(l)}$

(*smooth* trajectory, Equation 2)

Where:

- $\beta_{0(l)}, \beta_{1(l)}, \beta_{g(l)}$ and $\beta_{h(l)}$ denote, respectively, the fixed intercept, and baseline effects of
time, SNP and other covariate(s) on each longitudinal trait *l*.
- $b_i = ((b_{i0(l)}, b_{i1(l)})', \ldots, (b_{i0(L)}, b_{i1(L)})')'$ is the 2x*L* vector of random effects with
$b_i \sim N_{2L}(0, D)$.
- $D = \begin{pmatrix} D_{1,1} & \cdots & D_{1,L} \\ \vdots & \ddots & \vdots \\ D_{L,1} & \cdots & D_{L,L} \end{pmatrix}$ denotes the *overall* variance-covariance matrix for random effects,
accounting for serial dependencies *within* longitudinal traits ($D_{l,l}$, unstructured) and cross-
dependencies *between* longitudinal traits ($D_{l,m}$ for two traits $l \neq m$).

• $\varepsilon_{i(l)} = \left(\varepsilon_{i1(l)}, \dots, \varepsilon_{ij(l)}, \dots, \varepsilon_{in_i(l)}\right)'$ is the vector of error terms where $\varepsilon_{ij(l)} \sim N\left(0, \sigma^2_{(l)}\right)$ is
assumed independent and identically distributed.
• $\sigma^2_{(l)}$ is the residual variance of the QT *l*. We further assume that $\varepsilon_{ij(l)}$ and $b_{i(l)}$ are
independent[22].

(ii)   Time-to-event sub-model

The *time-to-event sub-model* is specified by a Proportional Hazards (PH) mixed effects model
(also known as frailty PH model) where the hazard function of each time-to-event trait *k* depends
on the SNP effect adjusted for a function of the subject's longitudinal trajectories $W_{i(k)}(t)$. We
introduce a subject-specific random effect (frailty term, u_i) to capture unexplained dependencies
(e.g. due to unmeasured shared factors) among the *K* time-to-event traits. Given the common
frailty term u_i, the *K* time-to-event traits are assumed independent[24].

Proportional Hazards mixed effects sub-model

$$\lambda_{i(k)}(t) = \lambda_{0(k)}(t) \times exp\{\gamma_{g(k)}SNP_i + W_{i(k)}(t) + \gamma_{v(k)}V_{i(k)} + u_i\} \qquad \text{(Equation 3)}$$

Where:

• $\lambda_{0(k)}(t)$ is a (parametric or non-parametric) baseline hazard function for the time-to-event
trait *k.*
• $\gamma_{g(k)}$ denotes the SNP effect on the time-to-event *k* adjusted for the indirect SNP effects
via longitudinal QT(s), taken into account by $W_{i(k)}(t)$.
• $\gamma_{v(k)}$ is the effect of the baseline covariate vector(s). Those covariates can be shared or
trait-specific covariates. They can also be time-dependent covariates for all or a subset of
the *K* time-to-event traits (in this case, $V_{i(k)} = V_{i(k)}(t)$).
• u_i and b_i, random effects from the time-to-event and longitudinal sub-models respectively,
are assumed independent. We assume $u_i \sim \Gamma(a, b)$, with *a*, *b*>0.

(iii)   Time-dependent association structures

The *time-dependent association structures*, $W_{i(k)}(t)$, that connect the *longitudinal* and *time-to-*
event sub-models (**Figure 1**), account for the specific temporal relationships between *each* set of
L_k $(1 \leq L_k \leq L)$ associated QT(s) and *each* time-to-event trait *k.*

$$W_{i(k)}(t) = \sum_{l=1}^{L_k} \alpha_{l(k)} f_{l(k)}(Y^*_{i(l)}(t)) \qquad\qquad \text{(Equation 4)}$$

Where:

• $1 \leq L_k \leq L$ is the set of longitudinal risk factors of the time-to-event trait *k.*
• $f_{l(k)}(Y^*_{i(l)}(t))$ denotes the *functional* form of the exposure effect on time-to-event trait *k*
corresponding to the *l*-th longitudinal risk. In the case of a *contemporaneous*
parametrization, the hazard of an event *k* at a time *t* depends on the longitudinal trait value

at the same time t (i.e. $f_{l(k)}\left(Y^*_{i(l)}(t)\right) = Y^*_{i(l)}(t)$). Other parametrizations have been

described in the literature[6,7,25].

• $\alpha_{l(k)}$ denotes the association parameter of the function of the *smooth* longitudinal

trajectory $Y^*_{i(l)}(t)$ with the time-to-event *k*.

• $Y^*_{i(l)}(t) = \{Y^*_{i(l)}(s), with\ 0 \le s \le t, 1 \le l \le L\}$ denotes the history of the underlying

longitudinal trajectory for trait *l* up to time *t.*

Interpretation

When the joint model is *correctly* specified, the parameters accurately represent multiple

relationships between each SNP and the correlated traits. Note that $\beta_{g(l)}$ is the SNP effect on the

longitudinal risk factor *l*, $\gamma_{g(k)}$ is the *direct* SNP effect on the time-to-event trait *k* via other

mechanisms than the indirect SNP association induced by the observed longitudinal risk factors.

In this context, $\theta_{k(l)} = \mu_{g(k,l)} + \gamma_{g(k)}$ is interpreted as the *overall* SNP effect on the time-to-event

trait *k,* where $\mu_{g(k,l)} = \alpha_{l(k)}\beta_{g(l)}$ represents the *indirect*[1] SNP effect on the time-to-event trait *k* via

the longitudinal QT *l.*

Under the proposed joint model formulation, when a longitudinal risk factor *l* is *associated* with a

time-to-event trait *k* ($\alpha_{l(k)} \ne 0$), a SNP that has an effect on the longitudinal QT risk factor *l* ($\beta_{g(l)} \ne$

0), but no effect on the time-to-event trait *k* ($\gamma_{g(k)} = 0$)*,* is interpreted as having an **indirect SNP**

association and its *overall* effect is equal to its *indirect* effect ($\theta_{k(l)} = \mu_{g(k,l)}$, with $\mu_{g(k,l)} =$

$\alpha_{l(k)}\beta_{g(l)}$). A SNP with an effect on the time-to-event trait *k* ($\gamma_{g(k)} \ne 0$)*,* but no effect on the

longitudinal QT *l* ($\beta_{g(l)} = 0$), is interpreted as having a **direct SNP association** and its *overall*

effect is equal to its *direct* effect ($\theta_{k(l)} = \gamma_{g(k)}$)*.* A SNP with an effect on the longitudinal risk factor

l ($\beta_{g(l)} \ne 0$) and an effect on the time-to-event trait *k* ($\gamma_{g(k)} \ne 0$) is interpreted as having both

direct and **indirect SNP associations** via distinct genetic pathways. In this case, its *overall* effect

is an aggregation of both *direct* and *indirect* effects ($\theta_{k(l)} = \mu_{g(k,l)} + \gamma_{g(k)}$, with $\mu_{g(k,l)} = \alpha_{l(k)}\beta_{g(l)}$)*.*

It is obvious that when an associated longitudinal risk factor is omitted from a time-to-event model,

as occurs in separate analyses of longitudinal and time-to-event traits, the marginal SNP effect

on the time-to-event trait will reflect partially the overall SNP effect $\theta_{k(l)}$. This is particularly the

case in GWAS when intermediate risk factors are ignored in the analysis models.

Effect estimation and test statistic construction

To mitigate the computational complexity involved in the maximization of the joint likelihood and

allow more flexibility in the model formulated, we estimate the parameters using a two-stage

approach (see **Appendix**). We work within the framework defined by Tsiatis and Davidian[26,27]

which guarantees that our estimators are consistent and asymptotically normal. Specifically, in

Stage 1 we fit the multivariate mixed model (Equation 1) using the *mvlme*() function from the R

package JoineRML[28] to estimate the parameters of the longitudinal trajectories of the risk factors

(Equation 2), and obtain fitted values of the smoothed trajectories. In Stage 2, we fit a Cox PH

frailty time-to-event model (Equation 3) adjusting for functions of the smoothed trajectories as

time-dependent covariates using the *coxph*() function from the R survival[29,30] package. We
assume a Gamma distribution for the frailty term (u_i) and a separate baseline hazard function for
each time-to-event trait using the *strata* argument in *coxph*(). To account for propagation of errors,
due to the uncertainty in Stage 1 estimates ignored during Stage 2 parameter estimation[16] and
empirically estimate the joint covariance matrix of SNP and trajectories effects we apply a
nonparametric bootstrap[31]. For each bootstrap sample *b* (1 ≤ *b* ≤ *B, B* is the total number of
bootstraps), we generate a new dataset by randomly sampling *N* individuals with replacement
and refitting the joint model on each new dataset *b*. We compute the empirical joint covariance
matrix of the estimated parameters using the *B* bootstrap parameter estimates. Wald statistics for
each $\beta_{g(l)}$ are computed as $S_{\beta_{g(l)}} = \widehat{\beta_{g(l)}}/se_{\beta_{g(l)}}$ using the empirical bootstrap standard errors
$se_{\beta_{g(l)}}$, and similarly for each $\gamma_{g(k)}$ as $S_{\gamma_{g(k)}} = \widehat{\gamma_{g(k)}}/se_{\gamma_{g(k)}}$.

Procedure to classify direct/indirect SNP associations

In **Table 1**, we present a practical hypothesis testing procedure to classify SNPs as having direct
and/or indirect association with a time-to-event trait *k* and each associated longitudinal risk factor
l. This procedure requires two significance thresholds, $P^*_{\beta_g}$ and $P^*_{\gamma_g}$ for $\beta_{g(l)}$ and $\gamma_{g(k)}$ respectively,
to be specified prior to the analysis and adjusted for the number of SNPs tested. Depending on
the research question, we can choose different values for P^*_β and P^*_γ, or the same one ($P^* = P^*_{\beta_g} =$
$P^*_{\gamma_g}$). The latter is applicable for instance, when we want to systematically classify direct/indirect
association among a set of *M* SNPs, and the former when assessing which SNPs, among the
ones reported to be associated with the longitudinal risk factor, have a direct effect on a time-to-
event trait.

[Insertion of Table 1]

Motivation: The DCCT Studies

The DCCT randomized-controlled trial was *pivotal* in demonstrating that intensive insulin therapy
prevents and delays progression of long-term T1DC[18]. Due to significant outcome differences
between the intensive and conventional treatment groups at interim analysis, the DCCT trial was
stopped early. At the closeout visit, patients were administratively censored and had a mean
follow-up time of 6.5 years (range 3 to 9), 99% had completed the study, and more than 95% of
all scheduled examinations were completed[18]. DCCT participants continued to be followed for
disease progression after the trial in a long-term epidemiologic cohort study and were genotyped
for GWAS in the DCCT Genetics Study. Because the goal of intensive therapy was to reduce
HbA1c into the non-diabetic range, which produced treatment differences in HbA1c values, we
focus on *N*=667 unrelated individuals of European descent ancestry from the Conventional
treatment group. Longitudinal measurements for HbA1c and SBP were collected at up to 39
quarterly visits during DCCT, while DR and DN events were diagnosed at annual and semi-annual
visits respectively. HbA1c and SBP were recorded irrespective of the occurrence of any
complication event(s). Although the DCCT implemented robust quality assurance procedures to
minimize potential sources of error during and after data collection[32,33], HbA1c and SBP are

subject to measurement error (e.g. individual variability). The first GWAS to analyze the DCCT
study phenotypes[19] identified two SNPs associated with cumulative HbA1c at genome-wide
significance in the Conventional treatment arm, namely rs10810632 (in *BNC2*, 9p22.2) and
rs1358030 (near *SORCS1,* 10q25.1), and reported consistent associations with secondary
outcomes of time-to-DR and/or time-to-DN. Subsequent association studies also reported
suggestive evidence for pleiotropy between DR and DN[34]. To preserve the inherent within-
individual variability, we designed simulation study evaluations of the joint modelling approach
that generates event times from HbA1c and SBP measurements in the DCCT Study data.

Simulation study

Design of the DCCT-data-based simulation study

To assess the performances of the methods, we simulate the data under a *genetic causal*
scenario that imitates the complex genetic architecture of T1DC (**Figure 2**). The genetic
association model involves *N*=667 DCCT subjects with *K*=2 non-independent *simulated* time-to-
T1DC traits (DR, DN) that depend on *M*=5 causal SNPs with *direct* effects on each time-to-T1DC
and/or *indirect* effects via *L*=3 longitudinal QTs: two as *measured* in DCCT (HbA1c, SBP) and
one other *simulated* (*U*) to induce some unexplained shared dependency between the T1DC
traits. We also introduce an effect of sex on SBP, and effects of T1D duration (at baseline) on
both T1DC traits, as observed in the original DCCT data.

[Insertion of Figure 2]

Algorithm for realistic data generation under a complex genetic architecture

To generate such a data structure that combines *observed* and *simulated* traits, we formulate a
data generating model including: *(i) L=3* linear mixed models linking each of the SNPs with an
indirect effect to each longitudinal risk factor, and *(ii) K=2* non-independent parametric time-to-
event models depending on SNPs with *direct* effects and on functions of the longitudinal QT
trajectories. For each DCCT individual *i* with observed HbA1c ($Y_{i(1)}$), SBP ($Y_{i(2)}$), visit times (t_i)
vectors at n_i time points, and covariates ($H_{i(l)}, V_{i(k)}$), we simulate longitudinal trait vector U_i, time-
to-event traits and genetic data at *M* causal SNPs as illustrated in **Figure 3** and detailed in
Supplementary Information SI-1. All SNPs are simulated under Hardy-Weinberg and linkage
equilibrium assumptions. Particularly, SNPs with indirect effects are simulated from the *observed*
(SNP1, SNP5) or *simulated* (SNP3) longitudinal QTs, while SNPs with direct effects (SNP2,
SNP4) on time-to-T1DC traits are simulated independently of the longitudinal QTs and are
included in the specified hazard function used to simulate each time-to-event trait *k* (**Figure 3**).

[Insertion of Figure 3]

For each SNP*m*, we specify MAF (p_m) and the SNP effects ($\beta_{g(l)}, \gamma_{g(k)}$), as well as the other
parameter values according to the DCCT Genetics Study and the T1DC literature (**Table 2**,

Tables S1). We assume contemporaneous association structures for HbA1c and SBP effects on
simulated time-to-T1DC traits. We simulate each time-to-event trait k under a Weibull model, with
shape and scale parameters specified to generate ~54% DR events and ~25% DN events in each
of the R=1000 replicated datasets generated under the *causal* scenario from **Figure 2**. Under the
global null genetic scenario, where none of the SNPs is associated with any traits, we simulate M
SNPs independently of the traits with the same MAF as for the causal SNPs.

Scenarios for DCCT-based complex genetic architecture

Overall, the simulated complex genetic architecture covers multiple types of SNP-trait
associations (**Figure 2** and **Table 2**): *direct* association with each T1DC trait (SNP2, SNP4),
indirect associations with both T1DC traits via *measured* (SNP1) and *unmeasured* (SNP3)
longitudinal QTs; and *direct and indirect* associations via a *measured* longitudinal QT (SNP5); all
longitudinal risk factors are subject to measurement error. Except for SNP3, all other SNP
scenarios represent GWAS discovery SNPs detectable by separate GWAS of a longitudinal risk
factor (SNP1, SNP5) or a time-to-event trait (SNP2, SNP4, SNP5).

[Insertion of Table 2]

SNP1, SNP3 and SNP5 have indirect effects on T1DC traits, such that their associations with the
T1DC traits are detectable with the marginal Cox PH time-to-event models (**Tables 2** and **S2**).
SNP1 corresponds roughly to rs10810632 and rs1358030 reported in the motivating DCCT
GWAS of HbA1c[19], while SNP5 represents a top hit detected by each GWAS (longitudinal and
time-to-event traits), analyzed separately. Because of their association with longitudinal and time-
to-event traits, SNP1 and SNP5 associations with time-to-event outcomes are conventionally
investigated using the Cox PH model adjusting for the *observed* longitudinal trait as a time-
dependent risk factor to determine if SNP association with the time-to-event trait is fully (or
partially) explained via the associated longitudinal QT(s).

Analysis of the simulated data

For the analysis of *each* SNP, done *separately*, we compare four analysis models on the
simulated data:

•   JM-cmp: a *completely* specified joint model that includes all other non-SNP variables used
for the data simulation.
•   JM-mis: includes the same variables as JM-cmp, except U, the unobserved longitudinal
QT.
•   JM-sep(k): includes the same variables as JM-mis but is fitted separately for the *two* time-
to-event traits. JM-sep(k) does not account for the shared dependency between the time-
to-event traits.
•   CM-obs: a Cox PH frailty model that includes the same variables as JM-mis, but adjusts
for the *observed* longitudinal QT values as time-dependent covariates.

Due to the latent nature of *U, JM-cmp* cannot be fitted in practice because *U* is unobserved, but
we include it as a benchmark for comparison against the *observable* models fitted without *U*. For
all these models, empirical covariance matrices are computed using 500 bootstrap iterations. We
assess the performance of the hypothesis testing procedure to classify direct/indirect association
for each of the 5 causal SNPs analyzed separately with each joint model (JM-cmp, JM-mis). To
assess the impact of longitudinal trait measurement errors on classification of SNPs, we substitute
test $P_{\gamma_{g(k)}}$ for $\gamma_{g(k)}$ estimated by the joint model by $P_{\gamma_{g(k)}}$ obtained with CM-obs. In our evaluations,
we assume the level of significance P^* such that $P^* = P^*_\beta = P^*_\gamma$, with P^* varying from 0.05 to 5x10^{-}
^{8}. Under the *causal* genetic scenario, we estimate for each SNP the proportion of replicates that
detect the correct direct and/or indirect association simulated (correct classification rate) and the
proportion of replicates where the causal SNP association is misclassified (misclassification rate).
Under the *global null* genetic scenario, we assess the classification test for each SNP and each
type of association at $P^* = 0.05$ and $P^* = 0.01$ to assess the type I error control of the hypothesis
testing procedure for each test of direct and/or indirect association.

Availability

DCCT data are available to authorized users at https://repository.niddk.nih.gov/studies/edic/
and https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000086.v3.p1.
Example R codes for DCCT-data-based simulation and analysis of the simulated data are
provided on GitHub (https://github.com/brossardMyriam/Joint-model-for-multiple-trait-genetics).
File S1 includes the Supplementary Information; File S2 includes the Supplementary Figures and
File S3 includes the Supplementary Tables.

Results

Simulation results

Estimation accuracy and single-parameter tests validity

When the longitudinal QT(s) are observed and measured with random *errors*, the simulation
results confirm that the proposed joint model reduces the bias of the parameter estimates for all
types of SNP association (**Figure S1** and **Tables S3-S6**), even when the analysis model is not
fully specified. This applies when: *(i)* two correlated time-to-event traits are analyzed *jointly* rather
than *separately* in the joint model (JM-cmp/JM-mis compared to JM-sep(*k*)); *(ii)* a SNP has a
direct effect on a time-to-event trait with a low event rate (SNP4/SNP5, JM-cmp/JM-mis compared
to CM-obs); *(iii)* a SNP has an *indirect* effect via a longitudinal QT measured with errors (for
example SNP5, JM-cmp/JM-mis compared to CM-obs). Particularly, for SNP5, CM-obs
overestimates the *direct* SNP5 effect on time-to-DN due to its inability to fully account for the
indirect SNP5 effect induced via SBP. Direct SNP2 and SNP4 effects on time-to-T1DC traits show
overall the larger relative bias reduction using joint models versus CM-obs (Ranges with JM-mis
0.702-0.983, with JM-cmp 0.934-0.966, see **Tables S4 and S6**)**.**

When a SNP has an indirect effect on both T1DC traits fully explained via an *unmeasured*
longitudinal QT (SNP3), all methods that ignore the longitudinal QT (JM-mis, JM-sep(*k*) or CM-
obs compared to JM-cmp) produce biased direct SNP3 estimates for both T1DC towards the
overall SNP3 effect ($\theta_{k(U)} = \alpha_{U(k)}\beta_{g(U)}$ =0.36, **Figure S1-c** and **S1-d, Table S5**). In this scenario,
the use of the frailty term in JM-mis does not reduce this bias (JM-mis *vs* JM-sep).

Classification of direct/indirect SNP associations with time-to-event traits

Under the *global null* genetic scenario of no genetic association with any of the traits, single-
parameter SNP test *P*-values $P_{\beta_{g(l)}}$ and $P_{\gamma_{g(k)}}$ from the joint model do not show departure from the
expected large sample distributions (χ^2 with 1 degree of freedom (*df)*, see **Figure S2**) and the
type I error of each test under the null is reasonably well controlled (**Table S7**). The hypothesis
testing procedure applied to single-parameter SNP tests, classifies each SNP as direct or indirect
association with each T1DC trait with rates close to the nominal level (**Table 3** and **Figure S3**).
However, the classification rates for test of *direct and indirect* SNP association test appears less
than nominal (see **Table 3** for SNP5 with SBP/DN traits and **Figure S3** for the other SNPs with
other QT/time-to-event trait pairs).

[Insertion of Table 3]

Under the *causal* scenario, the hypothesis testing procedure has high correct classification rates
for *indirect* (SNP1) or *direct* associations (SNP2, SNP4), as shown in **Figures 4-a, 4-b** and **S4**.
Conclusions are similar for JM-mis, with attenuated correct classification rates compared to JM-
cmp for direct SNP associations but still higher than classification rates based on CM-obs
(particularly for SNP4). Compared to results based on CM-obs, the largest relative classification
improvement is found for direct SNP4 association with DN (ranges: 1.18-251.9 with JM-cmp and
1.18-110.98 with JM-mis, **Figure 4**).

[Insertion of Figure 4]

On the other hand, SNP3 with an *indirect* effect via the unobserved longitudinal trait *U*, is most
frequently detected incorrectly as a *direct* association with each T1DC trait by both JM-mis and
CM-obs (**Figure 4-b**). SNP5, which has both *direct* and *indirect* effects on DN induced via SBP,
is most frequently misclassified by JM-cmp and JM-mis as an *indirect* association at significance
levels varying from $P^* = 0.01$ to 5x10⁻⁸ (**Figure 4-c** and **4-d**). SNP5 misclassification rate even
increases with P^* decreasing to 5x10⁻⁸, due to the direct SNP5 $P_{\gamma_{g(k)}}$. In comparison, the
procedure based on CM-obs results most frequently correctly classify SNP5 association in more
than 75% of the replicates at the same significance levels (**Figures 4-c** and **4-d**); this is explained
by the overestimated direct SNP5 effect, arising from measurement error in SBP as noted before.

In summary, our simulations show that by accounting for measurement errors in the longitudinal
QT risk factors and for dependencies between/within the traits, the proposed joint model improves
estimation accuracy and correct classification of associations of SNPs *directly* associated with

each time-to-event trait or *indirectly* associated via a measured longitudinal QT in comparison to
classification using the CM-obs approach. However, when a SNP has both *direct* and *indirect*
effects on a time-to-event trait, the proposed testing procedure can be conservative since it
requires the joint significance of the two SNP effects, $\beta_{g(l)}$ and $\gamma_{g(k)}$, where the power of each
test depends on effect size and the trait distribution. As a result, a SNP with a direct and an indirect
association can be misclassified as either a direct or an indirect association. Finally, when a SNP
has an indirect effect on both T1DC traits via an unmeasured QT, the testing procedure based on
JM-mis, that captures some of the unexplained dependency between time-to-event traits through
the frailty term, does not prevent misclassification as a direct association. This observation also
shows the importance of the correct specification of the joint model - including all the intermediate
QT(s) - to avoid misclassification of direct and/or indirect SNP associations.

Application in the DCCT Genetics Study data

We demonstrate the feasibility of the proposed approach by an application in the DCCT Genetics
Study data. We use time to mild DR and time to persistent microalbuminuria, for DR and DN
outcomes respectively, as previously defined in the motivating GWAS of HbA1c[19] (see
Supplementary information SI-2 for details). Genome-wide genotyping in DCCT subjects was
performed using HumanCoreExome Bead Array (Illumina, San Diego, CA, USA) and standard
quality controls procedures were applied to individuals and genetic markers[19,35]. Ungenotyped
autosomal SNPs were imputed using 1000 Genomes data phase 3[36] (v5) and minimac3[37]
(v.1.0.13), as previously described[35].

Out of the 667 DCCT individuals, we analyze *N*=516 subjects with genetic data, without mild to
moderate non-proliferative retinopathy or without DN event at DCCT baseline. By the time of the
DCCT close-out visit, 297 (57.6%) experienced a DR event, 61 (11.8%) a DN event, including 47
subjects (9.1%) that experienced both events. After SNP filtering and pruning on linkage
disequilibrium (see **Supplementary Information SI-2** for details), we analyze 307 candidate
SNPs reported as associated with HbA1c, SBP, and multiple definitions of DR and/or DN[19,34,38–42]
(see **Table S8** for the full list of SNPs).

Compared to the DCCT-based simulated datasets, the proportion of DN events observed in the
application is lower. We expect that the test power for direct SNP association with DN under the
joint model to be reduced; this implies reduce chances to correctly classify a SNP with either a
direct or a *direct and indirect* association with DN. We also observe larger effects of HbA1c on
T1DC traits ($\alpha_{1(k)}$, *k*=1, 2) and a smaller effect of SBP on DN ($\alpha_{2(2)}$), as shown in **Tables 2 vs**
S9. Larger $\alpha_{l(k)}$ values increase the contribution of the indirect effect ($\mu_{g(l,k)}$) to the overall SNP
effect ($\theta_{k(l)}$), which can result in a direct SNP effect ($\gamma_{g(k)}$) more severely biased if the longitudinal
QT is not accounted properly in the joint model, while lower $\alpha_{l(k)}$ values reduce $\mu_{g(l,k)}$ and thus
$\gamma_{g(k)}$ is less severely biased. Overall, in the DCCT application, we expect a SNP with an *indirect*
effect via HbA1c on any T1DC trait to be more subject to be misclassified as having both a *direct*
and *indirect* association, compared to the simulation results, while a SNP with an *indirect* effect
via SBP on DN is less subject to this misclassification.

Given prior evidence for cumulative effects of HbA1c on T1DC traits[20,21], we compare joint model
results obtained with contemporaneous, updated cumulative mean, and time-weighted
cumulative HbA1c effects on T1DC (**Supplementary Information SI-2** for details). We present
results under the latter association structure since it has a stronger prior association with T1DC
and in the DCCT individuals analyzed here (**Table S9**). We obtain similar conclusions with
alternative association structures (**Figure S5-a**). As shown in **Figure 5-a**, rs10810632 and
rs1358030 are classified as indirect associations with both T1DC traits via their association with
HbA1c shared risk factor ($P_{\beta_{g(l)}} \leq P^*$ and $P_{\gamma_{g(k)}} > P^*$, P^*=1.7x10⁻⁴ after Bonferroni correction for
the 289.02 effective SNPs tested[43]). These two SNPs have calculated indirect effects ($\mu_{g(1,k)}$)
and 95% bootstrap confidence intervals that do not include 0 (**Figure 5-b**). As noted before,
rs10810632 and rs1358030 were discovered previously in a GWAS of HbA1c in DCCT[19].
Classification of these two SNPs as direct and/or indirect associations may be affected by the
Winner's curse bias, due to the modest sample size in this analysis, and the same individuals
contributing to discovery and classification[44,45]. For the other candidate SNPs which were
discovered in larger and/or independent studies[19,34,38–42], we find little evidence of association with
the discovery traits in marginal analyses (**Table S8**).

Conclusions regarding classification of direct and/or indirect associations based on the procedure
using $P_{\gamma_{g(k)}}$ from CM-obs are similar to those based on the joint model results for all SNPs (**Figure
S5-b**), although we noticed some differences in $P_{\gamma_{g(k)}}$; that could be explained by the attenuated
association of HbA1c and SBP with T1DC traits due to measurement errors ignored from CM-obs
(**Tables S10 *vs* S9**).

Discussion

We present new methods for statistical genetic analysis under a joint model specification of
multiple time-to-event traits and multiple longitudinal risk factors designed to characterize the
complex genetic architecture of related traits in longitudinal studies of disease progression. The
proposed model accounts for dependencies within and between traits and can account for trait-
specific and shared covariates, measurement errors in the longitudinal traits, as well as effects of
unobserved baseline confounding factors between the time-to-event traits through the subject-
specific frailty term.

Evaluation by *realistic* data-informed simulation study of complex T1DC genetic architecture
shows that the proposed joint model improves estimation accuracy and efficiency compared to
the Cox PH frailty model adjusted for observed longitudinal trait values as time-dependent
covariates. This improvement holds particularly when the longitudinal risk factors are measured
with errors or the time-to-event outcome has a low event rate. Indeed, measurement error
attenuates the association of the longitudinal risk factor with the time-to-event trait, and as a result,
if a SNP has an indirect effect induced by the QT, the estimated direct SNP effect on a time-to-
event is biased away from zero. The magnitude of this bias tends to be worse with larger
measurement error[14,15]. Identification of direct and/or indirect SNP associations is also facilitated
under joint modelling in the presence of risk factor measurement error. When, however, a SNP
has an indirect effect via an unmeasured longitudinal risk factor, both the proposed and Cox PH

model approaches produce biased estimates of the direct SNP effects. The extent of this bias
depends on the magnitude of the effect sizes $\beta_{g(l)}$ and $\alpha_{l(k)}$. Although the frailty term in the joint
model time-to-event sub-model captures some of the unexplained shared dependency between
the time-to-event traits, we did not observe bias reduction of the direct SNP estimate in the case
of an unmeasured risk factor, which can lead to misclassification of the SNP as a direct
association. This may be a limitation of using a time-invariant frailty term that does not fully
account for the indirect SNP effect via the unmeasured longitudinal risk factor. Because joint
significance of both SNP effects is required to classify a SNP as having both direct and indirect
associations, a SNP with both direct and an indirect association can be misclassified as having
either a direct or an indirect association if one of the SNP effects is too small to be detected at
the specified level. The proposed procedure can be adapted to other research questions, for
example to identify SNPs that have direct association among the SNPs found associated with a
longitudinal risk factor. In this case, it may be desirable to apply a less stringent significance
threshold for the test of the direct SNP effect while controlling the overall type I error. Alternatively,
intersection-union tests, as based on the likelihood ratio test[46], could be used to assess joint
significance of $\beta_{g(l)}$ and $\gamma_{g(k)}$ but would require computation of the full joint likelihood.

Application of the proposed joint model approach in longitudinal studies of disease progression,
such as in the DCCT Genetics Study, improves classification of direct and/or indirect SNP
association which can help to elucidate the genetic architecture of complex traits. In the context
of mediation analysis, Liu et al[47] discussed various formulations and interpretations of joint
models, with shared-random-effects accounting for potential unmeasured baseline confounding
factors between *one* longitudinal and *one* time-to-event traits. Using applications in datasets from
two clinical trials, they illustrate interpretation of sensitivity analysis to unmeasured baseline
confounders. Adaptation of the joint model we propose for multiple longitudinal and multiple time-
to-event traits for mediation analysis requires extension of the mediation assumptions[48,49] to the
case of multiple mediators and multiple time-to-event traits. Specific evaluations of the proposed
model under these assumptions are also warranted.

Although our primary aim in this report is to develop statistical methods to accurately distinguish
among direct and/or indirect SNP associations with each time-to-event trait, the multi-trait aspect
of the joint model lends itself to development of multi-trait SNP association testing for SNP
discovery. In **Supplementary Information SI-3**, we present a joint-parameter test based on a
generalized Wald statistic. In application to the simulated DCCT-based complex genetic
architecture, we observe good type I error control under the global genetic null scenario, improved
power for SNP discovery when a SNP has multiple trait effects, and power maintenance in other
SNP association scenarios.

We acknowledge several features of the proposed joint model approach that warrant examination
in further work. *Firstly*, to reduce computational complexity and improve model flexibility, we used
two-stage parameter estimation. In some circumstances, this approach can produce biased
estimates and/or underestimated standard errors[16]. Biased estimates can result from non-random
censoring of the longitudinal trait values due to the occurrence of an event or from informative
dropout[50,51]. Our simulation results show minimal biases in the absence of informative censoring,

even when the joint model is mis-specified. In the DCCT application, characterized by
administrative censoring and a high completion rate, these biases are of minimal concern
because longitudinal trait values continued to be recorded regardless of the occurrence of any
T1DC events; we estimated the trajectories using all the available measurements. Furthermore,
we obtain robust bootstrap estimates of the covariance matrix, and simulation results under the
null did not show deviation from expected distributions. Use of the bootstrap also facilitates joint
testing of parameters from both stages. In the presence of informative censoring, we recommend
sensitivity analysis using existing implementations joint likelihood estimation. To our knowledge
those implementations only exist for simpler joint model formulation with either one longitudinal
and one time-to-event trait[52] or multiple longitudinal traits and one time-to-event outcome[28,53].
Secondly, because the joint model integrates longitudinal and time-to-event sub-models, model
misspecification can occur in multiple ways and lead to invalid inference[54]. Therefore, we
recommend a careful assessment of the model assumptions using joint model diagnostic tools[23].
Thirdly, patient visits were scheduled with high frequency in DCCT, so we ignored the modest
degree of interval censoring in the current implementation of the joint model; when there are
longer gaps between visits, extended methods are needed to account for interval censoring with
additional simulation studies to assess impact on joint model estimates. *Lastly*, joint models are
very computational demanding, particularly for genetic association studies that test millions of
variants. In DCCT application, it took ~60 seconds to fit the joint model for each SNP and
~1090.32 more seconds (~ 18 minutes) to estimate the covariance matrix with 500 bootstraps run
in parallel on 4 nodes (each node with 40 CPU and 202 GB RAM). While analysis at the genome
level involving, up to 85 million 1000G-imputed SNPs in DCCT, seems computationally
unrealistic, a screening approach without bootstrap to select the SNPs based on their association
P-values, followed by the bootstrap refinement would reduce the computational burden. Recent
computationally efficient algorithms have been developed to improve the efficiency of the linear
mixed model[55] and Cox PH model[56,57] for genetic association studies, but to date, they remain to
be implemented for multivariate outcomes.

With the increasing development of national biobanks[58–60] consisting of a large collection of
phenotypes, environmental factors, biomarkers and other risk factors collected over time and
linked with genetic data, we anticipate that joint model methods can be applied to characterize
the genetic architecture of complex traits using these data resources[61]. In addition, the results of
such analysis could contribute towards the translation of human genetic findings to personalized
medicine by providing more efficient SNP effect estimates for increased precision in polygenic
risk score development[62], and causal inference using mediation and mendelian randomization
studies. Lastly, the joint model framework can also enable dynamic prediction beneficial for
dynamic risk assessment[7,63] and optimization of intervention strategies[64,65].

Acknowledgements

This study uses the data provided by the Diabetes Control and Complications Trial / Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Research Group which is sponsored by the National Institute of Diabetes and Digestive and Kidney Diseases contract (#N01-DK-6-2204), National Institute of Diabetes and Digestive and Kidney Diseases grants (#R01-DK-077510, #R01-DK077489, and #P60-DK20595). Funding for genotyping by Illumina HumanCoreExome was provided by JDRF (#17-2013-9). The authors are grateful to the subjects in the DCCT/EDIC cohort for their long-term participation. A complete list of the individuals and institutions participating in the DCCT/EDIC Research Group can be found in **Supplementary Information**. This project was supported by: CIHR Operating/Project Grants (#MOP-84287, #PJT159509), CANSSI postdoctoral fellowship (MB), CIHR STAGE fellowships (MB and OEG, #GET-101831). Computations were performed on the Niagara supercomputer at the SciNet HPC Consortium and Galen, the HPC facility at the Lunenfeld-Tanenbaum Research Institute (LTRI). SciNet is funded by: the Canada Foundation for Innovation under the auspices of Compute Canada; the Government of Ontario; Ontario Research Fund - Research Excellence; and the University of Toronto. The LTRI HPC facility was also supported by the Canada Foundation for Innovation.

Contributions

All named authors affirm that authorship is merited based on the International Committee of Medical Journal Editors (ICMJE) authorship criteria. MB, SBB and RVC designed the study. MB conducted the simulation study and data analysis and drafted the manuscript. SBB, RVC and OEG contributed to the data analysis and interpretation of the results and reviewed/edited the manuscript. ADP contributed to the acquisition of the data and reviewed/edited the manuscript. All authors approve the final version to be published. SBB and MB are the guarantors of this work.

Abbreviations

DCCT: Diabetes control and complications trial;

DR: diabetic retinopathy;

DN: diabetic nephropathy;

GWAS: genome-wide association study;

HbA1c: Hemoglobin A1c;

LD: linkage disequilibrium;

MAF: minor allele frequency;

PH: proportional hazards;

590   QT(s): quantitative trait(s);

591   SBP: systolic blood pressure;

592   SNP: single nucleotide polymorphism;

593   T1DC: type 1 diabetes complications.

References

1. Ibrahim, J.G., Chu, H., and Chen, L.M. (2010). Basic concepts and methods for joint models of
longitudinal and survival data. J. Clin. Oncol. *28*, 2796–2801.

2. Chen, L.M., Ibrahim, J.G., and Chu, H. (2011). Sample size and power determination in joint
modeling of longitudinal and survival data. Stat. Med. *30*, 2295–2309.

3. Hogan, J.W., and Laird, N.M. (1998). Increasing efficiency from censored survival data by using
random effects to model longitudinal covariates. Stat. Methods Med. Res. *7*, 28–48.

4. Wu, L., Liu, W., Yi, G.Y., and Huang, Y. (2012). Analysis of longitudinal and survival data: joint
modeling, inference methods, and issues. J. Probab. Stat. *2012*, 1–17.

5. Asar, Ö., Ritchie, J., Kalra, P.A., and Diggle, P.J. (2015). Joint modelling of repeated
measurement and time-to-event data: An introductory tutorial. Int. J. Epidemiol. *44*, 334–344.

6. Hickey, G.L., Philipson, P., Jorgensen, A., and Kolamunnage-Dona, R. (2016). Joint modelling
of time-to-event and multivariate longitudinal outcomes: recent developments and issues. BMC
Med. Res. Methodol. *16*, 117.

7. Papageorgiou, G., Mauff, K., Tomer, A., and Rizopoulos, D. (2019). An overview of joint
modeling of time-to-event and longitudinal outcomes. Annu. Rev. Stat. Its Appl. *6*, 223–240.

8. Hickey, G.L., Philipson, P., Jorgensen, A., and Kolamunnage-Dona, R. (2018). Joint models of
longitudinal and time-to-event data with more than one event time outcome: a review. Int. J.
Biostat. *14*,.

9. Zhu, H., Ibrahim, J.G., Chi, Y.Y., and Tang, N. (2012). Bayesian influence measures for joint
models for longitudinal and survival data. Biometrics *68*, 954–964.

10. Tang, N.S., Tang, A.M., and Pan, D.D. (2014). Semiparametric Bayesian joint models of
multivariate longitudinal and survival data. Comput. Stat. Data Anal. *77*, 113–129.

11. Tang, A.M., and Tang, N.S. (2014). Semiparametric Bayesian inference on skew-normal joint
modeling of multivariate longitudinal and survival data. Stat. Med. *34*, 824–843.

12. Bycott, P., and Taylor, J. (1998). A comparison of smoothing techniques for CD4 data
measured with error in a time-dependent Cox proportional hazards model. Stat. Med. *17*, 2061–
2077.

13. Self, S., and Pawitan, Y. (1992). Modeling a marker of disease progression and onset of

disease. In AIDS Epidemiology, (Boston, MA: Birkhäuser Boston), pp. 231–255.

14. Tsiatis, A.A., Degruttola, V., and Wulfsohn, M.S. (1995). Modeling the relationship of survival
to longitudinal data measured with error. Applications to survival and CD4 counts in patients with
AIDS. J. Am. Stat. Assoc. *90*, 27–37.

15. Dafni, U.G., and Tsiatis, A.A. (1998). Evaluating Surrogate Markers of Clinical Outcome When
Measured with Error. Biometrics *54*, 1445.

16. Wulfsohn, M.S., and Tsiatis, A.A. (2006). A joint model for survival and longitudinal data
measured with error. Biometrics *53*, 330.

17. Paterson, A.D., and Bull, S.B. (2012). Does familial clustering of risk factors for long-term
diabetic complications leave any place for genes that act independently? J. Cardiovasc. Transl.
Res. *5*, 388–398.

18. The Diabetes Control and Complications Trial Research Group (1993). The effect of intensive
treatment of diabetes on the development and progression of long-term complications in insulin-
dependent diabetes mellitus. N. Engl. J. Med. *329*, 977–986.

19. Paterson, A.D., Waggott, D., Boright, A.P., Hosseini, S.M., Shen, E., Sylvestre, M.P., Wong,
I., Bharaj, B., Cleary, P.A., Lachin, J.M., et al. (2010). A genome-wide association study identifies
a novel major locus for glycemic control in type 1 diabetes, as measured by both A1C and glucose.
Diabetes *59*, 539–549.

20. Lind, M., Odén, A., Fahlén, M., and Eliasson, B. (1995). The relationship of glycemic exposure
(HbA(1c)) to the risk of development and progression of retinopathy in the diabetes control and
complications trial. Diabetes *53*, 1093–1098.

21. Lind, M., Odén, A., Fahlén, M., and Eliasson, B. (2010). The shape of the metabolic memory
of HbA1c: Re-analysing the DCCT with respect to time-dependent effects. Diabetologia *53*, 1093–
1098.

22. Laird, N.M., and Ware, J.H. (1982). Random-effects models for longitudinal data. Biometrics
38, 963–974.

23. Rizopoulos, D. (2012). Joint models for longitudinal and time-to-event data, with applications
in R (Chapman and Hall/CRC).

24. Hougaard, P. (1995). Frailty models for survival data. Lifetime Data Anal. *1*, 255–273.

25. Mauff, K., Steyerberg, E.W., Nijpels, G., van der Heijden, A.A.W.A., and Rizopoulos, D.
(2017). Extension of the association structure in joint models to include weighted cumulative
effects. Stat. Med. *36*, 3746–3759.

26. Tsiatis, A.A., and Davidian, M. (2001). A semiparametric estimator for the proportional
hazards model with longitudinal covariates measured with error. Biometrika *88*, 447–458.

27. Tsiatis, A.A., and Davidian, M. (2004). Joint modeling of longitudinal and time-to-event data:
An overview. Stat. Sin. *14*, 809–834.

28. Hickey, G.L., Philipson, P., Jorgensen, A., and Kolamunnage-Dona, R. (2018). JoineRML: A
joint model and software package for time-to-event and multivariate longitudinal outcomes. BMC
Med. Res. Methodol. *18*,.

29. Therneau, T.M. (2020). A Package for survival analysis in R. R package version 3.2-7, URL
http://CRAN.R-project.org/package=survival.

30. Terry M. Therneau, and Patricia M. Grambsch (2000). Modeling survival data: Extending the
Cox model (New York: Springer).

31. Efron, B. (1981). Nonparametric estimates of standard error: The jackknife, the bootstrap and
other methods. Biometrika *68*, 589.

32. Steffes, M., Cleary, P., Goldstein, D., Little, R., Wiedmeyer, H.-M.M., Rohlfing, C., England,
J., Bucksa, J., and Nowicki, M. (2005). Hemoglobin A1c measurements over nearly two decades:
sustaining comparable values throughout the Diabetes Control and Complications Trial and the
Epidemiology of Diabetes Interventions and Complications study. Clin. Chem. *51*, 753–758.

33. Lorenzi, G.M., Braffett, B.H., Arends, V.L., Danis, R.P., Diminick, L., Klumpp, K.A., Morrison,
A.D., Soliman, E.Z., Steffes, M.W., Cleary, P.A., et al. (2015). Quality control measures over 30
years in a multicenter clinical study: Results from the Diabetes Control and Complications Trial /
Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) study. PLoS One *10*,
e0141286.

34. Hosseini, S.M., Boright, A.P., Sun, L., Canty, A.J., Bull, S.B., Klein, B.E.K., Klein, R., and
Paterson, A.D. (2015). The association of previously reported polymorphisms for microvascular
complications in a meta-analysis of diabetic retinopathy. Hum. Genet. *134*, 247–257.

35. Roshandel, D., Gubitosi-Klug, R., Bull, S.B., Canty, A.J., Pezzolesi, M.G., King, G.L., Keenan,
H.A., Snell-Bergeon, J.K., Maahs, D.M., Klein, R., et al. (2018). Meta-genome-wide association
studies identify a locus on chromosome 1 and multiple variants in the MHC region for serum C-
peptide in type 1 diabetes. Diabetologia *61*, 1098–1111.

36. 1000 Genomes Project Consortium, Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P.,
Kang, H.M., Korbel, J.O., Marchini, J.L., McCarthy, S., McVean, G.A., et al. (2015). A global
reference for human genetic variation. Nature *526*, 68–74.

37. Das, S., Forer, L., Schönherr, S., Sidore, C., Locke, A.E., Kwong, A., Vrieze, S.I., Chew, E.Y.,
Levy, S., McGue, M., et al. (2016). Next-generation genotype imputation service and methods.
Nat. Genet. *48*, 1284–1287.

38. Wheeler, E., Leong, A., Liu, C.-T.T., Hivert, M.-F.F., Strawbridge, R.J., Podmore, C., Li, M.,
Yao, J., Sim, X., Hong, J., et al. (2017). Impact of common genetic determinants of Hemoglobin
A1c on type 2 diabetes risk and diagnosis in ancestrally diverse populations: A transethnic
genome-wide meta-analysis. PLoS Med. *14*, e1002383.

39. Evangelou, E., Warren, H.R., Mosen-Ansorena, D., Mifsud, B., Pazoki, R., Gao, H., Ntritsos,
G., Dimou, N., Cabrera, C.P., Karaman, I., et al. (2018). Genetic analysis of over 1 million people
identifies 535 new loci associated with blood pressure traits. Nat. Genet. *50*, 1412–1425.

40. Pollack, S., Igo, R.P., Jensen, R.A., Christiansen, M., Li, X., Cheng, C.-Y., Ng, M.C.Y., Smith,

A. V, Rossin, E.J., Segrè, A. V, et al. (2019). Multiethnic genome-wide association study of
diabetic retinopathy using liability threshold modeling of duration of diabetes and glycemic control.
Diabetes *68*, 441–456.

41. Grassi, M.A., Tikhomirov, A., Ramalingam, S., Below, J.E., Cox, N.J., and Nicolae, D.L.
(2011). Genome-wide meta-analysis for severe diabetic retinopathy. Hum. Mol. Genet. *20*, 2472–
2481.

42. Sandholm, N., Salem, R.M., McKnight, A.J., Brennan, E.P., Forsblom, C., Isakova, T., McKay,
G.J., Williams, W.W., Sadlier, D.M., Mäkinen, V.-P.P., et al. (2012). New susceptibility loci
associated with kidney disease in type 1 diabetes. PLoS Genet. *8*, e1002921.

43. Li, J., and Ji, L. (2005). Adjusting multiple testing in multilocus analyses using the eigenvalues
of a correlation matrix. Heredity (Edinb). *95*, 221–227.

44. Kraft, P. (2008). Curses—Winner's and Otherwise—in Genetic Epidemiology. Epidemiology
19, 649–651.

45. Sun, L., Dimitromanolakis, A., Faye, L.L., Paterson, A.D., Waggott, D., DCCT/EDIC Research
Group, and Bull, S.B. (2011). BR-squared: a practical solution to the winner's curse in genome-
wide scans. Hum. Genet. *129*, 545–552.

46. Schaid, D.J., Tong, X., Larrabee, B., Kennedy, R.B., Poland, G.A., and Sinnwell, J.P. (2016).
Statistical methods for testing genetic pleiotropy. Genetics *204*, 483–497.

47. Liu, L., Zheng, C., and Kang, J. (2018). Exploring causality mechanism in the joint analysis of
longitudinal and survival data. Stat. Med. *37*, 3733–3744.

48. Sobel, M.E. (1982). Asymptotic confidence intervals for indirect effects in structural equation
models. Sociol. Methodol. *13*, 290.

49. Mackinnon, D.P., Warsi, G., and Dwyer, J.H. (1995). A simulation study of mediated effect
measures. Multivariate Behav. Res. *30*, 41.

50. Albert, P.S., Shih, J.H., Albert, P.S., Shih, J.H., Albert1', P.S., and Shih2, J.H. (2010). On
estimating the relationship between longitudinal measurements and time-to-event data using a
simple two-stage procedure. Biometrics *66*, 983–991.

51. Ye, W., Lin, X., and Taylor, J.M.G. (2008). Semiparametric modeling of longitudinal
measurements and time-to-event data - A two-stage regression calibration approach. Biometrics
64, 1238–1246.

52. Rizopoulos, D. (2010). JM : An R Package for the joint modelling of longitudinal and time-to-
event data. J. Stat. Softw. *35*,.

53. Rizopoulos, D. (2016). The R package JMbayes for fitting joint models for longitudinal and
time-to-event data using MCMC. J. Stat. Softw. *72*,.

54. Arisido, M.W., Antolini, L., Bernasconi, D.P., Valsecchi, M.G., and Rebora, P. (2019). Joint
model robustness compared with the time-varying covariate Cox model to evaluate the
association between a longitudinal marker and a time-to-event endpoint. BMC Med. Res.

Methodol. *19*, 222.

55. Sikorska, K., Lesaffre, E., Groenen, P.J.F., Rivadeneira, F., and Eilers, P.H.C. (2018).
Genome-wide analysis of large-scale longitudinal outcomes using penalization - GALLOP
algorithm. Sci. Rep. *8*,.

56. Rizvi, A.A., Karaesmen, E., Morgan, M., Preus, L., Wang, J., Sovic, M., Hahn, T., and
Sucheston-Campbell, L.E. (2019). gwasurvivr: an R package for genome-wide survival analysis.
Bioinformatics *35*, 1968–1970.

57. Bi, W., Fritsche, L.G., Mukherjee, B., Kim, S., and Lee, S. (2020). A fast and accurate method
for genome-wide time-to-event data analysis and its application to UK Biobank. J. Clean. Prod.
107, 222–233.

58. Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., Motyer, A., Vukcevic,
D., Delaneau, O., O'Connell, J., et al. (2018). The UK Biobank resource with deep phenotyping
and genomic data. Nature *562*, 203–209.

59. Scholtens, S., Smidt, N., Swertz, M.A., Bakker, S.J.L., Dotinga, A., Vonk, J.M., Van Dijk, F.,
Van Zon, S.K.R., Wijmenga, C., Wolffenbuttel, B.H.R., et al. (2015). Cohort Profile: LifeLines, a
three-generation cohort study and biobank. Int. J. Epidemiol. *44*, 1172–1180.

60. Dummer, T.J.B., Awadalla, P., Boileau, C., Craig, C., Fortier, I., Goel, V., Hicks, J.M.T.,
Jacquemont, S., Knoppers, B.M., Le, N., et al. (2018). The Canadian Partnership for Tomorrow
Project: A pan-Canadian platform for research on chronic disease prevention. Cmaj *190*, E710–
E717.

61. Gasparini, A., Abrams, K.R., Barrett, J.K., Major, R.W., Sweeting, M.J., Brunskill, N.J., and
Crowther, M.J. (2020). Mixed-effects models for health care longitudinal data with an informative
visiting process: A Monte Carlo simulation study. Stat. Neerl. *74*, 5–23.

62. Young, A.I., Benonisdottir, S., Przeworski, M., and Kong, A. (2019). Deconstructing the
sources of genotype-phenotype associations in humans. Science *365*, 1396–1400.

63. Bull, L., Lunt, M., Martin, G., Hyrich, K., and Sergeant, J. (2020). Harnessing repeated
measurements of predictor variables for clinical risk prediction: a review of existing methods.
Diagnostic Progn. Res. *4*, 1–16.

64. Sweeting, M.J., and Thompson, S.G. (2011). Joint modelling of longitudinal and time-to-event
data with application to predicting abdominal aortic aneurysm growth and rupture. Biometrical J.
53, 750–763.

65. Yuen, H.P., Mackinnon, A., Hartmann, J., Amminger, G.P., Markulev, C., Lavoie, S., Schäfer,
M.R., Polari, A., Mossaheb, N., Schlögelhofer, M., et al. (2018). Dynamic prediction of transition
to psychosis using joint modelling. Schizophr. Res. *202*, 333–340.

66. Ibrahim, J.G., Chen, M.-H., and Sinha, D. (2001). Joint models for longitudinal and survival
data. In EMR-IBS Bi-Annual Meeting, pp. 262–289.

Appendix

Inference under the joint likelihood can be based on joint maximum-likelihood estimation, Bayesian estimation of the joint likelihood parameters, or a two-stage strategy[23]. Joint maximum-likelihood and Bayesian methods enable joint parameter estimation but can be computationally challenging, especially for multiple traits. Two-stage approaches[12–15] sequentially fit the longitudinal (Stage 1) and time-to-event (Stage 2) sub-models, with trajectory estimates from Stage 1 incorporated into Stage 2; this can be relatively computationally efficient and lends itself to flexible model formulation. However, as noted in introduction, inference can be mis-calibrated because of propagation of errors[16].

Joint likelihood function

Let Ω be the full parameter vector containing all fixed parameters from the longitudinal and time-to-event sub-models respectively. We assume:

(A1) $b_i \sim MVN(0, D)$

(A2) $\varepsilon_{ij(l)} \sim N\left(0, \sigma^2_{(l)}\right)$ and $\varepsilon_{ij(l)} \perp \varepsilon_{ip(l)}$ between visits $j \neq p$

(A3) $b_{i(l)} \perp \varepsilon_{ij(l)}$

(A4) $u_i \perp b_i$

Given (A1)-(A4), it is appropriate to assume that the random effects b_i account for: the association between the *L* longitudinal traits; the association between both the longitudinal and time-to-event outcomes[66]; and the serial correlation between the repeated measurements in each longitudinal process[22]; and that given (A4), the frailty term accounts for the residual dependency between the time-to-event traits[24]. Under these conditional independence assumptions, the joint likelihood function of the model parameters given the observed data is:

$$L(\Omega|Y_i, T_i, \delta_i, SNP_i, H_i, V_i) = \prod_{i=1}^{N} \int f_1(Y_i|\, b_i, \Omega) \times f_2(T_i, \delta_i \mid b_i, u_i, \Omega) \times f_3(b_i|\Omega) \times f_4(u_i|\Omega)\, du_i db_i$$

where:

- $(T_i, \delta_i) = \left((T_{i(1)}, \delta_{i(1)})', \ldots, (T_{i(k)}, \delta_{i(k)})', \ldots, (T_{i(K)}, \delta_{i(K)})'\right)'$
- $Y_i = (Y_{i(1)}', \ldots, Y_{i(l)}', \ldots, Y_{i(L)}')'$
- $f_1(Y_i|\, b_i, \Omega) = \prod_{l=1}^{L} \left(\dfrac{1}{\sqrt{2\pi\sigma^2_{(l)}}}\right)^{n_i} exp\left[-\dfrac{1}{2\sigma^2_{(l)}} \prod_{j=1}^{n_i} (Y_{ij(l)}(t_{ij}) - Y^*_{ij(l)}(t_{ij}))^2\right]$
- $f_2(T_i, \delta_i \mid b_i, u_i, \Omega) = \prod_{k=1}^{K} \left[\lambda_{i(k)}\left(T_{i(k)}|\, b_i, u_i, \Omega\right)\right]^{\delta_{ik}} S_{i(k)}(T_{i(k)}|\, b_i, u_i, \Omega)$
- $S_{i(k)}\left(T_{i(k)}|\, b_i, u_i, \Omega\right)$

$$801 \qquad = exp\left[- \int_0^{T_{i(k)}} \lambda_{i0}(s)exp\{\gamma_{g(k)}SNP_i + W_{i(k)}(s) + \gamma_{v(k)}V_{i(k)} + u_i\}ds\right]$$

• $f_3(b_i|\Omega) = \left(\frac{1}{\sqrt{2\pi}}\right)^{-q/2}|D|^{-1/2}exp\left\{-\frac{1}{2}b_i^T D^{-1}b_i\right\}$, where q is the dimension of the D matrix

• $f_4(u_i|\Omega) = \frac{u_i^{a-1}\exp{(-u_i/b)}}{\Gamma(a)b^a}$, i.e. we assume $u_i \sim \Gamma(a,b)$, with a, b>0.

Calculation of this full likelihood requires multivariate integration with respect to the random
effect's distribution, which can lead to demanding computation. When the random effect vector
b_i, has a small dimension, say less than 3, the integral can be evaluated via Gaussian quadrature
which approximates the integral by a weighted sum of the target function evaluated at pre-
specified sample points. However, when the dimension is larger, it is demanding to calculate the
integrals with satisfactory approximation accuracy. Although a full likelihood specification enables
rigorous study of asymptotic properties, its large sample approximation may not be accurate when
sample size is small. In comparison, the Bayesian paradigm does not require asymptotic
approximations, but the design of an efficient sampling algorithm to study the posterior distribution
is challenging.

Likelihood functions under the two-stage approximation

Let Ω and Γ be the vectors containing all fixed parameters from the longitudinal and time-to-event
sub-models respectively.

Stage1: Multivariate mixed model

$$818 \qquad L(\Omega|Y_i, X_i, C_i) = \prod_{i=1}^{N} \int_{b_i} f_1(Y_i| b_i,\Omega) \times f_3(b_i|\Omega)\, db_i$$

819 *Stage2:* Multivariate Cox PH model adjusted for fitted values of the trajectories $\widehat{Y_i^*(T_i)}$

$$820 \qquad L(\Gamma|T_i, \delta_i, C_i) = \prod_{i=1}^{N} \int_{u_i} f_2\left(T_i, \delta_i |\widehat{Y_{l(l)}^*}(T_i), u_i, \Gamma\right) \times f_4(u_i|\Gamma) \times du_i$$

Where:

• $f_2\left(T_i, \delta_i |\widehat{Y_i}(T_i), u_i, \Gamma\right) = \prod_{k=1}^{K}\left[\lambda_{i(k)}\left(T_{i(k)}|\widehat{Y_{l(l)}^*}(T_{i(k)}), u_i, \Gamma\right)\right]^{\delta_{ik}} S_{ik}(T_{i(k)}|\widehat{Y_{l(l)}^*}(T_{i(k)}), u_i, \Gamma)$

• $S_{i(k)}\left(T_{i(k)}|\widehat{Y_{l(l)}}(T_{i(k)}), u_i, \Gamma\right) = exp\left[-\int_0^{T_{i(k)}} \lambda_{i0}(s)exp\{\gamma_{g(k)}SNP_i + W_{i(k)}(s) + \gamma_{v(k)}V_{i(k)} + \right.$
$\left. u_i\}ds\right]$

• With $W_{i(k)}(s) = \sum_{l=1}^{L_k} \alpha_{l(k)}f_{l(k)}(\widehat{Y_{l(l)}^*}(s))$

Unlike the *joint likelihood function*, where the shared random effects b_i account for the
dependencies between the longitudinal and the time-to-event traits, the two-stage approach
accounts for the dependencies between the longitudinal and time-to-event traits via the fitted
values of the longitudinal trajectories. As mentioned previously, this approximation can produce
biased estimates and/or underestimated standard errors[16], particularly in presence of non-random
censoring of the longitudinal trait values due to the occurrence of an event or from informative
dropout[50,51] and/or because of propagation errors of Stage 1 parameters in Stage2[16]. Indeed,
informative missingness/dropouts lead in differential follow-up between patients with and without
an event, the time-to-event processes are related to the length of the follow-up, and thus the
random effects $b_{i(l)}$ can depend on the event times (e.g. patients who have an event early are
more likely to have positive random slopes). However, in absence of informative
dropouts/missingness, as we showed in our simulation studies, this approach has low bias and is
computationally feasible for genetic association studies.

Multivariate mixed-effects sub-model for L longitudinal QT(s) (and/or risk factors)

Measured longitudinal traits:

$$Y_{i(l)}(t_{ij}) = Y^*_{i(l)}(t_{ij}) + \varepsilon_{ij(l)} \sim N(0, \sigma^2_{(l)})$$

- *intermittently* (ie scheduled visits)
- measurement *errors* (ie biological variation)
- potentially *informatively* missing values

Longitudinal trajectories:

$$Y^*_{i(l)}(t_{ij}) = \beta_{0(l)} + b_{i0(l)} + (\beta_{1(l)} + b_{i1(l)})t_{ij} + \beta_{g(l)}SNP_i + \beta_{h(l)}H_{i(l)}$$

- *latent* and continuous processes
- various *shapes* (linear/non-linear)
- *smoothed*, account for *measurement* errors
- account for *shared* and/or trait-*specific* covariates
- $b_i \sim N_{2L}(0, D)$: subject-specific deviation from *average* trajectories
- D: dependencies *within/between* L traits

Time-dependent association structures

$$W_{i(k)}(t) = \sum_{l=1}^{L_k} \alpha_{l(k)} f_{l(k)}(Y^*_{i(l)}(t))$$

- account for the *indirect* SNP effects induced by each QT l
- $\alpha_{l(k)}$ association between each pair of traits l & k
- $f_{l(k)}(Y^*_{i(l)}(t))$, *functional dependence* between traits l & k, based on prior knowledge

Proportional Hazards frailty sub-model for K time-to-event traits

$$\lambda_{i(k)}(t) = \lambda_{0(k)}(t) \times exp\{\gamma_{g(k)}SNP_i + W_{i(k)}(t) + \gamma_{v(k)}V_{i(k)} + u_i\}$$

- possibly *informatively* censored
- $\gamma_{g(k)}$: *direct* SNP effect (other than via $W_{i(k)}(t)$)
- accounts for *shared* and/or trait-*specific* covariates
- $u_i \perp b_i$, subject-specific, accounts for *unexplained* dependencies

SNP associations with each time-to-event trait k and each longitudinal trait l ($\alpha_{l(k)} \neq 0$):

  a. Indirect: $\beta_{g(l)} \neq 0$ & $\gamma_{g(k)} = 0$
  b. Direct: $\beta_{g(l)} = 0$ & $\gamma_{g(k)} \neq 0$
  c. Direct & Indirect: $\beta_{g(l)} \neq 0$ & $\gamma_{g(k)} \neq 0$

Figure 1. Proposed joint modelling approach for characterization of complex genetic architecture of multiple disease progression

Figure 2. Realistic DCCT-data-based *causal* genetic scenario

The effect of gender on SBP (noted as $H_{i(2)}$), and effects of T1D duration at baseline (noted as $V_{i(k)}$) on both time-to-T1DC (k=1, 2) traits are not represented in this figure, but are included in the data generating model, see **Supplementary Information SI-1** for details. We simulated R=1000 replicates of N=667 DCCT individuals with M=5 causal variants and K=2 time-to-event traits simulated under this causal genetic scenario and R=1000 replicates of M=5 SNPs (with same MAF as the causal ones) under a global null genetic scenario where none of the SNPs is associated with any traits.

From DCCT: $t_i, n_i, Y_{i(1)}, Y_{i(2)}, H_{i(2)}, V_{i(1)}, V_{i(2)}$

Simulated risk factor: U_i, unexplained dependency between the K time-to-event traits

(repeated R times for each of individual i from DCCT)

Simulation of SNPm (p_m) with _indirect_ effect

$$SNPm_i \sim Mult(\pi_{0i}, \pi_{1i}, \pi_{2i})$$

Where:

- $\pi_{gi} = P(SNP_i | \psi_{i(s)}); \; \psi_i \sim N_{n_i}(X_i\beta, Z_i DZ_i' + \sigma^2 I_{n_i})$

- $\psi_i = \begin{cases} (Y_{i(1)} | SNP1_i = g_i, t_i, \Omega_{(1)}), \text{for SNP1}(MAF = p_1) \\ (Y_{i(2)} | SNP2_i = g_i, t_i, H_{i(2)}, \Omega_{(2)}), \text{for SNP2 } (MAF = p_2) \\ (U_i | SNP1_i = g_i, t_i, \Omega_{(3)}), \text{for SNP3 } (MAF = p_3) \end{cases}$

Trajectories from fitted mixed models

$Y_{i(l)}^*(t) \; \& \; U_i^*(t)$

Simulation of K time-to-event traits

- **_Censoring_ time** $C_i \sim \text{Uniform}(0, t_{n_i})$

- **_Uncensored_** $T_{i(k)}^*$ calculated from the inverse of the cumulative hazard function for each time-to-event k with specified parameters vector $\Gamma_{(k)}$

$$\lambda_{i(k)}(t) = \lambda_{0(k)}(t) \times exp\{\eta_{i(k)} + \alpha_{U(k)}U_i^*(t)\}$$

$$T_{i(k)} = min(T_{i(k)}^*, C_i)$$
$$\delta_{i(k)} = I(T_{i(k)}^* \le C_i)$$

Simulation of SNPm (p_m) with _direct_ effect

$$SNPm_i \sim Mult(\pi_0, \pi_1, \pi_2)$$
$$\pi_0 = (1 - p_m)^2; \; \pi_1 = 2p_m(1 - p_m)^2; \; \pi_2 = p_m^2$$

(m ϵ {4,5} for SNP4 &SNP5)

Figure 3. Illustration of the procedure developed for DCCT-based simulation study under the scenario from Figure 2

$\Omega_{(1)}$, $\Omega_{(2)}$ and $\Omega_{(U)}$ are specified parameter values for the longitudinal trait models, and $\Gamma_{(1)}$, $\Gamma_{(2)}$ for the time-to-event trait models, p is the specified vector of minor allele frequencies vector of the M SNPs. Parameters for the causal genetic scenario in **Figure 2** are summarized in **Table 2** and in **Table S1**. The DCCT-data-based simulation algorithm uses longitudinal trait/baseline covariates from DCCT and simulated genetic data and time-to-event traits. SNPs with indirect effects are simulated from a multinomial distribution with conditional genotype probabilities ($\pi_{0i}, \pi_{1i}, \pi_{2i}$) using longitudinal trait values for individual i. Here, X_i and Z_i denote the design matrices of the fixed and random effects of the mixed model assumed for each longitudinal trait and D is the specified covariance matrix for the random effects. SNPs with direct effects are simulated from the population probabilities, that depend on the MAF. Each time-to-event trait k is simulated by calculating the inverse of the cumulative specified hazard function using the Brent univariate root-finding method[34,35]. For DR, $\eta_{i(1)} = \gamma_{g(1)} SNP2_i + \alpha_{1(1)}Y_{i(1)}^*(t) + \gamma_{v(1)} V_{i(1)}$ and for DN, $\eta_{i(2)} = \gamma_{g(2)} SNP4_i + \gamma'_{g(2)} SNP5_i + \sum_{l=1}^{2} \alpha_{l(1)} Y_{i(l)}^*(t) + \gamma_{v(2)}V_{i(2)}$. We further added the effect of the longitudinal trajectory $U_i^*(t)$ to each $\eta_{i(k)}$ in the hazard function of each trait k to induce some unexplained dependencies between the simulated time-to-event traits. Details of the DCCT-based simulation procedure are presented in **Supplementary Information SI-1**.

Figure 4. **Performance of the testing procedure to recover underlying generating *causal* relationships from Figure 2, assessed using *R*=1000 replicates of *N*=667 DCCT subjects***.

The *Y* axis represents the proportion of replicates that accurately classify each causal SNP as: **(a.)** indirect; **(b.)** direct and **(c.)** direct and indirect causal associations for SNP1, SNP2, SNP4 and SNP5. SNP3 is tested as a direct association because the longitudinal risk factor *U* is assumed unmeasured. Results obtained for direct association of SNP3 and SNP4 with DN/SBP are similar to those presented for DN/HbA1c (**Figure S4**) and are not shown here. Plot **(d.)** shows that SNP5 is misclassified as an indirect association in a larger number of replicates compared to **(c.)** by the joint models compared to CM-obs**.** This misclassification rate tends to increase with decreasing P^*. Classification rates of all these SNPs with other time-to-event/longitudinal trait pairs are presented in **Figure S4.**

b.

Figure 5. **Classification of direct and/or indirect SNP associations in the DCCT Genetics Study data.**

(a.) Scatter plots of the P-values (-log10) for tests of $\beta_{g(l)}$ (X axis) and $\gamma_{g(k)}$ (Y axis) for DR/HbA1c, DN/HbA1c and DN/ SBP trait pairs. Significance levels $P^* = 1.7 \times 10^{-4}$ and $P^* = 0.05$ are indicated by red and grey horizontal and vertical dashed lines. **(b.)** Association results for the two SNPs that exibit indirect association with DR/DN outcomes at $P^* = 1.7 \times 10^{-4}$. Left panel present results from separate analysis of each trait; and right panel presents results from the joint model with bootstrap 95% confidence intervals for the direct and indirect SNP effects. Results are presented using the time-weighted cumulative HbA1c effects on T1DC traits.

		SNP association with the longitudinal risk factor l	
		$P_{\beta_{g(l)}} \leq P^*_{\beta_g}$	$P_{\beta_{g(l)}} > P^*_{\beta_g}$
SNP association with the time-to-event trait k	$P_{\gamma_{g(k)}} \leq P^*_{\gamma_g}$	Direct & Indirect	Direct
	$P_{\gamma_{g(k)}} > P^*_{\gamma_g}$	Indirect	No association

$P_{\beta_{g(l)}}$ and $P_{\gamma_{g(k)}}$ are the *P-values* from Wald test (1df) for SNP effects $\beta_{g(l)}$ and $\gamma_{g(k)}$.

$P^*_{\beta_g}$ and $P^*_{\gamma_g}$ are the significance thresholds (see methods for details). Note that thorough the paper, we use $P^* = P^*_{\beta_g} = P^*_{\gamma_g}$, but different levels can be specified for $P^*_{\beta_g}$ and $P^*_{\gamma_g}$.

<u>Table 1</u>. Hypothesis testing procedure to classify a SNP as having a direct and/or an indirect association with a time-to-event trait *k* and an associated longitudinal risk factor *l*.

Causal SNPs	Type of SNP association	Specified parameters[1]	Empirical power (separate analysis of each trait)[2]			
			Trait (*Tested parameter*)	$P^* =0.05$	$P^* =0.001$	$P^* =10^{-5}$	$P^* =5\text{x}10^{-8}$
SNP1 (MAF=0.30)	*Indirect* association with both T1DC traits via HbA1c	$\beta_{g(1)} = 0.70$	HbA1c ($\beta_{g(1)}$)	100%	100%	100%	99.8%
		$\alpha_{1(1)} = 0.20$	DR ($\gamma_{g(1)}$)	20.4%	1.2%	$<10^{-4}$	$<10^{-4}$
		$\mu_{g(1,k)} = 0.14$	DN ($\gamma_{g(2)}$)	8.70%	0.3%	$<10^{-4}$	$<10^{-4}$
SNP2 (MAF=0.10)	*Direct* association with DR	$\gamma_{g(1)} = 0.80$	DR ($\gamma_{g(1)}$)	100%	98.3%	87.4%	58.2%
SNP3 (MAF=0.40)	*Indirect* association with both T1DC traits via *U*	$\beta_{g(U)} = 0.80$	DR ($\gamma_{g(1)}$)	87.5%	42.8%	7.6%	0.9%
		$\alpha_{1(U)} = 0.40$ $\mu_{g(1,k)} = 0.36$	DN ($\gamma_{g(2)}$)	32.6%	2.6%	$<10^{-3}$	$<10^{-3}$
SNP4 (MAF=0.30)	*Direct* association with DN	$\gamma_{g(2)} = 0.70$	DN ($\gamma_{g(2)}$)	94.1%	56.5%	17.3%	4.0%
SNP5 (MAF=0.20)	*Direct* and *indirect* association with DN via SBP	$\beta_{g(2)} = 7.00$	SBP ($\beta_{g(2)}$)	100%	100%	100%	100%
		$\alpha_{2(2)} = 0.20$ $\mu_{g(2,2)} = 1.40$ $\gamma_{g(2)} = 0.40$	DN ($\gamma_{g(2)}$)	100%	100%	99.8%	98.0%

[1]$\beta_{g(l)}$ is the SNP effect on the longitudinal trait *l*, $\gamma_{g(k)}$ is the direct SNP effect on the time-to-event trait *k*; $\alpha_{l(k)}$ is the effect of each longitudinal trajectory for trait *l* on the time-to-event trait *k*. $\mu_{g(l,k)}$ is the *indirect* SNP effect, calculated as $\mu_{g(l,k)} = \beta_{g(l)}\alpha_{l(k)}$.

[2]For each scenario of direct/indirect SNP association, we present the empirical power to detect each SNP using *separate* analysis of each trait (linear mixed model for each longitudinal trait, Cox PH model for each time-to-event trait); as typically used for SNP discovery. Models are adjusted for all non-SNP baseline covariates, except *U,* as specified to simulate the data. The empirical power is calculated as the proportion of replicates in the causal scenario from Figure 2 with a significant SNP association detected at specified levels of P^*. As expected, SNP1, SNP3 and SNP5 appear associated with the T1DC outcomes due to their indirect effects induced by longitudinal risk factors absents from the time-to-event models. For comparisons, empirical power of the same tests based on parameters from the joint model are presented in **Table S2**.

Table 2. **Simulated scenarios of direct and/or indirect SNP associations, assessed using *R*=1000 replicates of *N*=667 DCCT subjects simulated under the *causal* genetic scenario.** This table presents parameters specified for SNPs and trajectory effects on time-to-event traits for the DCCT-based simulation study and illustrates how each type of SNP would be detected in GWAS discovery analysis based on separate analysis of each trait.

SNPs[1]	Trait pairs	$P^* =$0.05			$P^* =$0.01		
		JM-cmp	JM-mis	CM-obs	JM-cmp	JM-mis	CM-obs
Indirect association							
SNP1	DR/HbA1c	0.049	0.049	0.500	0.015	0.015	0.015
	DN/HbA1c	0.049	0.048	0.048	0.015	0.015	0.015
Direct association							
SNP2	DR/HbA1c	0.058	0.045	0.045	0.009	0.009	0.007
SNP3	DR/HbA1c	0.041	0.044	0.041	0.008	0.006	0.008
	DN/HbA1c	0.041	0.037	0.042	0.012	0.008	0.008
	DR/HbA1c	0.042	0.034	0.036	0.012	0.008	0.009
SNP4	DN/HbA1c	0.046	0.044	0.048	0.007	0.009	0.005
	DN/SBP	0.047	0.044	0.042	0.008	0.009	0.005
Direct and Indirect association							
SNP5	DN/SBP	<0.001	0.002	0.003	<0.001	<0.001	<0.001

[1]Each SNP is tested with the same direct/indirect association tests as applied to the causal SNPs, except SNP3 that is tested as a direct association because the longitudinal risk factor U is assumed unmeasured. Classification rates for all other possible QT/time-to-event trait pairs are presented in **Figure S3**.

[2] JM-cmp denotes the complete joint model fitted for each SNP separately, including all non-genetic covariates used for the data simulation (including U).

Table 3. Classification rates of the procedure for each SNP under the *global null* genetic scenario, assessed using *R*=1000 replicates of *N*=667 DCCT subjects.