Validation of a multi-gene qPCR-based pharmacogenomics panel across major ethnic groups in Singapore and Indonesia

Running Title

Multi-gene qPCR-based PGx panel for Asians

Authors

Anar Sanjaykumar KOTHARY[1,*], Caroline MAHENDRA[1,*], Mingchen TAN[1], Eunice Jia Min TAN[2], Jacelyn PHUA Hong Yi[1], Gabriella[1], TAN Xin Hui Jocelyn[1], Jessline Stephanie HARUMAN[1], Chun Kiat LEE[2], Benedict YAN[2,3], Astrid IRWANTO[1,4,#]

Affiliations

[1]Nalagenetics Pte Ltd, Singapore, Singapore

[2]Department of Laboratory Medicine, National University Health System, Singapore

[3]Stronghold Diagnostics Lab, Agency for Science, Technology and Research, Singapore

[4]Department of Pharmacy, Faculty of Science, National University of Singapore, Singapore

[*]equal contribution

[#]corresponding author: astrid@nalagenetics.com

Acknowledgements

We would like to thank Mr. Xie Zhicheng and Mr. Heng Khai Koon at the Genome Institute of Singapore for their assistance with performing the Agena VeriDose® Core and *CYP2D6* CNV Panel studies. We also thank Dr. Tan Zhihao for his feedback on the data analysis approach and manuscript, and PT Nalagenetik Riset Indonesia for their assistance in study recruitment and experimentation performed in Indonesia.

Abstract

Background – With up to 70% of adverse drug reactions (ADRs) having high genetic associations, the clinical utility of pharmacogenomics (PGx) has been gaining traction. Nala PGx Core™ is a multi-gene qPCR-based panel that comprises 18 variants and 2 *CYP2D6* Copy Number markers across 4 pharmacogenes – *CYP2C9*, *CYP2C19*, *CYP2D6* and *SLCO1B1*.

Objectives – In this study, we validated the performance of Nala PGx Core™ against benchmark methods, on the Singaporean and Indonesian populations. Additionally, we examined the allele and diplotype frequencies across 5 major ethnic groups present in these populations namely, Indonesians, Chinese, Malays, Indians and Caucasians.

Methods – Human gDNA samples, extracted from the buccal swabs of 246 participants, were tested on Nala PGx Core™ and two chosen benchmarks, Agena VeriDose® Core and *CYP2D6* Copy Number Variation (CNV) Panel, and TaqMan® DME Genotyping Assays. Performance was evaluated based on assay robustness, precision and accuracy at the genotype- and diplotype-level.

Results – Nala PGx Core™ demonstrated high genotype- and diplotype-level call rates of >97% and >95% respectively in *CYP2D6*, and 100% for *CYP2C9*, *CYP2C19* and *SLCO1B1*. A precision rate of 100% was observed on both intra- and inter-precision studies. Variant-level concordance to the benchmark methods was ≥96.9% across all assays, which consequently resulted in a diplotype-level concordance of ≥94.7% across *CYP2C9*, *CYP2C19* and *CYP2D6*. Overall, the allele frequencies of *CYP2D6*10* and *CYP2D6*36* were higher in our cohort as compared to previous records. Notably, *CYP2D6* copy number variation (CNV) analysis demonstrated a *CYP2D6* *10/*36* frequency of 26.5% amongst the Indonesian cohort.

Conclusion – Nala PGx Core™ produced robust and accurate genotyping when compared to other established benchmarks. Furthermore, the panel successfully characterized alleles of clinical relevance in the Singaporean and Indonesian populations such as *CYP2D6*10* and *CYP2D6*36*, suggesting its potential for adoption in clinical workflows regionally.

Keywords: Pharmacogenomics; *CYP2C9*; *CYP2C19*; *CYP2D6*; *SLCO1B1*; qPCR; adverse drug reactions; Singapore; Indonesia.

Abstract count: 290 words, 1764 characters
Main text count: 4337 words

Introduction

Pharmacogenomics (PGx) is a branch of medicine concerned with determining the response of an individual to therapeutic drugs using genetic and/or genomic information[1]. With up to 70% of adverse drug reactions (ADRs) having high genetic associations[2], and the sunk cost of trial-and-error prescriptions amounting to USD 30 billion[3], the clinical utility[4] of PGx appears to be gaining prominence.

Applications of PGx include the genotyping of Cytochrome P450 (P450) enzymes which are significantly involved in drug metabolism and chemical toxicity[5]. *CYP2C9*, *CYP2C19* and *CYP2D6* display common polymorphisms that produce altered enzyme activity, often resulting in deviant drug metabolism that leads to side effects from drugs metabolized by them[5].

Alterations in *CYP2C9* can disrupt the metabolism of warfarin, while *CYP2C19* is associated with the metabolism of a number of drugs including clopidogrel, omeprazole and phenytoin[6,7]. The highly polymorphic *CYP2D6* has more than 100 variants and is responsible for the metabolism of approximately 25% of all clinically used drugs, including codeine, fluoxetine and amitriptyline[7]. For this reason, pharmaceuticals are particularly cautious in the development of drugs that are known to interact with these three P450s[5]. Additionally, pharmacogenes, such as *SLCO1B1*, are responsible for encoding transporters of simvastatin and other statins into tissues also affect their response upon intake in individuals. Variants of this gene are associated with increased risk of muscle pain (myalgia) and damage, including myopathy and in severe cases rhabdomyolysis[8]. Previous studies suggest the role of specific *CYP2C19* and *SLCO1B1* variants in the bioactivation of drugs such as clopidogrel and atorvastatin in the Singaporean population[9,10]. The *CYP2C19* variant, rs4986893, which is implicated in reduced clopidogrel metabolism, has been observed to be 50 times more prevalent in the Singaporean Chinese population than in Europeans[10]. The *SLCO1B1* variant, rs4149056, which is implicated in statin metabolism, has been observed to occur less frequently in the Singaporean population[9]. Taken together, these cases highlight the need to harness PGx testing to account for population differences in the personalization of prescriptions[11].

Current methods on PGx testing for the identification of genetic polymorphisms revolve around variant genotyping that utilize technologies ranging from Next Generation Sequencing (NGS), Microarray, Mass Spectrometry to Quantitative PCR (qPCR)[1,6,12,13]. Main considerations in the adoption of a platform include its robustness, effectiveness and cost-saving, especially in localities with limited resources. qPCR-based assays satisfy these criteria and are a preferred platform for clinical laboratories[14–19]. However, some of the available commercial kits are limited in their ability to test the 4 genes of interest mentioned above in a single panel[20–25]. Nala PGx Core™ is a qPCR-based multi-gene panel consisting of 20 pharmacogenomic variants that are associated with altered *CYP2C9*, *CYP2C19*, *CYP2D6* and *SLCO1B1* activity in Asian geographies. Our study aims to evaluate the performance of Nala PGx Core™ in effectively identifying genetic variations that may contribute to PGx-related clinical outcomes in Singapore and Indonesia, with a secondary objective of highlighting the allele and diplotype frequencies in the geographies examined.

Methods and Materials

Ethics Approval

Institutional Review Board (IRB) approval was granted by the Parkway Independent Ethics Committee (Singapore) under IRB Reference Number PIEC/2019/021 and Siloam Hospitals Ethics Review Committee (Jakarta, Indonesia) under IRB Reference Number 001/EA/KEPKK/RSMRCCC/V/2019.

Study Recruitment

Patients were recruited on behalf of Nalagenetics Pte. Ltd. with written informed consent forms from recruitment sites in Singapore and Indonesia. A total of 251 samples were evaluated from 5 major ethnic groups to ensure the representation of the major ethnic groups residing in both countries – Chinese, Malays, Indians, Caucasians and Indonesians. Participants identifying as one or more of the following ethnicities were categorized as Indonesians: Ambon, Batak, Betawi, Jawa, Lampung, Manado, Minangkabau, Nusa Tenggara Timur, Palembang, Sulawesi, Sunda, Timor Leste, Tolaki and Toraja.

Buccal samples were collected using OraCollect (Cat No. DNA OCR-100 from DNA Genotek) and genomic DNA (gDNA) extracted using the Monarch Genomic DNA Purification Kit (Cat No. T3010 from NEB). The extraction procedure followed manufacturer's instructions with additional dry-spin step at maximum speed for 1 minute after the 2nd buffer washing step. The quality and concentration of gDNA extracts were quantified by NanoDrop 2000 Spectrophotometer (Singapore) and BioDrop-µLITE (Indonesia). Samples that failed to meet the DNA quality control criteria (n=5) of A260/280 >1.7 and DNA yield >500ng were excluded from the study. The remaining extracted gDNA samples (n=246) were stored at -20□ for downstream application.

Nala PGx Core™

The Nala PGx Core™ kit from Nalagenetics Pte. Ltd. consists of 20 qPCR-based variant assays across four genes – *CYP2C9*, *CYP2C19*, *CYP2D6* and *SLCO1B1*. Whilst assays for *CYP2C9*, *CYP2C19* and *CYP2D6* have been designed to enable the detection of specific star alleles, the *SLCO1B1* assay has been designed to detect the variant rs4149056, which is present in three reduced function haplotypes namely, *SLCO1B1*5*, *SLCO1B1*15* and *SLCO1B1*17*. The *SLCO1B1* assay is thus, unable to differentiate between each of the three aforementioned haplotypes. The variants covered by the kit are outlined in Table 1.

Assays were set up on a 96-well plate. Human gDNA was added at a concentration of 2 ng/µL as template for the qPCR reaction, which was then performed on the Bio-Rad CFX96 IVD Touch™ Real-Time PCR Detection System per the product insert. Run analysis was performed using the application CFX Manager 3.1 or CFX Maestro, and exported as raw .csv files. Exported files were uploaded into the companion software, Nala Clinical Decision Support™ (Nala CDS™) for further analysis of variant genotyping, diplotype determination and phenotype

105 translation. Genotyping using Nala PGx Core™ was performed at the Molecular Diagnosis Centre, National

106 University Health System, Singapore (NUHS MDC) and PT Nalagenetik Riset Indonesia.

Agena VeriDose® Core and *CYP2D6* Copy Number Variation (CNV) Panel

The VeriDose® Core and *CYP2D6* Copy Number Variation (CNV) Panel from Agena Bioscience® consists of 68

variant assays in 20 genes and 5 *CYP2D6* CNV assays, accompanied by a reporting software that automatically

analyzes each variation[26]. Genotyping using Agena VeriDose Core and *CYP2D6* CNV Panel was performed at the

Genome Institute of Singapore. Variants evaluated using this platform are listed in Table 1. The Agena VeriDose®

Panel has been utilized by the United States Centers for Disease Control and Prevention (CDC) as part of their

Genetic Testing Reference Material (GeT-RM) Coordination Program[27].

TaqMan® Drug Metabolism Enzyme (DME) Genotyping Assay

TaqMan® DME Genotyping Assays were utilized in the evaluation of *CYP2D6* rs769258 (TaqMan Assay ID

AH21B9N) and *CYP2D6* rs267608319 (TaqMan Assay ID C__27102444_F0). Assays were set up on a 384-well

plate[28] with a sample input of human gDNA at 2 ng/µL. The subsequent PCR reaction was performed on the

Applied Biosystems ViiA™ 7 Real-Time PCR System as per the recommended cycling conditions, at the Genome

Institute of Singapore. Post-PCR plate read was performed using the companion software, TaqMan® Genotyper™

Software for single nucleotide polymorphisms (SNP) genotyping. Similar to the Agena VeriDose® Panel,

TaqMan® DME Genotyping Assays were employed in the characterization of DNA samples as part of the CDC

GeT-RM program[27].

Robustness

125 Genotype- and diplotype-level call rates were defined as the percentage of samples that returned a genotype at the

126 variant-level or were assigned a distinct diplotype for the gene of interest, respectively. Failed tests were defined as

127 samples that did not return a genotype and/or diplotype call for the genes evaluated.

$$\text{Call Rates, \%} = \frac{\text{Total Sample Size-Failed Tests}}{\text{Total Sample Size}} \times 100\%$$

Precision

Three samples at 3 DNA concentrations were tested across 3 reagent lots on 2 machines. Each test condition was
repeated within the same plate for a triplicate. For variant assays that identified SNPs and indels, intra-precision was
performed within the same plate, run as triplicates across 47 tests. Inter-precision was assessed from 120 tests
performed across plate runs covering the 4 variables – samples, DNA concentration, reagent lots and machines.
Concordance rates across precision studies were calculated as the percentage of tests that returned a genotype call
concordant to the expected truth for each variant assay. Discordant genotype was defined as instances when the test
returned a genotype call that was different from the expected truth.

$$\text{Concordance Rate, \%} = \frac{\text{No. Of Tests Performed} - \text{No. Of Tests With Discordant Genotype}}{\text{No. Of Tests Performed}} \times 100\%$$

138 For *CYP2D6* CNV assays, copy number estimates for Intron 2 and Exon 9 of the three samples were derived based
139 on their cycle threshold (Ct) results across plate runs.

$$\text{Copy Number} = 2 \times 2^{-\Delta\Delta Ct}$$

$$\Delta\Delta Ct = \left(Ct_{\text{reference gene calibrator}} - Ct_{\text{CYP2D6 calibrator}}\right) - \left(Ct_{\text{reference gene}} - Ct_{\text{CYP2D6 sample}}\right)$$

142 Testing of the three samples was repeated for a number of plate runs, n (Table 5), and calculated for the average
143 copy number of each sample and their coefficient of variation (CV). The CV for each plate run was calculated by
144 finding the standard deviation (σ_{plate}) between triplicates within the same plate run, and divided by the triplicate
145 mean (μ_{plate}). The average of the individual CVs was reported as the intra-precision CV. For inter-precision CV,
146 standard deviation population ($\sigma_{plate\ means}$) was divided by the mean population, i.e. average of means.

$$\text{Intra- CV,\%} = \frac{\sum_n \frac{\sigma_{\text{plate}}}{\mu_{\text{plate}}}}{n} \times 100\%$$

$$\text{Inter- CV,\%} = \frac{\sigma_{\text{plate means}}}{\sum_n \mu_{\text{plate}} / n} \times 100\%$$

Accuracy

Variant-level Concordance

The accuracy of Nala PGx Core™ in genotyping at a variant-level was evaluated by comparing calls produced by
Nala PGx Core™ assay against benchmark methods as listed in Table 1. Samples that successfully produced
genotype calls for all variants tested on Nala PGx Core™ and its benchmarks were considered for the evaluation (n
= 225). Samples that failed to produce a genotype call on one or more of the platforms were excluded from the
concordance calculation (n = 21, Supplementary Table 8). Discordant calls were defined as instances in which Nala
PGx Core™ provided a genotype call that was different from that of a call made by the corresponding benchmark.
Percentage concordance to the benchmark was calculated per variant as follows –

$$\text{Concordance To Benchmark Per Variant, \%} = \frac{\text{Total Sample Size - Discordant Calls By Nala PGx Core™}}{\text{Total Sample Size}} \times 100\%$$

Diplotype-level Concordance

The accuracy of Nala PGx Core™ in assigning a diplotype call for *CYP2C9, CYP2C19,* and *CYP2D6*, was

evaluated by comparing calls against the Agena VeriDose® Core and *CYP2D6* CNV Panel. Samples that met the

following criteria were included in the sample size of each gene:

1.   Successful genotype-level calls on the relevant platforms for all variants covered by the gene of interest

2.   Successful assignment of a diplotype for the gene of interest on both Nala PGx Core™, and Agena

163   VeriDose® Core and *CYP2D6* CNV Panel

164   Discordant calls were defined as instances in which Nala PGx Core™ assigned a diplotype that differed from the

165   call made by the Agena VeriDose® Core and *CYP2D6* CNV Panel.

$$\text{Concordance, \%} = \frac{\text{Total Sample Size - Discordant Calls By Nala PGx Core™}}{\text{Total Sample Size}} \times 100\%$$

Frequencies By Ethnicity

Ethnicities were obtained based on participant self-identification across both the population cohorts as part of the

recruitment questionnaire. Participants that did not report an ethnic group in the recruitment form were excluded

from the frequency analysis. The following were further excluded from the frequency analysis:

1.   Samples in which participants did not report an ethnic group on the recruitment form (n=5)

2.   Samples with one or more variant level failures across the 4 genes evaluated in Table 1 (n=18)

3.   Samples with one or more diplotype-level failures ("No Call") across *CYP2C9, CYP2C19,* and *CYP2D6* on

173   Nala PGx Core™ and the Agena VeriDose® Core and *CYP2D6* CNV Panel (n=14)

174   4.   Samples with discordant diplotype calls for the gene of interest (n=variable, Supplementary Table 6)

176   The remaining samples were included in the allele-level frequency analysis of *CYP2C9* (n=206), *CYP2C19* (n=201),

177   *CYP2D6* (n=195) and *SLCO1B1* (n=203), as well as in the diplotype-level frequency analysis of *CYP2C9* (n=206),

178   *CYP2C19* (n=201) and *CYP2D6* (n=195). Allele and diplotype frequency values were derived using the following

179   formulae[29,30], for both the overall study cohort as well as for each ethnic group.

$$\text{Frequency Of Allele 'X' In A Given Population} = \frac{\text{Total Copies Of Allele 'X'}}{\text{Total Copies Of All Alleles For The Gene Of Interest}}$$

$$\text{Frequency Of Diplotype 'X' In A Given Population} = \frac{\text{Total Instances Of Diplotype 'X'}}{\text{Total Number Of Individuals In The Population}}$$

Results

Robustness

We first sought to evaluate the observed genotype- and diplotype-level call rates of the platforms evaluated in this study. 246 samples underwent variant genotyping and diplotype determination, across the four genes evaluated on the genotyping platforms (Tables 2, 3). The genotype-level call rates for Nala PGx Core™ were at 100% for *CYP2C9*, *CYP2C19* and *SLCO1B1*, and the diplotype-level call rates were at 100% for *CYP2C9* and *CYP2C19*. The benchmark platform, Agena VeriDose® Core Panel, demonstrated call rates of >98% at the genotype-level and >97% at the diplotype-level.

Most variants in *CYP2D6*, except for seven, achieved 100% call rates on Nala PGx Core™, while the corresponding call rates of the benchmark platforms were observed to be between 95.9 – 99.2% on Agena VeriDose® Core and *CYP2D6* CNV Panel, and 100% on TaqMan® DME Genotyping Assays. Out of the seven aforementioned variants, Nala PGx Core™ demonstrated higher call rates than the benchmark for the genotyping of rs1065852, Intron 2 and Exon 9 variants. For rs3892097, rs769258, rs5030865, and rs267608319, the accompanying benchmarks demonstrated higher call rates. At the diplotype-level, Nala PGx Core™ demonstrated a CYP2D6 call rate of 95.9% as compared to the benchmark, which was observed to be at 90.7% (Table 3).

Precision

A precision study was conducted to assess the consistency of Nala PGx Core™ for samples tested under the same conditions (intra-precision) and under different conditions (inter-precision). Both study resulted in 100% concordance for all assays across replicates, demonstrating consistent genotyping results across a range of DNA concentration, reagent lots and machine variations (Table 4). Precision of *CYP2D6* CNV assay was reported as the average copy number obtained for Intron 2 and Exon 9 of three samples, and their CV calculated across the test conditions (Table 5). The intra- CV ranged from 3-6% while inter- CV between 5-13%, demonstrating high precision of the assays across variables, where acceptable ranges were intra- CV below 10% and inter- CV below 15%[31].

Accuracy

Variant-level Concordance

To assess the accuracy of the panel, 18 SNPs and 2 *CYP2D6* Copy Number assays were genotyped on the panel, Nala PGx Core™, against benchmark methods as listed in Table 1. The 225 sample cohort consisted of DNA samples isolated from buccal swabs that had successfully produced genotype calls for all variants tested on Nala PGx Core™ and its benchmarks. The sample size per genotype displayed in Table 6 refers to the genotype calls made by the benchmark.

11 variants (*CYP2C9* rs1799853, rs1057910; *CYP2C19* rs12248560; *CYP2D6* rs5030655, rs3892097, rs35742686,
rs28371725, rs769258, rs5030656, rs59421388, rs267608319) were genotyped against Agena VeriDose® Core with
a resulting concordance rate of 100% (N = 225 samples). Discordance was observed for *CYP2C19* rs4244285 (n=7),
CYP2C19 rs4986893 (n=3), *CYP2D6* rs1065852 (n=1), *CYP2D6* rs16947 (n=7), *CYP2D6* rs1135840 (n=5),
CYP2D6 rs5030865 (n=1) and *SLCO1B1* rs4149056 (n=6), with a mismatch rate of 0.44% to 3.1% for the affected
assays. Overall, Nala PGx Core™ demonstrated >96% concordance to the benchmark, Agena VeriDose® Core, for
the 16 variants across 225 samples (Table 6).
222
223 Variants not present on Agena VeriDose® Core, *CYP2D6* rs769258 and *CYP2D6* rs267608319, were genotyped
224 using TaqMan® DME Genotyping Assays. Nala PGx Core™ demonstrated 100% concordance (N = 225) to the
225 benchmark for both SNPs.
226
227 For the *CYP2D6* Intron 2 and Exon 9 Copy Number assays, concordance was observed to be at 99.6% and 98.7%
228 respectively, against the Agena *CYP2D6* CNV Panel. Discordant calls were observed in samples with an Intron 2
229 copy number greater than 3 (n=1), and for samples with an Exon 9 copy number of one (n=1) and two (n=2). The
230 sample size per copy number displayed in Table 7 refers to the copy number calls made by the benchmark.
231
232 A summary of the concordance percentage as well as the distribution of genotypes of each variant are displayed in
233 Table 6. Similarly, the distribution of *CYP2D6* Intron 2 and Exon 9 copy numbers are displayed in Table 7.
234
Diplotype-level Concordance
Following successful genotyping at the variant level, we investigated the accuracy of Nala PGx Core™ in assigning
a diplotype call for *CYP2C9*, *CYP2C19* and *CYP2D6*, with reference to the Agena VeriDose® Core and *CYP2D6*
CNV Panel. Table 8 displays the percentage concordance after the further exclusion of samples that demonstrated
diplotype mismatches arising from technological differences, where technological differences refer to the varying
allele coverage of each platform (Supplementary Table 7). These differences were derived from the variant lists of
both Nala PGx Core™ (Table 1) and its benchmark, the Agena VeriDose® Core and *CYP2D6* CNV Panel[26].
242
243 Overall, a percentage agreement of 100% for *CYP2C9* (n=213), 96.4% for *CYPC219* (n=223) and 94.7% for
244 *CYP2D6* (n=209) was observed between Nala PGx Core™ and the benchmark. Discordance was observed at n=1
245 for all diplotypes listed in Table 8 except for the following with more than one discordant calls: *CYP2C19* *2/*2
246 (n=4), and *CYP2D6* *4/*36 (n=2).
247
Frequencies By Ethnicity
For samples that were concordant on Nala PGx Core™ and the benchmark platforms, we were able to observe the
allele frequencies amongst the populations residing in Singapore and Indonesia (Table 9). From the combination of
alleles present in individual's chromosome, we were able to observe both the diplotype and corresponding
phenotype frequencies amongst our study population (Table 10).

For *CYP2C9*, *3 allele was the most common amongst Chinese and Malay, and *2 allele amongst Caucasian, which
is in line with PharmGKB's reported distribution for the East Asian and European populations respectively. Our
study also reported *3 allele as the more common variant in Indian population than *2, as opposed to PharmGKB's
frequency. These allele frequencies translated to *1/*3 as a common diplotype observed in Chinese, Malay and
Indians, and *1/*2 in Caucasians.

For *CYP2C19*, we observed highest frequency of *CYP2C19*2* amongst Chinese, Malay and Indonesian which were
categorized as East Asian populations. This resulted into high frequency of *1/*2 heterozygous depicted as a
common diplotype amongst the population. The alleles *2 and *17 were observed as the common variants at equal
proportions of 0.229 in Indians and 0.138 in Caucasians. *CYP2C19*3* was a common minor allele least observed
amongst Indians and Caucasians, 0 and 0.017 respectively. As a result, *1/*2 and *1/*17 were common diplotypes
observed in Indian and Caucasian populations, and *2/*17 only seen in Indians.

Common polymorphisms of *CYP2D6* in our population were seen in *10 and *36 alleles, at almost two-fold higher
frequencies in Chinese, Malay and Indonesian than Indians. We noticed high frequencies of *36 in our East Asian
population. Additionally, two functional copies of *36 allele were present in 1.4% of Chinese population who
participated in our study (Figure 1). These alleles resulted in high frequency of *10/*36 as common diplotype
amongst Chinese, Malay and Indonesian population. Our Indian population observed equal distribution of alleles *2,
*10 and *36, although PharmGKB reported only *2 as the highest frequency in South Asians. The corresponding
common diplotypes observed in Indians were of equal proportions in *1/*2, *1/*36 and *1/*10 ranging from 0.12 to
0.16. The alleles *2 and *4 were most common amongst Caucasians as reported by PharmGKB in Europeans,
resulting in common diplotypes of *1/*2 and *1/*4 in the population.

For *SLCO1B1*, the frequencies of rs4149056 across all ethnicities were consistent with values reported in gnomAD,
with the variant being most common amongst Caucasians (0.167) and least amongst Indians (0.040). The frequency
amongst East Asians (0.125, gnomAD), as denoted by the Chinese and Indonesian ethnic groups in this study,
ranged between 0.074 and 0.125 respectively.

Discussion

In this study, we evaluated the performance of Nala PGx Core™, a qPCR-based panel that evaluates 18 variants and
2 *CYP2D6* Copy Number markers across 4 pharmacogenes with established relevance across major ethnic groups in
Singapore and Indonesia population. Nala PGx Core™ comes coupled with a reporting software that supports
variant detection, diplotype assignment, diplotype-to-phenotype translation and the generation of reports containing
clinical recommendations for each phenotype. The panel demonstrated high genotype-level call rates of >97% for
CYP2D6, and 100% for *CYP2C9*, *CYP2C19* AND *SLCO1B1*. Similarly, high diplotype-level call rates were
observed at >95% for *CYP2D6*, and 100% for *CYP2C9* and *CYP2C19*. A precision of 100% was observed under the

same conditions (intra) and across different conditions (inter). In comparison to other established platforms serving

as benchmarks during the study, Nala PGx Core™ had ≥96.9% concordance rate for all variant level assays, which

consequently resulted in ≥94.7% concordance at a diplotype level across *CYP2C9*, *CYP2C19* and *CYP2D6*.

Failures to produce a variant genotype call could be attributed to several reasons. Firstly, failures could potentially

stem from the quality of gDNA, despite the DNA quality checks (QC) performed prior to accepting a sample for

testing. Further QC that involves specific quantification of double-stranded non-fragmented DNA and traces of other

interfering materials like RNA, carryover carbohydrate, residual phenol, guanidine or other reagents could enhance

the call rate[32]. Regardless, the overall higher variant call rates on Nala PGx Core™ panel demonstrate high tolerance

of interfering substances, therefore alluding to the high robustness of the assay. Often, failures at variant genotyping

subsequently contribute to failures at determining diplotype, since an incomplete variant panel cannot translate into

a diplotype. Failures at diplotype calling could also arise from a combination of variants that do not map onto a

distinct diplotype, per the reference database, potentially indicating a novel combination.

We further sought to evaluate the allele frequency distribution in the study cohort across the 5 major ethnic groups

observed (Indonesian, Chinese, Malay, Indian and Caucasian). The data presented was limited strictly to the

geographical boundaries of Singapore and Indonesia, which could account for the difference in allele frequencies

observed in comparison to PharmGKB, which is representative of a more expansive and global cohort[33]. Whilst

dissimilar to database figures, our study demonstrated the distributions for the following to be concordant with

previous studies[7,34,35], suggesting a niche in the PGx landscape of Singapore and Indonesia –

1. *CYP2C9*3* allele as the more common variant within Indians than *CYP2C9*2*[7]

2. A *SLCO1B1* rs4149056 frequency of 12.5% amongst Indonesians[34]

3. High frequencies of the *CYP2D6*10* amongst Indonesians[34]

4. High frequencies of the *CYP2D6*36* allele as seen in the Indonesian, Chinese and Malay ethnicities[35]

5. Two functional copies of *CYP2D6* within our Chinese population[35]

Due to the lack of *CYP2D6* copy number references[34], it is our understanding that the frequency of *CYP2D6*36* in

Indonesia may not be well-represented. Our study revealed that the prevalence of *CYP2D6*36* to be approximately

seventeen times higher amongst Indonesians as compared to the corresponding East Asian allele frequency on

PharmGKB[36]. Furthermore, our study provides insight on the frequency of the *CYP2D6 *10/*36* diplotype in the

archipelago, which may help inform the adoption of population-specific PGx workflows in the region. Taken

together, the data presents a case for extending tailored PGx testing across the 4 pharmacogenes studied, *CYP2C9*,

CYP2C19, CYP2D6 and *SLCO1B1*, in South East Asia.

Limitations

The single largest reason for the diplotype-level discordance observed between Nala PGx Core™ and the Agena

VeriDose® Core and *CYP2D6* CNV Panel appears to be a difference in the technological offerings and hence, allele

coverage of each panel. From Supplementary Table 7, out of the 28 instances of discordance observed, 9 instances

were attributed to this cause (32.1%). This included two alleles each in *CYP2C9*, *CYP2C19* and *CYP2D6*.

The variants *CYP2C9*12*, *CYP2C9*13*, *CYP2C19*4A*, *CYP2C19*6* and *CYP2D6*13* were covered by Agena

VeriDose® Core but not by Nala PGx Core™. Due to lack of coverage for these alleles, the genotypes identified by

Nala PGx Core™ were inconsistent with the calls produced by Agena VeriDose® Core. It is important to note that

Nala PGx Core™ was curated specifically to include alleles that were relevant to Asian and European populations,

with a Minor Allele Frequency (MAF) >1% based on PharmGKB database for East Asians, South Asians and

Europeans[36]. Similarly, *CYP2D6*35*, an allele present in 5.5% of Europeans and 1.2% of South Asians, was

identified by Nala PGx Core™ but not by Agena VeriDose® Core. The technological differences and subsequent

inconsistencies in reporting suggest the need for further review from a product development standpoint.

For the highly polymorphic *CYP2D6* gene, Nala PGx Core™ was designed to only cover the variant alleles listed in

Table 1, that could lead to overestimation of *1 allele in our population. This means star alleles that may be present

in the population were identified as normal genotype, which could result in overestimation of *1 allele in our

population as compared to the PharmGKB's reported frequency (Supplementary Table 4).

The study also relied on self-reported ethnicity, which may have contributed to racial bias in summarizing the

observed allele and diplotype frequencies per ethnic group. Future studies may consider screening the ethnicities of

participants as well as their next-of-kin, up to two generations prior[10], to improve classification accuracy.

Finally, the study cohort consisted of 246 individuals, with each ethnic group consisting of approximately 24 to 75

individuals for the frequency analysis. Larger scale studies may be conducted to ensure that the trends observed as

part of this study may be generalized across the populations examined.

Future Work

Further investigation is required for the remaining samples that had discrepant genotypes across platforms. Nala

PGx Core™ was designed as a qPCR platform and genotyping was accompanied by Nala CDS™ that utilized

quantitative thresholds. There could be cases of samples displaying borderline or ambiguous qPCR amplification

curves and Ct-values, which could be misinterpreted by these thresholds. The benchmark platforms, Agena

VeriDose® Core and *CYP2D6* CNV Panel and TaqMan® DME Genotyping Assays, were also analyzed by their

accompanying software packages. Such ambiguous cases on either platform could lead to instances of discordance

and sequencing of the region of interest may be required to ascertain the truth. For gene variants caused by SNPs

and indels, the qPCR amplified product can be submitted for genome sequencing[13]. In the investigation of copy

number variants, a two-step approach can be adopted[37]. Firstly, CNV calls observed in this study may be checked

against an alternative benchmark such as digital droplet PCR (ddPCR)[38]. Secondly, where discordance in CNV

calling persists, methods such as XL-PCR can be adopted to identify the presence of *CYP2D6/CYP2D7* hybrid

states, with submitting samples for whole-genome sequencing[37] being an option. Investigation of copy number
variants may require amplification of region spanning across the gene of interest.

Conflict of Interest

Funding

All expenses for this study were funded by Nalagenetics Pte Ltd and PT Nalagenetik Riset Indonesia.

Data Availability Statement

The data that supports the findings of this study are available as supplementary materials. Any additional data is
available from the corresponding author upon reasonable request.

References

1. Berm EJJ, Looff M de, Wilffert B, et al. Economic Evaluations of Pharmacogenetic and Pharmacogenomic
Screening Tests: A Systematic Review. Second Update of the Literature. Bruns H, ed. *PLoS One*.
2016;11(1):e0146262. doi:10.1371/journal.pone.0146262
2. Chan SL, Ang X, Sani LL, et al. Prevalence and characteristics of adverse drug reactions at admission to
hospital: a prospective observational study. *Br J Clin Pharmacol*. 2016;82(6):1636-1646.
doi:10.1111/bcp.13081
3. Sultana J, Cutroneo P, Trifirò G. Clinical and economic burden of adverse drug reactions. *J Pharmacol
Pharmacother*. 2013;4(SUPPL.1). doi:10.4103/0976-500X.120957
4. Zhu Y, Lopes GS, Bielinski SJ, et al. Impact of Pharmacogenomic Information on Values of Care and
Quality of Life Associated with Codeine and Tramadol-Related Adverse Drug Events. *Mayo Clin Proc
Innov Qual Outcomes*. 2021;5(1):35-45. doi:10.1016/j.mayocpiqo.2020.08.009
5. Furge LL, Guengerich FP. Cytochrome P450 enzymes in drug metabolism and chemical toxicology: An
introduction. *Biochem Mol Biol Educ*. 2006;34(2):66-74. doi:10.1002/bmb.2006.49403402066
6. Malsagova KA, Butkova T V, Kopylov AT, et al. Pharmacogenetic Testing: A Tool for Personalized Drug
Therapy Optimization. *Pharmaceutics*. 2020;12(12):1-23. doi:10.3390/pharmaceutics12121240
7. Ling Goh L, Wei Lim C, Cheng Sim W, Xian Toh L, Pang Leong K. Analysis of Genetic Variation in
CYP450 Genes for Clinical Implementation. Published online 2017. doi:10.1371/journal.pone.0169233
8. Kisor DF, Hoefer C, Decker BS. Pharmacogenomics and precision medicine. In: *Clinical Pharmacy
Education, Practice and Research: Clinical Pharmacy, Drug Information, Pharmacovigilance,
Pharmacoeconomics and Clinical Research*. Elsevier; 2018:437-451. doi:10.1016/B978-0-12-814276-
9.00031-3
9. Ooi BNS, Raechell, Ying AF, et al. Robust Performance of Potentially Functional SNPs in Machine
Learning Models for the Prediction of Atorvastatin-Induced Myalgia. *Front Pharmacol*. 2021;12:605764.
doi:10.3389/fphar.2021.605764
10. Brunham LR, Chan SL, Li R, et al. Pharmacogenomic diversity in Singaporean populations and Europeans.
Pharmacogenomics J. 2014;14(6):555-563. doi:10.1038/tpj.2014.22
11. Bachtiar M, Ooi BNS, Wang J, et al. Towards precision medicine: interrogating the human genome to
identify drug pathways associated with potentially functional, population-differentiated polymorphisms.
Pharmacogenomics J. 2019;19(6):516-527. doi:10.1038/s41397-019-0096-y

12.   Perkel J. SNP genotyping: Six technologies that keyed a revolution. *Nat Methods*. 2008;5(5):447-453.
doi:10.1038/nmeth0508-447
13.   Yang W, Wu G, Broeckel U, et al. Comparison of genome sequencing and clinical genotyping for
pharmacogenes. *Clin Pharmacol Ther*. 2016;100(4):380-388. doi:10.1002/cpt.411
14.   S.A. Deepak, K.R. Kottapalli, R. Rakwal, et al. Real-Time PCR: Revolutionizing Detection and Expression
Analysis of Genes. *Curr Genomics*. 2007;8(4):234-251. doi:10.2174/138920207781386960
15.   Pratt VM, Zehnbauer B, Wilson JA, et al. Characterization of 107 genomic DNA reference materials for
CYP2D6, CYP2C19, CYP2C9, VKORC1, and UGT1A1: A GeT-RM and association for molecular
pathology collaborative project. *J Mol Diagnostics*. 2010;12(6):835-846. doi:10.2353/jmoldx.2010.100090
16.   Pratt VM, Everts RE, Aggarwal P, et al. Characterization of 137 Genomic DNA Reference Materials for 28
Pharmacogenetic Genes: A GeT-RM Collaborative Project. *J Mol Diagnostics*. 2016;18(1):109-123.
doi:10.1016/j.jmoldx.2015.08.005
17.   Syvänen AC. Accessing genetic variation: Genotyping single nucleotide polymorphisms. *Nat Rev Genet*.
2001;2(12):930-942. doi:10.1038/35103535
18.   Suo W, Shi X, Xu S, Li X, Lin Y. Towards low cost, multiplex clinical genotyping: 4-fluorescent
Kompetitive Allele-Specific PCR and its application on pharmacogenetics. Kalendar R, ed. *PLoS One*.
2020;15(3):e0230445. doi:10.1371/journal.pone.0230445
19.   Shen GQ, Abdullah KG, Wang QK. The TaqMan method for SNP genotyping. *Methods Mol Biol*.
2009;578:293-306. doi:10.1007/978-1-60327-411-1_19
20.   Davis BH, DeFrank G, Limdi NA, Harada S. Validation of the Spartan RX CYP2C19 Genotyping Assay
Utilizing Blood Samples. *Clin Transl Sci*. 2020;13(2):260-264. doi:10.1111/cts.12714
21.   Zhou Y, Armstead AR, Coshatt GM, Limdi NA, Harada S. Comparison of two point-of-care cyp2c19
genotyping assays for genotype-guided antiplatelet therapy. *Ann Clin Lab Sci*. 2017;47(6):738-743.
22.   Bergmeijer TO, Vos GJ, Claassens DM, et al. Feasibility and implementation of CYP2C19 genotyping in
patients using antiplatelet therapy. *Pharmacogenomics*. 2018;19(7):621-628. doi:10.2217/pgs-2018-0013
23.   Chae H, Kim M, Koh Y-S, et al. Feasibility of a Microarray-Based Point-of-Care CYP2C19 Genotyping
Test for Predicting Clopidogrel On-Treatment Platelet Reactivity. *Biomed Res Int*. 2013;2013:1-5.
doi:10.1155/2013/154073
24.   Lee CC, McMillin GA, Babic N, Melis R, Yeo K-TJ. Evaluation of a CYP2C19 genotype panel on the
GenMark eSensor® platform and the comparison to the Autogenomics Infiniti™ and Luminex CYP2C19
panels. *Clin Chim Acta*. 2011;412(11-12):1133-1137. doi:10.1016/j.cca.2011.03.001
25.   Vanwong N, Ngamsamut N, Hongkaew Y, et al. Detection of CYP2D6 polymorphism using Luminex
xTAG technology in autism spectrum disorder: CYP2D6 activity score and its association with risperidone
levels. *Drug Metab Pharmacokinet*. 2016;31(2):156-162. doi:10.1016/j.dmpk.2016.01.005
26.   Lois A, Everts RE, Nakorchevsky A, et al. The VeriDose Core Panel: Strong Performance When Analyzing
Challenging Pharmacogenetic Samples. Accessed April 2, 2021. https://agenabio.com/wp-
content/uploads/2020/06/Agena-Conference-Poster-Veridose-Core-V3-G020.pdf
27.   Gaedigk A, Turner A, Everts RE, et al. Characterization of Reference Materials for Genetic Testing of
CYP2D6 Alleles. *J Mol Diagnostics*. 2019;21(6):1034-1052. doi:10.1016/j.jmoldx.2019.06.007
28.   Technologies L. *TaqMan ® SNP Genotyping Assays TaqMan ® Predesigned SNP Genotyping Assays,*
TaqMan ® Custom SNP Genotyping Assays, and TaqMan ® Drug Metabolism Enzyme Genotyping Assays.;
2014.
29.   Population Genetics and the Hardy-Weinberg Principle. Accessed May 2, 2021.
https://www.cs.cmu.edu/~genetics/units/instructions/instructions-PGE.pdf
30.   Allele frequency. Accessed May 2, 2021. https://www.nature.com/scitable/definition/allele-frequency-298/
31.   Calculating Inter- and Intra-Assay Coefficients of Variability. Accessed May 1, 2021.
https://salimetrics.com/calculating-inter-and-intra-assay-coefficients-of-variability/
32.   Matlock B. Assessment of Nucleic Acid Purity. *Tech Bull NanoDrop Spectrophotometers*. Published online
2015:1-2.
33.   Huddart R, Fohner AE, Whirl□Carrillo M, et al. Standardized Biogeographic Grouping System for
Annotating Populations in Pharmacogenetic Research. *Clin Pharmacol Ther*. 2019;105(5):1256-1262.
doi:10.1002/cpt.1322
34.   Runcharoen C, Fukunaga K, Sensorn I, et al. Prevalence of pharmacogenomic variants in 100
pharmacogenes among Southeast Asian populations under the collaboration of the Southeast Asian
Pharmacogenomics Research Network (SEAPharm). *Hum Genome Var*. 2021;8(1):7. doi:10.1038/s41439-
021-00135-z

459 35. Chan W, Li MS, Sundaram SK, Tomlinson B, Cheung PY, Tzang CH. CYP2D6 allele frequencies, copy
460    number variants, and tandems in the population of Hong Kong. *J Clin Lab Anal*. 2019;33(1).
461    doi:10.1002/jcla.22634
462 36. Whirl-Carrillo M, McDonagh EM, Hebert JM, et al. Pharmacogenomics Knowledge for Personalized
463    Medicine. *Clin Pharmacol Ther*. 2012;92(4):414-417. doi:10.1038/clpt.2012.96
464 37. Pös O, Radvanszky J, Styk J, et al. Copy Number Variation: Methods and Clinical Applications. *Appl Sci*.
465    2021;11(2):819. doi:10.3390/app11020819
466 38. Gross AM, Ajay SS, Rajan V, et al. Copy-number variants in clinical genome sequencing: deployment and
467    interpretation for rare and undiagnosed disease. *Genet Med*. 2019;21(5):1121-1130. doi:10.1038/s41436-
468    018-0295-y

471 Tables

472 **Table 1** Genes and variants evaluated

Gene	Star Allele	Variant	Nucleotide Changes	Effect On Protein	Allele Clinical Function Status	Genotyping Methods Utilized		
						Nala PGx Core™	Agena VeriDose® Core and *CYP2D6* CNV Panel	TaqMan® DME Genotyping Assays
CYP2C9	*2	rs1799853	3608C>T	R144C	Decreased	Bi-allelic Assay	VeriDose Core	
	*3	rs1057910	42614A>C	I359L	None	Bi-allelic Assay	VeriDose Core	
CYP2C19	*2	rs4244285	19154G>A	Splicing Defect	None	Bi-allelic Assay	VeriDose Core	
	*3	rs4986893	17948G>A	W212X	None	Bi-allelic Assay	VeriDose Core	
	*17	rs12248560	-806C>T	5' Region	Increased	Bi-allelic Assay	VeriDose Core	
CYP2D6	*2	rs1135840	4181G>C	S486T	Normal	Bi-allelic Assay	VeriDose Core	
	*2	rs16947	2851C>T	R296C	Normal	Bi-allelic Assay	VeriDose Core	
	*3	rs35742686	2550delA	Frameshift	None	Bi-allelic Assay	VeriDose Core	
	*4	rs3892097, rs1065852	1847G>A, 100C>T	Splicing Defect, P34S	None	Bi-allelic Assay	VeriDose Core	
	*5	N/A	N/A	Gene Deletion	None	CNV Assay (Intron 2)	CYP2D6 CNV	
	*6	rs5030655	1708delT	Frameshift	None	Bi-allelic Assay	VeriDose Core	
	*8	rs5030865	1759G>T	G169X	None	Tri-allelic Assay	VeriDose Core	
	*9	rs5030656	2616delAAG	K281del	Decreased	Bi-allelic Assay	VeriDose Core	
	*10	rs3892097, rs1065852	1847G>A, 100C>T	Splicing Defect, P34S	Decreased	Bi-allelic Assay	VeriDose Core	
	*14	rs5030865	1759G>A	G169R	Decreased	Tri-allelic Assay	VeriDose Core	
	*21	rs72549352	2580_2581insC	Frameshift	None	Bi-allelic Assay	VeriDose Core	
	*29	rs59421388	3184G>A	V338M	Decreased	Bi-allelic Assay	VeriDose Core	
	*31	rs267608319	4043G>A	R440H	None	Bi-allelic Assay	NA	AH21B9N
	*35	rs769258	31G>A	V11M	Normal	Bi-allelic Assay	NA	C__27102444_F0
	*36	N/A	Recombination at Exon 9	CYP2D6-2D7 Hybrid	None	CNV Assay (Exon 9)	CYP2D6 CNV	
	*41	rs28371725	2989G>A	Splicing Defect	Decreased	Bi-allelic Assay	VeriDose Core	
SLCO1B1	NA†	rs4149056	g.52422T>C	V174A	Decreased	Bi-allelic Assay	VeriDose Core	

†Nala PGx Core™ detects the variant, rs4149056, which is associated with decreased enzymatic activity and is present in three known *SLCO1B1* haplotypes namely, *SLCO1B1*5*, *SLCO1B1*15* and *SLCO1B1*17*.

473
474
475

476   **Table 2** Observed genotype-level call rates per variant per gene per platform

Gene	Variant	Variant Call Rate, % (n=246)		
		Nala PGx Core™	Agena VeriDose® Core and *CYP2D6* CNV Panel	TaqMan® DME Genotyping Assays
CYP2C9	rs1799853	100.0	99.2	NA
	rs1057910	100.0	99.6	NA
CYP2C19	rs4244285	100.0	99.6	NA
	rs4986893	100.0	99.6	NA
	rs12248560	100.0	98.8	NA
CYP2D6	rs1065852	98.4	95.9	NA
	rs5030655	100.0	99.2	NA
	rs3892097	98.8	99.2	NA
	rs35742686	100.0	98.8	NA
	rs16947	100.0	99.6	NA
	rs28371725	100.0	99.2	NA
	rs1135840	100.0	99.6	NA
	rs769258	98.8	NA	100.0
	rs5030865	97.2	99.2	NA
	rs5030656	100.0	99.6	NA
	rs59421388	100.0	99.2	NA
	rs267608319	99.6	NA	100.0
	CNV Assay (Intron 2)	99.6	99.2	NA
	CNV Assay (Exon 9)	99.6	99.2	NA
SLCO1B1	rs4149056	100.0	99.2	NA

477
478   **Table 3** Observed diplotype-level call rates per gene per platform

Gene	Diplotype Call Rate, % (n=246)	
	Nala PGx Core™	Agena VeriDose® Core and *CYP2D6* CNV Panel
CYP2C9	100.0	97.2
CYP2C19	100.0	98.8
CYP2D6	95.9	90.7

479
480   **Table 4** Intra-precision and inter-precision concordance rates

Gene	Variant	Concordance rate, %	
		Intra-precision (47 tests in triplicate)	Inter-precision (120 tests across 4 variables)
CYP2C9	rs1799853	100%	100%
	rs1057910	100%	100%
CYP2C19	rs4244285	100%	100%
	rs4986893	100%	100%
	rs12248560	100%	100%
CYP2D6	rs1065852	100%	100%
	rs5030655	100%	100%
	rs3892097	100%	100%
	rs35742686	100%	100%

	rs16947	100%	100%
	rs28371725	100%	100%
	rs1135840	100%	100%
	rs769258	100%	100%
	rs5030865	100%	100%
	rs5030656	100%	100%
	rs59421388	100%	100%
	rs267608319	100%	100%
SLCO1B1	rs4149056	100%	100%

482 **Table 5** Intra-precision and inter-precision for *CYP2D6* Copy Number

Sample	Variant	Average Copy Number	Sample Size, n	Coefficient of Variation	
				Intra-precision	Inter-precision
221374	Intron 2	2	11	4%	6%
	Exon 9	1		5%	5%
222660	Intron 2	2	9	3%	5%
	Exon 9	2		4%	6%
222769	Intron 2	3	22	4%	10%
	Exon 9	2		6%	13%

484 **Table 6** Genotype concordance for genes *CYP2C9*, *CYP2C19*, *CYP2D6* and *SLCO1B1*

Gene	Variant	Genotype	Sample Size Per Genotype[†]	Concordance to Benchmark, %[‡]	Discordant Calls By Nala PGx Core™[§]
CYP2C9	**rs1799853**	C/C	212	100.0	0
	N = 225	C/T	13		0
		T/T	0		0
	rs1057910	A/A	204	100.0	0
	N = 225	A/C	21		0
		C/C	0		0
CYP2C19	**rs4244285**	G/G	121	96.9	1
	N = 225	A/G	89		2
		A/A	15		4
	rs4986893	G/G	206	98.7	2
	N = 225	A/G	19		1
		A/A	0		0
	rs12248560	C/C	0	100.0	0
	N = 225	C/T	0		0
		T/T	0		0
CYP2D6	**rs1065852**	G/G	77	99.6	0
	N = 225	A/G	76		1
		A/A	72		0
	rs5030655	A/A	224	100.0	0
	N = 225	delA/A	1		0
		delA/delA	0		0
	rs3892097	C/C	205	100.0	0
	N = 225	C/T	16		0
		T/T	4		0
	rs35742686	T/T	225	100.0	0
	N = 225	delT/T	0		0
		delT/delT	0		0
	rs16947	G/G	145	96.9	2
	N = 225	A/G	64		4
		A/A	16		1
	rs28371725	C/C	204	100.0	0
	N = 225	C/T	19		0
		T/T	2		0
	rs1135840	C/C	33	97.8	1

Gene	Variant	Genotype	Sample Size Per Genotype†	Concordance to Benchmark, %‡	Discordant Calls By Nala PGx Core™§
	N = 225	C/G	68		1
		G/G	124		3
	rs769258 N = 225	C/C	220	100.0	0
		C/T	5		0
		T/T	0		0
	rs5030865 N = 225	C/C	222	99.6	0
		C/T	3		1
		T/T	0		0
	rs5030656 N = 225	CTT/CTT	224	100.0	0
		delCTT/CTT	1		0
		delCTT/delCTT	0		0
	rs59421388 N = 225	C/C	224	100.0	0
		C/T	1		0
		T/T	0		0
	rs267608319 N = 225	C/C	225	100.0	0
		C/T	0		0
		T/T	0		0
SLCO1B1	rs4149056 N = 225	T/T	182	97.3	2
		C/T	42		4
		C/C	1		0

†"Sample Size Per Genotype" reflects the counts of each genotype as produced by the benchmark, and not by Nala PGx Core™.
‡Refer to Table 1 for an outline of the benchmark methods used per variant.
§Discordant calls refer to instances in which Nala PGx Core™ did not produce a genotype call concordant with the benchmark.

485
486 **Table 7** Genotype concordance for *CYP2D6* Intron 2 And Exon 9 Copy Number Variations

Gene	Variant	Copy Number	Sample Size Per Copy Number†	Concordance to Benchmark, %‡	Discordant Calls By Nala PGx Core™§
CYP2D6	Intron 2 N = 224	0	0	99.6	0
		1	12		0
		2	130		0
		>=3	82		1
	Exon 9 N = 224	0	0	98.7	0
		1	24		1
		2	193		2
		>=3	7		0

†"Sample Size Per Copy Number" reflects the counts of each copy number call as produced by the benchmark, and not by Nala PGx Core™
‡Refer to Table 1 for an outline of the benchmark methods used per variant.
§Discordant calls refer to instances in which Nala PGx Core™ did not produce a copy number call concordant with the benchmark.

487
488 **Table 8** Diplotype concordance for *CYP2C9*, *CYP2C19* and *CYP2D6* between Nala PGx Core™, and Agena
489 VeriDose® Core and *CYP2D6* CNV Panel

Genes	Concordance, %†	Discordant Diplotypes		
		Agena VeriDose® Core and *CYP2D6* CNV Panel	Nala PGx Core™	Instances
CYP2C9	100% (n = 213)	NA	NA	NA
CYP2C19	96.4% (n = 223)	*1/*1	*1/*3	1
		*1/*2	*1/*1	1
		*1/*2	*1/*3	1
		*1/*3	*2/*2	1
		*2/*2	*1/*1	2
		*2/*2	*1/*2	2
CYP2D6	94.7% (n = 209)	*1/*1	*2/*2	1
		*1/*10	*1/*10, CN >=3	1
		*1/*41	*39/*41	1
		*2/*2	*1/*1	1

		*2/*10	*1/*10	1
		*2/*36, CN >=3	*2/*10, CN >=3	1
		*4/*36	*4/*10	2
		*10/*36	*2/*36	1
		*13 *1/*10	*1/*10, CN >=3	1
		*13 *1/*41, CN = 1	*1/*41	1

[†]The concordance presented in this table excludes samples that have mismatches in diplotype calls arising from technological differences between platforms. Please refer to Supplementary Table 7 for a detailed breakdown of all diplotypes detected.

491 **Table 9** Observed allele frequencies by ethnicity

Gene	Allele or Variant	Allele Frequencies (Per This Study)						Allele Frequencies (PharmGKB)		
		Indonesian	Chinese	Malay	Indian	Caucasian	Overall	East Asian	Central/ South Asian	European
CYP2C9 (n = 206)	*2	0.000	0.000	0.000	0.060	0.172	0.032	0.002	0.114	0.127
	*3	0.000	0.040	0.039	0.100	0.069	0.044	0.038	0.110	0.076
CYP2C19 (n = 201)	*2	0.297	0.274	0.289	0.229	0.138	0.256	0.284	0.270	0.147
	*3	0.041	0.055	0.066	0.000	0.017	0.042	0.072	0.016	0.002
	*17	0.054	0.007	0.039	0.229	0.138	0.067	0.021	0.171	0.216
CYP2D6 (n = 195)	*2	0.088	0.118	0.167	0.140	0.196	0.136	0.121	0.295	0.277
	*4	0.000	0.035	0.069	0.020	0.161	0.051	0.005	0.091	0.185
	*5	0.029	0.028	0.028	0.080	0.000	0.031	0.049	0.046	0.030
	*6	0.000	0.000	0.000	0.000	0.018	0.003	0.000	0.000	0.011
	*9	0.000	0.000	0.000	0.000	0.018	0.003	0.002	0.003	0.028
	*10	0.338	0.299	0.222	0.160	0.143	0.251	0.436	0.087	0.016
	*14	0.000	0.014	0.000	0.000	0.000	0.005	0.003	ND	0.000
	*29	0.000	0.000	0.000	0.000	0.018	0.003	0.000	0.003	0.001
	*36	0.206	0.250	0.181	0.160	0.054	0.190	0.012	0.000	0.000
	*41	0.044	0.056	0.028	0.040	0.071	0.049	0.023	0.123	0.092
SLCO1B1 (n = 203)	rs4149056[†]	0.125	0.074	0.026	0.040	0.167	0.084	0.125[‡]	0.050[‡]	0.159[‡]

"ND" refers to instances in which no data is available for the given allele on PharmGKB.
[†]rs4149056 refers to the reduced function variant of *SLCO1B1* that is present in *SLCO1B1*5*, *SLCO1B1*15* and *SLCO1B1*17*.
[‡]Allele frequency values for rs4149056 have been obtained from gnomAD.

493 **Table 10** Observed diplotype frequencies by ethnicity

Gene	Diplotype	Phenotype[‡]	Diplotype Frequencies (Per This Study)[†]											
			Indonesian		Chinese		Malay		Indian		Caucasian		Overall	
			Obs	Freq	Obs	Freq	Obs	Freq	Obs	Freq	Obs	Freq	Obs	Freq
CYP2C9	*1/*1	NM	39	1.000	69	0.920	35	0.921	17	0.680	16	0.552	176	0.854
	*1/*2	IM	0	0.000	0	0.000	0	0.000	3	0.120	9	0.310	12	0.058
	*1/*3	IM	0	0.000	6	0.080	3	0.079	5	0.200	3	0.103	17	0.083
	*2/*3	PM	0	0.000	0	0.000	0	0.000	0	0.000	1	0.034	1	0.005
CYP2C19	*1/*1	NM	10	0.270	31	0.425	14	0.368	8	0.333	14	0.483	77	0.383
	*1/*2	IM	19	0.514	28	0.384	14	0.368	5	0.208	7	0.241	73	0.363
	*1/*3	IM	3	0.081	6	0.082	3	0.079	0	0.000	1	0.034	13	0.065
	*1/*17	RM	3	0.081	1	0.014	1	0.026	5	0.208	5	0.172	15	0.075
	*2/*2	PM	1	0.027	5	0.068	3	0.079	1	0.042	0	0.000	10	0.050
	*2/*3	PM	0	0.000	2	0.027	2	0.053	0	0.000	0	0.000	4	0.020
	*2/*17	IM	1	0.027	0	0.000	0	0.000	4	0.167	1	0.034	6	0.030
	*17/*17	UM	0	0.000	0	0.000	1	0.026	1	0.042	1	0.034	3	0.015
CYP2D6	*1/*1	NM	6	0.176	7	0.097	5	0.139	4	0.160	2	0.071	24	0.123
	*1/*2	NM	0	0.000	2	0.028	2	0.056	3	0.120	3	0.107	10	0.051
	*1/*2, CN >=3	UM	0	0.000	0	0.000	0	0.000	1	0.040	1	0.036	2	0.010
	*1/*4	IM	0	0.000	2	0.028	1	0.028	0	0.000	4	0.143	7	0.036
	*1/*5	IM	0	0.000	2	0.028	1	0.028	1	0.040	0	0.000	4	0.021

Genotype	Metabolizer	Obs	Freq	Obs	Freq	Obs	Freq	Obs	Freq	Obs	Freq	Obs	Freq
*1/*6	IM	0	0.000	0	0.000	0	0.000	0	0.000	1	0.036	1	0.005
*1/*10	NM	4	0.118	3	0.042	2	0.056	2	0.080	2	0.071	13	0.067
*1/*14	NM	0	0.000	1	0.014	0	0.000	0	0.000	0	0.000	1	0.005
*1/*29	NM	0	0.000	0	0.000	0	0.000	0	0.000	1	0.036	1	0.005
*1/*36	IM	3	0.088	3	0.042	5	0.139	4	0.160	0	0.000	15	0.077
*1/*41	NM	1	0.029	2	0.028	1	0.028	0	0.000	2	0.071	6	0.031
*1/*41, CN >=3	UM	0	0.000	0	0.000	0	0.000	1	0.040	0	0.000	1	0.005
*2/*2	NM	0	0.000	1	0.014	1	0.028	1	0.040	1	0.036	4	0.021
*2/*4	IM	0	0.000	0	0.000	1	0.028	0	0.000	2	0.071	3	0.015
*2/*5	IM	0	0.000	0	0.000	1	0.028	1	0.040	0	0.000	2	0.010
*2/*10	NM	4	0.118	4	0.056	4	0.111	0	0.000	1	0.036	13	0.067
*2/*14	NM	0	0.000	1	0.014	0	0.000	0	0.000	0	0.000	1	0.005
*2/*36	IM	2	0.059	4	0.056	2	0.056	0	0.000	1	0.036	9	0.046
*2/*41	NM	0	0.000	4	0.056	0	0.000	0	0.000	1	0.036	5	0.026
*4/*4	PM	0	0.000	1	0.014	1	0.028	0	0.000	1	0.036	3	0.015
*4/*5	PM	0	0.000	0	0.000	0	0.000	1	0.040	0	0.000	1	0.005
*4/*9	IM	0	0.000	0	0.000	0	0.000	0	0.000	1	0.036	1	0.005
*4/*10	IM	0	0.000	1	0.014	1	0.028	0	0.000	0	0.000	2	0.010
*5/*10	IM	0	0.000	2	0.028	0	0.000	1	0.040	0	0.000	3	0.015
*5/*41	IM	2	0.059	0	0.000	0	0.000	0	0.000	0	0.000	2	0.010
*10/*10	IM	3	0.088	1	0.014	2	0.056	0	0.000	2	0.071	8	0.041
*10/*10, CN >=3	IM	0	0.000	1	0.014	0	0.000	0	0.000	0	0.000	1	0.005
*10/*36	IM	9	0.265	28	0.389	5	0.139	4	0.160	1	0.036	47	0.241
*10/*41	IM	0	0.000	1	0.014	0	0.000	1	0.040	0	0.000	2	0.010
*36/*41	IM	0	0.000	1	0.014	1	0.028	0	0.000	1	0.036	3	0.015

[†]"Obs" and "Freq" are abbreviations for "Observations" and "Frequency" respectively.

[‡]"NM", "IM", "PM" and "UM" are abbreviations for "Normal Metabolizer", "Intermediate Metabolizer", "Poor Metabolizer" and "Ultrarapid Metabolizer" respectively.

Figure Legends

Figure 1. *CYP2D6*36* Frequency By Ethnicity. Distribution of individuals carrying one or two copies of the *CYP2D6*36* allele among the study cohort (n=205), grouped per ethnicity.

Supplementary Figure 1. Observed *CYP2D6* Functional Copy Number Frequencies By Ethnicity. Distribution of individuals carrying one, two or three or more copies of the *CYP2D6* functional gene among the study cohort (n=205), grouped per ethnicity.