SARS-CoV-2 mRNA vaccines induce a greater array of spike-specific antibody isotypes with more potent complement binding capacity than natural infection

Running title: Antibodies induced by COVID-19 vaccines

Jéromine Klingler[1,2], Gregory S. Lambert[1], Vincenza Itri[1], Sean Liu[1], Juan C. Bandres[1,2], Gospel Enyindah-Asonye[1], Xiaomei Liu[1,2], Kasopefoluwa Y. Oguntuyo[3], Fatima Amanat[3,4], Benhur Lee[3], Susan Zolla-Pazner[1,3], Chitra Upadhyay[1], Catarina E. Hioe[1,2,3*]

[1]Division of Infectious Diseases, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY USA.

[2]James J. Peters VA Medical Center, Bronx, NY, USA.

[3]Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[4]Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

*Corresponding author

Contact: catarina.hioe@mssm.edu, catarina.hioe@va.gov

Footnotes:

The authors do not have a commercial or other association that might post a conflict of interest.

This work was supported in part by the Department of Medicine of the Icahn School of Medicine at Mount Sinai Department of Medicine (to S.Z-P., C.E.H.); the Department of Microbiology and the Ward-Coleman estate for endowing the Ward-Coleman Chairs at the Icahn School of Medicine at Mount Sinai (to B.L.), the Department of Veterans Affairs [Merit Review Grant I01BX003860] (to C.E.H.) and [Research Career Scientist Award 1IK6BX004607] (to C.E.H.); the National Institutes of Health [grant AI139290] to C.E.H., [grants R01 AI123449, R21 AI1498033] to B.L, [grant R01 AI140909] to C.U. G.E-A and K.Y.O. were supported by Viral-Host Pathogenesis Training Grant T32 AI07647. K.Y.O was also supported by Training Grant F31 AI154739.

This manuscript has been deposited to medRxiv.

Corresponding author contact: catarina.hioe@mssm.edu, catarina.hioe@va.gov

Abstract (199 words)

Background: Mass vaccination against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is on-going in the United States with vaccines approved for emergency use by the FDA. Each vaccine can induce spike-specific antibodies (Abs) with virus-neutralizing activities; however, the Fc-mediated Ab activities have received little attention. Moreover, while plasma/serum Abs are commonly studied, scant information is available about Abs in the respiratory mucosa, the site of SARS-CoV-2 transmission.

Methods: Plasma and saliva were collected from mRNA vaccine recipients and convalescent coronavirus disease 2019 (COVID-19) patients. Antigen-specific total Ig and Ig isotypes were measured. In addition to virus neutralization, Fc-mediated activities were investigated, including antibody-dependent cellular phagocytosis (ADCP) and complement deposition.

Results: Similar to infection, vaccination stimulated spike-specific Ab responses detected in plasma and saliva, with IgG1 as the dominant isotype. Interestingly, vaccination produced greater IgG2, IgG3, and IgG4 responses and higher ratios of (IgG1+IgG3)/(IgG2+IgG4) than infection. Moreover, while plasma neutralization and ADCP potencies were comparable in vaccinated and convalescent individuals, vaccine-induced plasma Abs elicited stronger complement binding and activation.

Conclusion: Compared with natural infection, mRNA vaccines induced a greater array of IgG subtypes against spike in saliva and plasma. The vaccine-induced Abs were also more potent in mediating complement activation.

Keywords

SARS-CoV-2, COVID-19, vaccination, antibody isotypes, neutralization, ADCP, complement fixation, saliva

Text (3498 words)

Background

In the USA, three coronavirus disease-2019 (COVID-19) vaccines have received an authorization

for emergency use from the FDA: two messenger RNA (mRNA) vaccines from Pfizer-BioNTech

(BNT162b2) and Moderna (mRNA-1273), and one adenovirus-vectored vaccine from Johnson &

Johnson/Janssen (Ad26.CoV2.S). The two mRNA-based vaccines reported >90% efficacy in preventing

symptomatic SARS-CoV-2 infection [1,2], while the efficacy of Ad26.CoV2.S was ~70% [3]. All three

vaccines were designed to induce antibodies (Abs) targeting SARS-CoV-2 spike [4,5], a membrane-

anchored protein on the viral surface which contains the receptor-binding domain (RBD) necessary for

binding and entry into the host cells [6–8].

While the primary antiviral function of Abs is to neutralize the virions, Abs also mediate non-

neutralizing effector functions via their Fc fragments. In the case of SARS-CoV-2, the binding of anti-

spike Abs to virions, infected cells, or soluble spike proteins creates immune complexes capable of

engaging Fc receptors (FcRs) or complement via the Abs' Fc fragments [9,10]. The FcR engagement

triggers a cascade of intracellular signals critical for Fc-mediated activities, such as Ab-dependent cellular

phagocytosis (ADCP) and Ab-dependent cellular cytotoxicity (ADCC). The binding of C1q, the first

component in the classical complement pathway, to Fc fragments on immune complexes activates the

downstream complement cascade, resulting in the deposition of C3 and C4 degradation products and

leading to the generation of C5 convertase and the assembly of the membrane-attack complex responsible

for initiating complement-mediated lysis. Complement-opsonized immune complexes also interact with

complement receptors on leukocytes to trigger effector functions, such as complement-dependent cell-

mediated phagocytosis and cytotoxicity [11,12].

The Ab affinity for FcR and complement is determined by their immunoglobulin (Ig) isotypes.

Each of the Ig isotypes interacts with distinct FcRs, categorized into type I and type II [9,13]. The

canonical type I FcRs for IgG include FcγRI (CD64), FcγRIIA (CD32A), FcγRIIB (CD32B), FcγRIIIA

(CD16A), and FcγRIIIB (CD16B), which differ in cell distribution, affinity for IgG subtypes, and

functional effects. Among IgG subtypes, IgG1 and IgG3 have the strongest affinity for the activating

FcγRI, FcγRIIA, and FcγRIIIA [14]. On macrophages, the engagement of FcγRI and FcγRIIA stimulates

ADCP. On NK cells and some macrophages, FcγRIIIA serves as the key receptor for inducing ADCC

[15,16]. Each Ab isotype also activates the complement system with varying potency in the following

order: IgM=IgG3>IgG1>IgG2=IgA, while IgG4, IgD, and IgE do not engage complement [17].

The COVID-19 vaccines used for mass immunization demonstrate the capacity to elicit

neutralizing Ab responses, but the full properties of the elicited Abs are not yet known [2,18–21]. In

particular, Fc-mediated activities and the associated Ig isotypes elicited after vaccination have not been

studied. Moreover, while SARS-CoV-2-specific Ab profiles from convalescent individuals have been

shown to be similar in plasma and saliva [22], the Ab profiles in the respiratory mucosa, the initial site of

SARS-CoV-2 infection, from vaccinated individuals remains to be investigated. In this study, using a

multiplex bead assay developed in our previous study [23], we analyzed the spike- and RBD-specific Ig

isotypes in paired plasma/saliva specimens from seven mRNA vaccine recipients in comparison with eight

COVID-19-positive convalescent donors and four COVID-19-negative individuals. Additionally, the

vaccine-induced plasma Abs were examined for Fab-mediated virus neutralization and Fc-mediated

activities (ADCP, C1q binding and C3d deposition). This study provides the first evidence that SARS-

CoV-2 immunization induces a greater breadth of spike-specific Ig isotypes and more potent complement

binding capacities than natural infection.

Methods

Recombinant proteins. SARS-CoV-2 spike and RBD proteins were produced as described [24,25].

Human specimens. Plasma and saliva specimens were obtained from volunteers enrolled in IRB-approved protocols at the Icahn School of Medicine at Mount Sinai (IRB#17-00060, IRB#19-01243) and the James J. Peter Veterans Affairs Medical Center (IRB#BAN-1604): RN#1, RN#4 and RV#1-5 after immunization; RP#2-5, 7, 12-14 after infection; contemporaneous non-vaccinated COVID-19-negative and pre-pandemic plasma samples. The clinical data are summarized in the Supplementary Tables 1 (vaccinated individuals) and 2 (convalescent individuals). All participants provided written consent. All samples were heat-inactivated before use.

Multiplex bead Ab binding assay. Measurement of total Ig and Ig isotypes to SARS-CoV-2 antigen-coupled beads was performed as described [23]. Complement deposition onto plasma and saliva Abs reactive with spike and RBD were measured according to [26] with modifications. For the C1q assay, beads with spike-Ab or RBD-Ab complexes were incubated with C1q Component from Human Serum (Sigma, #C1740) for 1 hour at room temperature, followed by an anti-C1q-PE antibody (Santa Cruz, #sc-53544 PE). For the C3d assay, Complement Sera Human (33.3%, Sigma, #S1764) was added to the beads for 1 hour at 37°C, followed by a biotinylated monoclonal anti-C3d antibody (Quidel, #A702).

Virus neutralization. VSV-based SARS-CoV-2 pseudoviruses were used in neutralization assays as described [23,27].

ADCP. ADCP assays were performed using a reported protocol [28] with some modifications. FluoSpheres carboxylate-modified microspheres (Thermo Fisher, #F8823) were coupled with SARS-CoV-2 spike protein using the xMAP Antibody Coupling Kit (5 µg protein/~36.4×10^9 beads, Luminex #40-50016). Spike-conjugated microspheres were incubated with diluted plasma for 2 hours at 37°C in the dark. After washing and centrifugation (2,000 g, 10 minutes), the beads (~3×10^8 beads, 10 µL/well) were

130   incubated with THP-1 cells (0.25×10^5 cells, 200 µL/well) for 16 hours. The samples were analyzed on an

131   Attune NxT flow cytometer (Thermo Fisher, #A24858). Data analysis was performed using FCS Express

132   7 Research Edition (De Novo Software).

133   **Statistical analysis.** Statistical analyses were performed as designated in the figure legends using

134   GraphPad Prism 8 (GraphPad Software, San Diego, CA).

Results

COVID-19 mRNA vaccines induce Ab responses to SARS-CoV-2 spike, RBD, S1 and S2

antigens. Paired plasma and saliva specimens from seven healthy recipients of COVID-19 mRNA

vaccines were examined for total Ig and different Ig isotypes against the SARS-CoV-2 antigens. Two

individuals received the Pfizer vaccine, five received the Moderna vaccine, and samples were collected

15-37 days after the second vaccine dose (**Supplemental Table 1**). Four COVID-19-negative non-

vaccinated contemporaneous donors were tested for comparison and used to establish the cut-off values.

The plasma specimens were titrated for total Ig against spike, RBD, S1, S2, and nucleoprotein

antigens, with bovine serum albumin (BSA) as a control antigen (**Fig. 1a**). All vaccinated subjects

exhibited titrating amounts of plasma Ig against spike, RBD, S1 and S2, above the cut-off levels. As

expected, no reactivity was observed against the nucleoprotein, although one individual had weak

reactivity at the lowest dilution, reaching slightly above the cut-off.

Comparison with plasma samples from eight convalescent COVID-19-positive specimens

(collected 40-256 days post-symptom onset, **Supplemental Table 2**) showed differential Ig responses

against nucleoprotein. The area under the curve (AUCs) values of vaccinated and convalescent specimens

were not significantly different, although the median AUCs of the vaccinated group were 2.6-fold (for

spike) and 2.8-fold (for RBD) higher than those of the convalescent group (**Fig. 1b**).

COVID-19 mRNA vaccines induce varying levels of Ig isotypes against SARS-CoV-2 spike.

The plasma specimens were then tested for antigen-specific total Ig, IgM, IgG1-4, IgA1 and IgA2 at a

1:200 dilution (**Fig. 2a**). The percentage of responders for each isotype was determined using the cut-off

values calculated as mean+3 standard deviation (SD) of 12 pre-pandemic samples reported previously

[23]. All vaccinated plasma specimens had high levels of anti-spike and anti-RBD total Ig and IgG1. A

significant proportion (40-86%) of specimens also displayed IgG2-4 Abs against spike and RBD, although

the levels were low. Notably, 100% and 71% of individuals produced IgA1 Abs against spike and RBD,

respectively, while 14% and 57% exhibited spike- and RBD-specific IgA2. In addition, 71% generated

spike-specific IgM at varying levels. Only 14% had RBD-specific IgM mainly due to higher background,

consistent with our previous findings [23]. None of the four control specimens showed Ig isotype

reactivity with spike or RBD.

Linear regression analysis was performed to assess the relative contribution of each isotype to the

total Ig responses to spike and RBD (**Fig. 2b**). IgG1 had the highest and most significant r^2 values,

indicating that IgG1 is the major isotype induced by COVID-19 mRNA vaccines. In addition, IgG3 Abs

demonstrated a significant contribution, with an r^2 of 0.70 for spike and 0.64 for RBD with p values <0.05

for both.

Comparison of spike- and RBD-specific Ig isotypes generated by vaccination versus

infection. Linear regression data of plasma specimens from the seven vaccinated and eight convalescent

subjects tested in this study (**Supplemental Fig. 1**) were comparable to those from a previously reported

cohort of 29 convalescent subjects [23], demonstrating that IgG1 was the dominant isotype induced after

vaccination or infection. However, the association with IgG3 Abs was stronger in plasma from vaccinated

versus convalescent donors, where a significant association was observed with RBD-specific IgG3 only

(**Supplemental Fig. 1**, [23]).

Further comparison of Ig isotypes elicited by vaccination versus infection demonstrated that the

percentage of IgM responders in the vaccinated group was lower than those observed in our past study of

29 convalescent individuals [23] and the eight convalescent donors examined in parallel in this study

(88% for spike and 75% for RBD, **Supplemental Fig. 1**). In contrast, the percentages of IgG2, IgG3, and

IgG4 responders were higher (**Supplemental Fig. 1**), indicating that vaccination induces a broader range

of IgG subtypes. A heatmap is shown in **Fig. 3a** to illustrate these differences.

Interestingly, the ratios of IgG1+IgG3 mean fluorescence intensity (MFIs) to IgG2+IgG4 MFIs

were higher in the plasma of the seven vaccinated individuals than in those of the eight convalescent

individuals (p<0.05) (**Fig. 2c**), further distinguishing the two groups. A similar pattern was seen when

comparing these seven vaccinated specimens with the previously studied cohort of 29 convalescent

plasma specimens from <8 weeks post-symptom onset: the median ratios of spike-specific

(IgG1+IgG3)/(IgG2+IgG4) were 133 and 46 for the vaccinated and convalescent groups, respectively

(p=0.0019). For RBD-specific Abs, the median ratio of the vaccinated group was also two times higher

(p=0.06). These data demonstrate that, while vaccine-induced Ab responses are comprised mainly of

spike- and RBD-specific IgM, IgG1, and IgA1, vaccination elicited higher proportions of IgG1 and IgG3

over IgG2 and IgG4 than natural infection.

Vaccine-induced IgG1 and IgG3 against spike and RBD contribute the most to virus

neutralization activities. To test the neutralizing activity of vaccine-induced plasma Abs, specimens from

seven vaccinated individuals and two non-vaccinated COVID-19-negative individuals were examined in a

neutralization assay using a VSVΔG pseudovirus bearing the SARS-CoV-2 spike protein. All vaccinated

samples attained 90% neutralization, while none of the COVID-19-negative samples did (**Fig. 4a**). The

IC_{50} titers attained after vaccination were similar to those of convalescent samples collected <8 weeks

after symptom onset [23] or at later time points (>186 days) (**Fig. 4b, Fig. 3a**).

Linear regression analyses were performed to analyze the contribution of each isotype to the

neutralization activities (**Fig. 4c**). Similar to our past data with convalescent plasma [23], high r^2 values

(0.84-0.85) and significant p values were observed with total Ig and IgG1. Spike- and RBD-specific IgG3

Abs also demonstrated a significant association with neutralization, with r^2 values of 0.71 and 0.70,

respectively. Interestingly, IgM did not show any association. This result is contrary to our observation in

convalescent individuals [23] but is in line with lower IgM responses detected in these vaccinees after the

second vaccine dose. The association between IgM and neutralization was also not found in convalescent

samples collected at late time points (**Supplemental Fig. 2**), as the IgM response was lower in these

samples compared to specimens collected <8 weeks after symptom onset [23].

COVID-19 mRNA vaccines induce ADCP activities. Because Ig isotypes are key determinants

of Fc functions, we examined the Fc-mediated Ab activities in plasma specimens from the seven

vaccinated donors in comparison with convalescent plasma and negative controls. We evaluated 1) spike-

specific ADCP using THP-1 phagocytes and spike-coated fluorescent beads and 2) complement activation

based on C1q and C3d deposition on spike-Ab and RBD-Ab complexes.

All specimens from vaccinated subjects displayed ADCP activities above the negative control

(**Fig. 5a**), similar to those seen in convalescent specimens (**Supplemental Fig. 3, Fig. 3a**). The ADCP

levels were higher for samples of vaccinated donors versus convalescent donors (**Fig. 5b, Fig. 3a**),

corresponding to an ~2.6-fold difference in spike-binding total Ig median levels (**Fig. 1a**). To compare the

ADCP capacity of plasma Abs from vaccinated versus convalescent donors independent of the spike-

binding Ig levels, we calculated the AUC ratios of spike-specific ADCP over the spike-binding total Ig.

The ADCP capacity of Abs induced by vaccination versus infection was equivalent (**Fig. 5c**),

demonstrating that mRNA vaccines and SARS-CoV-2 infection induce plasma Abs with comparable

ADCP potency.

To assess the contribution of each isotype to ADCP in the plasma of vaccinated donors, regression

analyses showed the highest r^2 value and the most significant p value for spike-specific IgG1 Abs (**Fig.**

5d). Interestingly, spike-specific IgG2 and IgG3 Abs also demonstrated significant associations with

ADCP, with r^2 and p values comparable to those of total Ig but lower than those of IgG1. The ADCP

association with IgG1 and IgG2 Abs, but not IgG3 Abs, was also observed with convalescent plasma

(**Supplemental Fig. 3**).

COVID-19 mRNA vaccines induce Abs with greater C1q and C3d deposition activities. When

we examined C1q and C3d deposition, a high amount of variability was observed among the seven

vaccinated donors. One plasma sample did not show any C1q or C3d binding to spike- or RBD-binding

Abs (**Fig. 6a**), and the remaining samples exhibited a range of C1q and C3d binding levels. Regression

analyses demonstrated that, among the spike- and RBD-binding Ig isotypes tested, IgG1 and IgG3 Abs

contributed the most to the complement binding activities of the vaccinated plasma specimens (**Fig. 6b**).

This is consistent with the greater potency of IgG1 and IgG3 Abs, as compared with IgG2 and IgG4 Abs,

to activate the classical complement cascade [23]. C1q and C3d deposition was also detected with

convalescent plasma samples, albeit at lower levels (**Fig. 3a**, **Supplemental Fig. 4**). The binding levels of

C1q and C3d correlated with each other for spike- and RBD-specific Abs in vaccinated and convalescent

donors (**Fig. 6c**).

To compare the complement binding capacity of vaccine- versus infection-induced Abs, the ratios

of C1q or C3d binding AUCs over total Ig AUCs were calculated. C1q and C3d binding capacity was

higher for vaccine-induced Abs, although only the difference in C1q binding reached statistical

significance (**Fig. 6d**). Hence, the spike- and RBD-specific Abs induced by vaccination exhibited a greater

complement activation potency than the infection-induced counterparts.

We summarize the isotypes and functions of plasma Abs induced after vaccination versus natural

infection in **Fig. 3a**. The heatmap clearly shows a greater breadth of Ig isotypes against spike and RBD in

the plasma of vaccinated versus convalescent subjects. This was accompanied by more potent Fc-

dependent activities especially complement activation, even though neutralization potency was

indistinguishable.

COVID-19 mRNA vaccines induce spike-specific Ab responses in saliva. Since SARS-CoV-2

is a respiratory virus, we asked whether the COVID-19 mRNA vaccines also elicited Ab responses against

the spike antigen in the upper respiratory mucosa where the Abs would serve as the frontline defense

against this virus. To this end, we examined the levels of antigen-specific total Ig and Ig isotypes in saliva,

as a proxy for oral and upper respiratory mucosal secretion, following vaccination versus natural infection.

Saliva specimens were obtained from the same seven vaccinated and eight convalescent donors at the time

of blood collection and compared to four non-vaccinated COVID-19 negative saliva specimens.

262   The saliva specimens were first titrated for total Ig levels against spike, RBD, S1, S2, and

nucleoprotein antigens with BSA as a control (**Fig. 7a, Supplemental Fig. 5a**). All vaccinated subjects

exhibited titrating amounts of saliva Ig against spike and S1, similar to those detected in plasma (**Fig. 3b**),

although the saliva median effective concentration (EC_{50}) titers of spike-specific Ig were on average 229-

fold lower than the plasma titers. However, in contrast to 100% reactivity detected in plasma, only 5/7

saliva specimens from the vaccinated donors exhibited Ig reactivity against RBD and 2/7 against S2. No

reactivity was detected against nucleoprotein or BSA, similar to what was seen with plasma specimens. In

contrast, we detected Abs against spike, RBD, and nucleoprotein in saliva samples from convalescent

COVID-19 patients (**Fig. 7a**). The presence of nucleoprotein-specific Abs in the saliva samples again

differentiated convalescent from vaccinated subjects, similar to that seen in plasma. The AUCs of spike-

and RBD-specific total Ig in the saliva of vaccinated and convalescent subjects did not differ significantly,

although the median AUC of the vaccinated group was ~3-fold higher than that of the convalescent group.

Overall, a strong correlation was observed between spike- and RBD-specific total Ig levels in saliva

versus plasma (**Fig. 7b**).

276   The isotypes of spike- and RBD-specific Abs were subsequently examined in saliva at a 1:4

dilution. The COVID-19 mRNA vaccines induced anti-spike and anti-RBD IgG1 in the saliva of all

vaccine recipients (**Fig. 6c**). A significant proportion of saliva samples also displayed IgG2-4 Abs against

spike and RBD, even though the levels were lower than those of IgG1. Of the IgA isotypes in saliva, IgA1

responses against spike and RBD were more prevalent than their IgA2 counterparts. IgM responses were

weak; only 2/7 saliva samples had detectable spike-specific IgM and none had RBD-specific IgM. The

four control saliva specimens tested in parallel did not show any Ig isotype reactivity with spike or RBD.

283   Comparison of the plasma and saliva Ab profiles from the vaccinated versus convalescent groups

is summarized in **Fig. 3a,b**. The plasma and saliva Ig isotype profiles were comparable to each other but

differed between the vaccinated and convalescent group. The dominant isotype in saliva and plasma of

vaccinated and convalescent groups was IgG1. However, the saliva IgG2-4 levels were higher in the

vaccinated group, whereas the saliva IgM response was less prevalent in the vaccinated versus

convalescent groups as seen in plasma (**Supplemental Fig. 5a,b**, [23]). Moreover, the saliva

(IgG1+IgG3)/(IgG2+IgG4) ratios were higher in the vaccinated versus convalescent groups (p<0.01) (**Fig.**

7d), similar to that seen in plasma (**Fig. 2c**).

Discussion

294    This study provides the first evidence that COVID-19 mRNA vaccines induced plasma and saliva

295 Ab responses dominated by IgG1. As compared to natural infection, vaccination also induced Ab

296 responses marked by higher levels of IgG2-4 and greater ratios of (IgG1+IgG3)/(IgG2+IgG4). Vaccine-

297 induced plasma Abs also displayed stronger capacity to mediate complement binding and activation. The

298 differential potencies were apparent when the Ab levels were taken into account and when comparison

299 was made with infection-induced Abs from earlier or later time points. By contrast, neutralization and

300 ADCP potencies of vaccine- and infection-induced plasma Abs were comparable. The mRNA vaccine-

301 induced Ab responses were also reported in a recent work with >8000 finger stick blood specimens to

302 have higher seroconversion rates and greater cross-reactivity with SARS-CoV-1 and Middle Eastern

303 respiratory syndrome (MERS)-CoV RBDs [29], further indicating the superior quality of vaccine-induced

304 Ab responses.

305    We showed here that complement binding to spike-Ab immune complexes was stronger in plasma

306 from vaccinated versus convalescent groups. However, the individual levels were highly variable for both

307 groups, and no correlation was observed between the complement fixation capacity and the

308 (IgG1+IgG3)/(IgG2+IgG4) ratios. A parameter known to influence complement binding is Fc

309 glycosylation, as the removal of terminal galactose from IgG Fc glycans has been shown to decrease C1q

310 binding and downstream classical complement activation without affecting FcγR-mediated functions

311 [30,31]. Similarly, the sialylation of IgG Fc domains has been demonstrated to impair complement-

312 dependent cytotoxicity [32]. The Fc glycan compositions of vaccine- and infection-induced Abs are yet

313 unknown. The importance of complement binding/activation for protection against SARS-CoV-2 also

314 requires further investigation. It should be noted, however, that human neutralizing monoclonal antibodies

315 against SARS-CoV-2 require a functional Fc region capable of binding complement and engaging FcγR

316 for ADCP and ADCC, for optimal protection as therapy [33].

We noted that while all vaccinated and convalescent individuals displayed plasma neutralization

and ADCP activities, greater individual variability was seen in the levels of neutralization versus ADCP in

both vaccinated and convalescent groups. Indeed, no correlation was observed between neutralization and

ADCP activities, suggesting that these two functions are mediated by distinct Ig populations that may

target different epitopes. We did not examine binding and functional Ab activities against the SARS-CoV-

2 variants of concern such as B.1.1.7, B.1.351 and P.1. Most COVID-19 vaccines elicit Ab responses with

weaker neutralization activities against these variants [34,35], but the effects of these and other emerging

mutations on the non-neutralizing Fc-dependent functions are yet to be determined.

This study demonstrated that all vaccinated and convalescent individuals evaluated here had

detectable saliva and plasma spike-specific Abs. Plasma and saliva anti-spike Ig levels after natural

infection or vaccination correlated strongly. In a previous report, Abs against spike and nucleoprotein

were detected similarly in plasma and saliva following SARS-CoV-2 infection [22]. These data indicate

the potential use of saliva for monitoring of anti-spike Ab responses in vaccinated and convalescent

individuals. However, lower positive responses were detected in saliva against RBD, S1, and S2 as

compared to the intact spike antigen.

In conclusion, we demonstrate that, compared to natural infection, COVID-19 mRNA vaccines

induce spike-specific Abs of diverse Ig isotypes and with potent complement binding capacities. The data

highlight the superiority of vaccination in eliciting functional Ab responses against SARS-CoV-2.

References

1.   Baden LR, El Sahly HM, Essink B, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2

Vaccine. N Engl J Med. Massachusetts Medical Society; **2021**; 384(5):403–416.

2.   Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19

Vaccine. N Engl J Med. **2020**; 383(27):2603–2615.

3.   Sadoff J, Gray G, Vandebosch A, et al. Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine

against Covid-19. N Engl J Med [Internet]. Massachusetts Medical Society; **2021** [cited 2021 Apr 30]; .

Available from: https://www.nejm.org/doi/10.1056/NEJMoa2101544

4.   Amanat F, Krammer F. SARS-CoV-2 Vaccines: Status Report. Immunity. **2020**; .

5.   Le TT, Andreadakis Z, Kumar A, et al. The COVID-19 vaccine development landscape. Nat Rev

Drug Discov. Nature Publishing Group; **2020**; 19(5):305–306.

6.   Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of SARS-CoV-

2 by full-length human ACE2. Science. **2020**; 367(6485):1444–1448.

7.   Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2

and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell. **2020**; 181(2):271-280.e8.

8.   Walls AC, Park Y-J, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, Function, and

Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell. **2020**; 181(2):281-292.e6.

9.   Bournazos S, Ravetch JV. Diversification of IgG effector functions. Int Immunol. **2017**;

29(7):303–310.

10.   Goldberg BS, Ackerman ME. Antibody-mediated complement activation in pathology and

protection. Immunol Cell Biol. **2020**; 98(4):305–317.

11.   Kellner C, Otte A, Cappuzzello E, Klausz K, Peipp M. Modulating Cytotoxic Effector Functions

by Fc Engineering to Improve Cancer Therapy. Transfus Med Hemotherapy Off Organ Dtsch Ges

Transfusionsmedizin Immunhamatologie. **2017**; 44(5):327–336.

12.   Lee C-H, Romain G, Yan W, et al. IgG Fc domains that bind C1q but not effector Fcγ receptors

delineate the importance of complement-mediated effector functions. Nat Immunol. **2017**; 18(8):889–898.

13.   Pincetic A, Bournazos S, DiLillo DJ, et al. Type I and type II Fc receptors regulate innate and

adaptive immunity. Nat Immunol. **2014**; 15(8):707–716.

14.   Gillis C, Gouel-Chéron A, Jönsson F, Bruhns P. Contribution of Human FcγRs to Disease with

Evidence from Human Polymorphisms and Transgenic Animal Studies. Front Immunol. **2014**; 5:254.

15.   Temming AR, Taeye SW de, Graaf EL de, et al. Functional Attributes of Antibodies, Effector

Cells, and Target Cells Affecting NK Cell-Mediated Antibody-Dependent Cellular Cytotoxicity. J

Immunol Baltim Md 1950. **2019**; 203(12):3126–3135.

16.   Lewis GK, Ackerman ME, Scarlatti G, et al. Knowns and Unknowns of Assaying Antibody-

Dependent Cell-Mediated Cytotoxicity Against HIV-1. Front Immunol. **2019**; 10:1025.

17.   Noris M, Remuzzi G. Overview of complement activation and regulation. Semin Nephrol. **2013**;

33(6):479–492.

18.   Rogliani P, Chetta A, Cazzola M, Calzetta L. SARS-CoV-2 Neutralizing Antibodies: A Network

Meta-Analysis across Vaccines. Vaccines [Internet]. **2021** [cited 2021 Apr 13]; 9(3). Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7999682/

19.   Walsh EE, Frenck RW, Falsey AR, et al. Safety and Immunogenicity of Two RNA-Based Covid-

19 Vaccine Candidates. N Engl J Med. **2020**; 383(25):2439–2450.

20.   Sahin U, Muik A, Derhovanessian E, et al. COVID-19 vaccine BNT162b1 elicits human antibody

and TH1 T cell responses. Nature. **2020**; 586(7830):594–599.

21.   Jackson LA, Anderson EJ, Rouphael NG, et al. An mRNA Vaccine against SARS-CoV-2 —

Preliminary Report. N Engl J Med. Massachusetts Medical Society; **2020**; 0(0):null.

22.   Pisanic N, Randad PR, Kruczynski K, et al. COVID-19 Serology at Population Scale: SARS-CoV-

2-Specific Antibody Responses in Saliva. J Clin Microbiol [Internet]. **2020** [cited 2021 Apr 16]; 59(1).

Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7771435/

23.   Klingler J, Weiss S, Itri V, et al. Role of IgM and IgA Antibodies in the Neutralization of SARS-

CoV-2. J Infect Dis. **2020**; .

24.   Amanat F, Stadlbauer D, Strohmeier S, et al. A serological assay to detect SARS-CoV-2

seroconversion in humans. Nat Med. Nature Publishing Group; **2020**; :1–4.

25.   Stadlbauer D, Amanat F, Chromikova V, et al. SARS-CoV-2 Seroconversion in Humans: A

Detailed Protocol for a Serological Assay, Antigen Production, and Test Setup. Curr Protoc Microbiol.

2020; 57(1):e100.

26.   Perez LG, Martinez DR, deCamp AC, et al. V1V2-specific complement activating serum IgG as a

correlate of reduced HIV-1 infection risk in RV144. PloS One. **2017**; 12(7):e0180720.

27.   Oguntuyo KY, Stevens CS, Hung C-T, et al. Quantifying absolute neutralization titers against

SARS-CoV-2 by a standardized virus neutralization assay allows for cross-cohort comparisons of

COVID-19 sera. medRxiv. Cold Spring Harbor Laboratory Press; **2020**; :2020.08.13.20157222.

28.   Balasubramanian P, Williams C, Shapiro MB, et al. Functional Antibody Response Against V1V2

and V3 of HIV gp120 in the VAX003 and VAX004 Vaccine Trials. Sci Rep. **2018**; 8(1):542.

29.   Assis R, Jain A, Nakajima R, et al. Substantial Differences in SARS-CoV-2 Antibody Responses

Elicited by Natural Infection and mRNA Vaccination. bioRxiv. Cold Spring Harbor Laboratory; **2021**;

:2021.04.15.440089.

30.   Hodoniczky J, Zheng YZ, James DC. Control of recombinant monoclonal antibody effector

functions by Fc N-glycan remodeling in vitro. Biotechnol Prog. **2005**; 21(6):1644–1652.

31.   Boyd PN, Lines AC, Patel AK. The effect of the removal of sialic acid, galactose and total

carbohydrate on the functional activity of Campath-1H. Mol Immunol. **1995**; 32(17–18):1311–1318.

32.   Quast I, Keller CW, Maurer MA, et al. Sialylation of IgG Fc domain impairs complement-

dependent cytotoxicity. J Clin Invest. 125(11):4160–4170.

33.   Winkler ES, Gilchuk P, Yu J, et al. Human neutralizing antibodies against SARS-CoV-2 require

intact Fc effector functions for optimal therapeutic protection. Cell. **2021**; 184(7):1804-1820.e16.

34.   Bian L, Gao F, Zhang J, et al. Effects of SARS-CoV-2 variants on vaccine efficacy and response

strategies. Expert Rev Vaccines. :1–9.

35.   Bates TA, Leier HC, Lyski ZL, et al. Neutralization of SARS-CoV-2 variants by convalescent and

vaccinated serum. medRxiv. Cold Spring Harbor Laboratory Press; **2021**; :2021.04.04.21254881.

Acknowledgments

We thank Dr. Florian Krammer for providing spike and RBD antigens, and all the donors for their contribution to the research. We would like to thank the expertise and assistance of Dr. Christopher Bare and the Dean's Flow Cytometry CORE at Mount Sinai.

Author contributions

J.K. and C.E.H. wrote the manuscript. G.L., S.Z-P edited the manuscript. J.K., G.L., S.Z-P, C.U., and C.E.H. designed the experiments. J.K., G.L., V.I., and X.L. performed the experiments and collected the data. J.K., G.L., S.Z-P., C.U., and C.E.H. analyzed the data. K.Y.O., F.A., and B.L. provided protocols, antigens, cells and pseudovirus stocks. G.E-A., J.C.B., and S.L. obtained specimens. All authors read and approved the final manuscript.

a

Fig. 1. Titration of SARS-CoV-2-specific total Ig in plasma. (a) Titration curves are shown for total Ig against spike, RBD, S1, S2, nucleoprotein, and BSA in plasma specimens from seven vaccinated and four COVID-19-negative donors. Specimens were diluted at 4-fold dilutions from 1:100 to 1:6,400 or 102,400. The dotted gray lines indicated the cut-off calculated as the mean + 3SD of a 1:100 dilution of the four COVID-19 negative specimens. All data were generated using the multiplex bead antibody binding assay and plotted as mean fluorescent intensity (MFI). (b) Relative levels of plasma total Ig against spike, RBD, nucleoprotein (N), and BSA, as measured by area under the curves (AUC), in specimens from seven vaccinated donors and eight convalescent COVID-19 individuals tested in parallel. Statistical significance was calculated using a Kruskal-Wallis test.

a

b

	Plasma - vaccinated subjects			
	Linear regression spike MFI		Linear regression RBD MFI	
	r²	p	r²	p
Total Ig versus IgM	0.22	0.29	0.15	0.39
Total Ig versus IgG1	**0.86**	**0.003**	**0.88**	**0.002**
Total Ig versus IgG2	0.47	0.09	0.34	0.17
Total Ig versus IgG3	**0.70**	**0.02**	**0.64**	**0.03**
Total Ig versus IgG4	0.29	0.21	0.23	0.28
Total Ig versus IgA1	0.23	0.28	0.17	0.36
Total Ig versus IgA2	0.20	0.31	0.14	0.41

c

Fig. 2. SARS-CoV-2 immunization induces IgG, IgM and IgA Ab responses against spike and RBD in plasma. (a) Total Ig, IgM, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2 levels against spike and RBD were measured in plasma specimens from seven vaccinated and four COVID-19-negative individuals. The black line represents the median for each group. The percentages of responders above the cut-off for each spike- or RBD-specific Ig isotype are indicated. (b) Simple linear regression of spike-specific or RBD-specific plasma total Ig levels versus IgM, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2 levels from the seven vaccinated individuals. (c) Spike- or RBD-specific ratios of IgG1+IgG3 MFIs over IgG2+IgG4 MFIs for the vaccinated and convalescent groups. Statistical significance was calculated using a Mann-Whitney test.

Fig. 3. Heatmaps to show the relative levels of antibody binding, neutralization and Fc-mediated activities in (a) plasma and (b) saliva specimens from seven vaccinated donors, eight COVID-19 convalescent patients, and four COVID-19-negative controls. Y-axis represents each individual and X-axis represents each Ab binding or activity studied. Data were normalized by dividing each individual data by the highest value of the group.

a

● Vaccinated subjects
+ COVID-19-negative subjects

b

c

	Plasma - vaccinated subjects			
	Linear regression spike MFI		Linear regression RBD MFI	
	r^2	p	r^2	p
Total Ig versus neutralization	0.88	0.002	0.85	0.003
IgM versus neutralization	0.26	0.24	0.29	0.21
IgG1 versus neutralization	0.86	0.002	0.84	0.003
IgG2 versus neutralization	0.56	0.05	0.48	0.09
IgG3 versus neutralization	0.71	0.02	0.70	0.02
IgG4 versus neutralization	0.41	0.12	0.35	0.16
IgA1 versus neutralization	0.37	0.15	0.32	0.19
IgA2 versus neutralization	0.34	0.17	0.33	0.18

Fig. 4. Neutralization activities are detected in the plasma of all vaccinated individuals.

(a) Neutralization of SARS-CoV2 pseudovirus by plasma specimens from seven vaccinated individuals and two COVID-19-negative individuals. The dotted line indicates 50% neutralization. (b) Neutralization IC_{50} titers of plasma from seven vaccinated compared to eight convalescent donors. Statistical significance was calculated using a Mann-Whitney test. (c) Simple linear regression of neutralization IC_{50} titers versus spike- or RBD-specific total Ig, IgM, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2 levels from the seven vaccinated individuals.

Linear regression		
	r^2	p
Total Ig versus ADCP	0.72	0.02
IgM versus ADCP	0.41	0.12
IgG1 versus ADCP	0.90	0.001
IgG2 versus ADCP	0.75	0.01
IgG3 versus ADCP	0.71	0.02
IgG4 versus ADCP	0.38	0.14
IgA1 versus ADCP	0.16	0.38
IgA2 versus ADCP	0.45	0.1

Fig. 5. ADCP activities are detected in the plasma of all vaccinated individuals. (a) Spike-specific ADCP activities in plasma specimens from seven vaccinated individuals and two COVID-19-negative individuals. (b) ADCP AUC for the seven vaccinated and seven COVID-19 convalescent donors. Statistical significance was calculated using a Mann-Whitney test. (c) ADCP potency was calculated as ADCP AUC/total Ig AUC for the seven vaccinated and seven COVID-19 convalescent donors. Statistical significance was calculated using a Mann-Whitney test. (d) Simple linear regression of ADCP AUC versus spike-specific total Ig, IgM, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2 levels in the seven vaccinated individuals.

a

b

	Plasma - vaccinated subjects			
	Linear regression spike MFI		Linear regression RBD MFI	
	r²	p	r²	p
Total Ig versus C1q	0.74	0.01	0.83	0.05
IgM versus C1q	0.65	0.03	0.45	0.1
IgG1 versus C1q	0.97	< 0.0001	0.95	0.0002
IgG2 versus C1q	0.74	0.01	0.33	0.18
IgG3 versus C1q	0.86	0.003	0.84	0.004
IgG4 versus C1q	0.21	0.3	0.13	0.43
IgA1 versus C1q	0.07	0.56	0.04	0.66
IgA2 versus C1q	0.67	0.03	0.43	0.11

	Plasma - vaccinated subjects			
	Linear regression spike MFI		Linear regression RBD MFI	
	r²	p	r²	p
Total Ig versus C3d	0.64	0.03	0.75	0.01
IgM versus C3d	0.63	0.03	0.32	0.19
IgG1 versus C3d	0.92	0.0006	0.96	0.0001
IgG2 versus C3d	0.69	0.02	0.48	0.09
IgG3 versus C3d	0.79	0.008	0.73	0.01
IgG4 versus C3d	0.19	0.33	0.25	0.26
IgA1 versus C3d	0.06	0.6	0.07	0.56
IgA2 versus C3d	0.65	0.03	0.32	0.19

c

Spike — Spearman r = 0.90, p < 0.0001

RBD — Spearman r = 0.86, p < 0.0001

d

Fig. 6. Complement-binding activities are detected in the plasma of vaccinated individuals. (a) C1q or C3d binding to spike- or RBD-specific Abs in plasma specimens from seven vaccinated individuals and four COVID-19-negative controls. Specimens were diluted at 2-fold dilutions from 1:100 to 1:6,400 or 12,800. The dotted line represents the 100x dilution cut-off calculated as mean of the four control specimens + 3SD. ΔC1q and ΔC3d MFI values were calculated by subtracting background MFI from each assay. (b) Simple linear regression of C1q or C3d AUC values versus spike-specific total Ig, IgM, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2 levels from the seven vaccinated individuals. (c) Spearman correlation between spike- or RBD-specific Ab C1q AUC and C3d AUC for plasma specimens from the seven vaccinated (circles) and eight convalescent (diamonds) donors. (d) Complement binding and activation potencies of vaccine- versus infection-induced Abs were calculated by dividing the C1q or C3d AUC with spike-specific Ig AUC for each of the seven vaccinated and eight convalescent individuals. Statistical significance was calculated using a Kruskal-Wallis test.

Fig. 7. Titration of SARS-CoV-2-specific total Ig in saliva. (a) Relative levels of plasma total Ig against spike, RBD, nucleocapsid, and BSA, as measured by area under the curves, (AUC), with specimens from seven vaccinated donors and eight COVID-19 convalescent patients tested in parallel. Statistical significance was calculated using a Kruskal-Wallis test. All data were generated using a multiplex bead antibody binding assay and plotted as mean fluorescent intensity (MFI). (b) Spearman correlation between spike- or RBD-specific total Ig AUC of plasma specimens versus saliva specimens from vaccinated (circles) and convalescent (diamonds) individuals. (c) Total Ig, IgM, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2 levels against spike and RBD in saliva specimens from seven vaccinated and four COVID-19 negative individuals. The black line represents the median for each group. The percentages of responders above the cut-off for each spike- or RBD-specific Ig isotype are indicated. (d) Spike- or RBD-specific ratios of IgG1+IgG3 MFIs over IgG2+IgG4 MFIs for vaccinated and convalescent saliva specimens. Statistical significance was calculated using a Mann-Whitney test.