

1 **Characteristics of SARS-CoV-2 Positive Individuals in California From Two**
2 **Periods During Notable Decline in Incident Infection**

3 Lao-Tzu Allan-Blitz,¹ Isaac Turner,² Fred Hertlein,² Jeffrey D. Klausner³

4

5 1. Brigham and Women's Hospital, Boston, MA

6 2. Curative Inc. San Dimas, CA

7 3. Department of Preventive Medicine, Keck School of Medicine, University of
8 Southern California, Los Angeles, CA

9

10 Disclosures: Dr. Allan-Blitz served as a consultant for Curative Inc., Fred Hertlein is an
11 employee of Curative Inc., Isaac Turner is a Co-Founder and Chief Information Officer
12 of Curative Inc., and Dr. Klausner is the medical director of Curative Inc.

13

14 Funding and Acknowledgements: The authors would like to acknowledge the cities of
15 Los Angeles, Riverside, San Mateo, Berkeley, Menlo Park, Maywood, and Rancho
16 Mirage. There was no funding for this project.

17

18 Abstract Word Count: 99/100

19 Word Count: 1,323/1,500

20 References: 15/15

21 Tables: 1

22 Figures: 1

23

24 **Corresponding Author Contact Information:**

25 Lao-Tzu Allan-Blitz

26 Department of Medicine, Brigham and Women's Hospital

27 75 Francis Street

28 Boston, MA 02115, USA

29 Email: lallan-blitz@partners.org, phone: (805) 896-5313

30

31 **Alternative Author Information:**

32 Dr. Jeffrey D. Klausner

33 Department of Preventive Medicine

34 Keck School of Medicine

35 University of Southern California

36 Los Angeles, CA 90033, USA

37 Email: jdklausner@med.usc.edu

38

39

40 **Abstract**

41 Despite declining SARS-CoV-2 incidence, continued epidemic monitoring is warranted.
42 We collected SARS-CoV-2 test results from 150 drive-through testing centers across
43 California from two observation periods: February 23rd-March 3rd 2021 and April 15th-
44 April 30th 2021. We assessed SARS-CoV-2 positivity, stratified by Hispanic heritage
45 among sociodemographic characteristics and potential exposures. We analyzed
46 114,789 test results (5.1% and 2.6% positive during the respective observation periods).
47 Nearly half of all positive tests were among testers reporting a recent exposure (48.8%
48 and 45.3% during the respective observation periods). Those findings may provide
49 insight into evolving local transmission dynamics and support targeted public health
50 strategies.

51 **Key Words:** SARS-CoV-2; Epidemiology; California

52 **Introduction**

53 Over the last few months, the daily incidence of SARS-CoV-2 cases has declined
54 sharply in California (1), as well as in many parts of the United States. The cause of the
55 dramatic decline is likely multifactorial but greatly influenced by population-level
56 immunity due to prior infection and increasing vaccination coverage (see Figure) (2).

57 SARS-CoV-2, however, continues to be transmitted heterogeneously among
58 different subsets of the population (3), and over the last few months several states
59 reported resurging case counts (1). Fears that certain variants of SARS-CoV-2 (4) may
60 escape immunity warrant continued epidemic monitoring. We aimed to characterize
61 characteristics of those with SARS-CoV-2 positive test results during the recent decline
62 in cases in California to understand current viral transmission.

63

64 **Methods**

65 We evaluated SARS-CoV-2 polymerase chain reaction (PCR) test results from a
66 large testing program in California at two time periods: 1) February 23rd - March 3rd
67 2021, and 2) between April 15th - April 30th 2021. Individuals presented for testing to any
68 of 150 drive-through publicly accessible sites located in various counties and cities
69 including Los Angeles, Riverside, San Mateo, Berkeley, Menlo Park, Maywood, and
70 Rancho Mirage. Those who completed testing were asked via a confidential online
71 survey if in the last 14 days they had been contacted by local public health authorities
72 about a known SARS-CoV-2 exposure, they visited any of a list of public places, or
73 spent time with five or more strangers, as well as demographic information, employment
74 information, and report of any symptoms at the time of testing.

75 Self-collected oral swabs, which have been shown to perform similarly to
76 clinician-collected nasopharyngeal swabs (5), were processed with standard PCR
77 methods using a modified Center for Disease Control and Prevention testing protocol as
78 has been previously reported (6). We then conducted a cross-sectional analysis to
79 determine the frequency of infection among testers based on each of the above
80 characteristics. Further, we calculated positivity ratios by various characteristics to
81 evaluate trends in SARS-CoV-2 positivity during a period of notable decline in incident
82 infection. To account for confounding of Hispanic heritage, given the reports of the
83 disproportionate burden of infection among Hispanic persons (7), we stratified our
84 analysis by heritage. The Mass General Brigham institutional review board deemed the
85 analysis of de-identified data did not constitute human subjects' research
86 (2020P003530). All analyses were conducted using STATA 15.1 (StataCorp, College
87 Station, TX).

88

89 **Results**

90 We analyzed 114,789 test results (see Table). Of 52,9078 results between
91 February 23rd - March 3rd, 2021 2,679 (5.1%) were positive. Of 61,882 test results from
92 April 15th - 30th, 1,579 (2.6%) were positive. Testers included 47.9% who identified with
93 Hispanic heritage, and 54.6% reported female sex.

94 In the first period, the positivity of SARS-CoV-2 infection was higher among
95 Hispanic (7.6%) compared to non-Hispanic (2.8%) testers (p -value<0.001). In the
96 second period, the positivity of SARS-CoV-2 infection among Hispanic testers was 3.0%
97 compared to 2.0% among non-Hispanic Individuals (p -value=0.09). Of all who tested

98 positive in the first period 1,309 (48.8%) reported contact with an individual known to be
99 infected with SARS-CoV-2 in the last 14 days. In the second period, a similar
100 proportion, 715 (45.3%) of the 1,579 individuals who tested positive reported exposure
101 to a known contact in the last 14 days.

102 Among Hispanic testers during the first period, we found a high positivity of
103 infection among children (11.8%) and those who reported mixed Native Hawaiian or
104 Pacific Island and Hispanic heritage (6.5%). We found consistently elevated positivity
105 through both periods among individuals reporting any known exposure in the past 14
106 days (16.1% and 10.2%, respectively), individuals reporting employment as disability
107 care providers (5.5% and 5.5%, respectively), food service providers (8.6% and 3.7%,
108 respectively), and employment in retail or manufacturing (8.3% and 3.9%, respectively).

109 Among non-Hispanic testers, we found consistently elevated SARS-CoV-2
110 positivity among individuals reporting employment in retail or manufacturing in both
111 periods, periods (3.9% and 2.6%, respectively) as well as among testers reporting any
112 known exposure in the past 14 days (10.2% and 10.6%, respectively).

113

114 **Discussion**

115 We evaluated sociodemographic characteristics and potential exposures among
116 those presenting for SARS-CoV-2 testing during two periods of notable decline in
117 incident cases in California. The current trends in SARS-CoV-2 case rates across the
118 United States are at an uncertain point, with several states reporting continued declines,
119 while some are experiencing a resurgence (1). An epidemiologic understanding of those
120 most at risk for infection is essential as the transmission dynamics shift, in order to

121 guide future prevention efforts toward those at ongoing risk. Prevention programs
122 should include vaccination, and among those unvaccinated continued employment-
123 based regular testing, ventilation, indoor mask use and distancing. We identified notable
124 SARS-CoV-2 positivity among testers of Hispanic heritage and those reporting a recent
125 known exposure.

126 Nearly half of infections in our testing population were among individuals who
127 reported contact with someone known to be infected with SARS-CoV-2 in the last 14
128 days. A finding which persisted through both periods. Thus, contact tracing efforts,
129 perhaps now more than ever, are essential for identifying and isolating remaining cases,
130 and testing and quarantine of those exposed(8). Incorporation of detailed exposure
131 reporting into contact tracing efforts may also facilitate real-time monitoring of the
132 variations in risk exposures. The collection of more specific exposure data from both
133 infected and uninfected persons undergoing testing should be added as a crucial
134 strategy moving forward (9). Additionally, continuing to encourage testing and
135 vaccination among individuals with a known exposure will be essential.

136 We found high test positivity among Hispanic testers, and among children
137 specifically during the first period, supporting the key role of within household
138 transmission (10). Because schools were mostly closed during the initial observation
139 period in California, and the reopening of schools predominantly overlapped with the
140 second observation period, it is unlikely that school attendance is contributing to the
141 continued spread of infection. Household crowding among Hispanic individuals is
142 common with higher average number of individuals per room with multiple generations
143 (11), likely contributing more substantially substantial intra-familial transmission.

144 The disparities noted in SARS-CoV-2 positivity among testers identifying as
145 individuals of mixed minority heritage are consistent with recent studies (7, 12, 13),
146 which likely reflects systemic structural inequities that continue to put certain
147 populations at increased risk of exposure due to inadequate protections. Particularly
148 among individuals of Hispanic heritage, types of employment likely play a key role in
149 sustained transmission (14). Our findings suggest that prevention strategies may benefit
150 from focusing on employers in particular those employing food service workers and
151 disability care providers. Disability care providers are a particularly important population
152 as the morbidity and mortality of SARS-CoV-2 infection are substantial among
153 individuals of long-term care facilities (15). Thus, we encourage requiring vaccination
154 among such employment categories. Further research is still needed to both clarify the
155 specific exposures within different sub-populations that may contribute to sustained
156 transmission, as well as to address the underlying structural inequities that result in the
157 disparities identified.

158 Our study had several limitations. First, we analyzed laboratory-based testing
159 data and could not account for individuals with repeat testing. Second, we were unable
160 to collect detailed socioeconomic data like household size in order to control for
161 confounding factors. Data collection was also incomplete for some fields making further
162 statistical analyses and modeling not possible. However, the strengths of our study
163 were the very large sample size, thus improving precision of our results, the unbiased
164 collection of exposure data prior to receiving testing results, and the inclusion of
165 numerous testing sites across California, improving the generalizability of our results.

166

167 **Conclusions**

168 We report SARS-CoV-2 positivity by sociodemographic employment and various
169 exposure characteristics in California from two observation periods during a notable
170 decline in incident infection. We found higher SARS-CoV-2 positivity among Hispanic
171 testers as well as testers with a known recent exposure, as well as variations in
172 positivity by type of employment. Those findings are important because of the evolving
173 epidemiology of those at risk as the incidence of infection decreases, thus providing the
174 groundwork for future research and potentially informing public health strategies.

175

176 **References**

- 177 1. Centers for Disease Control and Prevention. COVID Data Tracker: United States
178 COVID-19 Cases and Deaths by State. Last updated April 2nd 2021. Available at:
179 https://covid.cdc.gov/covid-data-tracker/#cases_casesper100klast7days Accessed April
180 4th, 2021.
- 181 2. U.S. COVID Risk & Vaccine Tracker. Available at:
182 <https://covidactnow.org/?s=1793712> Last Accessed April 29th, 2021.
- 183 3. Centers for Disease Control and Prevention. COVID Data Tracker: Integrated
184 County View. Available at: [https://covid.cdc.gov/covid-data-](https://covid.cdc.gov/covid-data-tracker/?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fcases-updates%2Fcases-in-us.html#county-view)
185 [tracker/?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-](https://covid.cdc.gov/covid-data-tracker/?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fcases-updates%2Fcases-in-us.html#county-view)
186 [ncov%2Fcases-updates%2Fcases-in-us.html#county-view](https://covid.cdc.gov/covid-data-tracker/?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fcases-updates%2Fcases-in-us.html#county-view) Accessed 5/9/21. 2020.
- 187 4. Galloway SE, Paul P, MacCannell DR, et al. Emergence of SARS-CoV-2 B.1.1.7
188 Lineage - United States, December 29, 2020-January 12, 2021. MMWR Morb Mortal
189 Wkly Rep. 2021;70(3):95-9.
- 190 5. Food and Drug Administration. Accelerated Emergency Use Authorization
191 Summary: Curative SARS-CoV-2 Assay. Table 11. Available at:
192 <https://www.fda.gov/media/137089/download> Accessed March 24, 2021.
- 193 6. Kojima N, Turner F, Slepnev V, et al. Self-Collected Oral Fluid and Nasal Swab
194 Specimens Demonstrate Comparable Sensitivity to Clinician-Collected Nasopharyngeal
195 Swab Specimens for the Detection of SARS-CoV-2. Clin Infect Dis. 2020.
- 196 7. Macias Gil R, Marcelin JR, Zuniga-Blanco B, Marquez C, Mathew T, Piggott DA.
197 COVID-19 Pandemic: Disparate Health Impact on the Hispanic/Latinx Population in the
198 United States. J Infect Dis. 2020.

- 199 8. Hellewell J, Abbott S, Gimma A, et al. Feasibility of controlling COVID-19
200 outbreaks by isolation of cases and contacts. *Lancet Glob Health*. 2020;8(4):e488-e96.
- 201 9. Allan-Blitz L-T, Klausner JD. Response to: Seroprevalence of SARS-CoV-2
202 following the largest initial epidemic wave in the United States: Findings from New York
203 City, May 13-July 21, 2020. *The Journal of Infectious Diseases*. 2021.
- 204 10. Cerami C, Rapp T, Lin FC, et al. High household transmission of SARS-CoV-2 in
205 the United States: living density, viral load, and disproportionate impact on communities
206 of color. *medRxiv*. 2021.
- 207 11. Burr JA, Mutchler JE, Gerst K. Patterns of residential crowding among Hispanics
208 in later life: immigration, assimilation, and housing market factors. *J Gerontol B Psychol*
209 *Sci Soc Sci*. 2010;65(6):772-82.
- 210 12. Ogedegbe G, Ravenell J, Adhikari S, et al. Assessment of Racial/Ethnic
211 Disparities in Hospitalization and Mortality in Patients With COVID-19 in New York City.
212 *JAMA Netw Open*. 2020;3(12):e2026881.
- 213 13. Anand S, Montez-Rath M, Han J, et al. Prevalence of SARS-CoV-2 antibodies in
214 a large nationwide sample of patients on dialysis in the USA: a cross-sectional study.
215 *Lancet*. 2020.
- 216 14. Chen YH GM, Riley A, Balmes J, Duchowny Kate, Harrison R, Matthay E,
217 Bibbins-Domingo K. Excess mortality associated with the COVID-19 pandemic among
218 Californians 18–65 years of age, by
219 occupational sector and occupation: March through October 2020. *medRxiv*. Available
220 at: <https://www.medrxiv.org/content/10.1101/2021.01.21.21250266v1.full.pdf>. 2021.


221 15. Kaiser Family Foundation. State reports of long-term care facility cases and
222 deaths related to COVID-19 (as of October 16, 2020). Accessed October 17, 2020.
223 Available at: [https://www.kff.org/health-costs/issue-brief/state-data-and-policy-actions-](https://www.kff.org/health-costs/issue-brief/state-data-and-policy-actions-to-address-coronavirus/#top)
224 [to-address-coronavirus/#top](https://www.kff.org/health-costs/issue-brief/state-data-and-policy-actions-to-address-coronavirus/#top).

225

226

227

228 **Figure: New SARS-CoV-2 Infections Against Percent Vaccinated in California,**
229 **January - March 2021**
230



231
232
233 This is a figure shows percent vaccinated (by first and second dose) and new SARS-CoV-2 cases
234 per 100,000 individuals in California between January to March 2021. Data extrapolated from
235 U.S. COVID Risk & Vaccine Tracker (2).
236

237 **Table: SARS-CoV-2 Positivity Among Hispanics Individuals From California**
 238 **Comparing February 23rd - March 3rd and April 15th - April 30th, 2021**
 239

	Hispanic Ethnicity							Non-Hispanic Ethnicity						
	Total		No. SARS-CoV-2 Positive		SARS-CoV-2 Positivity		Positivity Ratio	Total		No. SARS-CoV-2 Positive		SARS-CoV-2 Positivity		Positivity Ratio
	Feb	April	Feb	April	Feb	April	February vs April	Feb	April	Feb	April	Feb	April	February vs April
Total	24929	30119	1907	907	7.6%	3.0%	2.5	27978	31763	772	672	2.8%	2.1%	1.3
Age														
<18 years	2,889	8,264	341	200	11.8%	2.4%	4.9	3,573	8,081	90	106	2.5%	1.3%	1.9
18-24 years	4,424	5,713	281	174	6.4%	3.0%	2.1	3,509	3,868	109	101	3.1%	2.6%	1.2
25-34 years	6,865	6,770	442	222	6.4%	3.3%	2.0	7,642	6,493	199	207	2.6%	3.2%	0.8
35-49 years	5,914	5,530	419	186	7.1%	3.4%	2.1	7,126	7,146	163	135	2.3%	1.9%	1.2
50-64 years	3,865	3,002	345	97	8.9%	3.2%	2.8	4,758	4,543	162	92	3.4%	2.0%	1.7
≥ 65 years	970	840	79	28	8.1%	3.3%	2.4	1,370	1,632	49	31	3.6%	1.9%	1.9
Gender														
Female	13,764	17,110	951	462	6.9%	2.7%	2.6	14,677	17,082	350	342	2.4%	2.0%	1.2
Male	11,111	12,947	954	444	8.6%	3.4%	2.5	13,212	14,610	419	328	3.2%	2.2%	1.4
Heritage														
American Indian or Alaska Native	245	338	16	4	6.5%	1.2%	5.5	59	88	5	1	8.5%	1.1%	7.5
Black or African American	203	251	11	10	5.4%	4.0%	1.4	3,322	4,486	62	145	1.9%	3.2%	0.6
Native Hawaiian or Another Pacific Island	73	70	10	4	13.7%	5.7%	2.4	354	442	16	12	4.5%	2.7%	1.7
Asian	95	134	3	0	3.2%	0.0%	*	6,199	6,812	159	101	2.6%	1.5%	*
Multiracial	401	514	18	15	4.5%	2.9%	1.5	805	1,022	17	19	2.1%	1.9%	1.1
White	5,365	6,756	330	195	6.2%	2.9%	2.1	14,920	16,071	429	316	2.9%	2.0%	1.5
Other/Prefer Not to Share	16,820	20,072	1,450	629	8.6%	3.1%	2.8	1,346	1,123	60	59	4.5%	5.3%	0.8
Employment														
Agricultural or Food Manufacturing Worker	183	323	12	10	6.6%	3.1%	2.1	143	211	4	1	2.8%	0.5%	*
Construction Worker	4	9	0	3	0.0%	33.3%	0.0	4	3	0	0	0.0%	0.0%	*
Correctional Facility	41	34	5	3	12.2%	8.8%	1.4	23	28	1	1	4.3%	3.6%	1.2
Delivery or Ride Share	2	6	0	1	0.0%	16.7%	0.0	8	1	1	0	12.5%	0.0%	*
Disability Care Provider	440	451	24	25	5.5%	5.5%	1.0	500	517	11	5	2.2%	1.0%	2.3
Education	1,160	3,812	44	23	3.8%	0.6%	6.3	1,886	3,694	24	36	1.3%	1.0%	1.3
Emergency Services	213	253	10	4	4.7%	1.6%	3.0	216	263	4	4	1.9%	1.5%	1.2
First Responder	5	4	0	0	0.0%	0.0%	*	3	2	0	0	0.0%	0.0%	*
Food Services	1,847	1,859	159	68	8.6%	3.7%	2.4	1,317	1,295	52	51	3.9%	3.9%	1.0
Government Personnel	5	4	0	0	0.0%	0.0%	*	2	2	0	0	0.0%	0.0%	*
Grocery Store Worker	8	3	0	0	0.0%	0.0%	*	3	3	0	0	0.0%	0.0%	*
Healthcare Worker	1,338	1,779	62	34	4.6%	1.9%	2.4	1,812	2,182	37	28	2.0%	1.3%	1.6
Members of the Media	0	0	0	*	*	*	*	2	1	0	0	0.0%	0.0%	*

Public Transportation	198	248	10	8	5.1%	3.2%	1.6	178	255	6	4	3.4%	1.6%	2.1
Retail or Manufacturing	1,698	1,480	141	58	8.3%	3.9%	2.1	1,116	890	44	23	3.9%	2.6%	1.5
Public Exposures														
Bars	136	338	0	2	0.0%	0.6%	0.0	311	625	3	11	1.0%	1.8%	0.5
Restaurants	1,269	3,232	13	32	1.0%	1.0%	1.0	3,168	5,446	17	29	0.5%	0.5%	1.0
Gas Stations	2,076	2,908	46	21	2.2%	0.7%	3.1	3,990	4,750	25	39	0.6%	0.8%	0.8
Public Park	1,006	2,704	13	24	1.3%	0.9%	1.5	3,337	4,877	12	18	0.4%	0.4%	*
Retail Store	1,785	3,862	34	36	1.9%	0.9%	2.0	3,739	5,744	17	31	0.5%	0.5%	0.8
Grocery Store	3,603	6,457	78	65	2.2%	1.0%	2.2	6,864	9,132	42	62	0.6%	0.7%	0.9
Place of Work	869	1,204	23	7	2.6%	0.6%	4.6	1,346	1,705	7	12	0.5%	0.7%	0.7
Public Transit	413	954	4	9	1.0%	0.9%	1.0	863	1,237	4	15	0.5%	1.2%	0.4
Place of Worship	243	549	4	3	1.6%	0.5%	3.0	236	512	1	1	0.4%	0.2%	*
Known COVID Contact in Last 14 days														
No	17,039	24,551	952	503	5.6%	2.0%	2.7	22,315	27,490	382	349	1.7%	1.3%	1.3
Yes	5,812	3,917	934	400	16.1%	10.2%	1.6	3,662	2,980	375	315	10.2%	10.6%	1.0