Transmission roles of symptomatic and asymptomatic COVID-19 cases: a modeling study

Authors:

Jianbin Tan[1]†, Yang Ge[2]†, Leonardo Martinez[3], Zhiping Chen[4], Changwei Li[5], Adrianna Westbrook[2], Enfu Chen[4], Jinren Pan[4], Yang Li[6], Feng Ling[4], Jimin Sun[4]‡, Hui Huang[1]‡, Ye Shen[2]‡.

Author affiliations:

[1]Sun Yat-Sen University, School of Mathematics, Guangzhou, China;
[2]University of Georgia, College of Public Health, Department of Epidemiology and Biostatistics, Athens, Georgia, United States;
[3]Stanford University, School of Medicine, Division of Infectious Diseases and Geographic Medicine, Stanford, California, United States;
[4]Zhejiang Provincial Center for Disease Control and Prevention, Hangzhou, China;
[5]Tulane University School of Public Health and Tropical Medicine, Department of Epidemiology, New Orleans, Louisiana, United States;
[6]Renmin University of China, School of Statistics, Beijing, China.

† Joint first authors; ‡ Joint senior authors.

Corresponding Author:

Dr. Ye Shen, University of Georgia, College of Public Health, Department of Epidemiology and Biostatistics, Athens, Georgia, United States

Word count: 3485.

Key words: COVID-19, SARS-CoV-2, age-dependent contact, asymptomatic case, transmission

Conflict of interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Abstract

Objective

Age-dependent asymptomatic and symptomatic transmission dynamics of COVID-19

have not been well quantified due to limited data.

Methods

Through a population-based surveillance network, we collected data on 1342 confirmed

cases with a 90-days follow-up for all asymptomatic cases.

Results

The difference in transmissibility of a symptomatic and asymptomatic case depended on

age and was most distinct for the middle-age groups. The asymptomatic cases had a

66.72% lower transmissibility rate than symptomatic cases, and 74.10% (95%CI: 65.85%

- 80.72%) of all asymptomatic cases were missed in detection. The average proportion

of asymptomatic cases was 28.22% (95%CI: 22.97% - 34.56%). Simulation showed that

the burden of asymptomatic transmission increased as the epidemic continued and

could potentially dominate the spreading.

Conclusion

Asymptomatic COVID-19 cases play a significant role in transmission. Vaccine Strategies

prioritizing the population between 30-60 years old are likely to have the most

population-level benefits.

Background

COVID-19, caused by the novel coronavirus (SARS-CoV-2),[1] is a great threat to

human health.[2] Nonpharmaceutical interventions such as social distancing, case

isolation and contact quarantines are the most common tools for suppressing the

pandemic in many countries where vaccine supplements are limited.[3–5] However, these

interventions are less effective in asymptomatic transmission prevention and control.[6]

COVID-19 patients may present and remain pre-symptomatic, asymptomatic, or

symptomatic and transmission may occur at each of these disease states.[7–9] Unlike the

transmission caused by symptomatic cases, pre-symptomatic and asymptomatic

transmission are hard to detect and difficult to measure as many surveillance systems

rely on symptom-based population screening.[7,10–12] Previous case studies suggested that

asymptomatic COVID-19 individuals are less infectious than symptomatic cases.[13,14]

However, asymptomatic cases may spread for a longer period due to reduced efficiency

in case detection.[15] Several studies investigated the silent transmission of SARS-CoV-2,

but presented contradictory conclusions with estimated burden ranged from 3% to

79%.[15–17]

Without sufficient follow-up time, asymptomatic and pre-symptomatic cases are

often indistinguishable. Consequently, studies using population-level data to estimate of

age-specific transmission and susceptibility parameters commonly falls short of accuracy

which potentially explains for the heterogeneous findings from different studies.[18–20]

Common issues were modeling without data on observed asymptomatic infection[15–17,21]

and inclusion of pre-symptomatic cases as part of an asymptomatic classification,[18–20] .

Meanwhile, few studies assessing asymptomatic infectiousness and viral load with

limited sample sizes fail to capture the transmission dynamics.[13,14,22–29] No studies to

date have attempted to combine reliable case symptom classification with

age-dependent transmissibility, social contact measures, and susceptibility parameters

at the population level to learn the transmission dynamics. However, a comprehensive

understanding of the age-specific symptomatic and asymptomatic transmission

dynamics at the population level is essential to the evaluation of an epidemic and the

creation of responding health policies.

In this study, we report on a longitudinal cohort of all diagnosed COVID-19

infections, between January 7[th] and February 22[nd], 2020, from Zhejiang province, China.

All patients without initial symptoms were followed by at least 90 days to distinguish

between asymptomatic and pre-symptomatic cases, an essential procedure rarely

implemented by previous studies to ensure reliable classification of case symptoms. We

then built age-stratified compartmental models to study the age-dependent

population-level transmission roles of symptomatic and asymptomatic COVID-19 cases.

Methods

Data sources

Zhejiang province is an eastern coastal province adjacent to Shanghai city with a population of approximately 54 million individuals.[30] The first and only major wave of the COVID-19 epidemic in Zhejiang began on January 7th, 2020 and continued until February 22nd, 2020 after which only sporadic single-case events were observed. We included information from all confirmed cases in this major wave (a total of 1342 cases), as well as a follow-up investigation related to all detected asymptomatic infections to distinguish between asymptomatic and pre-symptomatic cases. Individual-level data related to the symptom onset of symptomatic infections, as well as COVID-19 confirmation dates and ages of both symptomatic and asymptomatic cases were collected. For the contact data, we obtained contact matrices between age groups from surveys conducted in Shanghai.[31,32] Other primary sources of data used in our analyses include the population census data of Zhejiang province[30] and the estimated age-dependent susceptibility of COVID-19 from a previous study.[21]

On January 23rd, 2020, the provincial government changed its infectious disease alert category to the highest level and, on February 1st, began a comprehensive set of interventions.[33] As of April 10th, 2020, the date in which we restricted our data for this analysis, no additional outbreak had been observed. Trained health professionals investigated each confirmed case with a predefined questionnaire by which basic health and demographic information were collected.

Definition of symptomatic and asymptomatic cases

All confirmed cases and their close contacts were isolated or quarantined after

being identified through contact tracing. During the isolation/quarantine period, cases

and their contacts received regular testing and daily symptom screening for fever,

cough, and shortness of breath. Tests for case confirmation were conducted using

reverse transcription polymerase chain reaction (RT-PCR) or viral genome sequencing on

samples from throat swabs (oropharynx and nasopharynx). If a case or contact had a

positive test result but without any symptoms, they would be temporarily classified as

an asymptomatic/pre-symptomatic case at the time. All cases were followed for at least

90 days after their initial positive test to distinguish between asymptomatic and

pre-symptomatic cases. Among these subjects, those who developed symptoms later

would receive a final classification as a symptomatic case. Others who had never

developed any symptoms between their initial positive test and first subsequent

negative PCR test would be classified as asymptomatic cases.

Model structure

We divided the total population of Zhejiang province into seven age groups:

0-10, 10-20, 20-30, 30-40, 40-50, 50-60, 60+. To consider transmission related to

symptomatic and asymptomatic infections among different age groups, our model

contained 8 compartments for the i^{th} age group: susceptible population (S^i), exposed

contacts (E^i), pre-symptomatic cases (I_{ps}^i, infected but have not yet developed

symptoms), symptomatic cases (I_s^i), asymptomatic cases (I_a^i, infected but asymptomatic

till confirmed/recovery), and removed/recovery groups (R_{cs}^i, R_{ca}^i, R_h^i). We assumed

new infections were driven by transmission from compartments of I_{ps}^i, I_s^i and I_a^i in

all age groups. Asymptomatic cases (I_a^i) were infections without typical symptoms.

These infections were difficult to identify through symptom-based screening; therefore,

underreporting in this category of infections was notable. While some asymptomatic

cases may have been quarantined through contact tracing, others may have remained

unnoticed throughout the entirety of their transmissibility. As such, it is essential to

consider unconfirmed asymptomatic cases, also referred as the self-recovered

asymptomatic cases (R_h^i in Figure 1) in modeling. For all age groups, the details of the

compartments are given in Table 1, and the state transitions in the compartmental

model are given in Figure 1.

<u>Social contacts</u>

To identify age-varying transmissibility and susceptibility within the

compartmental model,[21] we assumed a time-varying curve for the average contact

numbers of i^{th} age group with j^{th} age group (c_t^{ij}), which is estimated with the

contact matrix between age groups through surveys conducted in Shanghai.[31,32] The

surveys provided two different strengths of contact (before and during the epidemic).

The first survey conducted in 2017 to 2018 in Shanghai was considered as a background

of the strength of contact.[16] The second survey, with the same design, was conducted

during the epidemic (from February 1 to 10, 2020). The details of the modeling of the

contact function (c_t^{ij}) are further described in the Supplementary Appendix.

<u>Susceptibility-contact-transmissibility transmission process</u>

When modeling transmission dynamics of infectious diseases, a key factor to

consider is the probability of infection for a susceptible individual given contact with an

infectious individual. Here, we separate this concept into two components:

transmissibility (T) and susceptibility (s). We define transmissibility (T) as the

infectiousness of one case. Theoretically, it is the secondary attack rate in their fully

susceptible contacts $(s = 1)$. Similarly, we define susceptibility (s) as the probability of

acquiring infection from an infectious case $(T = 1)$. Therefore, $s = 0$ corresponds to a

situation in which the susceptible individuals are immune to the disease (this setting

was later used to conduct a simulation study to assess possible age-dependent

vaccination strategies if a certain age group were to achieve 100% immunity by

vaccinations). We assumed that case transmissibility would depend on age and the

presence of symptoms. To better quantify transmissibility and susceptibility, we

introduced the estimated susceptibility (s_i) for i^{th} age group from a previous study as

a prior in our model.[21] In this setting, the occurrence of a transmission event depends

on both transmissibility of the index case and the susceptibility of the contact. The

number of secondary cases (ρ) from one infectious case at unit time can be calculated

as:

$$\rho = s \cdot \text{contact number} \cdot T$$

Let $T_{i,s}$ be the transmissibility of symptomatic infections in i^{th} age group. To

capture the age-dependent pattern, B splines basis functions were used to model the

variability in age-varying transmissibility smoothly. Given $T_{i,s}$, we assumed that the

age-varying transmissibility of asymptomatic infectious individuals was equivalent to

$T_{i,a} := \gamma_i T_{i,s}$, where $\gamma_i \leq 1$, i.e., the transmissibility of asymptomatic cases is no

stronger than that of symptomatic cases across different age groups, based on currently

available evidence[13,14,34]. The estimated mean of s_i and the B spline functions

mentioned above are shown in Table S3 and Figure S2 (in the Supplementary Appendix).

<u>Asymptomatic infections</u>

Our compartmental model (Figure 1) considered the transmission from

asymptomatic infections, which were often untraceable in the clinical survey and,

therefore, their contribution to population-level transmission would be underestimated.

To account for this, we assumed only a proportion of asymptomatic infections were

detected (R_{ca}^i), while others (R_h^i) would be unconfirmed (and not diagnosed) if

self-recovery occurred prior to diagnosis. We were able to observe disease confirmation

date but not the date of infection for the period from cases becoming infectious to the

diagnosis of COVID-19 ($\mathcal{A}_1^i$) in those with confirmed asymptomatic infection. Based on

the virus shedding pattern of asymptomatic infections reported in previous

studies,[12,14,35,36] we assumed that this period should be less than 30 days, after which

virus shedding generally ceases, and infection is no longer detectable through

pathogen-specific testing. As such, a uniform prior with an upper bound of 30 days was

adopted for $\mathcal{A}_1^i$. For the period from being infectious to becoming non-infectious

(negative conversion of virus shedding) in the undetected asymptomatic infections ($\mathcal{A}_2$),

we used a more informative log normal prior with a mean of 17 days and standard error

of 1.07 based on estimated parameters from a previous study.[12,14,35,36] The difference

between $\mathcal{A}_1^i \ and \ \mathcal{A}_2$ would reflect the intensity of the contact tracing for different

ages. More details are provided in the Supplementary Appendix.

<u>Model fitting</u>

Let N_i be the total population of i^{th} age group in Zhejiang province. Based on

the setting above, we defined a dynamic system for the 8×7 compartments, which is

shown in the equation (S3) (in the Supplementary). The state transitions in the

compartmental model are given in Figure 1. Based on our compartmental model, we

can estimate the secondary cases transmitted by symptomatic and asymptomatic

indexes respectively, which are used to evaluate the burden of both symptomatic and

asymptomatic transmission from different age groups. More details are given in the

equation (S4) and (S5) (in the Supplementary).

We built our model with features mentioned in above sections through a

Bayesian procedure for the estimation and inference of model parameters (listed in

Table S5).[37–39] Specifically, the compartmental model was fitted to the daily new

symptomatic and asymptomatic cases in Zhejiang province for each age group. The

Bayesian procedure with a Markov Chain Monte Carlo (MCMC) sampling algorithm also

utilized information relating to the COVID-19 transmission from other sources of

datasets, e.g. survey studies on the contact matrix, and estimated age-dependent

susceptibility from previous studies, therefore, the posterior distributions of parameters

could be elegantly extracted by the sufficient prior and sample evidence. More details of

the Bayesian model structure, assessment, and estimation of parameters are given in

the Supplementary Appendix. All analyses were implemented in R version 3.5.1.

Packages of deSolve,[40] extraDistr,[41] and splines[42] were used for model fitting.

Unless stated otherwise, the medians of the posterior distributions were used as the

point estimators of parameters and simulated numbers.

Results

Results

<u>Transmissibility</u>

The estimated transmissibility presented an age-dependent difference between

symptomatic and asymptomatic infections (Figure 2). While the transmission of

symptomatic cases monotonically increased with increasing age, the transmissibility of

asymptomatic infection remained low until age 40, after which point it significantly

increased with increasing age. The age-varying ratios of the two kinds of transmissibility

indicated asymptomatic cases were, on average, 66.72% lower in transmission than

symptomatic cases. However, the difference between the two types of infections was

not as big in those aged 0-20 and 60+ years old, but became more obvious in the

middle-aged group where the ratios were as low as 24.42% and 23.38% for those aged

30-40 and 40-50 years old, respectively.

<u>The proportion of asymptomatic cases</u>

In Figure 3, the proportion of asymptomatic cases ($\frac{R_h^i + R_{ca}^i}{R_h^i + R_{cs}^i + R_{ca}^i}$) estimated by our

model was much larger than what was observed in the data. The average proportion of

asymptomatic cases was 28.22% (95%CI: 22.97% - 34.56%) of the total counts of cases in

our model estimation, but was 9.24% in the observed data ($\frac{R_{ca}^i}{R_{cs}^i + R_{ca}^i}$). In our estimation

from the empirical data, the highest proportion of asymptomatic case was among 0-10

(60.18% (95%CI: 53.61% - 66.99%)) and 10-20 (57.64% (95%CI: 47.45% − 66.98%)) years

old groups. For asymptomatic cases, we further estimated the proportion of cases that

failed to be detected $(\frac{R_h^i}{R_h^i+R_{ca}^i})$. In the posterior samples, the average proportion of

unconfirmed cases in all asymptomatic infections was 74.10% (95%CI: 65.85% - 80.72%).

The maximum proportion of unconfirmed cases was observed in 20-30 years old at

86.59% (95%CI: 73.64% - 92.19%).

<u>Symptomatic and asymptomatic transmission</u>

To explore the impact of symptomatic and asymptomatic transmission, we

present several features of the estimated dynamic of the epidemic and the transmission

burden caused by symptomatic and asymptomatic cases in Figure 4. The estimated

number of daily new transmissions reached a peak around ten days prior to the peak of

the daily reported new confirmed cases (Figure 4A). We estimated a substantial number

of undetected asymptomatic cases (109 (95%CI: 73 - 164)) were infected before the first

asymptomatic case was diagnosed (January 27[th]) (Figure 4B). New transmissions were

nearly eliminated by February 1[st], 2020 (Figure 4A), when a comprehensive set of

restrictions[19] had been implemented. The peak of the two types of transmission both

occurred between January 17[th] to 21[st] (Figure 4C). The average burden of asymptomatic

transmission during the major outbreak period was estimated to be 12.86% (95%CI:

7.54% - 19.27%). The burden of asymptomatic transmission increased with time, ranging

from 7.77% to 16.03% (Figure 4D). Simulation studies were conducted to investigate the

dynamic changes in the transmission burden over time during a prolonged epidemic

(Figure S5). When the duration of the decreasing process of the contact function

(represented by "m" in Figure S1) was prolonged by two weeks and each individual's

daily contact number was increased by one person during the outbreak period, we

found a slower decreasing trend in daily new cases infected by asymptomatic cases

compared with that contributed by symptomatic cases (Figure S5, scenario 1).

Additional scenarios were generated demonstrating the possibility of asymptomatic

transmission dominating the total transmission under different conditions, especially

when the duration between symptom onset and disease confirmation for symptomatic

infections was shortened and the asymptomatic infections were not controlled (Figure

S5, scenarios 2 and 3).

<u>Age-depended transmission</u>

Within each age group, we observed heterogeneous transmission contributions

during different time periods (Figure 5A). Early on in the epidemic, the transmission

burden was dominated by persons of 50-60 years old (32.75% from January 7[th] to

January 11[th]), but the proportion of transmission contribution from people over 60 years

old significantly increased over time, surpassing the 50-60 years old and reaching 30.42%

by January 31[st], 2020. The proportion of transmission contribution among varying age

groups was distinct between symptomatic and asymptomatic cases (Figure 5B). The

majority of both symptomatic and asymptomatic transmissions were contributed by

persons over 30 years old (Table S9). Individuals below 30 years old only contributed

less than 5% of all symptomatic transmission and approximately 12% of all

asymptomatic transmission, respectively, despite representing almost 40% of the entire

population. Contributions to asymptomatic transmission among 20-30 and >60 year age

groups (9.44%, and 31.73%, respectively) were substantially higher than their

corresponding contributions to symptomatic transmission (3.77%, and 26.55%,

respectively). To further understand possible age-dependent vaccination strategies, a

simulation of seven scenarios was conducted to assess the percentage decline in

different age groups if one age group were to achieve 100% immunity by vaccinations

(Figure S6). The results suggested that vaccinations targeting age groups above 30 years

are likely to be more effective at the population level, with the most percentage decline

of cases from the entire population achieved by targeting the 50-60 years old group.

Meanwhile, vaccinating those younger than 30 years old are more likely to benefit their

own age groups.

Discussion

In our study, we found that asymptomatic cases were over 60% less infectious compared to symptomatic cases. While great efforts like mass screening and strict contact tracing were conducted, our results suggested that a large proportion of asymptomatic infections were still not detected.[43] The burden of asymptomatic transmission was inferior in the early outbreak but could become higher with the continuous spread of COVID-19. Under strained resources, age-specific prevention and control strategies aimed at the middle age population may return greater population-level benefits.

Current evidence suggests that asymptomatic COVID-19 cases are generally less infectious[13] than cases with symptoms. We found that this difference may partially be explained by patient age. Age may directly impact COVID-19 transmission through virus shedding patterns[14] as discussed in previous studies.[44] Symptoms are commonly mild in children[45] but severe in the elderly.[46] While still debatable,[47] higher severity has been associated with increased shedding of the virus.[48] In our study, symptomatic and asymptomatic cases were most infectious in individuals 60 years old or older. In contrary to the monotonic increasing association between age and transmission in symptomatic cases, there was a plateau of a low degree of transmission in young asymptomatic infections. We suspect older adults are not only the most vulnerable to succumb to COVID-19 but also may be more likely to transmit once infected, regardless of symptom status. Interventions attempting to suppress asymptomatic transmission,

such as mask-wearing, should primarily focus on older adults if interventions to the

whole community are not feasible.

Similar to previous studies, our results suggest a small proportion of

asymptomatic cases have been detected since the start of the COVID-19 pandemic in

our setting.[49–51] Symptom-based screening has limited capability in asymptomatic case

detection,[6] while mass pathogen or immunological-based testing at the population-level

consumes tremendous health resources, and thus is not feasible in most settings.

Considering these challenges, age-dependent screening strategies may be more

practical. We found that the highest number of undetected asymptomatic cases was

among young adults aged 20 to 30 years old (Table S7) and the corresponding

transmission contribution was significantly higher than that of symptomatic case (Figure

5B). Young people were less likely to adhere to social distancing guidelines,[52] often had

mild symptoms or were asymptomatic after infection,[53] and were not prioritized in

prevention and controlling strategies.[54] Meanwhile, younger asymptomatic patients

were also more likely to have normal CAT scan findings, which may further complicate

case detection.[11] Case detection of asymptomatic COVID-19 cases based on current

control strategies implemented in this study is alarming.

Based on the estimated transmission contributions from symptomatic and

asymptomatic infections, roughly 13% of infections were associated with asymptomatic

transmission and that percentage continuously increased with a prolonged period. The

overall burden of transmission was mainly contributed by symptomatic cases at the

beginning of the epidemic, but asymptomatic infections appeared to have increasing

percentages of subsequent cases later on. Additional simulations suggested that the

transmission burden could even be dominated by asymptomatic transmissions under

certain circumstances (Figure S5). Therefore, the spreading potential of asymptomatic

cases cannot be ignored, especially in the later stages of the epidemic and in regions

where social distancing has not been mandated. Meanwhile, potential differences in

transmission burden by age groups, as shown in Figure 5 and S6, supports prioritizing

age-dependent prevention and control strategies when facing strained resources. As the

larger contributor to the transmission of COVID-19, the older age population is not only

a highly vulnerable group but should also be the primary target for prevention

strategies. Vaccine Strategies prioritizing the population between 30-60 years old are

likely to have the most population-level benefits.

There are several limitations in this study. First, data collection likely missed

potential cases of the epidemic, despite intensified efforts devoted by the local

investigation team to trace contacts. Due to this, we introduced a compartment in our

model (R_h^i) to adjust for poor case ascertainment and missing cases. Second,

transmissibility and susceptibility were two factors related to symptomatic and

asymptomatic transmission estimation and can be difficult to capture simultaneously.

We used the susceptibility estimates from a previous study[21] as priors in our model to

account for this parameter identification problem. Third, the contact survey data we

used in our model were obtained in Shanghai, a city adjacent to Zhejiang province.

Although the two regions share a similar culture and modes of social activities, there

were potential uncertainties associated with the discrepancies in contact matrices. To

address this limitation, we introduced a correction parameter in our model, so these

uncertainties were partially adjusted for in the analyses.

In summary, our results suggest individual-level transmissibility of COVID-19

increases with patient age, therefore targeting older age groups with prevention and

intervention strategies is expected to be more efficient. While asymptomatic cases are

difficult to trace, the burden of asymptomatic transmission is still sizable and should not

be ignored. The results from our study can be used to inform policy decisions on

pandemic control and safe reopening.

<u>Ethics Statement</u>

The research protocol was approved by the institutional review board at the

Zhejiang Provincial Center for Disease Control and Prevention.

References

References

1   Guan W-J, Ni Z-Y, Hu Y, *et al.* Clinical Characteristics of Coronavirus Disease

2019 in China. *N Engl J Med* 2020; **382**: 1708–20.

2   World Health Organization. WHO Coronavirus Disease (COVID-19)

Dashboard (12/12/2020). https://covid19.who.int (accessed Dec 12, 2020).

3   Ferguson N, Laydon D, Nedjati Gilani G, *et al.* Report 9: Impact of

non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare

demand. Imperial College London, 2020 DOI:10.25561/77482.

4   Prem K, Liu Y, Russell TW, *et al.* The effect of control strategies to reduce

social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a modelling

study. *The Lancet Public Health* 2020; **5**: e261–70.

5   Lewnard JA, Lo NC. Scientific and ethical basis for social-distancing

interventions against COVID-19. *The Lancet Infectious Diseases* 2020; **20**: 631–3.

6   Gandhi M, Yokoe DS, Havlir DV. Asymptomatic Transmission, the Achilles'

Heel of Current Strategies to Control Covid-19. *N Engl J Med* 2020; **382**: 2158–60.

7   Rothe C, Schunk M, Sothmann P, *et al.* Transmission of 2019-nCoV Infection

from an Asymptomatic Contact in Germany. *N Engl J Med* 2020; **382**: 970–1.

8   Holshue ML, DeBolt C, Lindquist S, *et al.* First Case of 2019 Novel

Coronavirus in the United States. *N Engl J Med* 2020; **382**: 929–36.

9   Hoehl S, Rabenau H, Berger A, *et al.* Evidence of SARS-CoV-2 Infection in

Returning Travelers from Wuhan, China. *N Engl J Med* 2020; **382**: 1278–80.

10   Wölfel R, Corman VM, Guggemos W, *et al.* Virological assessment of

hospitalized patients with COVID-2019. *Nature* 2020; **581**: 465–9.

11   Kronbichler A, Kresse D, Yoon S, Lee KH, Effenberger M, Shin JI.

Asymptomatic patients as a source of COVID-19 infections: A systematic review and

meta-analysis. *International Journal of Infectious Diseases* 2020; **98**: 180–6.

12   Han MS, Seong M-W, Kim N, *et al.* Viral RNA Load in Mildly Symptomatic

and Asymptomatic Children with COVID-19, Seoul, South Korea. *Emerging Infectious*

Diseases 2020; **26**: 2497–9.

13   Gao M, Yang L, Chen X, *et al.* A study on infectivity of asymptomatic

SARS-CoV-2 carriers. *Respir Med* 2020; **169**: 106026.

14   Kim SE, Jeong HS, Yu Y, *et al.* Viral kinetics of SARS-CoV-2 in asymptomatic

carriers and presymptomatic patients. *Int J Infect Dis* 2020; **95**: 441–3.

15   Moghadas SM, Fitzpatrick MC, Sah P, *et al.* The implications of silent

transmission for the control of COVID-19 outbreaks. *PNAS* 2020; **117**: 17513–5.

16   Li R, Pei S, Chen B, *et al.* Substantial undocumented infection facilitates the

rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* 2020; **368**: 489–93.

17   Ferretti L, Wymant C, Kendall M, *et al.* Quantifying SARS-CoV-2

transmission suggests epidemic control with digital contact tracing. *Science* 2020; **368**.

DOI:10.1126/science.abb6936.

18   Hu Z, Song C, Xu C, *et al.* Clinical characteristics of 24 asymptomatic

infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life*

Sci 2020; **63**: 706–11.

19   Huang L, Zhang X, Zhang X, *et al.* Rapid asymptomatic transmission of

COVID-19 during the incubation period demonstrating strong infectivity in a cluster of

youngsters aged 16-23 years outside Wuhan and characteristics of young patients with

COVID-19: A prospective contact-tracing study. *J Infect* 2020; **80**: e1–13.

20   Meng H, Xiong R, He R, *et al.* CT imaging and clinical course of

asymptomatic cases with COVID-19 pneumonia at admission in Wuhan, China. *J Infect*

2020; **81**: e33–9.

21   Davies NG, Klepac P, Liu Y, *et al.* Age-dependent effects in the transmission

and control of COVID-19 epidemics. *Nat Med* 2020; **26**: 1205–11.

22   Zhou R, Li F, Chen F, *et al.* Viral dynamics in asymptomatic patients with

COVID-19. *International Journal of Infectious Diseases* 2020; **96**: 288–90.

23   Berlin DA, Gulick RM, Martinez FJ. Severe Covid-19. *N Engl J Med* 2020;

published online May 15. DOI:10.1056/NEJMcp2009575.

24   Verity R, Okell LC, Dorigatti I, *et al.* Estimates of the severity of coronavirus

disease 2019: a model-based analysis. *Lancet Infect Dis* 2020; **20**: 669–77.

25   CDC COVID-19 Response Team. Severe Outcomes Among Patients with

Coronavirus Disease 2019 (COVID-19) - United States, February 12-March 16, 2020.

MMWR Morb Mortal Wkly Rep 2020; **69**: 343–6.

26   Kong D, Zheng Y, Wu H, *et al.* Pre-symptomatic transmission of novel

coronavirus in community settings. *Influenza Other Respir Viruses* 2020; **14**: 610–4.

27   Furukawa NW, Brooks JT, Sobel J. Evidence Supporting Transmission of

Severe Acute Respiratory Syndrome Coronavirus 2 While Presymptomatic or

Asymptomatic. *Emerg Infect Dis* 2020; **26**. DOI:10.3201/eid2607.201595.

28   Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic

proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond

Princess cruise ship, Yokohama, Japan, 2020. *Eurosurveillance* 2020; **25**.

DOI:10.2807/1560-7917.ES.2020.25.10.2000180.

29   Bai Y, Yao L, Wei T, *et al.* Presumed Asymptomatic Carrier Transmission of

COVID-19. *JAMA* 2020; **323**: 1406–7.

30   Bureau of Statistics. Zhejiang Provincial Bureau of Statistics: Sixth Census

Data. 2014; published online Sept 3.

http://tjj.zj.gov.cn/art/2014/9/3/art_1530851_20980968.html (accessed July 30, 2020).

31   Zhang J, Litvinova M, Liang Y, *et al.* Changes in contact patterns shape the

dynamics of the COVID-19 outbreak in China. *Science* 2020; **368**: 1481–6.

32   Zhang J, Klepac P, Read JM, *et al.* Patterns of human social contact and

contact with animals in Shanghai, China. *Scientific Reports* 2019; **9**: 15141.

33   Chong KC, Cheng W, Zhao S, *et al.* Monitoring disease transmissibility of

2019 novel coronavirus disease in Zhejiang, China. *International Journal of Infectious*

Diseases 2020; **96**: 128–30.

34   Luo L, Liu D, Liao X, *et al.* Contact Settings and Risk for Transmission in 3410

Close Contacts of Patients With COVID-19 in Guangzhou, China : A Prospective Cohort

Study. *Ann Intern Med* 2020; **173**: 879–87.

441  35 Long Q-X, Tang X-J, Shi Q-L, *et al.* Clinical and immunological assessment of

442 asymptomatic SARS-CoV-2 infections. *Nat Med* 2020; **26**: 1200–4.

443  36 Lee S, Kim T, Lee E, *et al.* Clinical Course and Molecular Viral Shedding

444 Among Asymptomatic and Symptomatic Patients With SARS-CoV-2 Infection in a

445 Community Treatment Center in the Republic of Korea. *JAMA Internal Medicine* 2020;

446 **180**: 1447–52.

447  37 Bolstad WM, Curran JM. Introduction to Bayesian statistics. John Wiley &

448 Sons, 2016.

449  38 Wikle CK, Berliner LM. A Bayesian tutorial for data assimilation. *Physica D:*

450 *Nonlinear Phenomena* 2007; **230**: 1–16.

451  39 Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. Bayesian

452 Data Analysis, Third Edition. CRC Press, 2013.

453  40 Soetaert K, Petzoldt T, Setzer RW. Solving Differential Equations in R:

454 Package deSolve. *Journal of Statistical Software* 2010; **33**: 1–25.

455  41 Wolodzko T, Wolodzko MT. Package 'extraDistr'. 2017.

456  42 Perperoglou A, Sauerbrei W, Abrahamowicz M, Schmid M. A review of

457 spline function procedures in R. *BMC Medical Research Methodology* 2019; **19**: 46.

458  43 Flaxman S, Mishra S, Gandy A, *et al.* Estimating the effects of

459 non-pharmaceutical interventions on COVID-19 in Europe. *Nature* 2020; **584**: 257–61.

460  44 Esposito S, Principi N. To mask or not to mask children to overcome

461 COVID-19. *Eur J Pediatr* 2020; **179**: 1267–70.

45   Dong Y, Mo X, Hu Y, *et al.* Epidemiology of COVID-19 Among Children in

China. *Pediatrics* 2020; **145**: e20200702.

46   Zhou F, Yu T, Du R, *et al.* Clinical course and risk factors for mortality of

adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The*

Lancet 2020; **395**: 1054–62.

47   He X, Lau EHY, Wu P, *et al.* Temporal dynamics in viral shedding and

transmissibility of COVID-19. *Nature Medicine* 2020; **26**: 672–5.

48   Liu Y, Yan L-M, Wan L, *et al.* Viral dynamics in mild and severe cases of

COVID-19. *The Lancet Infectious Diseases* 2020; **20**: 656–7.

49   Fauci AS, Lane HC, Redfield RR. Covid-19 — Navigating the Uncharted. *New*

England Journal of Medicine 2020; **382**: 1268–9.

50   García LF. Immune Response, Inflammation, and the Clinical Spectrum of

COVID-19. *Front Immunol* 2020; **11**: 1441.

51   Oran DP, Topol EJ. Prevalence of Asymptomatic SARS-CoV-2 Infection⬚: A

Narrative Review. *Ann Intern Med* 2020; **173**: 362–7.

52   Czeisler MÉ, Tynan MA, Howard ME, *et al.* Public attitudes, behaviors, and

beliefs related to COVID-19, stay-at-home orders, nonessential business closures, and

public health guidance—United States, New York City, and Los Angeles, May 5–12, 2020.

Morbidity and Mortality Weekly Report 2020; **69**: 751.

53   Felsenstein S, Hedrich CM. COVID-19 in children and young people. *The*

Lancet Rheumatology 2020; **2**: e514–6.

54   Bedford J, Enria D, Giesecke J, *et al.* COVID-19: Towards controlling of a

pandemic. *The Lancet* 2020; **395**: 1015–8.

Tables and Figures

Table 1. The definitions of compartments.

Figure 1. Compartmental model for SARS-CoV-2 transmission, where "j" represents another age group different from "i" for the compartments.

Figure 2. (A) The estimated transmissibility and 95% credible intervals for each age group; (B) The ratios of asymptomatic transmissibility to symptomatic transmissibility for seven age groups.

Figure 3. The proportion of asymptomatic infections and unconfirmed asymptomatic infections until February 22nd, 2020 for seven age groups.

Figure 4. The estimated dynamics of the epidemic and the transmission burdens from symptomatic and asymptomatic cases. (A): The estimated numbers of daily new transmissions with 95% credible intervals and the observed numbers of daily reported new confirmed cases from January 7th to February 21st, 2020; (B) The observed numbers of daily reported new confirmed symptomatic (R_{cs}) and asymptomatic cases (R_{ca}) and the estimated numbers of daily new cases that failed to be detected (R_h) with 95% credible intervals; (C): The estimated numbers of infected individuals caused by symptomatic and asymptomatic transmission over time, with 95% credible intervals; (D): Corresponding proportions of symptomatic and asymptomatic transmissions over different time periods.

Figure 5. The burden of transmission caused by different ages. (A) The estimated (contribution) ratios of new transmissions from different ages over different time periods; (B) The estimated (contribution) ratios of symptomatic and asymptomatic transmission from different ages. The contribution ratio of each age group is calculated by the proportion of the transmissions caused by the corresponding age group to the number of all transmissions in each transmission type, from January 7th to February 1st, 2020.

Table 1. The definitions of compartments

Compartments	Meaning
S^i	Represents the individuals who are susceptible to the infectious disease.
E^i	Represents the individuals who are exposed to the infectious disease but are not infectious. The individuals of E would transfer to I_{ps} or I_a with proportion α_s^i and α_a^i for different ages after latent period $\mathcal{L}$.
I_{ps}^i	Represents the symptomatic infectious individuals before symptom onset, which would transfer to I_s^i after the period of pre-symptomatic transmission $\mathcal{S}_1$.
I_s^i	Represents the symptomatic infectious individuals after symptom onset, which would transfer to R_{cs}^i after the period of symptomatic transmission $\mathcal{S}_2^i$.
I_a^i	Represents the asymptomatic infectious individuals, which would transfer to R_{ca}^i (confirmed by testing after $\mathcal{A}_1^i$ days for different ages) or R_h^i (self-recovered after $\mathcal{A}_2$ days) eventually.
R_{cs}^i, R_{ca}^i and R_h^i	Represents the individuals assumed to be noninfectious and immune to the disease in the first wave of Zhejiang epidemic (roughly two months).

Figure 1. Compartmental model for SARS-CoV-2 transmission, where "*j*" represents
another age group different from "*i*" for the compartments.

Figure 2. (A) The estimated transmissibility and 95% credible intervals for each age
group; (B) The ratios of asymptomatic transmissibility to symptomatic transmissibility
for seven age groups.

Figure 3. The proportion of asymptomatic infections and unconfirmed asymptomatic
infections until February 22nd, 2020 for seven age groups. The estimated proportions of
asymptomatic cases, the proportions of cases that failed to be detected among
asymptomatic infections (unconfirmed proportions), and the observed proportions of
asymptomatic cases are defined as: $\frac{R_h^i + R_{ca}^i}{R_h^i + R_{cs}^i + R_{ca}^i}$, $\frac{R_h^i}{R_h^i + R_{ca}^i}$, and $\frac{R_{ca}^i}{R_{cs}^i + R_{ca}^i}$, respectively. The
95% credible intervals for the estimated proportions of asymptomatic cases are shown
for each age group.

Figure 4. The estimated dynamics of the epidemic and the transmission burdens from symptomatic and asymptomatic cases. (A): The estimated numbers of daily new transmissions with 95% credible intervals and the observed numbers of daily reported new confirmed cases from January 7[th] to February 21[st], 2020; (B) The observed numbers of daily reported new confirmed symptomatic (R_{cs}) and asymptomatic cases (R_{ca}) and the estimated numbers of daily new cases that failed to be detected (R_h) with 95% credible intervals; (C): The estimated numbers of infected individuals caused by symptomatic and asymptomatic transmission over time, with 95% credible intervals; (D): Corresponding proportions of symptomatic and asymptomatic transmissions over different time periods.

Figure 5. The burden of transmission caused by different ages. (A) The estimated
(contribution) ratios of new transmissions from different ages over different time
periods; (B) The estimated (contribution) ratios of symptomatic and asymptomatic
transmission from different ages. The contribution ratio of each age group is calculated
by the proportion of the transmissions caused by the corresponding age group to the
number of all transmissions in each transmission type, from January 7th to February 1st,
2020.