Cardiovascular vulnerability predicts hospitalisation in primary care clinically

suspected and confirmed COVID-19 patients: a model development and validation

study

Florien S. van Royen[1*], Linda P.T. Joosten[1], Maarten van Smeden[2], Pauline Slottje[3], Frans

H. Rutten[1], Geert-Jan Geersing[1], Sander van Doorn[1]

[1]Dept. General Practice, Julius Center for Health Sciences and Primary Care, University

Medical Center Utrecht, Utrecht University, Utrecht, Netherlands

[2]Dept. Epidemiology, Julius Center for Health Sciences and Primary Care, University

Medical Center Utrecht, Utrecht University, Utrecht, Netherlands

[3]Dept. General Practice, Academic network of general practice, Amsterdam UMC, Vrije

Universiteit Amsterdam, Amsterdam, Netherlands

*Corresponding author, e-mail: f.s.vanroyen-5@umcutrecht.nl, postal address:

Universiteitsweg 100, 3584 GX, Utrecht, Netherlands

Abstract

Objectives:

Cardiovascular conditions were shown to be predictive of clinical deterioration in hospitalised patients with coronavirus disease 2019 (COVID-19). Whether this also holds for outpatients managed in primary care is yet unknown. The aim of this study was to determine the incremental value of cardiovascular vulnerability in predicting the risk of hospital referral in primary care COVID-19 outpatients.

Design:

Analysis of anonymised routine care data extracted from electronic medical records from three large Dutch primary care registries.

Setting:

Primary care.

Participants:

Consecutive adult patients seen in primary care for COVID-19 symptoms in the 'first wave' of COVID-19 infections (March 1 2020 to June 1 2020) and in the 'second wave' (June 1 2020 to April 15 2021) in the Netherlands.

Outcome measures:

A multivariable logistic regression model was fitted to predict hospital referral within 90 days after first COVID-19 consultation in primary care. Data from the 'first wave' was used for derivation (n=5,475 patients). Age, sex, the interaction between age and sex, and the number of cardiovascular conditions and/or diabetes (0, 1, or ≥2) were pre-specified as candidate predictors. This full model was (i) compared to a simple model including only age and sex and its interaction, and (ii) externally validated in COVID-19 patients during the 'second wave' (n=16,693).

Results:

The full model performed better than the simple model (likelihood ratio test $p<0.001$). Older male patients with multiple cardiovascular conditions and/or diabetes had the highest

predicted risk of hospital referral, reaching risks above 15-20%, whereas on average this risk

was 5.1%. The temporally validated c-statistic was 0.747 (95%CI 0.729-0.764) and the model

showed good calibration upon validation.

Conclusions:

For patients with COVID-19 symptoms managed in primary care, the risk of hospital referral

was on average 5.1%. Older, male and cardiovascular vulnerable COVID-19 patients are

more at risk for hospital referral.

Introduction

Coronavirus disease 2019 (COVID-19), caused by SARS-CoV-2, has led to a global

pandemic ever since the first cases were described in late 2019. Despite clear improvement

in terms of vaccination efficacy and treatment options for hospitalised patients, COVID-19

remains a global public health burden. For instance, at present, global vaccination coverage

is still low in many countries (particularly in low- and middle-income countries). Moreover,

waning immunity after vaccination is already observed in countries with an early vaccination

program, notably in vulnerable individuals. In addition, new emerging virus variants may

escape protective immunity following vaccination, and finally, a substantial part (5-10% in

highly vaccinated countries) of the population refuse vaccination altogether for a variety of

reasons. Inevitably, (global) circulation of SARS-SoV-2 will remain with new (seasonal)

outbreaks likely to occur. Hence, COVID-19 will keep influencing the organisation of

healthcare worldwide in the upcoming years, perhaps decades.

Thereby, it is pivotal to increase our knowledge on this relatively new disease, for

instance in order to learn how to orchestrate the flow of patients depending on the expected

course or natural prognosis of the disease. Indeed, a fast-growing amount of prediction

models have been developed for prognosticating COVID-19 patients. However, most of

these models focus on an in-hospital population, with only few prediction models focusing on

outpatient management.(1) This is unfortunate as the clinical presentation of (suspected)

COVID-19 starts-off with initially mild to moderate symptoms in the first week of illness, and

only in some with progression to hypoxemia for which hospital (or even ICU) admission is
needed, typically occurring in the second week of illness.(2) If such deterioration occurs,
patients in the Netherlands, as in many countries worldwide, are seen in a primary care
setting. The primary care physician is therefore often the first to decide on the need and
optimal timing for more impactful measures, such as intensified monitoring or ultimately
referral for hospitalisation. Unfortunately, evidence based tools or knowledge to help primary
care physicians on deciding how to triage COVID-19 patients and detect patients in need for
referral from those in whom a relatively benign trajectory is to be expected are currently
lacking.

However, we know from hospitalised COVID-19 patients, that underlying
cardiovascular diseases are strong predictors for further disease deterioration towards ICU
admittance or death.(3–7) Moreover, we also know that in vaccinated populations, this
association still exists.(8) If indeed pre-existing cardiovascular disease and/or diabetes
increases the risk of clinical deterioration already in primary care, this likely is instrumental to
guide outpatient management in COVID-19 given that this clinical information is readily
available to primary care physicians. Intensified monitoring or even early treatment (e.g. with
budesonide inhalation) in high-risk patients might then prevent hospitalisation, thereby
lowering the burden on hospitals.

The aim of this study was therefore to determine the incremental value of
cardiovascular vulnerability – defined by the number of cardiovascular diseases and/or type
2 diabetes mellitus – in predicting the risk of escalation of care (i.e. hospital referral) in
primary care patients with clinically suspected or confirmed COVID-19. Results from this
analysis are intended to explore the incremental prognostic value of concurrent
cardiovascular disease in community COVID-19 outpatients, and to support primary care
physicians in applying a risk-tailored COVID-19 management in primary care.

Methods

Study design

This study involves an analysis of anonymised observational electronic medical record data

of community people registered by the primary care physician with either confirmed or

clinically suspected COVID-19. We assessed the incremental value of cardiovascular

disease and/or diabetes by developing a prognostic prediction model in a cohort of patients

from the 'first wave' of COVID-19 infections in the Netherlands (March 1 2020 to June 1

2020) that was temporally validated in a cohort of patients from the 'second wave' of

infections in the Netherlands (June 1 2020 to April 15 2021). Where appropriate for this

study, we adhered to the TRIPOD guideline for reporting prediction models.(9)

Databases

Patients were included from three similar ongoing and dynamic primary care databases run

by the academic hospitals of the cities and surrounding municipalities of Utrecht and

Amsterdam, containing pseudonymised medical data of approximately 850,000 patients in

total: the Julius General Practitioner's Network (JGPN) University Medical Center Utrecht,

the Academic Network of General Practice at VU University medical center in Amsterdam

(ANH VUmc), and the Academic General Practitioner's Network at Academic Medical Center

Amsterdam (AHA AMC).(10–12) Two databases (JGPN and ANH VUmc) were used to

identify patients for the development of the prediction model (i.e. development cohort) and all

three databases (JGPN, ANH VUmc and AHA AMC) were used to identify patients for the

temporal validation (i.e. the validation cohort).

Study population and data collection

Patients for the development cohort were included from March 1 2020 to June 1 2020 (the

'first wave' of the COVID-19 pandemic in the Netherlands). During this time period, very

limited polymerase chain reaction (PCR) testing for COVID-19 was available, and moreover

mainly restricted to more severe hospitalised cases. Consequently, many symptomatic

patients consulting their primary care physician were highly suggestive of COVID-19 and

were not tested. We therefore included all consecutive adult patients aged 18 years or older,

who visited their primary care physician with confirmed COVID-19 as well as those with

symptoms that were clinically suspected by their primary care physician.

For identification of the study population and data collection, the same methods were applied

in all three databases. Dutch primary care physicians record diagnoses and clinical

symptoms in the electronic medical records as diagnostic codes using the International

Classification of Primary Care (ICPC) coding system. The primary care physicians

participating in the JGPN, AHA AMC and ANH Vumc databases are trained in and

experienced with using ICPC codes. For the development cohort, COVID-19 suspected

patients were identified using the ICPC codes R74 (acute upper respiratory infection), R81

(pneumonia) and R83 (other respiratory infection). At the time, primary care physicians were

recommended to use R81 and R83 for indicating COVID-19 suspected and COVID-19

confirmed cases, respectively, for lack of COVID-19 specific ICPCs. Records of patients

labelled with ICPC R74 (unspecified acute upper respiratory infection) were manually

screened for COVID-19 suspicion in the consultation text by three (primary care) clinical

scientists (FSvR, LPTJ, and SvD) and cases of doubt were discussed until agreement was

reached. Patients with ICPC R74 yet without having a synonym of or reference to COVID-19

suspicion or related symptoms in the consultation text were excluded from the cohorts. Of all

included patients, baseline characteristics (i.e. age; sex; relevant diagnoses; and if available

Body Mass Index (BMI), oxygen saturation and C-Reactive Protein (CRP)) were collected.

(History) of relevant diseases (i.e. cardiovascular disease, type 2 diabetes mellitus,

hypercholesterolemia, hypertension, pulmonary diseases, cancer) were identified using

ICPC. Supplementary table 1 contains all ICPC codes that were used for this study.

For the validation cohort, consecutive adult JGPN patients were included from September 1

2020 until April 15 2021, and consecutive adult AHA AMC and ANH Vumc patients were

included from June 1 2020 until December 31 2020 (the 'second wave' of COVID-19

infections). At this point in time, the Dutch government made PCR COVID-19 tests freely

available and these were recommended for all symptomatic subjects in the Netherlands and

for those who were in close contact with a confirmed COVID-19 patient. Moreover, at that

time GPs were instructed to uniformly code confirmed cases in their medical records (i.e.

using ICPC codes either R83 or R83.03). Thus, only confirmed COVID-19 cases (i.e. with

ICPC R83 and R83.03) were included in the cohort for validation of the model.

Outcome

The primary outcome of the prediction model in this study was referral to an emergency ward

for intended hospital admission. This was defined as any clinical deterioration resulting in

hospital referral by the primary care physician that was recorded as such in the consultation

annotation (free text) of the medical record. To capture the full spectrum of complications of

COVID-19 resulting in hospitalisation, follow-up lasted 90 days after first consultation for

COVID-19 suspected symptoms. To this end, all anonymised consultation texts were

manually screened for any emergency hospital referral by (primary care) clinical scientists

(FSvR, LPTJ, SvD, and GJG) and cases of doubt were discussed, until consensus was

reached.

Candidate predictors

Based upon existing literature from hospitalised COVID-19 patients, we a-priori specified the

following candidate predictors prior to the analysis phase: age, sex, the interaction between

age and sex, and the number of cardiovascular diseases. The latter was defined as (history

of) type 2 diabetes mellitus, heart failure, coronary artery disease, peripheral arterial disease,

stroke/transient ischemic attack (TIA), venous thromboembolism (pulmonary embolism or

deep venous thrombosis; VTE), and/or atrial fibrillation (AF). Presence of these diseases was

based on the corresponding ICPCs (Supplementary table 1) at any point before the first

COVID-19 consultation in the patient's medical record. The number of cardiovascular

diseases were counted per patient and categorised into: no cardiovascular disease, one

cardiovascular disease, or two or more cardiovascular diseases.

Sample size

The model development cohort in JGPN and ANH Vumc yielded 5,475 eligible patients with an event fraction of 0.068 (6.8%, n=373) for the primary outcome referral to the hospital. Prior to prediction analysis, the number of allowed candidate predictors was determined. Based on the proposed calculation for sample size in prediction modelling by Riley et al.(13), the maximum number of candidate predictors that can be modelled was 30 with a R^2 Cox-Snell (R^2cs) of 0.0495. As this R^2cs was estimated in absence of a known value, varying R^2cs from 0.0395 to 0.0595 yielded a minimum of 24 and a maximum of 37 candidate predictors, including interaction terms. By using the candidate predictors age, sex, the interaction between age and sex, and the number of cardiovascular diseases with three categories, the sample size of 5,475 eligible patients was deemed sufficient and large enough for model development.

Missing data

Candidate predictors age, sex and cardiovascular disease had no missing data. Missing values in baseline of characteristics measurements of CRP, BMI and oxygen saturation level were not imputed as these determinants were not used further in predictive modelling.

Statistical analyses

Baseline characteristics were summarised using descriptive statistics with categorical variables as numbers with percentages and continuous variables as means with standard deviations or medians with interquartile ranges (IQR). A multivariable logistic regression modelling approach was used. All included patients were entered in a fixed model (full model) with the predictors age, sex, the interaction between age and sex, and the categorical number of cardiovascular diseases and/or diabetes. To determine the incremental value of the predictor number of cardiovascular diseases and/or, a second model (simple model) was fitted using only the predictors age, sex, and the interaction between age and sex. Age was

considered as a continuous variable and was studied using a restricted cubic spline function

to account for possible non-linearity with 4 knots on the percentiles 0.05, 0.35, 0.65 and

0.95.(14) Number of cardiovascular diseases and/or diabetes was taken as a dummy

variable with 'no cardiovascular disease' as reference category. The incremental value of the

number of cardiovascular diseases and/or diabetes was studied comparing the full and

simple model's c-statistics (ΔAUC), Cox-Snell R^2cs (ΔR^2cs), and a likelihood ratio test. To

determine incremental value, an alpha of 0.05 was used for the likelihood ratio test. The full

model was internally validated using Harrell's bootstrapping with 100 repetitions to obtain

optimism corrected estimates of the c-statistic, and R^2 and slope were calculated. For the

temporal external validation of the full model, calibration and discrimination were evaluated:

observed and predicted events were calculated and depicted in calibration plots and for

discrimination areas under the curve (AUC/c-statistic) were calculated. Other performance

measures for temporal external validation that were calculated are: calibration slope,

calibration intercept, calibration in the large, R^2cs, and Brier score. Brier scores assess the

overall goodness of fit of models, with smaller numbers indicating better performance.

Confidence intervals for c-statistics were obtained using the Delong method. For R^2cs and

Brier score confidence intervals, bootstrapping was used with repetitions set at 1000.

Validation was done in the whole validation dataset as well as separately in the JGPN, ANH

VUmc, and AHA AMC validation cohorts. All statistical analyses were performed in R version

4.0.3 with R base, rms, pROC, DescTools, and rmda packages.(15–19)

Ethics

This research was conducted in accordance with Dutch law and the European Union General

Data Protection Regulation and according to the principles of the Declaration of Helsinki. The

need for formal ethical reviewing was waived by the local medical research ethics committee

of the University Medical Center Utrecht, the Netherlands as the research did not require

direct patient or physician involvement. The JHN, ANH VUmc and AHA AMC databases may

be used for scientific purposes and contain pseudonymised routine care data from the EMRs

253 of all patients of the participating general practices, except those patients who objected to

254 this. Anonymised datasets were extracted from these databases by the respective data

255 managers for the purpose of this research.

257 **Results**

258 *Patient characteristics*

259 Patient characteristics of the (clinically suspected and confirmed COVID-19) development

260 cohort are described in table 1. There were 5,475 patients included in this cohort: 2,825 from

261 JGPN and 2,650 from ANH VUmc. In JGPN, 64.6% of the patients were coded as R74,

262 14.7% as R81, and 21.3% as R83. In ANH VUmc, 71.5% were coded as R74, 10.7% as

263 R81, and 19.2% as R83. Differences in patient characteristics between both datasets in the

264 development cohort were minor. Around a quarter of patients suffered from one or more

265 cardiovascular disease, most often type 2 diabetes and coronary artery disease.

267 Patient characteristics of the (confirmed COVID-19) validation cohort are described in table

268 2. From the total of 16,693 patients in the validation cohort 5,420 originated from JGPN,

269 4,989 from ANH VUmc, and 6,284 from AHA AMC. The patient characteristics in these three

270 datasets were very similar. Around 15-20% suffered from one or more cardiovascular

271 disease, again most often type 2 diabetes and coronary artery disease.

273 *Model development and internal validation*

274 All 5,475 patients in de development cohort were used for model development. 373 patients

275 (6.8%) had the outcome hospital referral. All predefined model regression coefficients of the

276 full and simple models with confidence intervals are shown in table 3. The apparent c-

277 statistic of the full model was 0.693 (95%CI 0.665-0.721) and the internally validated c-

278 statistic was 0.688 (95%CI 0.660-0.716). The apparent c-statistic of the simple model was

279 0.681 (95%CI 0.653-0.710) and the internally validated c-statistic was 0.680 (95%CI 0.652-

0.708). The full and the simple model are compared in table 4. The full model performed

significantly better than the simple model (p- value for likelihood ratio test, χ^2 = 19.5, df = 2,

p<0.001). Figure 1 gives a visual representation of the full model showing the predicted risks

of hospital referral as a function of (increasing) age, stratified by sex and by the number of

cardiovascular diseases and/or diabetes. Overall risks are higher for male patients and

increase with age. Furthermore, a higher risk is observed in patients with underlying

cardiovascular disease.

Temporal external validation

Predicted risks were overall slightly higher than the observed risk (6.2% versus 4.6%) and

the calibration slope was 1.36. Overall discrimination showed an AUC of 0.747 (95%CI

0.729-0.764). Performance measures based on the full validation cohort and stratified by

database are shown in table 5. The overall calibration plot and the calibration plots per

database separately are shown in Supplementary figure 1. The hospital referral prevalence

was lower in the validation datasets than in the development datasets (4.7% versus 6.8%).

Discussion

Cardiovascular vulnerability is a predictor of hospital referral in a population of 5,475

consecutive adult patients in primary care with confirmed or clinically suspected COVID-19 in

the 'first wave' of infections in the Netherlands. This finding was confirmed by temporal

validation in a population of 16,693 consecutive confirmed COVID-19 adult primary care

patients in the 'second wave', exemplifying the robustness of our inferences. On average, in

the combined data from the first and second wave (n=22,168 confirmed and clinically

suspected primary care COVID-19 patients), 5.1% was referred to the hospital for

considering admission. A model including the number of cardiovascular conditions and/or

diabetes (0, 1, or ≥2) in addition to age and sex and the interaction between age and sex,

showed moderate to good performance and demonstrated consistent and good

discrimination and calibration upon temporal external validation. The model showed a c-

statistic of 0.747 (95%CI 0.729-0.764).

Although most (vaccinated) COVID-19 patients experience a favourable prognosis without

the need for referral for hospital care, studies on COVID-19 are mainly focussed on those

seen in the hospital setting. While on average the overall risk for hospital referral in this adult

primary care cohort with COVID-19 was low (5.1%), it is much higher than the hospitalisation

rate for other lower respiratory infections in primary care which is estimated at approximately

1% of the adult population affected.(20) In our study, age, sex and the number of concurrent

cardiovascular conditions and/or diabetes predicted patients at far greater risk of hospital

referral. In fact, for female patients without cardiovascular diseases or diabetes, the risk of

hospital referral was well below 10% even in the eldest elderly (aged 80+). Contrastingly, in

the presence of cardiovascular diseases and/or diabetes, patients experience higher risks

already at younger ages, notably males. For instance, a male patient with two or more

underlying cardiovascular diseases and/or diabetes, had a predicted risk of 15% already at

the age of around 57 years and this predicted risk will even further increase to above 20%

from the age of 80 onwards. This indicates the incremental effect of cardiovascular diseases

and/or diabetes in addition to age and sex in predicting the risk for complicated COVID-19

disease trajectories in primary care patients.

Our findings overall confirm those from previous studies done in the hospital setting where

age and male sex are important predictors for disease progression towards the endpoints

ICU admission or death.(1,21–23) Social, behavioural, comorbidity and biological differences

(ACE2 expression, sex-hormones, X-chromosome exposure) between male and female

sexes all might contribute to the higher risks of COVID-19 progression observed in males,

although probably not all mechanisms have been fully elucidated yet.(24,25) Also, in

hospitalised patients, it has been demonstrated that there is an association between

cardiovascular disease and COVID-19 complicated disease trajectories, with higher

prevalence of cardiovascular disease and diabetes described in those with critical illness.(3–

7,26) Our study shows that this prognostically unfavourable effect is already present much

earlier on in the COVID-19 disease course, at the start of symptoms in primary care.

This is in line with previous research, where this additive effect of (cardiovascular)

comorbidities was also described by the 4C Mortality Score.(27) In this study, the authors

demonstrated that the number of comorbidities, importantly including cardiovascular

comorbidities, had a more predictive effect than taking only individual co-morbidities in

predicting in-hospital mortality of COVID-19 patients.(27) Furthermore, there are two large

community-based prediction studies also highlighting the importance of cardiovascular

comorbidities as a predictor in the community COVID-19 population. The QCOVID model

that was developed in the UK and recently validated in a vaccinated population, was based

on data from primary care and showed a c-statistic >0.9 for the primary outcome time to

death from COVID-19. The domain of that study, however, covered the whole general

population *regardless* of COVID-19 diagnosis and therefore this can best be interpreted as

the risk prediction of getting infected with COVID-19 *and* subsequently having complications

from COVID-19. Thus, the aim of this model was to inform UK health policy and support

interventions to manage COVID-19 related risks, rather than inform medical decision making

during patient consultations in confirmed or clinically suspected COVID-19 cases.(8,28) With

only 0.07% with the outcome death, and thus very low a priori chance, the c-statistic

'misleadingly' moves towards 1.0. Another similar public health based UK study in patients

with and without COVID-19 identified determinants that were associated with COVID-19

related death in the OpenSAFELY primary care database by linking primary care records to

reported COVID-19 related deaths. It found the most predictive clinical determinants to be

increasing age, male sex, type 2 diabetes mellitus, and cardiovascular disease, similar to our

findings.(29) While the domain notably differs between patients seeking primary care for

COVID-19 symptoms in our study and the adult community as a whole in these studies form

the UK, all draw similar conclusions on the increased risk of clinical deterioration in patients

with (multiple) cardiovascular disease and/or diabetes.

Strengths and limitations

This research contributes to the evidence-based prognostication of community COVID-19.

We were able to use routine primary care databases capturing both the 'first' and 'second'

wave of COVID-19 infections in the Netherlands. We used state-of-the-art methodology

including external temporal validation to predict clinical deterioration in a patient population

that is currently understudied. For full appreciation of our findings, however, some limitations

also need to be addressed. First, the model was developed in a dataset with a low event

fraction of the outcome hospital referral. Yet the number of hospital referral events did allow

us to perform robust multivariable regression techniques. Second, there are limitations to

using routine care registry data that could have resulted in misclassification of the study

population, predictors and outcome, and most importantly it has the risk of missing values.

For example, uncertainty concerning COVID-19 infection status may exist (primarily in the

first wave) as COVID-19 PCR test results were not automatically linked to the primary care

electronic medical records. However, the model proved its transportability in primary care

patients in a different time period with satisfactory calibration and discrimination, during a

time window where PCR testing was widely performed. Furthermore, the outcome hospital

referral was based upon a rigorous manual extraction of medical records by pairs of

researchers, albeit actual hospital *admittance* was not formally confirmed based upon linkage

to hospital records. Additionally, there are differences between our development and

validation population: the patients from the 'first wave' are all symptomatic patients that

visited their primary care physician for symptoms suggestive of COVID-19, while the patients

from the 'second wave' – due to government recommendation for individuals to get tested

even in the circumstance of only mild symptoms – also include more healthy people that just

informed their primary care physician of their positive COVID-19 PCR status. This could also

explain the lower event fraction in the validation set (4.7% versus 6.8% in the development

population). Furthermore, the model still has to show its robustness in the COVID-19

vaccinated population, although it is likely that existing risk factors will still be present even if

the risk of complications is lowered due to vaccination. Finally, the incremental value of the

number of cardiovascular diseases and diabetes on prognosticating COVID-19 was

assessed in different ways; although we did observe a highly significant change in the

likelihood ratio test, the delta in c-statistic and R^2cs was only small to modest. Possible

reasons for this include the overall low risk of hospital referral in most patients in our cohort,

as well as that most patients (80.2%) in fact in our cohort did not suffer from cardiovascular

diseases and/or diabetes. It has been widely acknowledged that, notably in such scenario's,

a change in e.g. the c-statistic is difficult to achieve.

Clinical implications

The readily availability of the chosen primary care predictors and the clinical applicability may

provide great advantages for risk profiling patient with suspected or confirmed COVID-19 in

the primary care and community setting. This can have several important clinical and public

health implications. First, it may be possible to identify patients that will benefit from closer

monitoring and frequent follow-up at home by predicting the risk of clinical deterioration early

on in the COVID-19 disease course. By intensified monitoring of higher risk patients, critical

illness may be detected earlier, potentially improving prognosis. Second, risk prediction could

also support advanced care planning. Informing both patients and physicians on the risk of

severe illness, may help in anticipating a more stringent or more lenient management. Last,

risk profiling may be used for targeting preventive measures. Additionally, experimental

regiments to *treat* symptomatic COVID-19 may be addressed to high-risk patients that may

benefit most. Examples include for instance treatment with budesonide or colchicine; both

treatment options likely benefit patients most at higher prior probability of having an adverse

prognosis.(30,31) Nevertheless, in the end, risk prediction in primary care has to prove its

value in daily practice at the background of changing characteristics of this challenging

COVID-19 pandemic and influences of vaccination and virus mutations. We however do

hope that prognostic studies, like ours, may aid physicians and policy makers by making

informed, evidence-based decisions and thereby improve patient outcomes.

Conclusion

In this primary care population-based study, risk of clinical deterioration leading to hospital

referral after suspected or confirmed COVID-19 was on average 5.1%. This risk increased

with age and was higher in males compared to females. Importantly, patients with concurrent

cardiovascular disease and/or diabetes had higher predicted risks. Identifying those at risk

for hospital referral could have clinical implications for COVID-19 early disease management

in primary care.

Funding statement

This work was supported by the Dutch Heart Foundation (grant number 2020T063). The

Dutch Heart Foundation had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript

Competing interests statements

The authors declare that they have no competing interests.

Author's contribution

All authors contributed to the conception and design of this study. FSVR, LPTJ, PS and SVD

collected the data. FSVR, LPTJ and SVD analysed the data and all authors contributed to

interpretation of the data. FSVR wrote the first draft and LPTJ, MVS, GJG, FHR, PS and

SVD critically revised and approved the final manuscript.

Acknowledgements

The authors would like to thank the data managers from the JGPN, ANH VUmc and AHA

AMC databases and all primary care physicians contributing data.

Data sharing statement

The data used for this study are available from the routine primary care registries JGPN,

ANH VUmc and AHA AMC. Restrictions apply to the availability of these data, which were

used under license for the current study, and so are not publicly available. Data will however

be available for scientific purposes from the authors upon reasonable request and with

permission from the individual registries.

References

1.   Wynants L, van Calster B, Collins GS, Riley RD, Heinze G, Schuit E, et al. Systematic

review and critical appraisal of prediction models for diagnosis and prognosis of

COVID-19 infection. 2020;369:m1328.

2.   Gandhi RT, Lynch JB, del Rio C. Mild or Moderate Covid-19. N Engl J Med.

2020;383(18):1757–66.

3.   Linschoten M, Peters S, Smeden M Van, Jewbali LS, Schaap J, Siebelink H, et al.

Cardiac complications in patients hospitalised with COVID-19. Eur Heart J Acute

Cardiovasc Care. 2020;9(8):817-23

4.   Li B, Yang J, Zhao F, Zhi L, Wang X, Liu L, et al. Prevalence and impact of

cardiovascular metabolic diseases on COVID-19 in China. Clin Res Cardiol.

2020;109(5):531–8.

5.   Bello-Chavolla O, Bahena-López J, Antonio-Villa N, Vargas-Vázquez A, González-

Díaz A, Márquez-Salinas A, et al. Predicting Mortality Due to SARS-CoV-2: A

Mechanistic Score Relating Obesity and Diabetes to COVID-19 Outcomes in Mexico.

J Clin Endocrinol Metab. 2020;105(8):2752-2761.

6.   Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, et al.

Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO

Clinical Characterisation Protocol: Prospective observational cohort study. BMJ.

2020;369:m1985.

7.   Amir E, Javanmardi F, Pirbonyeh N, Akbari A. Prevalence of Underlying Diseases in

Hospitalized Patients with COVID-19: a Systematic Review and Meta- Analysis. Arch

Acad Emerg Med. 2020;8(1):e35–e35.

8.   Hippisley-Cox J, Coupland CA, Mehta N, Keogh RH, Diaz-Ordaz K, Khunti K, et al.

Risk prediction of covid-19 related death and hospital admission in adults after covid-

19 vaccination: national prospective cohort study. BMJ. 2021;374:n2244

9.   Moons KGM, Altman DG, Reitsma JB, Ioannidis JPA, Macaskill P, Steyerberg EW, et

al. Transparent reporting of a multivariable prediction model for individual prognosis or

diagnosis (TRIPOD): Explanation and elaboration. Ann Intern Med. 2015;162(1):W1–

73.

10.   Julius General Practitioner's Network (JGPN).

Https://juliuscentrum.umcutrecht.nl/nl/julius-huisartsen-netwerk. Accessed 13

September 2021.

11.   Academisch Huisartsennetwerk AMC (AHA). Https://huisartsgeneeskunde-

amc.maglr.com/amsterdam_umc_huisartsgeneeskunde_amc_jaarbeeld_2018/acade

misch-huisartsennetwerk-amc. Accessed 13 September 2021.

12.   Academisch Netwerk Huisartsgeneeskunde VU (ANH).

Https://www.vumc.nl/anh/database-anh.htm. Accessed 13 September 2021.

13.   Riley RD, Ensor J, Snell KIE, Harrell FE, Martin GP, Reitsma JB, et al. Calculating the

sample size required for developing a clinical prediction model. BMJ. 2020;368:1–12.

14.   Harrell Jr FE. Regression modeling strategies: with applications to linear models,

logistic and ordinal regression, and survival analysis. [Internet]. 2[nd] edition. Springer.

2015. Accessed 11 May 2021. Available via:

https://link.springer.com/content/pdf/10.1007/978-3-319-19425-7.pdf.

15.   R Core Team. R: A language and environment for statistical computing. R Foundation

495   for Statistical Computing, Vienna, Austria. 2020. Available via: https://www.R-

496   project.org/.

497   16.   Harrell Jr FE. rms: Regression Modeling Strategies. R package version 6.2-0. 2021.

498   Available via: https://CRAN.R-project.org/package=rms.

499   17.   Xavier R, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J-C, et al. pROC: an

500   open-source package for R and S+ to analyze and compare ROC curves. MC

501   Bioinforma. 2011;12:77.

502   18.   Andri Signorell et mult. al. DescTools: Tools for descriptive statistics. R package

503   version 0.99.41. 2021.

504   19.   Brown M. rmda: Risk Model Decision Analysis. R package version 1.6. 2018, Available

505   via: https://CRAN.R-project.org/package=rmda.

506   20.   Little P, Stuart B, Smith S, Thompson MJ, Knox K, Van Den Bruel A, et al. Antibiotic

507   prescription strategies and adverse outcome for uncomplicated lower respiratory tract

508   infections: Prospective cough complication cohort (3C) study. BMJ. 2017;357.

509   21.   Liu K, Chen Y, Lin R, Han K. Clinical features of COVID-19 in elderly patients: A

510   comparison with young and middle-aged patients. J Infect. 2020;80:e14–8.

511   22.   Liu W, Tao ZW, Wang L, Yuan ML, Liu K, Zhou L, et al. Analysis of factors associated

512   with disease outcomes in hospitalized patients with 2019 novel coronavirus disease.

513   Chin Med J (Engl). 2020;133(9):1032–8.

514   23.   Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients

515   infected with 2019 novel coronavirus in Wuhan, China. Lancet. 2020;395(10223):497–

516   506.

517   24.   Haitao T, Vermunt J V., Abeykoon J, Ghamrawi R, Gunaratne M, Jayachandran M, et

518   al. COVID-19 and Sex Differences: Mechanisms and Biomarkers. Mayo Clin Proc.

519   2020;95(10):2189–203.

520   25.   Gebhard C, Regitz-Zagrosek V, Neuhauser HK, Morgan R, Klein SL. Impact of sex

521   and gender on COVID-19 outcomes in Europe. Biol of Sex Differ. 2020;11(1).

522   26.   Nishiga M, Wang DW, Han Y, Lewis DB, Wu JC. COVID-19 and cardiovascular

523   disease: from basic mechanisms to clinical perspectives. Nat Rev Cardiol.

524   2020;17(9):543-558.

525   27.   Knight SR, Ho A, Pius R, Buchan I, Carson G, Drake TM, et al. Risk stratification of

526   patients admitted to hospital with covid-19 using the ISARIC WHO Clinical

527   Characterisation Protocol: Development and validation of the 4C Mortality Score. BMJ.

528   2020;370: m3339.

529   28.   Clift AK, Coupland CAC, Keogh RH, Diaz-Ordaz K, Williamson E, Harrison EM, et al.

530   Living risk prediction algorithm (QCOVID) for risk of hospital admission and mortality

531   from coronavirus 19 in adults: national derivation and validation cohort study. BMJ.

532   2020;371 :m3731.

533   29.   Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. Factors

534   associated with COVID-19-related death using OpenSAFELY. Nature.

535   2020;584(7821):430–436.

536   30.   Ramakrishnan S, Nicolau D V, Langford B, Mahdi M, Jeffers H, Mwasuku C, et al.

537   Inhaled budesonide in the treatment of early COVID-19 (STOIC): a phase 2, open-

538   label, randomised controlled trial. Lancet Respir Med. 2021;19(21):1–10.

539   31.   Tardif JC, Bouabdallaoui N, L'Allier PL, Gaudet D, Shah B, Pillinger MH, et al.

540   Colchicine for community-treated patients with COVID-19 (COLCORONA): a phase 3,

541   randomised, double-blinded, adaptive, placebo-controlled, multicentre trial. Lancet

542   Respir Med. 2021;9(8):924–32.

Table 1 - Baseline characteristics of the development cohort of 5,475 community patients with confirmed or clinically suspected COVID-19 from the 'first wave' in the Netherlands

Characteristics	JGPN (n=2,825)	ANH VUmc (n=2,650)
Age in years (IQR)	48 (34-62)	49 (36-62)
Male sex	1144 (40.1%)	1068 (40.3%)
Cardiovascular and metabolic diseases		
Heart failure	109 (3.9%)	80 (3.0%)
Coronary artery disease	230 (8.1%)	189 (7.1%)
Atrial fibrillation	139 (4.9%)	177 (6.7%)
Peripheral arterial disease	49 (1.7%)	37 (1.4%)
History of stroke/TIA	136 (4.8%)	117 (4.4%)
History of VTE	123 (4.4%)	100 (3.8%)
Type 2 diabetes mellitus	300 (10.6%)	262 (9.9%)
0 CVD/Diabetes	2132 (75.5%)	2053 (77.5%)
1 CVD/Diabetes	440 (15.6%)	386 (14.6%)
≥2 CVD/Diabetes	253 (9.0%)	211 (8.0%)
Other comorbidities		
History of any cancer	198 (7.0%)	207 (7.8%)
Hypertension	721 (25.5%)	616 (23.2%)
Hypercholesterolemia	261 (9.2%)	331 (12.5%)
All pulmonary disease	575 (20.4%)	432 (16.3%)
COPD	234 (8.3%)	155 (5.8%)
Asthma	434 (15.4%)	315 (11.9%)
BMI in kg/m^2 (IQR)	27 (24-31) (n=1,091)	27 (24-31) (n=1,072)
Median oxygen saturation in % (IQR)	98 (95-98) (n=191)	98 (96-99) (n=87)
CRP in mg/L (IQR)	6 (2-23) (n=480)	3 (1-9) (n=646)
Hospital referrals	185 (6.5%)	188 (7.1%)

 BMI = Body Mass Index; CRP = C-reactive protein; CVD = cardiovascular disease; IQR =

552 interquartile rage; TIA = transient ischemic attack; VTE=venous thromboembolism.

556 **Table 2** - Baseline characteristics of the validation cohort of 16,693 community patients with

557 confirmed COVID-19

Characteristics	JGPN (n=5,420)	ANH VUmc (n=4,989)	AHA AMC (n=6,284)
Age in years (IQR)	43 (30-56)	47 (34-59)	49 (36-60)
Male sex	2400 (44.3%)	2121 (42.5%)	2462 (39.2%)
Cardiovascular diseases			
Heart failure	86 (1.6%)	64 (1.3%)	101 (1.6%)
Coronary artery disease	262 (4.8%)	270 (5.4%)	346 (5.5%)
Atrial fibrillation	122 (2.3%)	139 (2.8%)	176 (2.8%)
Peripheral arterial disease	39 (0.7%)	44 (0.9%)	39 (0.6%)
History of stroke/TIA	123 (2.3%)	151 (3.0%)	229 (3.6%)
History of VTE	124 (2.3%)	134 (2.7%)	226 (3.6%)
Type 2 diabetes mellitus	476 (8.8%)	495 (9.9%)	715 (11.4%)
0 CVD/Diabetes	4566 (84.2%)	4059 (81.4%)	4964 (79.0%)
1 CVD/Diabetes	586 (10.8%)	663 (13.3%)	955 (15.2%)
≥2 CVD/Diabetes	268 (4.9%)	267 (5.4%)	365 (4.8%)
Other comorbidities			
History of any cancer	207 (3.8%)	331 (6.6%)	304 (4.8%)
Hypertension	883 (16.3%)	1,009 (20.2%)	1,486 (23.6%)
Hypercholesterolemia	309 (5.7%)	493 (9.9%)	527 (8.4%)
All pulmonary diseases	648 (12.0%)	610 (12.2%)	867 (13.8%)
COPD	174 (3.2%)	186 (3.7%)	184 (2.9%)
Asthma	534 (9.9%)	477 (9.0%)	726 (11.6%)
BMI in kg/m^2 (IQR)	28 (24-32) (n=1,685)	27 (24-32) (n=1,823)	29 (25-33) (n=2,178)
Median oxygen saturation in % (IQR)	98 (96-98) (n=176)	98 (97-99) (n=134)	98 (97-99) (n=75)
CRP in mg/L (IQR)	3 (2-12) (n=545)	2 (1-5) (n=936)	3 (1-10) (n=1,055)
Hospital referrals	219 (4.0%)	187 (3.7%)	357 (5.7%)

BMI = Body Mass Index; CRP = C-reactive protein; CVD = cardiovascular disease; IQR =

interquartile range; TIA = transient ischemic attack; VTE=venous thromboembolism.

565 **Table 3** - Model development and internal validation using logistic regression

	Full model		Simple model	
Predictor	**Regression coefficient**	**95% confidence interval**	**Regression coefficient**	**95% confidence interval**
Intercept	-4.110	-6.338; -2.149	-3.988	-6.228; -2.038
Age	0.026	-0.034; 0.092	0.022	-0.037; 0.088
Age'	0.045	-0.170; 0.251	0.085	-0.127; 0.289
Age''	-0.172	-0.719; 0.389	-0.268	-0.809; 0.286
Female sex	0.324	-2.358; 3.105	0.214	-2.450; 2.974
Interaction sex and age	-0.019	-0.102; 0.062	-0.016	-0.099; 0.065
Interaction sex and age'	0.032	-0.247; 0.314	0.015	-0.262; 0.296
Interaction sex and age''	-0.056	-0.805; 0.686	-0.014	-0.758; 0.726
CVD 1	0.564	0.280; 0.844		
CVD ≥2	0.636	0.282; 0.983		
Apparent c-statistic	0.693	0.665; 0.721	0.681	0.653; 0.710
R^2cs	0.030	0.021; 0.039	0.026	0.018; 0.035
Internal validation c-statistic	0.688	0.660; 0.716	0.680	0.652; 0.708
Internal validation R^2	0.070		0.061	
Internal validation slope	0.965		0.957	

Regression coefficients with 95% confidence intervals, c-statistic, and internal validation

performance measures of the full model with predictors age, sex and number of

cardiovascular diseases and the simple model with only age, sex and the interaction

between age and sex as predictors. Age was divided into three subgroups (shown as age,

age' and age'') using restricted cubic spline function to account for non-linearity. CVD =

cardiovascular disease; R^2cs = R^2 Cox-Snell.

577   **Table 4** – Comparing the full and the simple model

Δ**AUC**	0.012
Δ**R^{2}cs**	0.004
Likelihood ratio, χ^2	19.531, df=2, p = 5.740e-05

579   ΔAUC, ΔR^{2}cs and Deviance (likelihood ratio) in comparing model with and model without

580   number of cardiovascular diseases (0, 1 or ≥2) modelled. ΔAUC is calculated by subtracting

581   model 2 unadjusted c-statistic from model 1 unadjusted c-statistic. ΔR^{2}cs is calculated by

582   subtracting model 2 R^{2}cs from model 1 R^{2}cs.

584   **Table 5** - performance temporal validation

	Total validation population (n=16,693)	Total patients JGPN (n=5,420)	Total patients AHA AMC (n=6,284)	Total patients ANH Vumc (n=4,989)
Area under the curve	0.747 (CI 0.729; 0.764)	0.780 (CI 0.750; 0.810)	0.718 (CI 0.691; 0.745)	0.751 (CI 0.714; 0.789)
Calibration slope	1.358 (CI 1.246; 1.471)	1.510 (CI 1.306;1.718)	1.218 (CI 1.050; 1.387)	1.415 (1.190; 1.645)
Calibration intercept	-0.333 (CI -0.408; -0.261)	-0.396 (CI -0.535; -0.262)	-0.143 (CI -0.253; -0.036)	-0.561 (CI -0.712; -0.416)
Calibration in the large: Mean observed risk Mean predicted risk	0.046 0.062	0.040 0.058	0.057 0.065	0.038 0.063
R^{2}cs	0.035 (CI 0.029;	0.040 (CI 0.030;	0.033 (CI 0.024;	0.032 (CI 0.021;

		0.041)	0.050)	0.042)	0.042)
Brier score		0.042 (CI 0.039; 0.045)	0.037 (CI 0.033; 0.041)	0.052 (CI 0.047; 0.056)	0.035 (CI 0.031; 0.040)

586   Temporal validation performance measures with 95% confidence intervals (CI). $R^2cs = R^2$

587   Cox-Snell.

590   **Figure 1** – Predicted risk of hospitalisation

600   Plotted risk of hospital referral in (clinically suspected and confirmed) COVID-19 primary care

601   patients at a certain age and stratified by sex and by the number of cardiovascular diseases

602   and/or diabetes (i.e. 0, 1 or ≥2). Confidence intervals are shown in grey.