# A novel deterministic epidemic model considering mass vaccination and lockdown against Covid-19 spread in Israel: Numerical study

**Motoaki Utamura[1], Makoto Koizumi[2], Seiichi Kirikami[3]**

[1]    PhD, BSc, Former Professor Research Laboratory for Nuclear Reactors, Tokyo Institute of Technology

[2]    PhD, MSc, BSc, Former Researcher of Hitachi Research Laboratory, Hitachi Ltd.

[3]    BSc, Former Engineer of Hitachi Works, Hitachi Ltd.

2-5-2-612 Kamiyoga Setagaya-ku Tokyo 158-0098 Japan

## ABSTRACT

Why public health intervention by Israel government against COVID-19 spread has been successful while the most countries in the world is still coping with it? To give the answer, a simple numerical epidemic model is prepared to simulate an entire trend of various infection related variables considering vaccination campaign and simultaneous lockdown. The model is an extension of the deterministic physical model ATLM previously published by the authors that aims to predict an entire trend of variables in a single epidemic. The time series data of both vaccine dose ratio and lockdown period are employed in the model. Predictions have been compared with observed data in terms of daily new cases, isolated people, infectious at large and effective reproductive number and the model is verified. Moreover, parameter survey calculations for several scenarios have clarified a synergy effect of vaccination and lockdown have existed. In particular, it is suggested the key element of Israel success lies in a high dose rate of vaccination that avoids the onset of the rebound of daily new cases on the rescission of the lockdown.

_**Key words**_: Covid-19, Israel, vaccination, ATLM, epidemic model, discrete delay differential equation

## INTRODUCTION

The entire world is still in a crisis of COVID-19 pandemic. The introduction of a lockdown cannot prevent completely prevent the development of an epidemic. Mass vaccination of the population is the most radical method of limiting the growth of the epidemic. Israel government authority has started vaccination operation against SARS-CoV-2 virus on December 19th 2020 earlier than most other country. Cumulative doses exceeded 100% toward the end of March 2021, which was fastest in the world. In parallel the lockdown was practiced. As a result daily new cases drastically decreased from a peak of 10 thousand to 5 hundred at the end of March 2021. This successful experience should be shared in other countries. To do so it is essential to have a prediction model to count the number of people infected given the data of

vaccination dose rate and the period of lockdown. Few deterministic models based on standard SIR model [1] have been reported[2] because it is usually applied to an initial stage of epidemic when the number of susceptible individuals is close to collective population. Conversely full simulation of the entire epidemic is needed to take efficacy of vaccine into account. To this end Agent Based Model (ABM) have been recently investigated. The ABM [3][4][5][6][7] is a kind of stochastic approach and differs from mathematical models such as SIR and SEIR in that it can set up a virtual society consisting of a large number of autonomous decision-making and action actors (Agent) with various attributes and can analyze complex social events involving interactions between actors. Based on the rules we set on a computer, agents act spontaneously while interacting in a given environment. There are reports that simulated the comparative and joint impact of COVID-19 vaccine efficacy and coverage together with non-pharmaceutical interventions(NPI) by taking population of US cities as an example[6],[7]. They suggested non-pharmaceutical interventions such as masking, social distancing etc. are still requisite to reduce infections against SARS-CoV-2 even if higher vaccine coverage and efficacy are attained. In this method, detailed setting is possible, but it needs many empirical constants because of its stochastic approach and the calculation may take time depending on the scale. Moreover comparison was made only in terms of cumulative basis. Prediction need to be examined with factors such as inpatients, daily new cases etc. by available observations

In view of these difficulties in the past literature, a deterministic approach different from SIR was attempted in the present study. Apparent Time Lag Model (ATLM)[8] was extended to include observed vaccine dose rate and NPI in Israel and the validity of the present model was demonstrated using publicly available data. This model is featured in small number of empirical constants most of which can be measured.

# METHODS

## Data

Infectious data of COVID-19 in Israel was referred to worldometers [9] and statistics of vaccination and reproductive number of COVID-19 were cited by Our World in Data.[10]. Israel's vaccination started on December 19 2020 and total dose rate per habitant exceeded 100% toward the end of March 2021. Lockdown was issued from December 27 2020 till February 5 2021[11][12]. Data for time lags H(delay in PCR reporting date from infection), V(period of isolation), and T(period from infection till isolation) were set 14days[13], 12days[14] and 14days respectively.
We adopted SARS-CoV-2 infection data in Israel from November 8 till March 31 for analysis. Various data such as daily new cases, cumulative infections, currently isolated, currently infected, reproductive number and were used to examine the numerical model. We assumed all the vaccine dosed is Pfizer-BioNTech vaccine(BNT162b2 mRNA COVID-19 vaccine). Its effectiveness in real world was assumed 80% for single dose recipient at ten days or later after injection and similarly 95% for double dose recipient in consideration of reports [15][16][17][18][19] [20] .

## ATLM with vaccine model

Coronavirus SARS-CoV-2 causing COVID-19 pandemic includes several time delay events in the

spread of infection as follows:

(1) Lose infectious capability and extinguish within a month if no second infection occurs.

(2) Incubation time is about 5 to7days after infected in which last 2 or 3 days are infectious period.

(3) Infected individuals are detected and recognized as PCR positives in delay of about two weeks.

(4) PCR positives are mandatorily isolated and released after certain time according to administrative guidance.

(5) Effectiveness of mass vaccination is revealed in about ten days.

Therefore the mathematics of SARS-CoV-2 epidemic can be understood as demography of progressive events with time delay. Apparent Time Lag Model (ATLM) establishes a single discrete delay differential equation(DDE) in natural manner as a governing equation. It is different from conventional deterministic approach such as SIR model where delay is implicitly included in transmission rate whose reciprocal implies 1st order delay. ATLM provides a full simulation of a whole wave of the epidemic while SIR model expresses an entire solution as a combination of plural piecewise solutions in a small period. To do so ATLM introduces virtual collective population(M) instead of actual one(N) of the area in concern. M is not known a priori and to be determined by parameter fitting. M is much smaller than N, which matches the observation that action range of ordinary person is limited and thus consecutive infections may likely happen in the same area. As we can regard M as susceptible at time t=0, we assume that mass vaccinations play the role to reduce M. We place following assumption in the analysis: Every infected person onsets being PCR positive and eventually isolated T days after infection. Then it is possible to count infectious at large at time t as x(t)-x(t-T) where x(t) is cumulative infected persons until t and infected individuals before t-T are all isolated. The portion of susceptible may be expressed by 1-x(t)/M(t). We assume the rate of new infection dx/dt is proportional to the product of infectious at large and the portion of susceptible. Hence, governing equation of ATLM becomes

$$\frac{dx(t)}{dt} = \alpha(t)\{x(t) - x(t-T)u(t-T)\}(1 - \frac{x(t)}{M(t)}) \quad , \qquad (1)$$

$$M(t) = M0\left(1 - \frac{\mu(t)}{100}\right), \alpha(t) = \alpha0(1 - \rho(t)) \quad , \qquad (2)$$

where, x(t): cumulative infected individuals, dx(t)/dt: rate of new infection, α0: transmission rate(1/day), M0: virtual collective population, μ(t): effectiveness of vaccination(%), ρ(t): mobility of people, u(t): step function. As the influence of vaccination corresponds to the reduction of susceptible, we expressed net collective population of a group M(t) with effectiveness of vaccination. Also, the effect of lockdown is the decrease in people's mobility and reduction of transmission rate as in equation (2). Given initial value x(0), Eqs. (1) and (2) were solved numerically using 4th order Runge Kutta Gill method with a time step of half day and obtained x(t).

Then the rest of infection related variables were calculated by use of x(t) as follows:

Cumulative positive number Y(t) is reported in delay of H days then

$$Y(t) = x(t - H) \qquad .\qquad (3)$$

This also means the number of cumulative isolated patients. H: period from infection till PCR test result report reaches public health authority. H was assumed to be 14days[13].

Similarly daily new cases reported by public health authority, are actually daily new positive cases dY(t)/dt expressed by

$$dY(t)/dt = dx(t - H)/dt \qquad .\qquad (4)$$

Infectious at large W(t) is the sum of new infection in the period [t-T, t]

$$W(t) = x(t) - x(t - T) \qquad ,\qquad (5)$$

where T is a period during which infected keep infectious at large(or until isolated). T was assumed to be 14days. It is because T=L+F+G, where L:latent period[21], F:period of getting tired 5~7days, G: go to hospital and quarantined 2~3days, which yields 12(=5+5+2) <T<15(=5+7+3) whose average gives T=14.

The number of isolated people P(t) consisting of hospitalizations, self-isolated at hotels and home and people under adjusting can be counted by

$$P(t) = x(t - T) - x(t - S) \qquad ,\qquad (6)$$

where x(t-S) is the number of recovered or deaths. Denoting isolation period as V, S can be written as the sum of T and V i.e. S=T+V. Israel's Health Ministry decided to shortened the mandatory coronavirus isolation period from 14days to 12days, provided the patient received two negative tests from November 12 2020[14]. Then, V=12 in Israel. Therefore, S=T+V=14+12=26.

ATLM defines effective reproductive number Rt(t) as

$$Rt(t) = \alpha(t)T(1 - \frac{x(t)}{M(t)}) \qquad (7)$$

In subsequent sections, we will compare predictions by Eqs. (3)-(7) with available data from Worldometer and Our World in Data and examine the present analysis method.

In summary, the values of delay parameters adopted in the prediction are H=T=14, V=12, S=26 which will be used throughout this paper.

**Modeling of vaccination in Israel**

Let the number of people vaccinated per 100 habitants A, those vaccinated twice, B and cumulative doses per 100 habitants C, then those vaccinated only once becomes A-B and A+B=C holds at any instance. When vaccination was completed i.e. everyone is two times recipient, then C would be 200%. Trends of data for A, B and C in Israel[10] are shown in Fig. 1. The target value of B is said to be 5.5million among 9 million citizens which has been almost accomplished before the end of March.

Let efficacy of vaccine for a single dose recipient $\beta$ and that of double dose one $\gamma$, then average efficacy $\mu$ (t) of a group consisting of both A and B is written for a single type vaccine

$$\mu \text{ (t)} = \beta \text{ (A(t)} - \text{B (t))} + \gamma \text{ B(t)} \tag{8}$$

Virus type of 3rd wave epidemic in Israel is mainly Britain type variant[15]. Efficacy of BNT162b2 mRNA COVID-19 vaccine (Pfizer-BioNTech COVID-19 vaccine) in a nation-wide mass vaccination campaign in Israel is reported in peer reviewed articles [18],[19],[20]. Referring to these we adopted $\beta$ =0.8 and $\gamma$ =0.95 and assumed vaccine activated after ten days of vaccination for both single and double injection recipients.

Data plot of μ(t) (Eq. (8)) and its polynomial approximation Eq.(9) is presented in Fig.2.

$$\mu \text{ (t)} = -0.000034\, t^3 + 0.007073 t^2 + 0.297875\, \text{t} - 33.064707 \tag{9}$$

Since the effectiveness of vaccine becomes activated in ξ days after injection, μ(t) in Eq. (9) was replaced by μ(t-ξ). Referring to literatures [16] [17], we assumed ξ=10 in the computation.

**Modeling of lockdown**

Israel government executed a national lockdown as a necessary step to block the increase of morbidity [10] on December 27, 2020(63days) for two weeks with an option to extend it. Actually it was extended until February 5, 2021(103days) because it saw delay in turnaround from the health and economic crisis on January 31, 2021(98days)[12]. We modelled lockdown effect by a sudden decrease in transmission rate as shown in Fig. 3. Its duration was from 63days through 103days.

$\rho_{min}$ was set to 0.72 as a result of parameter fitting. Other parameter values determined after fitting were x(0)=650, α0=0.117 and M0=1,500,000 respectively. They were used throughout subsequent calculations.

# RESULS

**Daily new cases and total cases**

Daily new cases were taken from worldometer data source[9] first for comparison with prediction by ATLM equipped with vaccine and lockdown models. Examined period of time was from November 8 through March 31 which covers a whole span of third wave of COVID-19 epidemic in Israel in which mass vaccination and a nation-wide lockdown are involved. We regarded 'daily new cases' in worldometer as daily new positives that are observed behind new infection by H days(14 days). Computation started ahead of November 8 by H days i.e. October 25 that is the origin of time.

Figure 4 compares prediction with observed data. Both are expressed as 7 day average basis to avoid fluctuation caused by the difference in the number of PCR tests in a week. The prediction is 7 day

average of dY/dt (Eq. (4)). There are important findings in the following:

- Effects of lockdown start and release are seen apparent at 77days and 117days in data respectively while actual dates the interventions were executed 63days and 103days respectively. The former reduces rate of infection resulting in the appearance of a peak, the latter producing plateau as a result of antagonist between rebound of lockdown and efficacy of vaccine. This fact implies the assumption H=14(infection becomes apparent in delay of 14 days) is proper. Both observations in data were simulated by the prediction within 10% precision in the aspect of both time and magnitude.

- In early phase of the epidemic the prediction is seen to underestimate the observed data. This fact can be interpreted as the observed data contain tailing effect of the 2nd wave if we note November 8(14days) is the minimum cases between 2nd and 3rd waves. The plateau appearing around 90 to 100 days were not reproduced in computation. As a whole the simulation of the entire trend of daily new cases was satisfactory.

Figure 5 presents cumulative positives. Excellent agreement was attained between prediction and observed data. Underestimation in the early phase and toward the end comes from the same reason mentioned above.

**The number of isolated patient**

The number of isolated patients is shown in Fig. 6. Although calculation by Eq.(6)　overestimates actual data by 15% at a peak around 85days, overall trend is well simulated. In particular, the trend after 130days was excellently simulated. This proves the assumption T=14 in the model is correct.

**Reproductive number**

Figure 7 compares Eq.(7) with data[11] with respect to reproductive number(Rt). Apart from an early phase, predicted Rt(t) coincides with Rt(t) reduced from observed data. Discrepancy in early phase could be accounted for by the existence of the tail of 2nd wave of epidemic.

**The number of infectious at large**

It is important to note infectious at large is an important index to tell when epidemic will end, much more accurate than daily new cases. It is because the time to give W(t)=0 indicates the end of epidemic. We will show the infectious at large may be obtained through data processing as follows: Substitution of daily new cases Y(t) to RHS of Eq.(5) would yield W(t-T). W(t) would be obtained by shifting function W(t-T) leftward by T along time axis. To confirm the validity of newly obtained W(t), agreement with numerical prediction which has been already verified through arguments described above. As shown in Fig. 8, excellent agreement was obtained between two, which suggests the correctness of the data processing mentioned above. W(t) peak to dY(t)/dt ratio was found to be about 13times.

**Estimation of separate effect of vaccination and lockdown**

Based on the confirmed epidemic model, parameter survey calculations were conducted for following four cases.

(1) Without vaccination, without lockdown: case C

$\mu = \rho = 0$

(2) With vaccination, without Lockdown: case D

$\mu \neq 0, \rho = 0$

(3) Without vaccination, with lockdown: case B

$\mu = 0, \rho \neq 0$

(4) With vaccination, with lockdown: case A(real intervention)

$\mu \neq 0, \rho \neq 0$

Results are presented in Fig. 9. Without lockdown, peaking value would grow 2~2.5 times larger than real one. Without vaccination, the epidemic would decelerate in a later stage and even worse on earlier release of lockdown. Without lockdown, blue line(A) would have shifted to red(D). If vaccination dose rate had been slower, blue(A) would have approached grey(B). In general, lockdown decreases rate of transmission, while vaccination accelerate to converge epidemic. Next, separate effect of either vaccination or lockdown was investigated by taking ratios D/C and B/C. Figure 10 presents both ratios together with combined effect. Here, the ratio means relative infection rate to C. The smaller the ratio, the lower the infection rate. As vaccinations proceed, the ratio D/C monotonously decreases. To the contrary, the ratio B/C of lockdown has a minimum due to rebound on the release of lockdown. It is seen until 120days the effect of lockdown exceeds that of vaccination. After 120days the effect of vaccination is superior to lockdown. Combined effect A/C is presented in a broken line, which shows the suppression of rebound to some extent owing to vaccination. There exists a period (90days~125days) when the green broken line has the smallest ratio among three. It is seen synergy effect was created by interaction between vaccination and lockdown. It is inferred the faster the dose rate, the smaller the magnitude of the rebound, which would contribute to earlier ending of the epidemic.

# DISCUSSION

**Primary findings**

ATLM equipped with newly developed vaccination and lockdown model simulated a whole trend of various infection variables for COVID-19 3rd wave epidemic in Israel within 15% error. In particular, the number of isolated PCR positives and infectious at large were separately evaluated with high resolution. Moreover the equation of effective reproductive number Eq.(7) derived from ATLM showed a good agreement with existing data trend[10] that reflects the influence of mass vaccination

and lockdown. These facts affirm the present vaccination model whose concept is to reduce virtual collective population by acquired effectiveness of the vaccination. To the contrary, most previous works using ABM disclosed so far were limited to total and daily new cases[4][5][6].

Separate effect of vaccination and lockdown was explored. Although lockdown is most effective to suppress infection rate in ascending phase of epidemic, it is inevitable to induce rebound on its release. Vaccination is found to be a strong tool to suppress it. To make this effective, it was inferred from parameter survey calculations that vaccination dose rate should be fast enough to achieve at least 40% effectiveness on the rescission of lockdown.

Additionally data processing method to extract infectious at large from total cases was clarified. The infectious at large $W(t)$ is a decisive index to estimate the end of the epidemic when $W(t)$ becomes zero which occurs behind the time when daily new cases become zero. It yields a conservative measure to plan public health prevention and control of epidemic.

In the present study ATLM ignored asymptomatic patient. Nonetheless prediction agreed with observation, especially for the number of isolated individuals and infectious at large. This fact implies most of symptomatic patients, if any, were non-infectious in Israel.

**Limitations and future work**

Limitations of the present method follow. First, the present vaccination model is established based on efficacy data of BNT162b2 mRNA COVID-19 vaccine (Pfizer-BioNTech COVID-19 vaccine). It should be extended to accommodate other COVID-19 vaccines. Also, it is assumed activation time needed after injection $\xi$ =10. Since another observations exist, wide parameter survey is requisite to get a conservative solution for policy making.

In summary, this study is the first work of a full simulation of 3rd wave of epidemic in Israel accompanied by mass vaccination campaign and simultaneous lockdown. Existing data and policy making elsewhere would be aided by inclusion of the present model. On its occasion potential users of the present model are advised to reassess the values of the delay constants T,H,V, $\xi$ in the model if the application to other countries is attempted because they are influenced by the difference in medical regulations or practice.

# CONCLUSION

A simple numerical epidemic model is proposed to simulate an entire trend of various infection related variables considering vaccination campaign and simultaneous lockdown. The model is an extension of the deterministic physical model ATLM previously published by the authors which is described by a single discrete delay differential equation with explicit inclusion of time lag of isolation of infectious. It gives full simulation of various infection variables in the entire span from the initial infection through endpoint with small number of fitting parameters. The time series data of

both vaccine dose ratio and lockdown period were employed in the model. The model was examined within 15% of errors compared with Israel's COVID-19 various epidemic trend data such as daily new cases, the number of isolated people and infectious at large, and effective reproductive number. Moreover, parameter survey calculations for several scenarios have clarified there exists a synergy effect between vaccination and lockdown. In particular, it is suggested the key element of Israel's success in drastic reduction of infection rate should lie in a high dose rate in mass vaccination campaign that avoids the onset of the rebound of daily new cases on the rescission of the lockdown.

## DATA AVAILABILITY

We used time-series data of COVID-19 for November 8, 2020 through March 31, 2021 in Israel [9],[10].

## AUTHOR CONTRIBUTIONS

M. Utamura developed and verified the epidemiological model. S. Kirikami developed vaccination model. M. Koizumi examined calculation results by another solution algorithm. All authors have read and agreed to the published version of manuscript.

## FUNDING

This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

## CONFLICT OF INTEREST

The authors declare that they have no conflict of interest related to this report or the study it describes.

## ACKNOWLEDGMENTS

MU is the former professor of Research Laboratory for Nuclear Reactors of Tokyo Institute of Technology. MK is the former researcher of Hitachi Research Laboratory, Hitachi Ltd. SK is the former engineer of Hitachi Works, Hitachi Ltd. Authors are grateful to the editor for kind guidance.

## ABBREVIATIONS

ABM: Agent Based Model

ATLM: Apparent Time Lag Model (present epidemic model)

DDE: Delay Differential Equation

NPI: Non-Pharmaceutical Intervention

PCR: polymerase chain reaction

SIR: Susceptible, Infectious and Removed model

# REFERENCES

[1] Kermack WO, Mckendrick AG. A contribution to the mathematical theory of epidemics. *Proc. The Royal Society A* (1927) 115(772) pp.700-721.

[2]M.D. Patel et al., The Joint Impact of COVID-19 Vaccination and Non-Pharmaceutical Interventions on Infections, Hospitalizations, and Mortality: An Agent-Based Simulation | medRxiv **doi:** https://doi.org/10.1101/2020.12.30.20248888.

[3] Erik Cuevas,(2020) An agent-based model to evaluate the COVID-19 transmission risks in facilities, comput Biol Med. 2020 Jun; 121: 103827.

[4]S. Pageaud et al.(2021), Adapting French COVID-19 vaccination campaign duration to variant dissemination | medRxiv **doi:** https://doi.org/10.1101/2021.03.17.21253739.

[5]E. Chang, K.A. Moselle(2021) Agent-Based Simulation of Covid-19 Vaccination Policies in CovidSIMVL | medRxiv **doi:** https://doi.org/10.1101/2021.01.21.21250237.

[6]O. Alagoz et al.(2021), The Impact of Vaccination to Control COVID-19 Burden in the United States: A Simulation Modeling Approach | medRxiv **doi:** https://doi.org/10.1101/2021.03.22.21254131.

[7]A.Truszkowska et al.(2021), High-Resolution Agent-Based Modeling of COVID-19 Spreading in a Small Town - Truszkowska - 2021 - Advanced Theory and Simulations - Wiley Online Library.

[8] Utamura M, Koizumi M, Kirikami S, An Epidemiological Model Considering Isolation to Predict COVID-19 Trends in Tokyo, Japan: Numerical Analysis. JMIR Public Health Surveill. 2020 Dec 16;6(4):e23624. doi: 10.2196/23624. PMID: 33259325; PMCID: PMC7746226.

[9] Israel COVID: 836,926 Cases and 6,334 Deaths - Worldometer (worldometers.info).

[10] Coronavirus (COVID-19) Vaccinations - Statistics and Research - Our World in Data.

[11]Israel government press release(23.12.2020), Joint Prime Minister's Office-Health Ministry Statement | Prime Minister's Office (www.gov.il).

[12]Egypt independent(February1, 2021), Israel extends lockdown, sees delay in COVID-19 turnaround - Egypt Independent .

[13] Kazuo Maki(2020), A delayed SEIQR epidemic model of COVID-19 in Tokyo area, MedRxiv, doi:https://doi.org/10.1011/2020.8.18.20177709.

[14]The Jerusalem post(November 8, 2020), COVID-19 isolation period to be shortened to 12 days starting Thursday - The Jerusalem Post (jpost.com).

[15]CDC Newsroom(March 29, 2021), CDC Real-World Study Confirms Protective Benefits of mRNA COVID-19 Vaccines | CDC Online Newsroom | CDC.

[16]A.J. Tande et al., Impact of the COVID-19 Vaccine on Asymptomatic Infection Among Patients

Undergoing Pre-Procedural COVID-19 Molecular Screening | Clinical Infectious Diseases | Oxford Academic (oup.com)  https://doi.org/10.1093/cid/ciab229, 10 March 2021.

[17]Pfizer press release(March 11, 2021), Pfizer Inc. - Real-World Evidence Confirms High Effectiveness of Pfizer-BioNTech COVID-19 Vaccine and Profound Public Health Impact of Vaccination One Year After Pandemic Declared.

[18]S. Amit et al.,( Feb. 18, 2021) Early rate reductions of SARS-CoV-2 infection and COVID-19 in BNT162b2 vaccine recipients - The Lancet

DOI:https://doi.org/10.1016/S0140-6736(21)00408-7.

[19]N. Dagan et al., BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting | NEJM N Engl J Med 202 1; 384:1412-1423

DOI: 10.1056/NEJMoa2101765 April 15, 2021.

[20]F.P.Polack et al., Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine | NEJM N Engl J Med 2020; 383:2603-2615

DOI: 10.1056/NEJMoa2034577, December 31, 2020.

[21] Lauer, S.A. et al., The incubation period of coronavirus disease 2019(COVID-19) from publicly reported confirmed cases: estimation and application, Ann Intern Med. 172(9), 577-582(2020).

Fig. 1 The number of people of A, B per 100 habitants and total vaccine dose per 100 habitants C in Israel. Abscissa represents time(days) and its origin is October 25 2020 where calculation starts. A, B and C are observed data in which A and B represent the number of people vaccinated per 100 habitants and those vaccinated twice in Israel. C is total doses per 100 habitants. Axis

shows the number of people or doses per 100 habitants. Vaccination started on December 19 (55days) onward. Total doses exceeded 110 at the end of March 2021(157days).

Fig.2 Presumed effectiveness of mass vaccination( in case of no delay). Inferred from observed data by Eq.(8) for collective population composed of recipients mixed up with single and double dose vaccinated. Axis represents expected number of people per 100 habitants and abscissa time(days). Vaccination began on December 19(55days), 2020.

Fig. 3 Modeling of lockdown. Axis represents 1-ρ(t) in Eq. (2). Lockdown was operated during 63days and 103days. Its effect is anticipated to become apparent on 77days and 117days because of delay in reporting.

Fig. 4 Daily new cases by observation and prediction. Red line :data, blue :prediction, yellow:

effectiveness of vaccination 1-μ(t)/100 referring to right axis. Left axis: the number of new infected, Abscissa: time(days) origin of time coincides with October 25, 2020, 14 days ahead of November 8, the start of 3$^{rd}$ wave of epidemic investigated in this study. Both data are displayed as 7 day average. The 3$^{rd}$ wave of COVID-19 epidemic in Israel was regarded from November 8, 2020 to March 31, 2021. Nation-wide mass vaccination started on December19, 2020(55days). Its effectiveness was revealed behind ten days(65days). Lockdown was operated between December 27, 2020(63days) and February 5, 2021(103days). Calculation shows dY(t)/dt which was carried out using Eq.(4).

Fig. 5 Trends of cumulative positives. Equation (3) was used to calculate cumulative positives, Y(t)

Fig. 6 Trend of currently isolated patients. Orange dots are data cited from worldometer whose notation is 'currently infected'[9]. Blue line is prediction by ATLM. Coincidence in timing implies the assumption T=14 is proper. Time to give peak is behind that of daily new cases(Fig. 4) by a week.

Fig. 7 Effective reproductive number. Observed data was cited from our world in data[10] and Eq.(7) was used for calculation.

Fig. 8 Infectious at large. Red line is processed from data and blue line from prediction.

Fig. 9 Predicted daily new cases for various combinations of public health interventions

with or without vaccination and lockdown. Axis: daily new cases, Abscissa: time（days）
Case A: With vaccination, with lockdown, Case B: Without vaccination, with lockdown,
Case C: Without vaccination, without lockdown, Case D: With vaccination, without
Lockdown.

Fig. 10 Separate effect of vaccination and lockdown on reduction of infection. Ratio B/C(blue
line) denotes lockdown effect without vaccination. Ratio D/C(red line) represents vaccination
effect without lockdown. Ratio A/C(green line) shows real combined effect.