Mask mandate and use efficacy for COVID-19 containment in US States

Damian D. Guerra[1,*] and Daniel J. Guerra[2]

1.   Department of Biology, University of Louisville, Louisville, Kentucky, United States of America

2.   Authentic Biochemistry, VerEvMed, Clarkston, Washington, United States of America

* Corresponding author (damian.guerra@louisville.edu)

Abstract

Background: COVID-19 pandemic mitigation requires evidence-based strategies. Because COVID-19 can spread

via respired droplets, most US states mandated mask use in public settings. Randomized control trials have not

clearly demonstrated mask efficacy against respiratory viruses, and observational studies conflict on whether mask

use predicts lower infection rates. We hypothesized that statewide mask mandates and mask use were associated

with lower COVID-19 case growth rates in the United States.

Methods: We calculated total COVID-19 case growth and mask use for the continental United States with data from

the Centers for Disease Control and Prevention and Institute for Health Metrics and Evaluation. We estimated post-

mask mandate case growth in non-mandate states using median issuance dates of neighboring states with mandates.

Results: Earlier mask mandates were not associated with lower total cases or lower maximum growth rates. Earlier

mandates were weakly associated with lower minimum COVID-19 growth rates. Mask use predicted lower

minimum but not lower maximum growth rates. Growth rates and total growth were comparable between US states

in the first and last mask use quintiles during the Fall-Winter wave. These observations persisted for both natural

logarithmic and fold growth models and when adjusting for differences in US state population density.

Conclusions: We did not observe association between mask mandates or use and reduced COVID-19 spread in US

states. COVID-19 mitigation requires further research and use of existing efficacious strategies, most notably

vaccination.

Keywords: COVID-19, SARS-CoV-2, face covering, medical mask, mask mandate, nonpharmaceutical intervention

Introduction

The COVID-19 pandemic has increased mortality and induced socioeconomic upheaval worldwide (1). Evidence-based containment strategies are warranted, given that age, obesity, cardiovascular disease, and diabetes are common comorbidities associated with severe COVID-19 symptoms (e.g., pneumonia, blood clots, cytokine storm), hospitalization, and death (2, 3). Respired droplets and aerosols containing SARS-CoV-2 are intuitive modes of community transmission (4). To reduce viral spread, governments have issued mandates to wear medical masks or cloth face coverings in public settings. From April to December 2020, 40 States of the United States issued mask mandates. Mask mandates have limited precedent, making efficacy unclear. Our first objective was to evaluate the efficacy of mask mandates in attenuating COVID-19 growth in US states.

Prior studies have conflicted on whether masks reduce COVID-19 spread. For USS Theodore Roosevelt crew, mask use was lower among COVID-19 cases compared with non-infected (56% vs. 81%) (5). There were no infections for 48% of universally masked patrons exposed to COVID-19 positive hair stylists (6), but PCR tests were not obtained for the other 52% of patrons (6), and first wave COVID-19 hospitalizations were no higher in public schools (high density with minimal masking) than elsewhere in Sweden (7). A randomized controlled trial (RCT) of Danish volunteers found no protective benefit of medical masks against COVID-19 infection (8). In RCTs before COVID-19, viral infections were not lower in Vietnamese clinicians who wore cloth or medical masks than in the control arm (9), and N-95 respirators (but not medical masks) protected Beijing clinicians from bacterial and viral diseases compared to no masks (10). Mask compliance in RCTs is not always clear (11). Mask use was 10% and 33% for Beijing households with and without intrahousehold COVID-19 case growth, respectively (12). This suggests greater mask use may reduce COVID-19 spread. Our second objective was to assess if mask use predicts lower COVID-19 case growth.

We assessed if mask mandates and compliance in US States predict statewide COVID-19 growth during

the second and third infection waves (1 June 2020-1 March 2021). Controlling for infection wave timing

with logarithmic and linear relative growth models, we found limited association between COVID-19

case growth and mask mandates or mask use before 1 October 2020, and no association during the

subsequent and largest third wave. These findings do not support the hypothesis that statewide mandates

and enhanced mask use slow COVID-19 spread. Pharmaceutical interventions (including recently

available COVID-19 vaccines) provide alternative, evidence-based strategies to minimize COVID-19

related morbidity and mortality.

Materials and methods

Data Sources and Terms

We obtained total (confirmed and probable) COVID-19 cases up to 6 March 2021 for the 49 continental

US states, normalized per 100,000 residents, from the Centers for Disease Control and Prevention (CDC)

(13). To reduce reporting lag effects, we used 7-day simple moving means (e.g., the 7-day simple moving

mean of cases on 31 March is the mean of daily cases between 28 March and 3 April). Hawaii was

excluded because COVID-19 growth patterns deviated from those of continental US states. Confirmed

and probable cases are defined by the Council of State and Territorial Epidemiologists. Confirmed cases

require PCR amplification of SARS-CoV-2 RNA from patient specimens. Probable cases require one of

the following: clinical and epidemiologic evidence, clinical or epidemiologic evidence supported by

SARS-CoV-2 antigen detection in respiratory specimens, or vital records listing COVID-19 as

contributing to death. Total PCR tests for each state were obtained from Worldometers on 25 May 2021

(14).

Mask mandates are statewide emergency executive public health orders requiring nose and mouth

coverings in public settings in more than 50% of counties within a state (15, 16). We assigned US states

to one of five quintiles based on when mandates went into effect (effective dates): 18 April-16 May 2020

(Q1), 29 May-3 July 2020 (Q2), 8 July-27 July 2020 (Q3), 1 Aug-9 Dec 2020 (Q4), or no statewide

mandate as of 6 March 2021 (Q5). Effective dates were obtained from US state executive and health

departments and press releases (available upon request).

We assessed mask use with the University of Washington Institute for Health Metrics and Evaluation

(IHME) COVID-19 model site (17), which estimates daily compliance from Premise, the Facebook

Global Symptom Survey (University of Maryland), the Kaiser Family Foundation, and the YouGov

Behavior Tracker Survey. Mask use is the percentage of people who always wear masks in public

settings. We assigned US states to mask quintiles based on the mean percent mask use from 1 Jun-1 Oct

2020 (Summer) or from 1 Oct 2020-1 Mar 2021 (Fall-Winter).

To assess geographic differences, we assigned each US state to one of five regions: Northeast

(Connecticut, Delaware, Massachusetts, Maryland, Maine, New Hampshire, New Jersey, New York,

Pennsylvania, Rhode Island, and Vermont); Midwest (Illinois, Indiana, Iowa, Kentucky, Kansas,

Michigan, Minnesota, Missouri, Ohio, West Virginia, Wisconsin); Mountains-Plains (Colorado, Idaho,

Montana, Nebraska, New Mexico, North Dakota, Oklahoma, South Dakota, Utah, Wyoming); South

(Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina,

Tennessee, Texas, Virginia); and Pacific (Alaska, Arizona, California, Nevada, Oregon, Washington).

Growth Rate Calculation

COVID-19 growth has been modeled logarithmically (15, 18, 19) and linearly (19, 20). Therefore, we

calculated COVID-19 case growth for each US state by measuring percent natural logarithmic (Ln

Growth) and percent linear (Fold Growth) relative growth rates:

$$\text{Ln Growth:} = 100 * ln\frac{C_t}{C_{t-1}}$$

$$\text{Fold Growth:} = 100 * (-1 + \frac{C_t}{C_{t-20}})$$

Where C_t, C_{t-1}, and C_{t-20} are total normalized cases on a day, the prior day, and 20 days prior, respectively.

We determined adjusted population density by calculating the weighted mean of each state's urban (U)

and rural (R) population density using the following formulas:

$$\text{Urban Density (U)} = u1p1 + u2p2 + u3p3$$

$$\text{Mean Rural Density (R)} = \frac{State\ Population * (1-F)}{State\ Area - a1 + a2 + a3}$$

For the three most populous urban regions in each US state, we obtained urban population density (u;

people/square mile in an urban area), urban land area (a; size of urban area in square miles), and

population proportion (p; fraction of combined urban population) via 2010 US Census Bureau estimates

(21). For some states, two rather than three urban regions were used. The proportion of urban (F) and

rural (1-F) population of each state was similarly obtained (21). We thus calculated adjusted population

density of each state as:

$$\text{Adjusted Population Density (APD)} = UF + R(1 - F)$$

To assess association between population density and growth rates, we multiplied Fold Growth by the

inverse of normalized APD:

$$\text{Adj. Fold Growth} = \text{Fold Growth} * \left(\frac{Mean\ APD\ of\ all\ US\ States}{State\ APD}\right)$$

We defined minima and maxima (extrema) as the lowest and highest growth rates between the end of the

Summer wave and the height of the Fall-Winter wave. Ln Growth extrema comprised 20-day windows

when Ln Growth rates were lowest or highest:

$$\text{Ln Growth extremum} = \max/\min_{\substack{20\ Jul\ 2020 \\ \leq t \leq 1\ Mar\ 2021}} \frac{\sum_{t=1}^{t=20} Ln\ Growth\ on\ Day\ t}{20}$$

Fold Growth extrema similarly comprise the lowest and highest Fold Growth:

$$\text{Fold Growth extremum} = \max/\min_{\substack{20\ Jul\ 2020 \\ \leq t \leq 1\ Mar\ 2021}} Fold\ Growth\ (t\text{-}20,\ t)$$

For each state, surges are differences between maxima and minima (relative growth rate increase), masks

at extrema are the 20-day mean mask use at minima and maxima, and Δ Masks is the percent change in

mask use between maxima and minima.

To assess association between mandates and growth rates in the 48 contiguous states (excluding Alaska

and Hawaii), we determined Ln, Fold, and Adj. Fold Growth between 1 March 2021 (C_{301}) and the

mandate effective date (C_M) for US states in mandate quintiles 1-3:

$$\text{Post-Mandate Ln Growth} = 100 * \ln \frac{C_{301}}{C_M}$$

$$\text{Post-Mandate Fold Growth} = 100 * (-1 + \frac{C_{301}}{C_M})$$

For states in quintiles 4-5, *modeled effective dates* are medians of actual dates among bordering states of

mandate quintiles 1-3. For example, the modeled effective date of Tennessee (10 July) is the median of

effective dates of Arkansas (20 July), Alabama (16 July), Kentucky (10 July), North Carolina (26 June),

and Virginia (29 May).

For each state, Summer 2020 (1 June-1 Oct) and Fall-Winter 2020-21 (1 Oct-1 Mar) mask use is mean

mask use between these dates. Cases on 1 June or 1 Oct were the 20-day mean total normalized cases on

these two dates. We likewise defined Summer and Fall-Winter case growth using Ln and Fold Growth

formulas:

$$\text{Summer Ln Growth} = \ln \frac{C_{1001}}{C_{601}} \qquad \text{Summer Fold Growth} = 100 * (-1 + \frac{C_{1001}}{C_{601}})$$

$$\text{Fall-Winter Ln Growth} = \ln \frac{C_{301}}{C_{1001}} \qquad \text{Fall-Winter Fold Growth} = 100 * (-1 + \frac{C_{301}}{C_{1001}})$$

Where c_{601}, c_{1001}, and c_{301} are total normalized cases on 1 June 2020, 1 October 2020, and 1 March 2021,

respectively.

Statistics

We used Prism 9.2 (GraphPad; San Diego, CA) to construct figures and perform null hypothesis

significance tests, for which the significance threshold was $p < \alpha = 0.05$ (**Worksheet D in S1 Table**).

Error bars denote standard deviations, 95% confidence intervals, or interquartile ranges as indicated in

figure legends. We performed D'Agostino-Pearson tests to assess normality of residuals.

To evaluate mask mandate and use efficacy among categories (mandate effective date or mask use

quintiles), we performed ordinary one-way ANOVA with Tukey posttests. For non-normal data, we

performed Kruskal-Wallis with Dunn posttests. For two sample comparisons (e.g., **Fig. 3G, J**), we

conducted two-tailed t tests or Mann-Whitney tests for normal and non-normal data, respectively.

This decision tree conforms with recommended practices for datasets of N > 5 (22).

For interval variable associations, we performed ordinary least squares (OLS)-simple linear regression

with null hypotheses of zero slope. Infectious disease research has employed OLS previously (23, 24),

with linear and ln-linear models reported in recent COVID-19 studies (25, 26). For the Summer wave,

Northeast states were excluded because they deviated from other states with respect to total cases and

growth covariation. We used weighted least squares (WLS) for heteroscedastic data, as determined by the

GraphPad Prism Test for Homoscedasticity. Regardless of statistical significance, R^2 values denote

coefficients of determination for lines of best fit with unconstrained slopes.

Results

COVID-19 growth rates vary with time

With the aim of reducing COVID-19 case growth, 40 US states enacted mask mandates in 2020. We

wondered if mask mandate timing affected COVID-19 growth patterns. To identify patterns of COVID-

19 growth, we graphed natural logarithmic (Ln) Growth of COVID-19 in US states as a function of time

(**Worksheet A in S1 Table**). We observed six phases of COVID-19 growth up to 6 March 2021: first

wave (before May 2020), Spring minimum (May-June 2020), Summer wave maximum (June-August

2020), post-Summer minimum (August-October 2020), Fall-Winter wave maximum (October-January

2020), and third minimum (March 2021) (**Fig. 1A and S1 Fig**). Hawaii growth patterns deviated from

those of continental US states and was thus excluded from further analysis. Regardless of mask mandate

effective date quintile, Ln growth patterns were comparable for all continental US states, and there was no

association between normalized total cases and PCR tests (**S1 Fig**).

Earlier mask mandates are not consistently associated with COVID-19 growth rates in US states

A recent study reported time-enhanced negative association between mask mandates and Ln Growth of

COVID-19 (15), but simple Fold-Growth (an alternative COVID-19 metric (19, 20)) may be preferred for

post-exponential, linear pandemic spread. PCR testing for COVID-19 was limited before Summer 2020

(27). Thus, to determine if US states with earlier mask mandates exhibited less COVID-19 spread, we

examined both Ln Growth and Fold Growth at the post-Summer wave minimum and the Fall-Winter

wave maximum (**Fig 1A-B**)—periods of low and high transmission, respectively. We assigned US states

to one of five quintiles (MQ1-5), with MQ1 including states with the earliest mandates, MQ4 the latest

mandates, and MQ5 states without mandates. Ln minima (p=0.07; **Fig. 1C**) and Fold minima (p=0.047;

Fig. 1D) trended lower for earlier mandates. Fold minima was 3-fold higher in MQ4 than MQ1 after

adjusting for population density (p=0.04), but all other pairwise comparisons were not significant (**Fig.**

1E). This suggested that mask mandate duration was a weak predictor of lower minimum growth. Ln

maxima (p=0.23; **Fig. 1F**) and Fold maxima (p=0.19; **Fig. 1G**) did not differ among quintiles. Adjusting

for population density, Fold maxima were 2.8-fold higher in MQ4 than MQ2 (p=0.01), but MQ1, 3, and 5

were indistinguishable (**Fig. 1H**), suggesting mask mandate duration was not associated with lower

maximum growth. Likewise, surges (growth rate increases from minima to maxima) were MQ-

independent for Ln (p=0.08) and Fold (p=0.13) models, and only MQ2 and MQ4 exhibited significantly

different Fold surges with population density adjustment (p=0.03; **S2 Fig**). Most MQ4 states exhibited

lower initial and Summer 2020 infection waves than Q1-2 (**S1 Fig**), suggesting high MQ4 growth rates

could be an artifact of lower total cases. While there was strong positive association between earlier

mandates and Fall-Winter mask use (p<0.001; **Fig. 1I**), total cases on 1 March 2021 were MQ-

independent (p<0.07; **Fig. 1J**). Direct MQ1 vs. MQ5 comparison by t test uncovered a small (1.2-fold)

and non-significant (p=0.078) difference in total cases. Taken together, these findings suggest that US

state mask mandates were not associated with slower spread of COVID-19.

Early mask mandates do not predict lower post-mandate COVID-19 growth in contiguous US states

Most US states enacted mandates during infection waves, which confounds assessment of effectiveness.

To assess association between mandate effective date and post-mandate case growth, we compared

growth after actual MQ1-3 mandates with growth after modeled MQ4-5 mandates up to 1 March 2021.

For a MQ4-5 state, the modeled date was the effective date median of contiguous, common-border MQ1-

3 states (**Fig. 2A**). Post-mandate growth in total cases was MQ-independent by Ln (p=0.43), Fold (p=41),

and population density-adjusted Fold (p=0.15) models (**Fig. 2B**). Direct MQ1 vs. MQ5 comparison by

Mann-Whitney test uncovered a small (1.3-fold) and non-significant (p=0.86) difference in adjusted fold-

growth. Overall, we did not obtain an association between mandates and lower COVID-19 growth.

Mask use is not associated with most state COVID-19 case growth

We speculated that statewide mask use, rather than mask mandates per se, may predict lower COVID-19

growth rates. The Institute of Health Metrics and Evaluation (IHME) provides robust estimates for mask

use (defined as the percentage of people who always wear masks in public settings) (17). By simple linear

regression, mask use was associated with lower Ln, Fold, and adjusted Fold minima (p<0.0001; **Fig 3A-B**

and **S3A Fig**). To better understand this trend, we assigned US states to one of five mask use quintiles

(UQ1-5), with UQ1 including states with the highest mask use and UQ5 states with the lowest mask use.

UQ5 exhibited a 3.4-fold greater adjusted Fold minimum than UQ1 (p=0.002; **Fig 3C**), suggesting

potential association between mask use and COVID-19 spread at minima. By contrast, mask use was not

associated with Ln (p=0.071), Fold (p=0.058), or adjusted Fold (p=0.076) maxima (**Fig 3D-E** and **S3D**

Fig). Adjusted Fold maxima were also UQ-independent (p=0.56; **Fig 3F**), with direct UQ1 vs. UQ5

comparison by Mann-Whitney test uncovering a modest (1.5-fold) and non-significant (p=0.16)

difference in maxima. This suggests that mask use is not associated with COVID-19 spread at maxima.

We wondered why mask use was associated with lower minimum but not lower maximum growth rates.

Mask use was not associated with total cases at Ln minima (p=0.54) or maxima (p=0.086; **S3C-D Fig**),

indicating potential confounders in the mask-minimum growth relationship. Excluding Northeast states,

which exhibited the largest first waves and July 2020 seroprevalence (13, 28), total cases predicted lower

Ln minima (p=0.001; **S3E Fig**). This suggested that the link between mask use and lower minima may be

an artifact of the tendency for faster case growth to occur at lower case prevalence. In support of this, for

1 June – 1 Oct. 2020 mask use quintiles, normalized cases grew from 400 to 1350 at similar rates for UQ1

(which includes eight Northeast states) and UQ5 (p=0.22; **Fig. 3G**). UQ5 exhibited exponential growth

and reached these case totals ~50 days after UQ1 (**Fig 3H-I**), further implying that higher growth rates

may reflect lower total cases in low mask use states before minima. 0-80 days after Ln maxima, when

total case differences were smaller among states, UQ1 and UQ5 exhibited indistinguishable growth rates

(p=0.78; **Fig 3J**; 1 Oct. 2020 – 1 March 2021 mask use quintiles). Growth was post-exponential for both

UQ1 and UQ5 during this period (**Fig 3K-L**), and total cases predicted lower Ln maxima in all

continental US states (p<0.0001; **S3F Fig**). Together, these data suggest that mask use is an unreliable

predictor of COVID-19 growth in US states.

Mask use does not predict Summer or Fall-Winter COVID-19 cumulative growth in US states.

As expected, total cases were negatively associated with Ln growth in non-Northeast states for 1 June-1

Oct. 2020 and all continental US states for 1 Oct. 2020 – 1 March 2021 (p<0.0001; **S4A-B Fig**). We

reasoned that even if mask use could not predict growth rate, mask use may be negatively associated with

cumulative case growth. 1 June-1 Oct. 2020 (Summer) 1 Oct. 2020 – 1 March 2021 (Fall-Winter)

represent two distinct COVID-19 growth waves (**Fig 4A-B**). Excluding Northeast states, masks were not

associated with lower Summer growth using Ln (p=0.11; **Fig 4C**) or Fold (p=0.18; **S4C Fig**) models.

Mask use trended with lower adjusted Fold Summer growth (p=0.05; **S4D Fig**). While adjusted Fold

Summer growth was 3-fold higher in UQ4 than UQ1 (p=0.009), all other pairwise comparisons were not

significant (**Fig. 4D**). Likewise, mask use was not associated with lower Fall-Winter growth using Ln

(p=0.94; **Fig 4E**), Fold (p=0.91; **S4E Fig**), or adjusted Fold (p=0.71; **S4F Fig**) models, and adjusted Fold

Fall-Winter growth was not significantly different among mask use quintiles (p=0.38; **Fig. 4F**). These

data suggest that mask use is not consistently associated with Summer wave growth and not associated

with Fall-Winter wave growth in US states. Furthermore, low Summer growth did not protect Northeast

states from subsequent Fall-Winter growth. In summary, statewide SARS-CoV-2 transmission waves

appear independent of reported mask use (17).

Discussion

Our main finding is that mask mandates and use likely did not affect COVID-19 case growth. Mask

mandates were associated with greater mask use but ultimately did not influence total normalized cases or

post-mandate case growth. Higher mask use (rather than mandates per se) has been argued to decrease

COVID-19 growth rates (11). While compliance varies by location and time, IHME estimates are derived

from multiple sources and densely sampled. Even when accounting for population density, higher mask

use was not associated with lower Ln or fold maximum growth rates or lower Fall-Winter case growth

among continental US states. By contrast, mask use-growth rate association was highly significant at

minima. This antinomy warrants consideration. Mask use did not predict normalized cases at growth

minima or maxima, whereas there were more cases in the highest than the lowest mask use quintile before

minima. Northeast states exhibited the highest seroprevalence by July 2020 (28) and comprised 80% of

the highest mask use quintile, suggesting that mask use may be a lagging indicator of case growth. At

maxima, when case prevalence was similar among states, COVID-19 growth rates were also similar for

the highest and lowest mask use quintiles. Thus, initial association between masks and lower COVID-19

growth rates that dissipated during the Fall-Winter 2020-21 wave is likely an artifact of fewer normalized

cases begetting faster growth in states with coincidental low mask use.

There is inferential but not demonstrable evidence that masks reduce SARS-CoV-2 transmission. Animal

models (29), small case studies (6), and growth curves for mandate-only states (16) suggest that mask

efficacy increases with mask use (11). However, we did not observe lower growth rates over a range of

compliance at maximum Fall-Winter growth (45-83% between South Dakota and Massachusetts during

maxima) (17) when growth rates were high. This complements a Danish RCT from 3 April to 2 June

2020, when growth rates were low, which found no association between mask use and lower COVID-19

rates either for all participants in the masked arm (47% strong compliance) or for strongly compliant

participants only (8). While N-95 respirators offer some protection against respiratory viruses (10), there

is limited evidence for cloth and medical masks. Higher self-reported mask use protected against SARS-

CoV-1 in Beijing residents (30), but RCTs found no differences in PCR confirmed influenza among Hong

Kong households assigned to hand hygiene with or without masks (mask use 31% and 49%, respectively)

(31). Medical and cloth masks did not reduce viral respiratory infections among clinicians in Vietnam (9)

or China (10), and rhinovirus transmission increased among universally masked Hong Kong students and

teachers in 2020 compared with prior years (32). These findings are consistent with a 2020 CDC meta-

analysis (33) and a 2020 Cochrane review update (34).

Our study has implications for respiratory virus mitigation. Public health measures should ethically

promote behaviors that prevent communicable diseases. The sudden onset of COVID-19 compelled

adoption of mask mandates before efficacy could be evaluated. Our findings do not support the

hypothesis that greater public mask use decreases COVID-19 spread. As masks have been required in

many settings, it is prudent to weigh potential benefits with harms. Masks may promote social cohesion

during a pandemic (35), but risk compensation can also occur (36). By obscuring nonverbal

communication, masks interfere with social learning in children (37). Likewise, masks can distort verbal

speech and remove visual cues to the detriment of individuals with hearing loss; clear face-shields

improve visual integration, but there is a corresponding loss of sound quality (38, 39). Prolonged mask

use (>4 hours per day) promotes facial alkalinization and inadvertently encourages dehydration, which in

turn can enhance barrier breakdown and bacterial infection risk (40). British clinicians have reported

masks to increase headaches and sweating and decrease cognitive precision (41). Survey bias

notwithstanding, these sequelae are associated with medical errors (42). Future research is necessary to

assess risks of long-term daily mask use (34). As COVID-19 remains a public health threat, it is also

appropriate to emphasize interventions with demonstrated efficacy against COVID-19, most notably

vaccination (43) and vitamin D repletion (44).

In conclusion, we found mask mandates and use to be poor predictors of COVID-19 spread in US states.

Strengths of our study include assessing COVID-19 association with both mandates and reported use;

evaluating both Ln and Fold growth models; accounting for population density differences; and

measuring case growth after modeled mandate effective dates in states with late or no mandates. Our

study also has key limitations. We did not assess counties or localities, which may trend independently of

state averages. While dense sampling promotes convergence, IHME masking estimates are subject to

survey bias. We only assessed one biological quantity (confirmed and probable COVID-19), but the

ongoing pandemic warrants assessment of other factors such as hospitalizations and mortality.

Importantly, our study does not disprove the efficacy of all masks in limited and controlled

circumstances, such as properly worn N95 respirators. A recent study found that at typical respiratory

fluence rates, medical masks decrease airway deposition of 10-20μm SARS-CoV-2 particles but not 1-

5μm SARS-CoV-2 aerosols (45). Aerosol expulsion increases with COVID-19 disease severity in non-

human primates, as well as with age and BMI in humans without COVID-19 (46). Together with

enhanced vaccination rates, aerosol treatment with improved ventilation and air purification could help

reduce the size of COVID-19 outbreaks.

Acknowledgments

The authors thank Brandy Jesernik, Ashley Tracey, Jay Bhattacharya, Scott Atlas, and Erik Fostvedt for

manuscript input.

References

1.   Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with
Pneumonia in China, 2019. N Engl J Med. 2020;382(8):727-33.
2.   Hojyo S, Uchida M, Tanaka K, Hasebe R, Tanaka Y, Murakami M, et al. How COVID-19 induces
cytokine storm with high mortality. Inflamm Regen. 2020;40:37.
3.   Chu Y, Yang J, Shi J, Zhang P, Wang X. Obesity is associated with increased severity of disease in
COVID-19 pneumonia: a systematic review and meta-analysis. Eur J Med Res. 2020;25(1):64.
4.   Jayaweera M, Perera H, Gunawardana B, Manatunge J. Transmission of COVID-19 virus by
droplets and aerosols: A critical review on the unresolved dichotomy. Environ Res. 2020;188:109819.
5.   Payne DC, Smith-Jeffcoat SE, Nowak G, Chukwuma U, Geibe JR, Hawkins RJ, et al. SARS-CoV-2
Infections and Serologic Responses from a Sample of U.S. Navy Service Members - USS Theodore
Roosevelt, April 2020. MMWR Morb Mortal Wkly Rep. 2020;69(23):714-21.
6.   Hendrix MJ, Walde C, Findley K, Trotman R. Absence of Apparent Transmission of SARS-CoV-2
from Two Stylists After Exposure at a Hair Salon with a Universal Face Covering Policy - Springfield,
Missouri, May 2020. MMWR Morb Mortal Wkly Rep. 2020;69(28):930-2.
7.   Ludvigsson JF, Engerström L, Nordenhäll C, Larsson E. Open Schools, Covid-19, and Child and
Teacher Morbidity in Sweden. N Engl J Med. 2021;384(7):669-71.
8.   Bundgaard H, Bundgaard JS, Raaschou-Pedersen DET, von Buchwald C, Todsen T, Norsk JB, et al.
Effectiveness of Adding a Mask Recommendation to Other Public Health Measures to Prevent SARS-CoV-
2 Infection in Danish Mask Wearers : A Randomized Controlled Trial. Ann Intern Med. 2021;174(3):335-
43.
9.   MacIntyre CR, Seale H, Dung TC, Hien NT, Nga PT, Chughtai AA, et al. A cluster randomised trial
of cloth masks compared with medical masks in healthcare workers. BMJ Open. 2015;5(4):e006577.
10.   MacIntyre CR, Wang Q, Rahman B, Seale H, Ridda I, Gao Z, et al. Efficacy of face masks and
respirators in preventing upper respiratory tract bacterial colonization and co-infection in hospital
healthcare workers. Prev Med. 2014;62:1-7.
11.   Howard J, Huang A, Li Z, Tufekci Z, Zdimal V, van der Westhuizen H-M, et al. An evidence review
of face masks against COVID-19. Proceedings of the National Academy of Sciences.
2021;118(4):e2014564118.
12.   Wang Y, Tian H, Zhang L, Zhang M, Guo D, Wu W, et al. Reduction of secondary transmission of
SARS-CoV-2 in households by face mask use, disinfection and social distancing: a cohort study in Beijing,
China. BMJ Glob Health. 2020;5(5).
13.   United States COVID-19 Cases and Deaths by State over Time. Centers for Disease Control and
Prevention (CDC); 2021.
14.   Worldometers.info. Worldometer Dover, DE USA2021 [updated 25 May 2021; cited 2021.
Available from: https://www.worldometers.info/coronavirus/country/us/.
15.   Guy GP, Jr., Lee FC, Sunshine G, McCord R, Howard-Williams M, Kompaniyets L, et al.
Association of State-Issued Mask Mandates and Allowing On-Premises Restaurant Dining with County-
Level COVID-19 Case and Death Growth Rates - United States, March 1-December 31, 2020. MMWR
Morb Mortal Wkly Rep. 2021;70(10):350-4.
16.   Lyu W, Wehby GL. Community Use Of Face Masks And COVID-19: Evidence From A Natural
Experiment Of State Mandates In The US. Health Aff (Millwood). 2020;39(8):1419-25.
17.   Institute for Health Metrics and Evaluation COVID-19 model: University of Washington; 2021
[Available from: https://covid19.healthdata.org/.
18.   Liu Z, Fang CT. A modeling study of human infections with avian influenza A H7N9 virus in
mainland China. Int J Infect Dis. 2015;41:73-8.
19.   Perneger T, Kevorkian A, Grenet T, Gallée H, Gayet-Ageron A. Alternative graphical displays for
the monitoring of epidemic outbreaks, with application to COVID-19 mortality. BMC Medical Research
Methodology. 2020;20(1):248.

20.   De Flora S, La Maestra S. Growth and decline of the COVID-19 epidemic wave in Italy from
March to June 2020. J Med Virol. 2021;93(3):1613-9.
21.   Bureau USC. 2010 Census Urban and Rural Classification and Urban Area Criteria. In: Commerce
UDo, editor. 2010.
22.   Curtis MJ, Alexander S, Cirino G, Docherty JR, George CH, Giembycz MA, et al. Experimental
design and analysis and their reporting II: updated and simplified guidance for authors and peer
reviewers. British Journal of Pharmacology. 2018;175(7):987-93.
23.   Khan A, Waleed M, Imran M. Mathematical analysis of an influenza epidemic model,
formulation of different controlling strategies using optimal control and estimation of basic
reproduction number. Mathematical and Computer Modelling of Dynamical Systems. 2015;21(5):432-
59.
24.   Samaras L, Sicilia M-A, García-Barriocanal E. Predicting epidemics using search engine data: a
comparative study on measles in the largest countries of Europe. BMC Public Health. 2021;21(1):100.
25.   Ghosal S, Sengupta S, Majumder M, Sinha B. Linear Regression Analysis to predict the number of
deaths in India due to SARS-CoV-2 at 6 weeks from day 0 (100 cases - March 14th 2020). Diabetes Metab
Syndr. 2020;14(4):311-5.
26.   Chu J. A statistical analysis of the novel coronavirus (COVID-19) in Italy and Spain. PloS one.
2021;16(3):e0249037.
27.   Jha AK, Tsai T, Jacobson B. Why we need at least 500,000 tests per day to open the economy —
and stay open: Brown School of Public Health; 2020 [Available from:
https://globalepidemics.org/2020/04/18/why-we-need-500000-tests-per-day-to-open-the-economy-
and-stay-open/.
28.   Anand S, Montez-Rath M, Han J, Bozeman J, Kerschmann R, Beyer P, et al. Prevalence of SARS-
CoV-2 antibodies in a large nationwide sample of patients on dialysis in the USA: a cross-sectional study.
Lancet (London, England). 2020;396(10259):1335-44.
29.   Chan JF, Yuan S, Zhang AJ, Poon VK, Chan CC, Lee AC, et al. Surgical Mask Partition Reduces the
Risk of Noncontact Transmission in a Golden Syrian Hamster Model for Coronavirus Disease 2019
(COVID-19). Clin Infect Dis. 2020;71(16):2139-49.
30.   Wu J, Xu F, Zhou W, Feikin DR, Lin CY, He X, et al. Risk factors for SARS among persons without
known contact with SARS patients, Beijing, China. Emerg Infect Dis. 2004;10(2):210-6.
31.   Cowling BJ, Chan KH, Fang VJ, Cheng CK, Fung RO, Wai W, et al. Facemasks and hand hygiene to
prevent influenza transmission in households: a cluster randomized trial. Ann Intern Med.
2009;151(7):437-46.
32.   Fong MW, Leung NHL, Cowling BJ, Wu P. Upper Respiratory Infections in Schools and Childcare
Centers Reopening after COVID-19 Dismissals, Hong Kong. Emerg Infect Dis. 2021;27(5).
33.   Fong MW, Gao H, Wong JY, Xiao J, Shiu EYC, Ryu S, et al. Nonpharmaceutical Measures for
Pandemic Influenza in Nonhealthcare Settings-Social Distancing Measures. Emerg Infect Dis.
2020;26(5):976-84.
34.   Jefferson T, Del Mar CB, Dooley L, Ferroni E, Al-Ansary LA, Bawazeer GA, et al. Physical
interventions to interrupt or reduce the spread of respiratory viruses. Cochrane Database of Systematic
Reviews. 2020(11).
35.   Goh Y, Tan BYQ, Bhartendu C, Ong JJY, Sharma VK. The face mask: How a real protection
becomes a psychological symbol during Covid-19? Brain Behav Immun. 2020;88:1-5.
36.   Yan Y, Bayham J, Richter A, Fenichel EP. Risk compensation and face mask mandates during the
COVID-19 pandemic. Scientific Reports. 2021;11(1):3174.
37.   Spitzer M. Masked education? The benefits and burdens of wearing face masks in schools during
the current Corona pandemic. Trends Neurosci Educ. 2020;20:100138.

38.   Corey RM, Jones U, Singer AC. Acoustic effects of medical, cloth, and transparent face masks on
speech signals. J Acoust Soc Am. 2020;148(4):2371.
39.   Atcherson SR, Finley ET, Renee McDowell BR, Watson C. More Speech Degradations and
Considerations in the Search for Transparent Face Coverings During the COVID-19 Pandemic. Audiology
Today. 2020;Nov/Dec.
40.   Hua W, Zuo Y, Wan R, Xiong L, Tang J, Zou L, et al. Short-term skin reactions following use of N95
respirators and medical masks. Contact Dermatitis. 2020;83(2):115-21.
41.   Davey SL, Lee BJ, Robbins T, Randeva H, Thake CD. Heat stress and PPE during COVID-19: impact
on healthcare workers' performance, safety and well-being in NHS settings. J Hosp Infect. 2021;108:185-
8.
42.   Rosenberg K. The joint commission addresses health care worker fatigue. Am J Nurs.
2014;114(7):17.
43.   Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of
the BNT162b2 mRNA Covid-19 Vaccine. N Engl J Med. 2020;383(27):2603-15.
44.   Radujkovic A, Hippchen T, Tiwari-Heckler S, Dreher S, Boxberger M, Merle U. Vitamin D
Deficiency and Outcome of COVID-19 Patients. Nutrients. 2020;12(9).
45.   Xi J, Si XA, Nagarajan R. Effects of mask-wearing on the inhalability and deposition of airborne
SARS-CoV-2 aerosols in human upper airway. Phys Fluids (1994). 2020;32(12):123312.
46.   Edwards DA, Ausiello D, Salzman J, Devlin T, Langer R, Beddingfield BJ, et al. Exhaled aerosol
increases with COVID-19 infection, age, and obesity. Proceedings of the National Academy of Sciences of
the United States of America. 2021;118(8).

476 **Figure Legends**

Fig 1. Earlier mask mandates are not consistently associated with lower COVID-19 growth rates in continental US States. A-B. Natural logarithmic (A) and Fold (B) COVID-19 growth in continental US states. Red horizontal lines denote growth rate minima (Min) and maxima (Max) between Summer and Fall-Winter waves. Surge: growth rate increase between Min and Max. C. Ln minima were not associated with the time quintile of a state's mandate effective date (MQ). D. Fold minima trended lower in MQ1 than MQs 3 and 5. E. Adjusted Fold minima were lower in MQ1 than MQ4 and indistinguishable for all other pairwise comparisons. F-G. Ln (F) and Fold (G) maxima were not associated with mandate effective date time quintiles. H. Adjusted Fold maxima were lower in MQ2 than MQ4 and indistinguishable for all other pairwise comparisons. I. States with earlier mask mandates exhibited greater mask use between Oct. 2020 and March 2021. J. Cases per 100,000 by 1 March 2021 were not associated with mandate effective date time quintiles. Different letters denote p<0.05 by Tukey tests after one-way ANOVA (C, F, G) or all pairwise comparison Dunn tests after Kruskal-Wallis (D, E, H-J). *: p<0.05 by Kruskal-Wallis. n.s.: not significant. Error bars: 95% confidence intervals (A-B), standard deviations (C, F, G), and interquartile ranges (D, E, H-J).

Fig 2. Earlier mask mandates are not associated with lower post-mandate COVID-19 growth rates in contiguous US states. A. Effective and modeled effective (***bold, italicized***) dates in 2020 for mask mandates in contiguous US states. State colors denote effective date time quintiles. Modeled dates of MQ4-5 states (late or no actual mandates) are medians of effective dates among bordering states of MQ1-3 (earlier mandates). Dashed lines denote MQ1-3 states that border a given MQ4-5 state. B-D. Between actual or modeled mandate effective dates and 1 March 2021, Ln Growth (B), Fold Growth (C), and population density-adjusted Fold Growth (D) were not associated with mandate effective date time quintiles. n.s.: not significant by one-way ANOVA (B) or Kruskal-Wallis (C-D). Error bars: standard deviations (B) and interquartile ranges (C-D).

Fig 3. Mask use does not consistently predict COVID-19 case growth in continental US states. A-C.

At minima, mask use was associated with lower ln (A), fold (B), and population density-adjusted fold (C)

growth rates. D-F. At maxima, mask use was not associated with ln (D), fold (E), or population density-

adjusted fold (F) growth rates. G. States in June-Oct. 2020 mask use quintiles (UQ) 1 and 5 grew from

400 to 1350 normalized cases at indistinguishable rates before minima. H-I. Ln cases (H) and cases (I) vs.

time for UQ1 and UQ5. J. States in Oct. 2020-March 2021 mask use UQ1 and UQ5 exhibited

indistinguishable Fold Growth 80 days after maxima. K-L. Ln cases (K) and cases (L) vs. time for UQ1

and UQ5. Simple linear regression used weighted (A-B) or ordinary (D-E) least squares. R^2 values refer

to unconstrained lines of best fit. Different letters denote $p < 0.05$ by all pairwise comparison Dunn tests

after Kruskal-Wallis (C, F). n.s.: not significant. Error bars: Interquartile ranges (C, F, J, K) and 95%

confidence intervals (G-H).

Fig 4. Mask use does not predict lower COVID-19 growth during the Summer or Fall-Winter

waves. A-B. Ln Growth rate (A) and total COVID-19 cases (B) for continental US states from 20 April

2020 to 6 March 2021. Red vertical lines denote Summer (Jun-Oct 2020) and Fall-Winter (Oct 2020-Mar

2021) waves. C. Mask use does not predict Summer Ln Growth in non-Northeast states. D. In the

Summer wave, population-adjusted Fold Growth was lower in Summer mask use UQ1 than UQ4 and

indistinguishable among UQ2-5. E. Mask use does not predict Fall-Winter Ln Growth in continental US

states. F. In the Fall-Winter wave, population-adjusted Fold Growth was indistinguishable among Fall-

Winter mask use quintiles. Simple linear regression used ordinary least squares (C, E). R^2 values refer to

unconstrained lines of best fit. Different letters denote $p < 0.05$ by all pairwise comparison Dunn tests after

Kruskal-Wallis (D, F). n.s.: not significant. Error bars: 95% confidence intervals (A-B) and interquartile

ranges (D, F). Solid circles (●): All continental US states. Hollow circles (○): Excluded Northeast states.

Supporting Information Legends

S1 Fig. A. Natural logarithmic COVID-19 growth rates in continental US states by mask mandate time quintile. Y-axis values are differences between natural logarithms of total cases on a day and natural logarithms of total cases on the prior day. States are listed by time quintiles at which mask mandates became effective statewide. Thin black and wide gray denote mean and 95% confidence intervals, respectively. Red vertical lines denote dates of mask mandate issuance. Red horizontal lines indicate growth rate minima (phase 4) and maxima (phase 5) between the Summer and Fall-Winter waves. B. There is no association between cases and COVID-19 PCR tests among US states (case and test totals as of 25 May 2021).

S2 Fig. Earlier mask mandates are not consistently associated with lower COVID-19 growth surges in continental US States. Ln (A) and Fold (B) surges were not associated with the time quintile of a state's mandate effective date. C. Adjusted Fold maxima were lower in MQ2 than MQ4 and indistinguishable for all other pairwise comparisons. Different letters denote $p<0.05$ by all pairwise comparison Dunn tests after Kruskal-Wallis (C). n.s.: not significant by one-way ANOVA. Error bars: standard deviations (A, B) and interquartile ranges (C).

S3 Fig. **Inverse association between normalized cases and COVID-19 growth rates in continental US states**. A. At minima, mask use predicts lower population density-adjusted Fold Growth. B. At maxima, mask use does not predict population density-adjusted Fold Growth. C. Normalized cases do not predict mask use at minima. D. Normalized cases do not predict mask use at maxima. E. At minima, greater total cases predict lower growth rates in non-Northeast continental US states. F. At maxima, greater total cases predict lower growth rates in continental US states at maxima. Black circles (●): All continental US states. Hollow blue circles (○): Excluded Northeast states. Solid blue circles (●): Included Northeast states. Red squares (■): Midwest states. Green triangles (▲): Mountain-Plains States. Grey

triangles (▼): South states. Gold diamonds (♦): Pacific states. Simple linear regression used ordinary (A-

D, F) or weighted (E) least squares and included all continental states except E (which excluded Northeast

states). ε: Non-normal residuals by D'Agostino-Pearson test. R^2 values refer to unconstrained lines of best

fit.

S4 Fig. **Limited association between mask use and COVID-19 case growth during Summer and**

Fall-Winter waves. A. Mask use does not predict Summer Fold Growth in non-Northeast states. B. Mask

use is marginally associated with lower population-adjusted Fold Growth in non-Northeast states. C-D.

Normalized cases were associated with lower Ln Growth during Summer (C) and Fall-Winter (D) waves

in non-Northeast states (C) and all continental US states (D), respectively. E-F. Mask use does not predict

Fall-Winter Fold Growth (E) or population-adjusted Fold Growth (F) in continental US states. Simple

linear regression used ordinary least squares and excluded Northeast states (A, C, D) or included all

continental US states (B, E, F). ε: Non-normal residuals by D'Agostino-Pearson test. R^2 values refer to

unconstrained lines of best fit.

S1 Table. **Supplemental Data Tables**. Worksheet A. 7-day rolling mean of total normalized cases (cases

per 100,000 residents of each US state from CDC), 31 March 2020 - 1 March 2021. Worksheet B. Natural

logarithmic daily case growth, 1 April 2020 – 1 March 2021. 20-day minima and maxima are denoted in

red and gold, respectively. Mask mandate effective dates and modeled effective dates are provided for

each state. Worksheet C. Population density calculations. See Materials and Methods for explanation of

formulas. Worksheet D. 20-day Fold Growth, 12 August 2020 – 1 March 2021. Minima and maxima are

denoted in red and gold, respectively. Population density-adjusted Fold Growth data are provided for each

state. Worksheet E. Mask use for each US state at Ln and Fold extrema and during the Summer and Fall-

Winter waves. Date range mask use values are simple arithmetic means of daily mask use over the

specified date range. Numeric daily mask use (obtained from IHME) are provided.

A

Mandate Effective Date Quintile
- MQ1
- MQ2
- MQ3
- MQ4
- MQ5

C_M : Cases on effective (MQ_{1-3}) or modeled effective (MQ_{4-5}) mandate day.

B — **Ln Growth** After MQ_{1-3}

% ln (C_{301}/C_M)

n.s.

Mandate Quintile

C — **Fold Growth** After MQ_{1-3}

% $C_{301}/C_M - 1$

n.s.

Mandate Quintile

D — **Adj. Fold Growth** After MQ_{1-3}

% $C_{301}/C_M - 1$

n.s.

Mandate Quintile

A Daily %Ln Case Growth (Continental US States)

Ln Growth ($\ln C_t/C_{t-1}$)

Days since 20 April 2020

Summer

Fall-Winter

B Total Cases (Continental US States)

Cases/100k

Fall-Winter

Summer

Days since 20 April 2020

C ○ Excluded Northeast States

Ln Summer Growth vs. Masks

$\ln C_{1001}/C_{601}$

$R^2 = 0.07$
$p = 0.11$

% Masks

D Adj. Summer Fold Growth by 1Jun-1Oct Mask Quintile

Fold-Growth

a $\quad ab$ $\quad ab$ $\quad b$ $\quad ab$

% Masks 64-79 59-64 53-58 49-51 33-48

E Ln Fall-Winter Growth vs. Masks

$\ln C_{301}/C_{1001}$

$R^2 = 0.0001$
$p = 0.94$

% Masks

F Adj. Fall-Winter Fold Growth by 1Oct-1Mar Mask Quintile

Fold-Growth

n.s.

% Masks 78-82 74-77 69-74 64-69 48-64