1

A feasibility study to test a novel approach to dietary weight loss with a focus on

assisting informed decision making in food selection

Mindy H. Lee[1], Catherine C. Applegate[1], Annabelle L. Shaffer[1], Abrar Emamaddin[2],

John W. Erdman Jr.[1,2], Manabu T. Nakamura[1,2*]

[1]Division of Nutritional Sciences, University of Illinois at Urbana-Champaign, Urbana, IL

[2]Department of Food Science and Human Nutrition, University of Illinois at Urbana-
Champaign, Urbana, IL
13

14 * Corresponding author

15 E-mail: mtnakamu@illinois.edu

# Abstract

Obesity is a significant contributor to the development of chronic diseases, some of which can be prevented or reversed by weight loss. However, dietary weight loss programs have shortcomings in the success rate, magnitude, or sustainability of weight loss. The Individualized Diet Improvement Program's (iDip) objective was to test the feasibility of a novel approach that helps individuals self-select a sustainable diet for weight loss and maintenance instead of providing weight loss products or rigid diet instructions to follow.

The iDip study consisted of 22 dietary improvement sessions over 12 months with six months of follow-up. Daily weights were collected, and a chart summarizing progress was provided weekly. Six 24-hour dietary records were collected, and dietary feedback was provided in the form of a protein-fiber plot, in which protein/energy and fiber/energy of foods were plotted two-dimensionally together with a target box specific to weight loss or maintenance. An exit survey was conducted at 12 months.

Twelve (nine female, 46.3±3.1 years (mean±SE)) of the initial 14 participants (BMI>28 kg/m$^2$) completed all sessions. Mean percent weight loss (n=12) at six and 12 months was -4.9%±1.1 (p=0.001) and -5.4%±1.7 (p=0.007), respectively. Weight loss varied among individuals at 12 months; top and bottom halves (n=6 each) achieved -9.7%±1.7 (p=0.0008) and -1.0%±1.4 weight loss, respectively. The 24-hour records showed a significant increase in protein density from baseline to final (4.1g/100kcal±0.3 vs. 5.7g/100kcal±0.5; p=0.008). Although mean fiber density showed no significant change from the first month (1.3g/100kcal±0.1), the top half had significantly higher

fiber/energy intake than the bottom half group. The survey suggested that all

participants valued the program and its self-guided diet approach.

In conclusion, half of the participants successfully lost >5% and maintained the

lost weight for 12 months without strict diet instructions, showing the feasibility of the

informed decision-making approach.

# **Introduction**

With the advancement of food production globally, obesity has become a

worldwide epidemic, increasing the risk of morbidity and mortality [1, 2]. In the United

States, 70.2% of adults are obese (body mass index, BMI>30 kg/m$^2$) or overweight

(BMI>25 kg/m$^2$), with 7.7% having extreme obesity (BMI>40 kg/m$^2$) [3]. Overweight and

obesity are risk factors for several comorbidities, including type 2 diabetes mellitus,

cardiovascular disease, and cancers. Based on 2013 US medical spending, obesity and

its complications cost $342.2 billion annually (28.2% of all medical expenditures) [4].

Modest weight losses of 3-5% initial body weight are capable of lowering blood

triglycerides, fasting blood glucose, and improving glycemic control. Further weight

losses enhance these benefits, and can also reduce blood pressure, cholesterol, and

need for medications [2, 5].

Currently available weight loss methods do not reliably achieve medically

significant weight loss as they are insufficient in success rate, sustainability, or

magnitude of weight loss. Exclusion of certain food groups is a common approach of

popular dieting programs, which can result in some weight loss, but high attrition rates

often result due to a lack of dietary adherence [6-8]. For example, commonly used

dieting programs, including Atkins (carbohydrate restriction), Zone (substituting

carbohydrate with protein), and Ornish (reduction of animal foods), can induce modest

weight losses of <5 kg but lack a high success rate and long-term sustainability [9-12].

Participants grow fatigued with the strict recipes and are unable to adapt the plan to

their daily lives [13, 14]. This translates to low adherence, unsuccessful weight loss, and

lack of diet sustainability.

A negative energy balance is a requirement for weight loss. Calorie monitoring is

often proposed to achieve this and is recommended by health professionals [15, 16].

However, it is laborious and frequently inaccurate [17, 18]. Often, programs utilizing

calorie counting or food journals lead to high attrition and thus, insignificant long-term

weight loss [19-21]. More importantly, the protein requirement increases during periods

of negative energy balance compared with the requirement during weight maintenance

[22, 23]. Focusing only on energy reduction may result in exacerbating the loss of lean

body mass and decreased efficiency in weight loss [23].

Successful, clinically significant weight losses can be achieved using meal

replacement products; however, weight maintenance remains a substantial challenge

for this approach [5]. Very-low-calorie diets (VLCD) in the form of meal replacements

have shown clinically significant weight loss in the short term, but VLCD participants are

less successful in weight maintenance compared to a restricted-energy whole-food diet

[24-26]. Even when a VLCD was combined with intensive dietary and lifestyle

intervention programs using counseling, education, and exercise, participants regained

most or all of the lost weight during the follow-up periods [5, 27].

Pharmaceuticals are available for treating obesity, such as orlistat and

phentermine/topiramate, and work by reducing fat absorption and suppressing appetite,

respectively. However, these medications result in only 3-6% weight loss and can

include several side effects such as reduced fat-soluble vitamin absorption or

gastrointestinal upset [28]. As a result of the low efficacy in medicine and lifestyle

interventions, bariatric surgery remains the most reliable weight-loss treatment and can

reduce obesity complications, such as hyperglycemia [29]. Bariatric surgery does have

significant drawbacks, including a mean cost of $14,389 per surgery, surgical

complications, nutritional deficiencies, and the need for drastic dietary changes [30].

In summary, although dietary weight loss can treat comorbidities associated with

obesity, existing programs have shortcomings in sustainability, magnitude, adherence,

and success rate. As such, a dietary weight loss program that can be used as a reliable

treatment option is yet to be developed. Thus, we tested the feasibility of a new

approach to dietary weight loss that does not rely on traditionally used methods.  First,

in place of common methods such as exclusion of food groups, strict weight-loss meal

plans, and dieting products, iDip increases nutritional knowledge allowing participants to

create their own weight loss diet by making informed and individualized food choices

based on their preferences. Second, we monitored a target range of protein density per

energy for participants to create a weight loss diet with increased protein while reducing

the overall energy intake to meet an increased protein requirement during weight loss.

Third, we replaced calorie counting/food journaling with daily weight monitoring and

provision of a weekly progress chart to monitor energy balance.

# Materials and methods

This study was approved by the University of Illinois Institutional Review Board (#18069) on August 30[th], 2017. The study was registered at the US National Institutes of Health (ClinicalTrial.gov) #NCT04605653. Registration was delayed due to a lack of knowledge on required clinical trial registration. The authors confirm that all ongoing and related trials for this intervention are registered. Participant recruitment started on September 1[st], 2017, and the follow-up period ended on July 28[th], 2018. Informed consent was obtained from all participants included in the study.

## Participant recruitment

Adults aged 18 to 64 years old and with a BMI >28 kg/m$^2$ were recruited via posters, flyers, and word of mouth in Urbana-Champaign, Illinois, USA, and surrounding regions from September 2017 through January 2018. A power calculation was performed using an average standard deviation 7.0 for percentage weight loss at 12 months from similar dietary weight loss studies [31]. We set 5% weight loss as a primary outcome to be detected. The minimum sample number required with 95% confidence interval was 7.5. With n=10, we will be able to detect 5% difference when actual SD is 8.0. Alternatively, we will be able to detect 4.3% difference if SD is 7.0 as estimated. Based on a mean attrition rate of 22.1% from 60 studies [32], we recruited 14 participants assuming 30% attrition rate.

Inclusion criteria in addition to age and BMI were access to Wi-Fi at home, working email and smartphone, fluent English communicator, and willingness to self-weigh daily with a provided Wi-Fi scale for a minimum of 18 months. Exclusion criteria were the following: adults with self-reported severe metabolic, cardiovascular or

musculoskeletal disease, insulin use, pregnancy, planned pregnancy, or lactating; or

unwillingness to attend any of the educational sessions. Among eligible applicants,

invitation was prioritized to best reflect the gender and racial composition of the

community. Selected applicants were invited for an individual meeting for an additional

screening process at the University of Illinois at Urbana-Champaign (UIUC). After

receiving explanations about the study, a Food Frequency Questionnaire (FFQ) was

administered, and anthropometric measurements were collected. Informed consent was

obtained, and a Wi-Fi scale was provided. Other than a Wi-Fi scale, no monetary

compensation was provided to the participants for participating in the study.

## Study design

The iDip study utilized a non-randomized, single-arm, before-and-after study

design with a cohort of participants and comprised of 12 months of intervention followed

by six months of follow-up. The objective was to test the feasibility of the informed

decision-making approach to dietary weight loss by utilizing two forms of quantitative

visual feedback for dietary improvement and weight loss progress. The iDip consisted of

four weekly sessions (month one), 12 biweekly sessions (months two-six), and six

monthly sessions (months seven-12). Nineteen 60-minute group-based diet

improvement sessions and three 30-minute individual advising sessions were delivered

by graduate students with registered dietitian credentials at UIUC from January 2018 to

January 2019. All group sessions were focused on building knowledge and skills to

make informed food selections. Weekly sessions in month one focused on the basic

strategy of weight loss; bi-weekly sessions in months two-six focused on protein, fiber,

physical activity, weight loss plateauing, and weight maintenance; and months seven-12

of monthly sessions focused on healthy eating and micronutrients beyond weight

management (S1 Appendix). All participants received two types of feedback: 1) weekly

weight chart and 2) protein-fiber (PF) plot (Fig 1) of their 24-hour diet record analysis at

one, two, three, five, seven and 12 months and pre-and-post FFQs to assist individuals

in monitoring their weight and dietary pattern, respectively. PF plot was the sole diet

feedback and analysis that participants received. No food groups were excluded from

participants' diets, and even no dieting products nor recipes were distributed. Weight

and dietary changes were evaluated during individual sessions, and participants were

advised using our new approach.

## Development of novel learning tools

### A quantitative nutrient display method

The PF plot (Fig 1) is a two-dimensional data-visualization method for easy

comparison of quantitative nutrition values in multiple foods designed to improve

food selection based on protein and fiber density of foods [33]. The PF plot was

developed by our lab, and its efficacy in aiding restaurant customers in selecting

healthier menus has been demonstrated previously [33]. Protein and fiber are critical

nutrients for weight management as adequate protein intake protects muscle mass

from degradation during weight loss [22, 34], and consuming sufficient protein and

fiber decreases excessive calorie consumption [35-37]. The fat to carbohydrate ratio

was disregarded in this study as previous randomized-controlled trials have not

shown that long-term weight loss favors a low-carbohydrate and high-fat diet or vice

versa [38-40].

Protein and fiber targets of the PF plot are derived from the Acceptable

Macronutrient Distribution Ranges (AMDR) and Adequate Intakes (AI) for weight

maintenance in adults, respectively, as previously described [33]. Briefly, for

protein, the AMDR is 10-36% daily intake; and for fiber, the AI is 14 g/1000 kcal

daily [15]. For the weight maintenance box (blue): using 16-32% kcals from

protein, we set 4-8 g/100 kcal as protein target; and for fiber, 1.4-2.8 g/100 kcal

(Fig 1). The weight loss target (green) was set assuming a caloric reduction of

minimum 25% of the daily requirement as weight loss requires a negative energy

balance. By adjusting for 25% calorie reduction and 1.25x increase in protein

requirement [22, 23], we set a protein range for weight loss of 7-11 g/100 kcal.

Using 14 g fiber/1000 kcals and factoring 25% reduction of caloric intake, the

resulting range was set as 1.8-3.2 g/100 kcals (Fig 1). The PF plot with a target

box of protein and fiber density for weight loss was used as a key tool for food

selection by participants throughout the study in both educational sessions and

individual diet analysis.

**Fig 1. Protein and fiber plot with example food items.** Foods with green dots are

dense in protein or fiber and can be easily fit in the target box when combined, whereas

foods with red dots cannot be due to low protein and fiber densities. A yellow food has

protein and fiber density between green and red foods.  A blue dot with a shadow

shows where a meal falls when all foods are combined.  Blue box, weight maintenance

target; green box, weight loss target.

## Energy balance monitoring

In the iDip, calorie counting and daily food journaling were replaced with daily self-weighing and weekly progress summary charts to monitor energy balance. As mentioned in the Introduction, calorie counting and food log methods are conventional in weight management, but these methods are time-consuming and prone to error, resulting in high attrition rates [19-21]. Daily self-weighing using a Wi-Fi-enabled scale (Weight Gurus, Greater Goods LLC, MO) provided a reliable energy balance estimation that requires little time. Participants were provided with a weekly weight feedback chart showing weight, a six month goal weight and an intended weight loss rate in the format shown in Fig 2.

**Fig 2. A weekly weight chart feedback.** The red line represents the target weight loss slope (1 lb. loss (0.45 kg) per week) over time, the blue line represents actual daily weights, and the green line represents the target of 25 lbs. (11.3 kg) weight loss (based on a six month weight loss goal of 1 lb. (0.45 kg) loss per week).

# Outcome measures

## Anthropometrics measures

Anthropometrics measurements were the primary outcomes, and entailed height, weight, waist, and hip circumferences. These were obtained at baseline and at 12 months. Height was recorded to the nearest 0.25 inches with shoes removed via stadiometer (Seca 700, Hanover, MD), and weight was measured to the nearest 0.1 lbs in light clothing on the same instrument. BMI was calculated using measured height and

204 weight. In addition, weights were monitored and obtained daily via a Wi-Fi scale. Waist

205 and hip circumferences were measured using a Retractable Tape Measure (Gulik II,

206 Gay Mills, WI) and recorded to the nearest 0.1 cm. Waist to hip ratio was obtained

207 based on measured waist and hip circumferences.

## Dietary assessment

Dietary changes were the secondary outcomes assessed in this study. A paper

version of the validated EPIC-Norfolk FFQ questionnaire (S2 Appendix) was used at

baseline and at 12 months to assess dietary patterns [41]. Twenty-four-hour dietary

records were administered at one, two, three, five, seven and 12 months to evaluate

dietary changes. The results were returned to the participants in the form of a PF plot.

The US Department of Agriculture National Nutrient Database for Standard Reference

(https://fdc.nal.usda.gov/) and manufacturer information were used to calculate protein

and fiber density.

## Survey measures

An exit survey was conducted at the end of the 12 month intervention. A

questionnaire containing 14 items was developed to explore personal gain, program

content, program materials and feedback, engagement, and overall evaluation of the

program. Possible responses for each question were "Definitely Yes," "Yes," "A little bit,"

"Neutral," "Not really," "No," "Definitely Not." A score was provided for each response

from one to seven, with seven being "Definitely Yes".

## Data analysis

Descriptive analysis was used to summarize demographic characteristics.

Weight loss of ≥5% from the baseline was considered meaningful as it has been shown

to reduce risks associated with obesity [42]. Paired t-test analysis determined the pre-

and post-intervention differences in outcome measures for the program completers.

Pearson correlation analyses were performed to determine associations between body

weight, protein density, and fiber density. Cohen's d was computed by single sample z-

test. Normality of data was tested using Shapiro-Wilk. Repeated measure ANOVA was

used to compare 24-hour dietary records. All statistical analyses were performed using

Microsoft Office Excel 2016 and R Computing (Version 3.6.1 © 2019), and $p<0.05$ was

considered statistically significant.

# 235 Results

## 236 Participant characteristics

A flow diagram of participant enrollment is shown in Fig 3. Fourteen participants

started the study. The mean age was 44.6±3.0 years, and mean BMI was 36.7±1.7

kg/m$^2$. The demographics and characteristics of participants are presented in Table 1.

Enrolled participants had medical histories of hypertension (36%), hypercholesterolemia

(21%), thyroid issues (21%), skeletal problems (14%), sleep apnea (14%), and

hyperlipidemia (7%).  Participants had prior dieting experiences including Weight

Watchers (71%), Jenny Craig (14%), liquid diets (14%), Ideal Protein (14%), low

carbohydrate (14%), South Beach (7%), low fat (7%), low calories (7%), cabbage soup

diet (7%), Atkins (7%), 21-day fix (7%), Human Chorionic Gonadotropin (HCG) diet

(7%), and Transformation diet (7%). No significant differences were found in any

demographic data between top and bottom groups in weight loss.

**Fig 3. Flow diagram of the participants**

**Table 1. Demographic data of participants**

|  | Enrollees (%) N=14 | Completers (%) N=12 |
|---|---|---|
| **Age in years** | | |
| **18-34** | 14.3 | 8.3 |
| **35-50** | 50.0 | 50.0 |
| **51-65** | 35.7 | 41.7 |
| **Gender** | | |
| **Male** | 21.4 | 25.0 |
| **Female** | 78.6 | 75.0 |
| **Marital Status** | | |
| **Married** | 64.3 | 75.0 |
| **Single** | 35.7 | 25.0 |
| **Ethnicity** | | |
| **White** | 57.1 | 50.0 |
| **African American** | 42.9 | 50.0 |
| **Educational level** | | |
| **Graduate degree** | 50.0 | 58.3 |
| **Bachelor's degree** | 42.9 | 41.7 |
| **High school** | 7.1 | 0.0 |

253    **Table 2. Characteristics of participants who completed dietary improvement**

254    **sessions**

|  | Baseline (n=12) | 6 months (n=12) | 12 months (n=12) | Follow-up (n=6) |
|---|---|---|---|---|
| **Height (cm)** | 167.4±1.2 | 167.4±1.2 | 167.4±1.2 | 166.5±1.3 |
| **WC (cm)** | 112.9±1. | | 109.9±1.5(0.18) | |
| **HC (cm)** | 125.5±1.2 | | 123.5± 1.6(0.45) | |
| **WHR** | | | | |
| **Female** | 0.88±0.02 | | 0.85±0.02(0.17) | |
| **Male** | 1.00±0 | | 0.98±0.04(0.54) | |
| **Weight (kg)** | 103.3±6.5 | 98.1±6.2(0.001) | 98.4±7.4(0.007) | 98.2±11.9(0.07) |
| **BMI (kg/m$^2$)** | 36.7±1.7 | 34.9±1.7(0.001) | 34.9±2.0(0.007) | 35.0±2.9(0.07) |

255    Values are presented as mean±SE for baseline and mean±SE (p-value) for

256    subsequent months. P-values by paired t-test are shown in parenthesis. WC: Waist

257    circumference, HC: Hip circumference, WHR: Waist to hip ratio

## Retention

Fourteen participants started the program. Participants who were unable to attend the regular sessions received make-up sessions. One participant dropped out after four weeks, and the other participant dropped out after eight weeks, yielding 85.7% retention. The reasons for leaving the study were as follows: too busy to focus on weight loss (n=1) and unsatisfying weight loss (n=1). During the six-month follow-up period, six out of 12 participants continued to weigh daily.

## Change in anthropometric measures

Anthropometric changes following the intervention are presented in Table 2 and Figures 4 and 5. At 12 months, six participants (50%) successfully lost >5% of their initial body weights (Fig 4). Of the unsuccessful participants (n=6), four participants did not reach 5% weight loss during the 12 months, whereas two participants regained lost weight after six months (Fig 4). On average, participants (n=12) achieved a significant weight loss of -5.4%±1.7% at 12 months (p <0.01) from the baseline (Fig 5). Cohen's d for 12 month weight loss was -0.21 indicating a small, negative change in weight. A large difference in weight loss outcome was observed among participants at 12 months (p<0.001); the top half (n=6) showed a significant loss from the baseline reaching -9.7±1.7%, whereas the bottom half (n=6) lost -1.0±1.4% (Fig 5). The difference between the two groups widened as time progressed from the start to 12 months, particularly during the last five months (Fig 5).

**Fig 4. Weight changes (%) of each participant at six, 12, and 18 months.** The data are ordered by magnitudes of weight loss at 12 months.

282  **Fig 5. Weight changes (%) of all, top-half and bottom-half groups over 12 months.**

283  Participants were divided into two groups of equal size (n=6 each) based on the weight

284  changes from baseline at 12 months. **p<0.01, ***p<0.001 at 12 months compared with

285  baseline.

## Change in dietary components

Changes in diets over 12 months based on FFQ and 24-hour dietary records are

shown in Fig 6. A significant increase in fiber density (p=0.02) but not protein density

was found from the FFQ. Twenty-four-hour records showed a significant improvement in

protein density in all 5 subsequent records when compared with the first month record,

whereas fiber density increased only at three months (p=0.02). Next, 24-hour records

were compared between the groups of the bottom and top half in weight loss at 12

months (Fig 7). While only one out of six records of the bottom half was inside the

weight maintenance box, four out of six records of the top half stayed within the weight

maintenance box. A significant difference in fiber density was observed between the top

half and bottom half (p=0.04) although there was no difference in protein density

(p=0.19) (Fig 7). There was no significant difference in protein density between the two

groups at any time point.

**Fig 6. Changes in protein and fiber density of the whole group assessed by FFQ**

**and 24-hour records**. Data are presented as mean. FFQ were collected at the baseline

(n=12) and the end of the trial (n=11). Twenty-four-hour records were collected at one –

seven months (n=12) and at 12 months (n=10). Changes in protein density and fiber

density were compared between baseline and 12 months in FFQ, and between the first

month and two-12 months in 24-hour records.  *p=0.02, ***p<0.001 in protein density;

§p=0.02 in fiber density by paired t-test.

**Fig 7. Changes in protein and fiber density of top and bottom half groups in**

**weight loss at 12 months.** Twenty-four-hour records conducted at one, two, three, five,

seven and 12 months. Dietary records were divided to two groups based on top half

(n=6) and bottom half (n=6) of weight loss at 12 months. Data are presented as mean.

§p=0.04 significant difference in fiber density between two groups by repeated measure

ANOVA when all time points of 24-hour records were combined (n=35 for each group).

# Correlation among weight loss and protein and fiber density

Correlations between diet and weight changes were tested to identify possible

determinants of weight loss success (Fig 8). A significant correlation was found between

weight loss at three months and weight loss at 12 months, suggesting participants who

lost weight during the initial three months continued to lose more (p=0.03, $r_p$=0.64). A

positive correlation was observed between protein and fiber densities at three months

(p=0.02, $r_p$=0.64). A positive correlation between protein density at three months and its

density at six months was also observed (p=0.002, $r_p$=0.80). No significant associations

were found between weight loss and either protein or fiber density at any months tested.

**Fig 8. Correlation matrix showing the interrelationship among weight loss and**

**protein and fiber density at three months, six months, and 12 months.** Positive

correlations are displayed in blue and negative correlations in red. *p<0.05, **p<0.01.

## Exit survey

The results of the exit survey at 12 months are shown in S3 Appendix. Questions one and three suggested that participants who completed the program found it beneficial for applying skills such as interpreting nutrition information to maintain healthier eating behaviors. A high score was received in program content with instructors being knowledgeable (question 5) and available to offer support (question 8). Finally, participants reported that the visual feedbacks, including the PF plot and weekly weight chart, were easy to understand and helpful (questions 9 and 10). The overall findings from the survey suggested that iDip equipped participants with valuable nutrition knowledge and skills, which enabled some individuals to adhere to dietary changes. However, the scores of questions 2 and 11 showed that the program was lacking in some aspects of satisfaction in overall engagement and individual goal achievements.

## Discussion

This non-randomized and single-arm trial demonstrated the feasibility of a new approach to weight loss that utilized two key visual feedback systems: the PF plot and the weekly weight chart. These quantitative feedbacks allowed participants to self-select dietary modifications and monitor their energy balance without calorie counting. iDip uniquely used no calorie counting, meal plans or recipes, meal replacements, or food exclusions. At 12 months, mean weight loss was -5.4±1.7% (p <0.01), and 50% of participants achieved >5% weight loss.

## Innovations

iDip was aimed to increase dietary flexibility and easily monitor energy balance, two shortcomings of many existing weight loss programs. To increase dietary flexibility, iDip provided nutrition education and individualized dietary feedback in the form of a PF plot to empower participants to develop a weight loss diet suitable to their taste preferences and lifestyle. Participants were able to create and advance a weight loss diet based on the PF plot. Participants were not given specific meal plans, meal replacement products, or recipes; rather, all dietary modifications were selected and implemented by individuals with guidance from individual counseling and educational sessions delivered by dietitians.  Importantly, no macronutrient or food group was excluded or limited in contrast to the practice commonly used in dieting programs, such as the Atkins diet. Also, iDip did not place any focus on the fat and carbohydrate ratios in a diet as modifying fat and carbohydrate ratios has been reported to be ineffective for long-term weight loss [10, 13]. Participants were encouraged to self-select to limit intake of foods that contain minimal fiber and protein, but they were not instructed to exclude any particular foods from their diet. Thus, all diets and macronutrient distributions differed by participants.

The iDip study participants significantly lost weight by creating their weight-loss diet based on feedback in the form of a PF plot. Previously in a 12-week university cafeteria study, we tested if providing nutrient information in a PF plot format could assist customers in making an informed, healthier choice [33]. In the study, no nutritional information was provided during weeks one-three and seven-nine, and a Nutrition Facts Panel and a PF plot were posted on identical menu

items, during weeks four-six and weeks 10-12, respectively. Sales data, the

primary outcome, showed that when the PF plot was displayed, the protein

content and protein + fiber content of purchased meals was significantly higher

than no posting periods, whereas the Nutrition Facts Panel had no effect. These

results have shown the PF plot's ease of understanding and its ability to improve

consumer food choices [33]. In the current study, all educational session

materials and individual diet feedback were provided in the form of a PF plot,

demonstrating the applicability of the PF plot to creating a weight-loss diet by

individual participants.

iDip utilized daily self-weighing on a Wi-Fi-enabled scale to provide easy and

reliable energy balance monitoring without calorie counting or daily food journals. Self-

monitoring is a critical factor in behavior changes for weight loss [43, 44]. Bertz et al.

showed daily self-weighing and weekly summary feedback can limit weight gain [45].

We provided a cumulative weight loss progress chart every week, eliminating the need

for calorie counting entirely and providing means for self-monitoring. The purpose of the

six 24-hour records administered was to evaluate progress in dietary improvement, not

to monitor energy balance.

iDip had a retention rate of 85.7%, indicating likeability of our program and its

flexible dietary approach. This high retention was achieved without monetary

compensation. Participants were given the Wi-Fi scale (value ~ $60) to keep after

completing the study. The dietary flexibility, self-selection of foods, and no need for daily

calorie counting may have at least in part contributed to the low attrition.

## Limitations

We chose the minimum number of participants to achieve the study objective of testing the feasibility of a novel dietary weight loss approach, not to test the robustness of the program as a clinical treatment option. Despite achieving a clinically significant mean percent weight loss of >5%, we observed some limitations. First, a success rate of 50% of participants achieving >5% weight loss at 12 months is not sufficient to be used as a reliable treatment program. Also, a greater magnitude of weight loss will be needed for iDip to become a clinical treatment option. Second, the follow-up period (six months) was too short to assess long-term sustainability. At 18 months, 50% of the cohort was lost to follow-up and were no longer weighing regularly, suggesting that half of the participants regarded daily weighing as a requirement of the program, not an essential part of weight management. Among the remaining participants, not all were successful in maintaining their weight between 12 and 18 months, indicating the dietary changes made during the program were not sustained in some. iDip aimed to improve long-term sustainability through non-exclusion of certain food groups and self-selected approach; thus, future studies will include a longer follow-up period with better implementation of sustainable dietary changes and daily weighing for monitoring energy balance. Third, body composition changes were not measured or tracked throughout the study. However, with the focus on increased protein density, we expect little muscle mass loss. Lastly, changes in health parameters, such as blood lipid and glucose levels and changes in medications associated with obesity and other comorbidities, were not tracked. Although a randomized and controlled trial is standard for clinical studies, a large-scale, randomized, controlled weight-loss trial has shown that the weight of the

placebo group was unchanged during one year, whereas the metformin and lifestyle

modification groups lost significant weight [46]. Therefore, this landmark study suggests

that the lack of a no-treatment control group would not pose a major limitation in

interpreting the weight loss outcome of our study.

**Future directions**

This study showed proof of concept for the non-exclusion of certain food groups

and self-selected approach to dietary weight loss. Based on this success, steps of

future studies will include: 1) increase the magnitude of mean percent weight loss; 2)

improve the overall success rate of participants achieving >5% weight loss; 3) monitor

body composition to minimize loss of skeletal muscle mass; 4) calculate intervention

cost; and 5) collaborate with other health professionals to track health markers, such as

blood lipids, and address medical and psychological barriers to weight loss. In the long

term, a large-scale, multi-center trial with a comparable control group is required to test

general applicability and effectiveness of the program as a treatment option for obesity.

**Conclusions**

In conclusion, this non-randomized and single-arm trial has shown the feasibility

of the informed decision-making approach for dietary weight loss by utilizing two forms

of quantitative visual feedback. One was the PF plot to monitor protein per energy and

fiber per energy intake, and the other was the daily weight chart to monitor energy

balance. Participants understood these visual weight management tools well.

Participants significantly improved fiber density from the baseline FFQ to the final FFQ,

and protein density significantly increased from the baseline 24-hour record at several

time points in the study. The participants achieved a mean percent weight loss of -

5.4±1.7%, and half of them successfully lost ≥5% baseline weight. The dietary flexibility

and self-selection of foods may have contributed to a low attrition rate. Future studies

are warranted to improve success rate, magnitude of weight loss and long-term

sustainability.

# References

1. Popkin BM. Global nutrition dynamics: the world is shifting rapidly toward a diet

linked with noncommunicable diseases. Am J Clin Nutr. 2006;84(2):289-98. Epub

2006/08/10. doi: 10.1093/ajcn/84.1.289. PubMed PMID: 16895874.

2. Jensen MD, Ryan DH, Apovian CM, Ard JD, Comuzzie AG, Donato KA, et al.

2013 AHA/ACC/TOS Guideline for the Management of Overweight and Obesity

in Adults A Report of the American College of Cardiology/American Heart

Association Task Force on Practice Guidelines and The Obesity Society. Journal

of the American College of Cardiology. 2014;63(25):2985-3025. doi:

10.1016/j.jacc.2013.11.004. PubMed PMID: WOS:000338435300004.

3. National Institute of Diabetes and Digestive and Kidney Diseases. Overweight &

Obesity Statistics 2019. Available from: https://www.niddk.nih.gov/health-

information/health-statistics/overweight-obesity.

4. Biener A, Cawley J, Meyerhoefer C. The Impact of Obesity on Medical Care

Costs and Labor Market Outcomes in the US. Clin Chem. 2018;64(1):108-17.

Epub 2017/11/04. doi: 10.1373/clinchem.2017.272450. PubMed PMID:

29097513.

5. Look AHEAD Research Group. Cardiovascular effects of intensive lifestyle

intervention in type 2 diabetes. N Engl J Med. 2013;369(2):145-54. Epub

2013/06/26. doi: 10.1056/NEJMoa1212914. PubMed PMID: 23796131; PubMed

Central PMCID: PMCPMC3791615.

6. Finley CE, Barlow CE, Greenway FL, Rock CL, Rolls BJ, Blair SN. Retention

rates and weight loss in a commercial weight loss program. Int J Obes (Lond).

2007;31(2):292-8. Epub 2006/06/07. doi: 10.1038/sj.ijo.0803395. PubMed PMID:

16755283.

7. Mitchell C, Stuart RB. Effect of self-efficacy on dropout from obesity treatment. J

Consult Clin Psychol. 1984;52(6):1100-1. Epub 1984/12/01. doi: 10.1037//0022-

006x.52.6.1100. PubMed PMID: 6520280.

8. Volkmar FR, Stunkard AJ, Woolston J, Bailey RA. High attrition rates in

commercial weight reduction programs. Arch Intern Med. 1981;141(4):426-8.

Epub 1981/03/01. PubMed PMID: 7212881.

9. Gardner CD, Kiazand A, Alhassan S, Kim S, Stafford RS, Balise RR, et al.

Comparison of the Atkins, Zone, Ornish, and LEARN diets for change in weight

and related risk factors among overweight premenopausal women: the A TO Z

Weight Loss Study: a randomized trial. JAMA. 2007;297(9):969-77. Epub

2007/03/08. doi: 10.1001/jama.297.9.969. PubMed PMID: 17341711.

10. Dansinger ML, Gleason JA, Griffith JL, Selker HP, Schaefer EJ. Comparison of

the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart

disease risk reduction: a randomized trial. JAMA. 2005;293(1):43-53. Epub

2005/01/06. doi: 10.1001/jama.293.1.43. PubMed PMID: 15632335.

11. Lowe MR, Miller-Kovach K, Phelan S. Weight-loss maintenance in overweight

individuals one to five years following successful completion of a commercial

weight loss program. Int J Obes Relat Metab Disord. 2001;25(3):325-31. Epub

2001/04/25. doi: 10.1038/sj.ijo.0801521. PubMed PMID: 11319628.

12. Thomas JG, Raynor HA, Bond DS, Luke AK, Cardoso CC, Foster GD, et al.

Weight loss in Weight Watchers Online with and without an activity tracking

device compared to control: A randomized trial. Obesity (Silver Spring).

2017;25(6):1014-21. Epub 2017/04/25. doi: 10.1002/oby.21846. PubMed PMID:

28437597.

13. Foster GD, Wyatt HR, Hill JO, Makris AP, Rosenbaum DL, Brill C, et al. Weight

and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet:

a randomized trial. Ann Intern Med. 2010;153(3):147-57. Epub 2010/08/04. doi:

10.7326/0003-4819-153-3-201008030-00005. PubMed PMID: 20679559;

PubMed Central PMCID: PMCPMC2949959.

14. Hall KD, Kahan S. Maintenance of Lost Weight and Long-Term Management of

Obesity. Med Clin North Am. 2018;102(1):183-97. Epub 2017/11/21. doi:

10.1016/j.mcna.2017.08.012. PubMed PMID: 29156185; PubMed Central

PMCID: PMCPMC5764193.

15. U.S. Department of Health and Human Services and U.S. Department of

Agriculture. 2015–2020 Dietary Guidelines for Americans. Washington, DC:: US

Dept of Health and Human Services; Dec 2015 [April 2020]. 8th

Edition:[Available from: http://health.gov/dietaryguidelines/2015/guidelines/.

16. Raynor HA, Champagne CM. Position of the Academy of Nutrition and Dietetics:

Interventions for the Treatment of Overweight and Obesity in Adults. J Acad Nutr

Diet. 2016;116(1):129-47. Epub 2016/01/01. doi: 10.1016/j.jand.2015.10.031.

PubMed PMID: 26718656.

17. Archundia Herrera MC, Chan CB. Narrative Review of New Methods for

Assessing Food and Energy Intake. Nutrients. 2018;10(8). Epub 2018/08/15. doi:

10.3390/nu10081064. PubMed PMID: 30103401; PubMed Central PMCID:

PMCPMC6116053.

18. Carels RA, Harper J, Konrad K. Qualitative perceptions and caloric estimations of

healthy and unhealthy foods by behavioral weight loss participants. Appetite.

2006;46(2):199-206. Epub 2006/02/10. doi: 10.1016/j.appet.2005.12.002.

PubMed PMID: 16466830.

19. Cordeiro F, Epstein DA, Thomaz E, Bales E, Jagannathan AK, Abowd GD, et al.

Barriers and Negative Nudges: Exploring Challenges in Food Journaling. Proc

SIGCHI Conf Hum Factor Comput Syst. 2015;2015:1159-62. Epub 2016/02/20.

doi: 10.1145/2702123.2702155. PubMed PMID: 26894233; PubMed Central

PMCID: PMCPMC4755274.

20. Kong A, Beresford SAA, Alfano CM, Foster-Schubert KE, Neuhouser ML,

Johnson DB, et al. Self-monitoring and eating-related behaviors are associated

with 12-month weight loss in postmenopausal overweight-to-obese women. J

Acad Nutr Diet. 2012;112(9):1428-35. Epub 2012/07/17. doi:

10.1016/j.jand.2012.05.014. PubMed PMID: 22795495; PubMed Central PMCID:

PMCPMC3432675.

21. Hwang KO, Ning J, Trickey AW, Sciamanna CN. Website Usage and Weight

Loss in a Free Commercial Online Weight Loss Program: Retrospective Cohort

Study. Journal of Medical Internet Research. 2013;15(1). doi: ARTN

e1110.2196/jmir.2195. PubMed PMID: WOS:000315113200018.

22. Hoffer LJ, Bistrian BR, Young VR, Blackburn GL, Matthews DE. Metabolic effects

of very low calorie weight reduction diets. J Clin Invest. 1984;73(3):750-8. Epub

1984/03/01. doi: 10.1172/JCI111268. PubMed PMID: 6707202; PubMed Central

PMCID: PMCPMC425077.

23. Wycherley TP, Moran LJ, Clifton PM, Noakes M, Brinkworth GD. Effects of

energy-restricted high-protein, low-fat compared with standard-protein, low-fat

diets: a meta-analysis of randomized controlled trials. Am J Clin Nutr.

2012;96(6):1281-98. Epub 2012/10/26. doi: 10.3945/ajcn.112.044321. PubMed

PMID: 23097268.

24. Ard JD, Lewis KH, Rothberg A, Auriemma A, Coburn SL, Cohen SS, et al.

Effectiveness of a Total Meal Replacement Program (OPTIFAST Program) on

Weight Loss: Results from the OPTIWIN Study. Obesity (Silver Spring).

2019;27(1):22-9. Epub 2018/11/14. doi: 10.1002/oby.22303. PubMed PMID:

30421863; PubMed Central PMCID: PMCPMC6587830.

25. Franz MJ, VanWormer JJ, Crain AL, Boucher JL, Histon T, Caplan W, et al.

Weight-loss outcomes: a systematic review and meta-analysis of weight-loss

clinical trials with a minimum 1-year follow-up. J Am Diet Assoc.

2007;107(10):1755-67. Epub 2007/10/02. doi: 10.1016/j.jada.2007.07.017.

PubMed PMID: 17904936.

26. Marinilli Pinto A, Gorin AA, Raynor HA, Tate DF, Fava JL, Wing RR. Successful

weight-loss maintenance in relation to method of weight loss. Obesity (Silver

Spring). 2008;16(11):2456-61. Epub 2008/08/23. doi: 10.1038/oby.2008.364.

PubMed PMID: 18719680; PubMed Central PMCID: PMCPMC2666007.

27. Diabetes Prevention Program Research Group. The 10-year cost-effectiveness

of lifestyle intervention or metformin for diabetes prevention: an intent-to-treat

analysis of the DPP/DPPOS. Diabetes Care. 2012;35(4):723-30. Epub

2012/03/24. doi: 10.2337/dc11-1468. PubMed PMID: 22442395; PubMed Central

PMCID: PMCPMC3308273.

28. Jones BJ, Bloom SR. The New Era of Drug Therapy for Obesity: The Evidence

and the Expectations. Drugs. 2015;75(9):935-45. Epub 2015/05/20. doi:

10.1007/s40265-015-0410-1. PubMed PMID: 25985865; PubMed Central

PMCID: PMCPMC4464860.

29. Schauer PR, Bhatt DL, Kirwan JP, Wolski K, Brethauer SA, Navaneethan SD, et

al. Bariatric Surgery versus Intensive Medical Therapy for Diabetes-3-Year

Outcomes. New Engl J Med. 2014;370(21):2002-13. doi:

10.1056/NEJMoa1401329. PubMed PMID: WOS:000336125500008.

30. Doble B, Wordsworth S, Rogers CA, Welbourn R, Byrne J, Blazeby JM, et al.

What Are the Real Procedural Costs of Bariatric Surgery? A Systematic

Literature Review of Published Cost Analyses. Obes Surg. 2017;27(8):2179-92.

Epub 2017/05/28. doi: 10.1007/s11695-017-2749-8. PubMed PMID: 28550438;

PubMed Central PMCID: PMCPMC5509820.

31. Johns DJ, Hartmann-Boyce J, Jebb SA, Aveyard P, Behavioural Weight

Management Review G. Diet or exercise interventions vs combined behavioral

weight management programs: a systematic review and meta-analysis of direct

comparisons. J Acad Nutr Diet. 2014;114(10):1557-68. Epub 2014/09/27. doi:

10.1016/j.jand.2014.07.005. PubMed PMID: 25257365; PubMed Central PMCID:

PMCPMC4180002.

32. Borek AJ, Abraham C, Greaves CJ, Tarrant M. Group-Based Diet and Physical

Activity Weight-Loss Interventions: A Systematic Review and Meta-Analysis of

Randomised Controlled Trials. Appl Psychol Health Well Being. 2018;10(1):62-

86. Epub 2018/02/16. doi: 10.1111/aphw.12121. PubMed PMID: 29446541.

33. Pratt NS, Ellison BD, Benjamin AS, Nakamura MT. Improvements in recall and

food choices using a graphical method to deliver information of select nutrients.

Nutrition Research. 2016;36(1):44-56. doi: 10.1016/j.nutres.2015.10.009.

PubMed PMID: WOS:000368755600006.

34. Atkinson RL, Dietz WH, Foreyt JP, Goodwin NJ, Hill JO, Hirsch J, et al. Very-

Low-Calorie Diets. Jama-Journal of the American Medical Association.

1993;270(8):967-74. doi: DOI 10.1001/jama.1993.03510080071034. PubMed

PMID: WOS:A1993LT56700028.

35. Larsen TM, Dalskov SM, van Baak M, Jebb SA, Papadaki A, Pfeiffer AFH, et al.

Diets with High or Low Protein Content and Glycemic Index for Weight-Loss

Maintenance. New England Journal of Medicine. 2010;363(22):2102-13. doi: DOI

10.1056/NEJMoa1007137. PubMed PMID: WOS:000284587800005.

36. Liu SM, Willett WC, Manson JE, Hu FB, Rosner B, Colditz G. Relation between

changes in intakes of dietary fiber and grain products and changes in weight and

development of obesity among middle-aged women. American Journal of Clinical

Nutrition. 2003;78(5):920-7. PubMed PMID: WOS:000186146500006.

37. Koh-Banerjee P, Franz MV, Sampson L, Liu SM, Jacobs DR, Spiegelman D, et

al. Changes in whole-grain, bran, and cereal fiber consumption in relation to 8-y

weight gain among men. American Journal of Clinical Nutrition. 2004;80(5):1237-

45. PubMed PMID: WOS:000225036000019.

38. Nordmann AJ, Nordmann A, Briel M, Keller U, Yancy WS, Jr., Brehm BJ, et al.

Effects of low-carbohydrate vs low-fat diets on weight loss and cardiovascular

risk factors: a meta-analysis of randomized controlled trials. Arch Intern Med.

2006;166(3):285-93. Epub 2006/02/16. doi: 10.1001/archinte.166.3.285. PubMed

PMID: 16476868.

39. Stern L, Iqbal N, Seshadri P, Chicano KL, Daily DA, McGrory J, et al. The effects

of low-carbohydrate versus conventional weight loss diets in severely obese

adults: one-year follow-up of a randomized trial. Ann Intern Med.

2004;140(10):778-85. Epub 2004/05/19. doi: 10.7326/0003-4819-140-10-

200405180-00007. PubMed PMID: 15148064.

40. Foster GD, Wyatt HR, Hill JO, McGuckin BG, Brill C, Mohammed BS, et al. A

randomized trial of a low-carbohydrate diet for obesity. N Engl J Med.

2003;348(21):2082-90. Epub 2003/05/23. doi: 10.1056/NEJMoa022207. PubMed

PMID: 12761365.

41. Mulligan AA, Luben RN, Bhaniani A, Parry-Smith DJ, O'Connor L, Khawaja AP,

et al. A new tool for converting food frequency questionnaire data into nutrient

and food group values: FETA research methods and availability. Bmj Open.

2014;4(3). doi: 10.1136/bmjopen-2013-004503. PubMed PMID:

WOS:000333921300056.

42. Blackburn G. Effect of Degree of Weight-Loss Benefits. Obesity Research.

1995;3:S211-S6. doi: DOI 10.1002/j.1550-8528.1995.tb00466.x. PubMed PMID:

WOS:A1995RZ53500018.

43. Van Dorsten B, Lindley EM. Cognitive and behavioral approaches in the

treatment of obesity. Med Clin North Am. 2011;95(5):971-88. Epub 2011/08/23.

doi: 10.1016/j.mcna.2011.06.008. PubMed PMID: 21855703.

44. Brown JD, Buscemi J, Milsom V, Malcolm R, O'Neil PM. Effects on

cardiovascular risk factors of weight losses limited to 5-10 %. Translational

Behavioral Medicine. 2016;6(3):339-46. doi: 10.1007/s13142-015-0353-9.

PubMed PMID: WOS:000385261500002.

45. Bertz F, Pacanowski CR, Levitsky DA. Frequent Self-Weighing with Electronic

Graphic Feedback to Prevent Age-Related Weight Gain in Young Adults. Obesity

(Silver Spring). 2015;23(10):2009-14. Epub 2015/09/29. doi: 10.1002/oby.21211.

PubMed PMID: 26414563; PubMed Central PMCID: PMCPMC4751025.

46. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA,

et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or

metformin. N Engl J Med. 2002;346(6):393-403. Epub 2002/02/08. doi:

10.1056/NEJMoa012512. PubMed PMID: 11832527; PubMed Central PMCID:

PMCPMC1370926.

## Supporting information

**S1 Appendix. iDip educational session topics**

**S2 Appendix. Food Frequency Questionnaire**

**S3 Appendix. Exit survey outcomes**

**S4 Appendix. CONSORT for pilot and feasibility study**

**S5 Appendix. TIDieR Checklist**

**S6 Appendix. Clinical trial protocol**