Independent and combined effects of nutrition and sanitation interventions on enteric pathogen carriage and child growth in rural Cambodia: a factorial cluster-randomised controlled trial

Amanda Lai[1,2], Irene Velez[3], Ramya Ambikapathi[4], Krisna Seng[3], Karen Levy[5], Erin Kowalsky[1,2], David Holcomb[1], Konstantinos T. Konstantinidis[2], Oliver Cumming[6], Joe Brown[1*]

[1] Department of Environmental Sciences and Engineering, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA
[2] School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, Georgia, USA
[3] MSI, A Tetra Tech Company, Arlington, Virginia, USA
[4] Department of Public Health, Purdue University, West Lafayette, Indiana, USA
[5] Department of Environmental and Occupational Health Sciences, University of Washington, Seattle, Washington, USA
[6] Department of Disease Control, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, UK

* Corresponding author: Joe Brown, joebrown@unc.edu

Summary

Background

Childhood exposure to enteric pathogens associated with poor sanitation contributes to
undernutrition, associated with adverse effects later in life. This trial assessed the independent
and combined effects of nutrition and sanitation interventions on child growth outcomes and
enteric pathogen infection in rural Cambodia, where the prevalence of childhood stunting
remains high.

Methods

We conducted a factorial cluster-randomised controlled trial of 4,015 households with 4,124
children (1-28 months of age at endline) across three rural provinces in Cambodia. Fifty-five
communes (clusters) were randomly assigned to a control arm or one of three treatments: a
nutrition-only arm, a sanitation-only arm, and a combined nutrition and sanitation arm receiving
both treatments. The primary outcome was length-for-age Z-score (LAZ); other outcomes
included weight-for-age Z-score (WAZ), weight-for-length Z-score (WLZ), stunting, wasting,
underweight, and caregiver-reported diarrhoea. We assayed stool specimens from a subset of all
children (n = 1,620) for 27 enteric pathogens (14 bacteria, 6 viruses, 3 protozoa, and 4 soil-
transmitted helminths) and estimated effects of interventions on enteric pathogen detection and
density. Analysis was by intention-to-treat. The trial was pre-registered with ISRCTN Registry
(ISRCTN77820875).

Findings

Self-reported adherence was high for the nutrition intervention but uptake was low for sanitation.
Compared with a mean LAZ of -1.04 (SD 1.2) in the control arm, children in the nutrition-only
arm (LAZ +0.08, 95% CI -0.01-0.18) and combined nutrition and sanitation arm (LAZ +0.10,
95% CI 0.01-0.20) experienced greater linear growth; there were no measurable differences in
LAZ in the sanitation-only arm (LAZ -0.05, 95% CI -0.16-0.05). We found no effect of any
intervention (delivered independently or combined) on either enteric pathogen frequency or
pathogen load in stool. Compared with a mean WAZ of -1.05 (SD 1.1) in the control arm,
children in the nutrition-only arm (WAZ +0.10, 95% CI 0.00-0.19) and combined intervention
arm (WAZ +0.11, 95% CI 0.03-0.20) were heavier for their age; there was no difference in WAZ
in the sanitation-only arm. There were no differences between arms in prevalence of stunting,
wasting, underweight status, one-week period prevalence of diarrhoea, pathogen prevalence, or
pathogen density in stool.

Interpretation

Improvements in child growth in nutrition and combined nutrition and sanitation arms are
consistent with previous efficacy trials of combined nutrition and sanitation interventions. We
found no evidence that the sanitation intervention alone improved child growth or reduced
enteric pathogen detection, having achieved only modest changes in access and use.

Funding

United States Agency for International Development (USAID), contracts AID-OAA-M-13-
00017 and AID-OAA-TO-16-00016. The contents of this publication are the sole responsibility
of the authors and do not necessarily reflect the views of USAID or the United States
Government.

Introduction

Childhood undernutrition is associated with higher susceptibility to infectious disease, reduced cognitive function, and various adverse outcomes later in life[1]. Growth faltering is an effect of chronic undernutrition and tends to manifest in a child's first two years[2]. Many studies have focused on improving infant and child nutrition to achieve better growth outcomes[3,4]. However, nutrition interventions alone have not been successful in eliminating stunting, suggesting that broader interventions addressing other important factors are needed alongside exclusive breastfeeding and improved nutritional intake[5].

Reducing early childhood exposure to enteric pathogens through safe water, sanitation, and hygiene (WASH) may complement other interventions by reducing diarrhoeal diseases and environmental enteric dysfunction (EED)[6]—both of which can impact early childhood growth and development[7]. Observational studies have found strong associations between child growth faltering and poor access to sanitation[8]. However, recent randomised controlled trials (RCTs) in Zimbabwe[9], Bangladesh[3], and Kenya[4] that delivered standalone household-level sanitation interventions (not coupled with other nutrition or hygiene interventions) were not found to improve child growth.

The community-led total sanitation (CLTS) framework is an approach to ending open defecation (OD) through behavioural change and collective action rather than through the provision of hardware and materials. CLTS and other rural promotion-based interventions shift the focus from individual and household sanitation practices to a community-level concern over OD by triggering collective behaviour change through powerful emotional drivers such as shame and disgust, as well as positive motivators such as improved health, dignity, and pride. Observational studies in Cambodia[10] and Ecuador[11] found higher community-level sanitation coverage to be associated with reduced prevalence of stunting. Despite this, recent RCTs employing promotion-based interventions have found mixed effects on child growth. One trial was found to be successful in improving child growth in Mali[12], but this effect was not observed in other trials elsewhere[13,14].

This study contributes to a growing body of literature on the impact of combined nutrition and sanitation interventions on early child growth, caregiver-reported diarrhoea, and detection and quantification of enteric pathogens in stool as a proxy for enteric infection. While diarrhoea has been widely used as a primary outcome measure in WASH studies[3,4,12,15], recent studies have used stool-based detection of enteric pathogens[15] and anthropometry measurements[3,4,9] as primary outcomes that are more objectively measurable and may also broadly indicate health status by capturing cumulative effects of exposures via EED[6]. We used a factorial cluster-randomised controlled trial (cRCT) to assess the independent and combined effects of nutrition and sanitation interventions delivered in the context of a large-scale, USAID-funded rural nutrition and sanitation/hygiene program in Cambodia. We hypothesised that children receiving both sanitation and nutrition interventions would have increased linear growth compared with children from control areas lacking these interventions. We further hypothesised that combined nutrition and sanitation interventions would lead to synergistic improvements in linear growth beyond what was realised in either standalone intervention arm. The hypothesised pathway for

106   these effects, consistent with secondary outcome measures, was reduced enteric pathogen
107   frequency and enteric pathogen load in stools (Figure 1).

Methods
108

Study design and participants
109
We implemented a two-by-two factorial cRCT in rural communes in three provinces in
Cambodia: Battambang, Pursat, and Siem Reap. The communes targeted by the program were
selected based on two criteria: communes where at least 30% of the population was living below
the poverty line according to the 2011 Cambodia Ministry of Planning's Commune Database;
and communes where latrine subsidies were not then in place. This study is reported per the
Consolidated Standards of Reporting Trials (CONSORT) guideline (see Supplementary Material
for CONSORT checklist).
117

Randomisation and masking
118
In 2015, prior to the start of project activities, we randomly assigned communes to one of three
treatment arms (nutrition only, sanitation/hygiene only, combined nutrition and
sanitation/hygiene) or control arm using a random number generator with reproducible seed in
Stata 13 (Stata, College Station, TX). Randomisation was conducted at the commune level to
limit the risk of contamination between study arms and all villages within each commune
received the assigned intervention. Following randomisation, three communes were dropped
from the trial due to objections from the local governments of overlap with other current
programming. This resulted in 55 communes with treatment arms of different sizes: 11
communes in nutrition-only arm; 13 in sanitation-only arm; 12 in combined-intervention arm;
and 19 in control arm (Figure 2). The trial enrolled primary caregivers with a child who was born
after intervention implementation began (up to 28 months prior) and who had lived in the
commune during the child's entire life, resulting in a participant population of children 1-28
months old. Neither participants nor field staff were masked to treatment status due to the nature
of the interventions, but data collection teams were blinded to the arm assignment and number of
treatment arms.
134

Procedures
135
The interventions were delivered in the 36 intervention communes over the course of two years,
between 2015-2017, while the remaining 19 control communes were unexposed to the
programmes. Two international non-governmental organizations—Save the Children and SNV—
provided programmatic implementation and coordinated activities with local governments. The
nutrition interventions included complementary feeding activities and education through
community-based growth promotion sessions; caregiver groups; home visits; and conditional
cash transfers (CCTs) linked to the utilization of key health and nutrition services focusing on
first 1,000 days of life. The sanitation interventions consisted primarily of CLTS as it was
delivered here, latrine vouchers coupled with supply-side support for sanitation and hygiene
products, and social behaviours change communications (SBCC). Intervention activities and
frequency are summarized in Table 1, and additional details about the interventions are described
in the Supplementary Material.
148

The survey was communicated in the Khmer language to assess household and child-level risk
factors of children under 28 months of age. Enumerators completed in-home interviews with the

primary caregiver of children in the household about basic household member information;
breastfeeding and nutrition of children up to age 28 months; number of pregnancies and child
births of the caregiver; intervention exposure and participation; household WASH conditions and
practices; and household assets/characteristics to construct wealth scores (excluding WASH
variables). We also documented process evaluation (PE) indicators based on self-reported receipt
of, and participation in, intervention activities to assess intervention fidelty and adherence,
respectively. We attempted to collect a stool sample from each child and randomly selected a
subset of stools for analysis by reverse-transcription quantitative polymerase chain reaction (RT-
qPCR) of 30 enteric pathogen genes using a custom-developed TaqMan Array Card (TAC;
ThermoFisher Scientific, Waltham, VA), as described in the Supplementary Material.
161

Outcomes
The primary outcome was length-for-age Z-score (LAZ). For children 1-24 months in age, we
measured recumbent length; for children 24-28 months in age, we measured standing height.
Herein, "length" will be inclusive of both recumbent length and standing height. Secondary
outcomes included weight-for-age Z-score (WAZ); weight-for-length Z-score (WLZ); proportion
of children stunted (LAZ<-2), underweight (WAZ<-2), and wasted (WLZ<-2); caregiver
reported diarrhoea; all-cause mortality; and enteric pathogen detection and quantification in
stool. Child length and weight were measured by trained paired enumerators following
guidelines from the National Health and Nutrition Examination Survey (NHANES)[16]. Final
measurements took place in August 2019, 28 months after the end of the roll-out period. Data
collection was completed by KHANA Centre for Population Health Research, with oversight and
support from Management Systems International (MSI). Data collection details, measurement
protocols, and PE indicators are further described in Supplementary Material.
175

We assessed enteric pathogens as the prevalence of individual gene targets, the number of co-
detected pathogens, and enteric pathogen-associated gene copies per gram of stool based on PCR
quantification cycle (Cq) and standard curves. *E.coli* pathotypes were defined as: EAEC (*aaiC*,
or *aatA*, or both), atypical EPEC (*eae* without *bfpA*, *stx1*, and *stx2*), typical EPEC (*bfpA*), ETEC
(*STh*, *STp*, or *LT*), and STEC (*eae* without *bfpA* and with *stx1*, *stx2*, or both). Details on nucleic
acid extraction and molecular assaying are described in Supplementary Material.
182

Statistical analysis
We performed an intention-to-treat (ITT) analysis for all outcomes using generalised estimating
equations (GEE) with robust standard errors to account for clustering at the village level. We did
not consider pre-intervention covariate balance[17] but present secondary analyses adjusted for pre-
specified pre-intervention covariates in the Supplementary Material. Outcomes in each treatment
arm were compared to the control arm and between standalone treatment arms and the combined
treatment arm. We used linear regression to estimate mean differences in LAZ, WAZ, WLZ, and
$\log_{10}$-transformed pathogen gene target densities and used log-linear Poisson regression to
estimate the prevalence ratio (PR) between arms for nutritional status (stunting, wasting, and
underweight), diarrhoea, and overall mortality. Enteric pathogen gene outcomes were
dichotomised, with positive detections defined by a Cq <35[18], and Poisson regression was used
to estimate PRs for individual pathogens detected in stool. We further estimated the incidence
rate ratio (IRR) of co-detected pathogens (total and in subgroups by bacteria, viruses, protozoa,
and STHs) using negative binomial regression. We did not adjust for multiple comparisons for

growth, diarrhoea, or mortality outcomes[3,19], but we did apply the Benjamini-Hochberg
procedure to control the false discovery rate within analyses of multiple enteric pathogen
outcomes[20]. Details on power calculations are included in Supplementary Material.
200
Ethics
The study received approval from the National Ethics Committee for Health Research in the
Cambodian Ministry of Health, Georgia Institute of Technology, and New England Institutional
Review Board. Prior to any data collection, the trial was explained to participants in the Khmer
language. Written and verbal consent were obtained prior to administering the surveys and
anthropometry measurements. The trial was pre-registered with ISRCTN Registry
(ISRCTN77820875).

Results
Among 82 presumptively eligible communes, the provincial governments in 27 declined to
participate. Ultimately, the evaluation included 55 communes randomly assigned to one of three
treatment arms (n=36 communes) or control arm (n=19 communes); the control arm was
relatively oversized to enhance statistical efficiency of multiple hypothesis testing[21]. Figure 2
shows the trial profile by intervention subgroups. 4,015 households participated in endline
surveys; 4,005 households were included in these analyses (10 were excluded due to incomplete
surveys), and 4,124 children had anthropometry measures taken.
216
Household and caregiver characteristics were mostly similar across treatment and control groups
(Table 2). Primary caregivers in the control group reported lower levels of primary school
attendance compared to the treatment groups, but paternal primary school attendance was
similar. Households in the nutrition-only and sanitation-only groups had higher wealth index
scores compared to households in the combined intervention and control groups. The control
group had a higher prevalence of improved water source as their main source of drinking water
compared to the treatment groups.
224
Nutrition intervention fidelity was high, with households in the nutrition-only and combined-
intervention arms reporting significantly higher participation in these activities compared to the
sanitation-only and control groups (Table 3). Approximately 60% of households in the nutrition-
only and combined-intervention arms reported participating in at least four of the eight nutrition
intervention activities, compared to 4% in the sanitation-only and control arms. Conversely,
sanitation intervention fidelity was very low, with only 6% of households in the sanitation-only
and control arms reporting participation in any CLTS activity, compared to 14% of households
in the nutrition-only arm and 25% in the combined-intervention arm.
233
More households in the control arm (70%) had an improved water source as their main source of
drinking water, an indicator of nutrition intervention adherence, compared to other arms
(approximately 60% in other arms; Table 4). The combined intervention arm had greater access
to improved sanitation facilities (61%) compared to the nutrition-only (55%), sanitation-only
(51%), and control (52%) arms. OD (self-reported) was practiced less in the combined
intervention arm (7%) compared to the nutrition-only (14%), sanitation-only (16%), and control
(16%) arms. Notably, the sanitation-only arm experienced a significantly larger increase in
sanitation coverage (+25 percentage points [pp]) compared to all other arms (+14pp in nutrition-

only arm, +19pp in combined and control arms), though sanitation gains across all arms were
evident in the intervention period, reflecting a strong secular trend of sanitation expansion that
has been widely documented in rural Cambodia[8,10,22,23]. Additional intervention adherence
indicators related to environmental hygiene are reported in the Supplementary Material.
246
247   Primary and secondary outcomes
248   Mean LAZ in the control arm was -1.04 (SD 1.20). Compared with control, children in the
249   nutrition-only arm were longer by a mean of 0.08 LAZ (95% CI -0.01, 0.18), and children in the
250   combined-intervention arm were longer by 0.10 LAZ (95% CI 0.01, 0.20), although these
251   differences were not observed in the adjusted analyses (
252

253   Table 5, Figure 3, Supplementary Material). Children in the nutrition-only arm and combined-
254   intervention arm were heavier than children in the control arm by a mean of 0.10 WAZ (95% CI
255   0.00, 0.19) and 0.11 (95% CI 0.03, 0.20), respectively. These differences were slightly attenuated
256   in the adjusted analyses (
257

Table 5, Supplementary Material). No differences were observed between the control arm and intervention arms in terms of WLZ. Children in the combined intervention arm were also longer and heavier, on average, than children in the sanitation-only arm by 0.16 LAZ (95% CI 0.04, 0.27) and 0.10 WAZ (95% CI 0.01, 0.20), respectively. LAZ and WAZ were similar between children in the nutrition-only and combined intervention arms.

Compared with the control arm, none of the intervention arms differed in the prevalence of children who experienced stunting, wasting, diarrhoea (7-day recall), or mortality (Table 6). However, the combined intervention reduced underweight prevalence by 18% (PR 0.82, 95% CI 0.68, 0.99) relative to the control arm. Although the combined intervention did not significantly impact stunting prevalence compared with the control arm or the nutrition-only arm, the sanitation-only arm was associated with a 20% increase (PR 1.2, 95% CI 1.0, 1.5) in the prevalence of both stunting and underweight status when compared to the combined intervention. All associations with stunting and underweight were attenuated in adjusted analyses (Supplementary Material).

Enteric pathogen results

We assessed enteric pathogen-associated gene targets in 1,620 randomly selected stools that demonstrated acceptable amplification (of 4,114 stools total, see Supplementary Material): 305 from the nutrition arm, 333 from the sanitation arm, 438 from the combined-intervention arm, and 544 from the control arm. We detected at least one bacterial gene in 87% of all samples, at least one viral gene in 49% of samples, at least one protozoan gene in 20% of samples, and at least one STH gene in 2% of samples. Enteroaggregative *E. coli* (EAEC), enteric pathogenic *E. coli* (EPEC), enterovirus, *Campylobacter* spp., and enterotoxigenic *E. coli* (ETEC) were the most prevalent pathogens (Table 7). We detected a mean 2.2 bacterial genes (out of 9), 0.59 viral genes (out of 6), 0.21 protozoan genes (out of 4), and 0.03 STH genes (out of 4) in each sample. We found no differences in the rate of bacterial, viral, protozoan, or STH gene co-detection between the control arm and any treatment arm or between the combined arm and the standalone intervention arms (Table 8). Prevalence increased with age for many pathogens (aEPEC, ETEC, *Shigella*/EIEC (*ipah*), STEC, adenovirus, *Giardia*), while prevalence peaked for children 9-17 months for other pathogens (*Campylobacter* spp., *C.diff*, EAEC, *Salmonella* spp.; Supplementary Material).

Examining prevalence of specific targets compared to the control arm, the nutrition-only arm demonstrated increased prevalence of any bacterial gene (PR 1.06, 95% CI 1.01, 1.11), adenovirus (PR 1.88, 95% CI 1.41, 2.51) and heat-labile/heat-stable ETEC (PR 2.00, 95% CI 1.19, 3.36) and reduced prevalence of EIEC/*Shigella* spp. (PR 0.60, 95% CI 0.39, 0.94). Children in the sanitation-only arm had less EPEC (PR 0.88, 95% CI 0.78, 1.00) compared to control. In the combined-intervention arm, atypical-EPEC prevalence decreased (PR 0.85, 95% CI 0.73, 0.98) while the prevalence increased for heat-stable ETEC (PR 1.42, 95% CI 1.01, 2.00), heat-labile/heat-stable ETEC (PR 1.74, 95% CI 1.06, 2.86), and any viral gene (PR 1.16, 95% CI 1.02, 1.31). We found similar mixed effects when comparing pathogen gene prevalence in individual treatment arms compared to the combined arm; there was slightly lower combined prevalence of any bacterial target (PR 0.96, 95% CI 0.91, 1.02) and enterovirus (PR 0.82, 95% CI 0.67, 1.00) in the sanitation-only arm, and we found higher prevalence of adenovirus (PR 1.42, 95% CI 1.07, 1.87) in the nutrition-only arm (Table 9).

Generally, differences in mean gene quantities were consistent with prevalence differences

(Table 9;

Table 10). We detected lower concentrations of pathogen-associated genes in the nutrition-only
and sanitation-only arms compared with the control arm; children in the nutrition-only arm
carried lower quantities of STEC (-1.46 $\log_{10}$-copies, 95% CI -2.97, 0.06) and *Giardia* (-1.73
$\log_{10}$-copies, 95% CI -3.02, -0.44), and children in the sanitation-only arm carried lower
quantities of EPEC (-0.54 $\log_{10}$-copies, 95% CI -1.17, 0.09) and STEC (-1.71 $\log_{10}$-copies, 95%
CI -3.07, -0.34). There was no measurable difference in mean gene quantities between the
combined and control arm. There was no significant difference in quantity of pathogen genes
between treatment arms after adjusting for multiple comparisons[20].

Discussion

We found a modest effect on growth from the nutrition-only intervention and a greater effect in
the combined intervention arm that was likely attributable to the nutrition intervention alone. By
contrast, the sanitation intervention alone was not associated with growth improvements, relative
to control conditions, and demonstrated significantly poorer growth than the combined
intervention arm. The similar impacts on linear growth between the nutrition-only and combined
intervention arms were consistent with the observed linear growth improvements being
attributable primarily to the nutrition intervention alone, further suggesting that the addition of
this sanitation intervention to the nutrition intervention did not produce synergistic effects.
Intermediate outcomes of meal frequency and dietary diversity were similar between arms, so the
observed effects may have been attributable to other elements of the nutrition or combined
intervention not captured by these measures. We observed no meaningful differences between
arms with respect to secondary outcome measures of WAZ, WLZ, stunting, wasting,
underweight status, diarrhoea, mortality, pathogen prevalence, pathogen co-detection rate, or
pathogen gene copy quantity. Although molecular detection of a specific pathogen in stool does
not necessarily signal active enteric infection, potential for disease, or direct effects on the
individual, it does unambiguously indicate prior exposure to that pathogen. Our specific
pathogen targets were selected *a priori* based on a range of globally observed enteric pathogens
and may not fully capture the relevant enteric pathogens in rural Cambodia; however, the
consistently high pathogen prevalence across all treatment arms suggests the suite of
interventions assessed in this trial did not prevent environmental exposure to enteric pathogens.
336
Our findings are consistent with results from several recent randomised factorial WASH and
nutrition efficacy trials reporting protective effects of combined/integrated interventions and null
effects of WASH alone on child growth outcomes[3,4,9]. A small number of experimental trials[24]
and many observational studies[8,10] have reported increases in child growth and reductions in
stunting prevalence with improvements in sanitation coverage and commensurate reductions in
OD; among the latter, unmeasured confounding is a likely explanation for observed effects[24].
343
While gains in the proportion of the population self-reporting access to sanitation were highest in
the sanitation only arm (+25 pp), these were only modestly higher than the gains for the control
(+19 pp) and nutrition arms (+14 pp); furthermore, sanitation coverage gains for the combined
intervention matched the control arm at +19 pp. Comparable secular trends of increasing
sanitation coverage in Cambodia have been documented previously: the percentage of children
younger than five years of age with access to an improved sanitation facility increased from 5%
in 2000 to 17% in 2005, 29% in 2010, and 54% in 2014, the most recent nationally
representative DHS survey[8,23]. Correspondingly swift improvements have been documented

specifically in rural areas, where access to any sanitation facility increased from 30% in 2010 to
44% in 2014 and improved sanitation coverage rose from 27% to 43%[8,23]. The rapid pace of
WASH development in rural Cambodia makes it challenging to measure the impact of specific
programs. However, the lack of differences in other sanitation intervention adherence indicators
suggests low overall uptake of the sanitation intervention and only modest increases in sanitation
coverage attributable to the intervention, which were likely insufficient to reduce community
exposure and transmission. Zoonotic transmission from domestic animals, for instance, was not
addressed by this or many other WASH trials, as indicated by nearly 80% of households across
all treatment arms lacking access to an area free of animals for children to safely play[25].
361
The frequency and intensity of contact from program promoters was much greater in the
nutrition intervention than the sanitation intervention. Recipients of the nutrition intervention
participated in monthly activities, whereas the sanitation intervention consisted of one triggering
session with infrequent follow-up visits. The lower contact frequency may explain the
discrepancy in intervention adherence. Both arms receiving the nutrition programming reported
higher levels of participation in the key intervention activities—including sanitation intervention
activities—suggesting higher adherence to the nutrition intervention than the sanitation
intervention. Self-reported CLTS participation rates were equally low in both the sanitation-only
and control arms at 6%, while 14% of nutrition-only recipients and 25% of combined
intervention recipients reported CLTS participation. The comparatively elevated CLTS
participation in the nutrition-only arm may reflect biases embedded in the self-reporting process;
given the 28+ months that had elapsed since the initial CLTS triggering session and the
infrequency of CLTS follow-up visits, households that only received the sanitation intervention
may have been less likely to recall programming of any kind than households that participated in
the more frequent and intense nutrition intervention activities. Furthermore, the "Growth
Together" SBCC campaign, which promoted 13 core health, nutrition, sanitation, and hygiene
practices, was fully incorporated into all intervention activities across the three intervention
arms, meaning households receiving the higher intensity nutrition programming also encountered
the associated SBCC sanitation messaging much more frequently than households in the
sanitation-only arm. The SBCC campaign was also promoted nationally on television, such that
households in the control arms may have been nearly as exposed to its content as sanitation-only
households, while the nutrition-only and combined intervention arms received substantial in-
person promotion of the SBCC campaign messaging.
385
Due to the nature of the interventions and resource considerations, all trial outcomes were
assessed during a single survey round conducted 28 months after initiating the intervention
programming, which introduced some limitations. Growth and pathogen outcomes were assessed
in children from one to 28 months of age, meaning that older children received the treatments for
a longer duration but were initially exposed to less mature intervention conditions than the
younger children born later. The timing of outcome ascertainment also precluded detecting
effects that may only be realized later in childhood, such as potentially rapid catch-up growth
after 24 months of age that may reverse earlier growth faltering[26]. While a focus on the first
1,000 days is justified[2], investigation of growth and growth-promoting factors after this window
may provide additional insight on improved WASH practices and their role in supporting long-
term development and health.
397

While molecular detection of a specific pathogen in stool unambiguously indicates prior
exposure to that pathogen, our data do not necessarily indicate active enteric infection, potential
for disease, or direct effects on the individual. We were limited by the suite of pathogen targets
selected on our custom TAC assay; we selected these targets *a priori* based on a range of
globally observed enteric pathogens, but we cannot know whether these were the most important
enteric pathogens in rural Cambodia. There is also evidence that the *invA* gene, which was
selected for *Salmonella* spp. detection, is not specific to *Salmonella enterica* and suggest the
consideration of other genes, such as *ttrA/C*, for reliable detection of *S. enterica*[27].
It is highly plausible that this sanitation intervention simply failed to sufficiently reduce
environmental exposure to enteric pathogens. For example, only 22% of households in our
survey were observed to have a child play environment free of animals, with little difference
between treatment and control arms; this is a transmission pathway that our trial and many other
WASH trials have not addressed[25]. The trial design is predicated on the theory that gains in
sanitation coverage may lead to improved growth outcomes in children via reductions in the
transmission of enteric infection and disease, though links between sanitation coverage and
specific outcomes are poorly understood in high-burden settings. The change in community
coverage in this trial was limited and likely insufficient to reduce community exposure and
transmission.
416
There are a few key observations from this study that should be considered in future
interventions and effectiveness trials of comparable interventions. Increased frequency, duration,
and intensity of CLTS programming could have resulted in greater uptake of sanitation in target
communes. Despite the sanitation coverage gains observed in the sanitation-only arm, much of
which may have been as a result of the sanitation intervention, we are unable to attribute
beneficial effects—i.e., measurable differences in prespecified outcomes—to the sanitation
intervention due to the high sanitation gains also observed in the control arm. There may have
been other benefits of sanitation gains that were not measured, including in safety and broader
measures of well-being[28]. Future trials may also include additional objective outcome measures,
including intermediate measures of environmental contamination that are on the causal pathway
between interventions and exposures.
428
429 Our work is consistent with a growing body of research reporting high prevalence of enteric
430 pathogen exposure in early childhood, which may lead to long-term effects on health[15,18,29,30].
431 Reducing these exposures in high-burden settings requires transformative interventions that have
432 the potential to dramatically reduce direct and indirect contact with all faeces, including animal
433 faeces[24], across multiple pathways.
434

Other Information

Trial Registry

The trial is registered with ISRCTN Registry ([ISRCTN77820875](#)).

Protocol

The National Ethics Committee for Health Research in the Cambodian Ministry of Health reviewed and approved the protocols (NECHR #110) prior to the start of data collection. The study also received approvals from the Institutional Review Board at Georgia Institute of Technology (Ref: H19286) and from New England IRB (IRB#: 120190186).

Funding

This study was funded by the United States Agency for International Development (USAID) under contract number OAA-M-13-00017. The contents of this publication are the sole responsibility of the authors and do not necessarily reflect the views of USAID or the United States Government.

Acknowledgements

We thank the staff and participants of the study for their important contributions. We also acknowledge contributions by Caroline Akerley, Farran Bush, Isabelle du Plessis, and Juliann Pham.

Author contributions

Conceptualisation	IV, OC, JB
Data Curation	IV, KS, AL
Formal Analysis	AL, RA, IV
Funding Acquisition	IV, JB
Investigation	AL, KS, IV, JB
Methodology	AL, IV, OC, JB
Project Administration	IV, KS, JB
Resources	IV, JB
Software	N/A
Supervision	IV, KS, OC, JB
Validation	N/A
Visualisation	N/A
Writing – Original Draft Preparation	AL
Writing – Review & Editing	AL, IV, RA, EK, DH, KS, OC, JB, KL

455
456 **Figure 1: Theory of change diagram**

457
458
459
460

461
462 **Figure 2: Trial profile**

464
465 **Figure 3: Unadjusted intervention effects on LAZ. Estimates are mean differences (point)**
466 **with 95% CIs (line)**

Figure 4: Impact of interventions on unadjusted prevalence ratio of individual pathogens. Point estimates and 95% confidence intervals were determined using log-linear Poisson models with generalized estimating equations.

473 **Table 1: Summary of intervention activities**

	Intervention activity	Frequency
Nutrition-only		
	Community dialogues led by village chief and VHSG to support children's growth	Quarterly
	Caregiver group sessions led by local women trained by staff to promote 13 key stunting prevention behaviours	Monthly
	GMP sessions led by VHSGs to monitor growth and refer children who were sick or not growing well to health centers	Monthly
	Home visits to pregnant women, caregivers of children 9-11 months old, and caregivers of children not growing well to promote childcare and feeding, home hygiene, and handwashing	Monthly
	Village fair help twice per year to offer hands-on learning experiences (health/nutrition, WASH and agricultural using games, latrine marketing and sales	Twice per year
	CCT (cash for antenatal and postnatal care visits and adherence to handwashing stations), vouchers for water filters and food baskets	Up to six payments over first 1,000 days of child life
Sanitation-only		
	CLTS triggering session	Once
	Door-to-door visits to provide information about sanitation/latrines	At least five times per village
	Latrine vouchers to subsidise poor households in villages that reached 75% sanitation coverage to achieve sufficient open defecation free (ODF) coverage	Once, as needed
	Promoted supply-side support by connecting small- and medium-sized enterprises (SMEs) with communes after triggering event	Continuously
Combined		
	All activities described in NUTR and SAN groups above	See above
Control		
	None	N/A

474
475

476   **Table 2: Household and caregiver characteristics**

	Nutrition-only		Sanitation-only		Combined		Control	
	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI
Buddhist	0.98	(0.97, 0.99)	0.97	(0.96, 0.98)	0.96	(0.95, 0.97)	0.99	(0.98, 0.99)
Married or living together	0.95	(0.93, 0.96)	0.94	(0.92, 0.95)	0.94	(0.92, 0.95)	0.95	(0.94, 0.96)
Maternal age (years)	32.0	(31.2, 32.7)	31.1	(30.4, 31.9)	31.9	(31.2, 32.5)	31.0	(30.5, 31.5)
Primary caregiver has attended primary school	0.87	(0.85, 0.89)	0.87	(0.84, 0.89)	0.88	(0.85, 0.89)	0.81	(0.79, 0.83)
Spouse has attended primary school	0.90	(0.88, 0.92)	0.86	(0.83, 0.88)	0.87	(0.85, 0.89)	0.87	(0.85, 0.89)
Household size	5.63	(5.48, 5.78)	5.54	(5.40, 5.69)	5.36	(5.24, 5.47)	5.52	(5.42, 5.62)
Number of children in HH	2.61	(2.52, 2.70)	2.61	(2.51, 2.70)	2.43	(2.36, 2.50)	2.43	(2.37, 2.49)
Has electricity	0.73	(0.70, 0.76)	0.74	(0.71, 0.77)	0.76	(0.73, 0.78)	0.75	(0.72, 0.77)
Owns a mobile phone	0.93	(0.92, 0.95)	0.90	(0.88, 0.92)	0.87	(0.85, 0.89)	0.90	(0.89, 0.92)
Has a finished floor [1]	0.92	(0.90, 0.94)	0.93	(0.91, 0.95)	0.96	(0.94, 0.97)	0.96	(0.95, 0.97)
Wealth index score, excluding WASH variables	0.06	(-0.06, 0.19)	0.19	(0.05, 0.33)	-0.02	(-0.14, 0.09)	-0.12	(-0.21, -0.03)
Improved drinking water source [2]	0.59	(0.56, 0.62)	0.56	(0.52, 0.59)	0.61	(0.58, 0.64)	0.70	(0.68, 0.72)
Has water source on site	0.68	(0.64, 0.71)	0.58	(0.55, 0.61)	0.60	(0.57, 0.63)	0.65	(0.62, 0.67)
Water source is <5 min, roundtrip	0.20	(0.16, 0.25)	0.23	(0.19, 0.28)	0.24	(0.19, 0.29)	0.34	(0.29, 0.39)
Reported minutes to fetch water, roundtrip	12.42	(10.6, 14.2)	14.14	(12.5, 15.8)	14.50	(13.1, 15.9)	13.56	(12.3, 14.8)

477 [1] Finished floor defined as floor made of wood plans, palm/bamboo, parquet or polished wood, vinyl or asphalt strips, ceramic
478 tiles, cement tiles, or cement. Floor materials were classified by enumerator observation. [2] Improved sources of drinking water
479 include: piped water into dwelling/yard/plot, public tap or standpipe, tube well or borehole, protected dug well, protected spring,
480 bottled water, and rainwater.
481
482

483 **Table 3: Intervention fidelity indicators**

	Nutrition-only			Sanitation-only			Combined			Control		
Nutrition	N	n	%	N	n	%	N	n	%	N	n	%
Participated in any "First 1,000 Days" type activity [1]	817	615	75%	792	145	18%	1,055	813	77%	1,460	383	26%
Participated in any GMP	817	641	78%	792	181	23%	1,055	935	89%	1,460	482	33%
Received home visit VHSG	817	518	63%	792	227	29%	1,055	661	63%	1,460	490	34%
Enrolled in any CCT program for health and nutrition [2]	817	224	27%	792	19	2%	1,055	228	22%	1,460	31	2%
Received any voucher for food basket [3]	817	440	54%	792	1	0%	1,055	554	53%	1,460	6	0%
Received any voucher for water filter [3]	817	41	5%	792	40	5%	1,055	149	14%	1,460	100	7%
Aware of *Grow Together* campaign [4]	817	353	43%	792	93	12%	1,055	471	45%	1,460	149	10%
Participation in nutrition intervention activities: none (0 of 8)	817	65	8%	792	404	51%	1,055	44	4%	1,460	585	40%
Participation in nutrition intervention activities: low (1-3 of 8)	817	262	32%	792	359	45%	1,055	367	35%	1,460	809	55%
Participation in nutrition intervention activities: med (4-6 of 8)	817	387	47%	792	29	4%	1,055	501	47%	1,460	65	4%
Participation in nutrition intervention activities: high (7-8 of 8)	817	103	13%	792	0	0%	1,055	143	14%	1,460	1	0%
Sanitation												
Any CLTS participation [5]	817	115	14%	792	46	6%	1,055	261	25%	1,460	81	6%
Received any voucher to build latrine [6]	817	66	8%	792	51	6%	1,055	123	12%	1,460	84	6%
[1] "First 1,000 Days" activities were administered in nutrition-only and combined arms and include: community dialogues, caregiver group education sessions, and village fairs. [2] CCT program ended in Jan 2019 and the Government of Cambodia started a new CCT program in July 2019 across study area. CCT program administered in nutrition-only and combined arms only. [3] Vouchers for water filter and food baskets were targeted subsidies distributed to CCT participants in nutrition-only and combined arms. [4] *Grow Together* campaign was part of the nutrition programming (nutrition-only and combined arms). However, three TV spots were seen across all four arms. [5] In sanitation-only and combined arms, the Ministry of Rural Development confirmed that the project was the only CLTS campaign active in those areas. [6] Latrine vouchers were targeted subsidies given to households in villages that reached 75% sanitation coverage in sanitation-only and combined arms.												

484
485

Table 4: Intervention adherence indicators (28-months after intervention)

Nutrition	Nutrition-only			Sanitation-only			Combined			Control		
	N	n or mean	% or SD	N	n or mean	% or SD	N	n or mean	% or SD	N	n or mean	% or SD
Visited health facility for at least four antenatal care check-ups	697	632	91%	712	646	91%	910	819	90%	1,257	1116	89%
Brought child for monthly GMP at community or health center	817	641	78%	792	181	23%	1,055	935	89%	1,460	482	33%
Breastfeeding exclusively for children <6 months	171	109	64%	161	110	68%	205	140	68%	272	182	67%
Ever breastfed (all children)	817	797	98%	792	765	97%	1,048	1021	97%	1,457	1420	97%
Solid and semi-solid foods eaten for children >6 months	646	609	94%	631	611	97%	843	792	94%	1,185	1134	96%
Dietary diversity score (0-7)	817	2.24	1.61	792	2.19	1.58	1,048	2.20	1.62	1,457	2.33	1.59
Achieved minimum dietary diversity	817	202	25%	792	177	22%	1,048	247	24%	1,457	350	24%
Achieved minimum meal frequency	817	537	66%	792	518	65%	1,048	680	65%	1,457	977	67%
Achieved minimum acceptable diet	817	170	21%	792	159	20%	1,048	205	20%	1,457	310	21%
Treated drinking water	817	548	67%	792	471	59%	1,055	765	73%	1,460	1,037	71%
Treated drinking water with filter	817	151	18%	792	160	20%	1,055	302	29%	1,460	548	40%
Sanitation												
Had improved sanitation facility [1]	816	452	55%	791	400	51%	1,054	638	61%	1,459	759	52%
Open defecation (OD)	817	112	14%	792	126	16%	1,055	73	7%	1,460	231	16%
Used shared toilet	817	252	31%	791	264	33%	1,054	343	33%	1,459	468	32%
Caregiver reported adults in HH openly defecating	697	92	13%	658	116	18%	973	118	12%	1,208	213	18%
Time to get to toilet, one way (minutes)	171	4.22	4.11	166	3.92	3.83	219	4.74	8.17	291	5.05	8.27
Reported latrine built as a result of CLTS activity	115	51	44%	46	15	33%	261	91	35%	81	28	35%
Reported latrine built using latrine voucher	50	10	20%	15	4	27%	91	37	41%	28	12	43%
Main reason for not constructing latrine: lack of funds	20	17	85%	18	14	78%	60	55	92%	14	14	100%
Main reason for construction latrine: privacy	51	7	14%	15	6	40%	91	6	7%	28	7	25%
Main reason for construction latrine: security	51	10	20%	15	2	13%	91	20	22%	28	4	14%
Main reason for construction latrine: hygiene	51	17	33%	15	5	33%	91	43	47%	28	10	36%
Main reason for construction latrine: OD is harmful	51	5	10%	15	1	7%	91	9	10%	28	5	18%

Child stools properly disposed of [2]	817	581	71%	792	515	65%	1,055	781	74%	1,460	993	68%
Community-level open defecation before intervention	817	28%	27%	792	41%	33%	1,055	28%	28%	1,460	32%	31%
Community-level open defecation after intervention	817	14%	21%	792	16%	21%	1,055	9%	17%	1,460	13%	20%
Environmental hygiene												
Child stools left in the open	817	147	18%	792	170	21%	1,055	160	15%	1,460	315	22%
Child play environment observed to be free of animals	817	182	22%	792	187	24%	1,055	261	25%	1,460	294	20%
Child play environment observed to be free of garbage/HH waste	817	298	36%	792	290	37%	1,055	419	40%	1,460	567	39%
Child play environment observed to be free of sharp objects	817	449	55%	792	427	54%	1,055	639	61%	1,460	818	56%
Child play environment observed to be free of faeces	817	313	38%	792	304	38%	1,055	448	42%	1,460	555	38%
[1] Improved sanitation facilities include: flush/pour flush toilet to a piped sewer system, septic tank or pit latrine, a ventilated improved pit latrine, a pit latrine with slab, and a composting toilet. [2] Proper disposal of children faeces consist of putting or rinsing stool into a sanitation facility or burying it; improper disposal of children faeces includes putting or rinsing stool into a drain or ditch, throwing it into garbage or leaving it in the open.												

489 **Table 5: Effects of interventions on length and weight (Primary outcome (LAZ) and**
490 **secondary outcomes (WAZ, WLZ)), comparing intervention arms to control and single**
491 **intervention arms to combined intervention**

	N	Mean	SD	Unadjusted mean difference (95% CI)	
				Compared to control arm	Compared to combined intervention arm
LAZ					
Nutrition-only	798	-0.95	1.16	0.08 (-0.01, 0.18)	-0.02 (-0.12, 0.09)
Sanitation-only	777	-1.09	1.23	-0.05 (-0.16, 0.05)	-0.16 (-0.27, -0.04)
Combined	1037	-0.94	1.16	0.10 (0.01, 0.20)	--
Control	1443	-1.04	1.20	--	--
WAZ					
Nutrition-only	815	-0.95	1.29	0.10 (0.00, 0.19)	-0.02 (-0.12, 0.08)
Sanitation-only	792	-1.04	1.13	0.01 (-0.07, 0.09)	-0.10 (-0.20, -0.01)
Combined	1044	-0.94	1.11	0.11 (0.03, 0.20)	--
Control	1452	-1.05	1.10	--	--
WLZ					
Nutrition-only	814	-0.60	1.04	0.06 (-0.03, 0.15)	-0.02 (-0.12, 0.08)
Sanitation-only	790	-0.59	0.98	0.06 (-0.02, 0.14)	-0.02 (-0.11, 0.07)
Combined	1043	-0.58	1.03	0.08 (0.00, 0.16)	--
Control	1452	-0.65	0.98	--	--

492
493
494

Table 6: Effects of intervention on child health outcomes, comparing intervention arms to control and single intervention arms to combined intervention.

	N	Mean	SD	Compared to control arm	Compared to combined-intervention arm
				PR (95% CI)	PR (95% CI)
Stunted					
Nutrition-only	801	0.15	0.36	0.84 (0.69, 1.03)	0.93 (0.74, 1.15)
Sanitation-only	782	0.21	0.40	1.12 (0.94, 1.33)	1.23 (1.02, 1.49)
Combined	1046	0.17	0.37	0.91 (0.76, 1.09)	--
Control	1449	0.18	0.39	--	--
Wasted					
Nutrition-only	815	0.07	0.26	0.87 (0.65, 1.17)	1.12 (0.80, 1.57)
Sanitation-only	790	0.07	0.26	0.84 (0.62, 1.14)	1.08 (0.76, 1.53)
Combined	1052	0.07	0.25	0.78 (0.58, 1.04)	--
Control	1457	0.08	0.28	--	--
Underweight					
Nutrition-only	816	0.15	0.35	0.85 (0.71, 1.03)	1.04 (0.84, 1.29)
Sanitation-only	792	0.17	0.38	1.00 (0.85, 1.19)	1.22 (1.00, 1.49)
Combined	1053	0.14	0.35	0.82 (0.68, 0.99)	--
Control	1457	0.17	0.38	--	--
Diarrhoea (7-day recall)					
Nutrition-only	788	0.19	0.39	0.89 (0.74, 1.06)	0.95 (0.78, 1.14)
Sanitation-only	752	0.21	0.41	0.99 (0.84, 1.17)	1.05 (0.88, 1.25)
Combined	1018	0.20	0.40	0.94 (0.80, 1.11)	--
Control	1411	0.21	0.41	--	--
All-cause mortality					
Nutrition-only	1574	0.03	0.16	1.55 (0.71, 3.39)	1.61 (0.68, 3.82)
Sanitation-only	1636	0.03	0.16	1.09 (0.50, 2.40)	1.13 (0.48, 2.68)
Combined	1932	0.03	0.16	0.96 (0.44, 2.10)	--
Control	2688	0.03	0.16	--	--

495
496
497
498

Table 7: Enteric pathogen gene prevalence among treatment arms

	All samples (N=1620)	Nutrition-only (N=305)	Sanitation-only (N=333)	Combined (N=438)	Control (N=544)
Bacteria					
Campylobacter spp.	551 (34%)	104 (34%)	114 (34%)	162 (37%)	171 (31%)
Clostridium difficile	139 (9%)	33 (11%)	25 (8%)	41 (9%)	40 (7%)
EAEC	1029 (64%)	204 (67%)	207 (62%)	281 (64%)	337 (62%)
EPEC	899 (55%)	172 (56%)	173 (52%)	234 (53%)	320 (59%)
aEPEC	703 (43%)	139 (46%)	137 (41%)	173 (39%)	254 (47%)
tEPEC	109 (7%)	15 (5%)	17 (5%)	32 (7%)	45 (8%)
ETEC	422 (26%)	86 (28%)	79 (24%)	114 (26%)	143 (26%)
ETEC-LT	342 (21%)	75 (25%)	68 (20%)	86 (20%)	113 (21%)
ETEC-ST	194 (12%)	39 (13%)	37 (11%)	63 (14%)	55 (10%)
ETEC-LT/ST	114 (7%)	28 (9%)	26 (8%)	35 (8%)	25 (5%)
Salmonella spp.	134 (8%)	28 (9%)	19 (6%)	39 (9%)	48 (9%)
Shigella spp.	186 (11%)	24 (8%)	39 (12%)	52 (12%)	71 (13%)
STEC	132 (8%)	27 (9%)	22 (7%)	45 (10%)	38 (7%)
Vibrio cholerae	10 (1%)	1 (0%)	6 (2%)	1 (0%)	2 (0%)
At least 1 bacterium detected	1410 (87%)	276 (90%)	282 (85%)	386 (88%)	466 (86%)
Mean number of bacteria detected	2.48 (2.42, 2.55)	2.46 (2.32, 2.60)	2.43 (2.28, 2.57)	2.51 (2.39, 2.63)	2.51 (2.40, 2.62)
Viruses					
Adenovirus	287 (18%)	77 (25%)	59 (18%)	78 (18%)	73 (13%)
Astrovirus	7 (0%)	2 (1%)	1 (0%)	3 (1%)	1 (0%)
Enterovirus	558 (34%)	97 (32%)	105 (32%)	169 (39%)	187 (34%)
Norovirus	54 (3%)	8 (3%)	9 (3%)	20 (5%)	17 (3%)
Rotavirus	17 (1%)	2 (1%)	2 (1%)	6 (1%)	7 (1%)
Sapovirus	24 (1%)	4 (1%)	5 (2%)	9 (2%)	6 (1%)
At least 1 virus detected	788 (49%)	152 (50%)	157 (47%)	231 (53%)	248 (46%)
Mean number of viruses detected	1.20 (1.17, 1.23)	1.25 (1.17, 1.33)	1.15 (1.09, 1.21)	1.23 (1.17, 1.30)	1.17 (1.12, 1.23)
Protozoa					
Cryptosporidium	17 (1%)	4 (1%)	1 (0%)	6 (1%)	6 (1%)
Entamoeba	13 (1%)	1 (0%)	1 (0%)	8 (2%)	3 (1%)
Giardia	306 (19%)	52 (17%)	68 (20%)	84 (19%)	102 (19%)
At least 1 protozoan detected	328 (20%)	56 (18%)	69 (21%)	93 (21%)	110 (20%)
Mean number of protozoa detected	1.02 (1.01, 1.04)	1.02 (0.98, 1.05)	1.01 (0.99, 1.04)	1.05 (1.01, 1.10)	1.01 (0.99, 1.03)
STH					
Ascaris lumbricoides	3 (0%)	0 (0%)	0 (0%)	3 (1%)	0 (0%)
Trichuris trichiura	3 (0%)	1 (0%)	1 (0%)	0 (0%)	1 (0%)
Ancylostoma duodenale	17 (1%)	0 (0%)	4 (1%)	4 (1%)	9 (2%)
Necator americanus	20 (1%)	3 (1%)	6 (2%)	5 (1%)	6 (1%)
At least 1	37 (2%)	4 (1%)	10 (3%)	9 (2%)	14 (3%)
Mean number of STH detected	1.16 (1.04, 1.29)	1.00 (1.00, 1.00)	1.10 (0.90, 1.30)	1.33 (1.00, 1.67)	1.14 (0.95, 1.34)

500
501

502 Table 8: Unadjusted incidence rate ratios of co-detected bacteria, viruses, protozoa, and
503 STHs, comparing intervention arms to control and single intervention arms to combined
504 intervention.

	N	Compared to control arm			Compared to combined arm	
		Nutrition-only	Sanitation-only	Combined	Nutrition-only	Sanitation-only
Bacteria	1620	1.04 (0.95, 1.13)	0.96 (0.87, 1.05)	1.03 (0.95, 1.12)	0.01 (-0.08, 0.10)	-0.07 (-0.17, 0.02)
Viruses	1620	1.16 (0.99, 1.37)	1.02 (0.86, 1.19)	1.22 (1.05, 1.41)	-0.04 (-0.21, 0.12)	-0.18 (-0.34, -0.02)
Protozoa	1620	0.92 (0.68, 1.23)	1.03 (0.79, 1.35)	1.10 (0.85, 1.41)	-0.18 (-0.48, 0.12)	-0.06 (0.66, -0.34)
STHs were omitted because <5% of samples had detectable STH genes.						

506 Table 9: Unadjusted prevalence ratios (PR) of detected bacteria, viruses, protozoa, and
507 STHs, comparing intervention arms to control and single intervention arms to combined
508 intervention.

	Compared to control arm			Compared to combined arm	
	Nutrition-only	Sanitation-only	Combined	Nutrition-only	Sanitation-only
Any bacterium	1.06 (1.00, 1.11)	0.99 (0.93, 1.05)	1.03 (0.98, 1.08)	1.03 (0.98, 1.08)	0.96 (0.91, 1.02)
Camploybacter spp.	1.08 (0.89, 1.32)	1.09 (0.90, 1.32)	1.18 (0.99, 1.40)	0.92 (0.76, 1.12)	0.93 (0.76, 1.12)
C.diff	1.47 (0.95, 2.28)	1.02 (0.63, 1.65)	1.27 (0.84, 1.93)	1.16 (0.75, 1.78)	0.80 (0.50, 1.29)
EAEC	1.08 (0.97, 1.20)	1.00 (0.90, 1.12)	1.04 (0.94, 1.14)	1.04 (0.94, 1.16)	0.97 (0.87, 1.08)
EPEC	0.96 (0.85, 1.08)	0.88 (0.78, 1.00)	0.91 (0.81, 1.02)	1.06 (0.93, 1.20)	0.97 (0.85, 1.11)
aEPEC	0.98 (0.84, 1.14)	0.88 (0.75, 1.03)	0.85 (0.73, 0.98)	1.15 (0.97, 1.37)	1.04 (0.88, 1.24)
tEPEC	0.59 (0.34, 1.05)	0.62 (0.36, 1.06)	0.88 (0.57, 1.37)	0.67 (0.37, 1.22)	0.70 (0.39, 1.24)
ETEC	1.07 (0.85, 1.35)	0.90 (0.71, 1.15)	0.99 (0.80, 1.22)	1.08 (0.85, 1.38)	0.91 (0.71, 1.17)
ETEC-LT	1.18 (0.92, 1.53)	0.98 (0.75, 1.29)	0.95 (0.74, 1.21)	1.25 (0.95, 1.65)	1.04 (0.78, 1.38)
ETEC-ST	1.26 (0.86, 1.86)	1.10 (0.74, 1.63)	1.42 (1.01, 2.00)	0.89 (0.61, 1.29)	0.77 (0.53, 1.13)
ETEC-LT/ST	2.00 (1.19, 3.36)	1.70 (1.00, 2.89)	1.74 (1.06, 2.86)	1.15 (0.71, 1.85)	0.98 (0.60, 1.59)
Salmonella spp.	1.04 (0.67, 1.62)	0.65 (0.39, 1.08)	1.01 (0.67, 1.51)	1.03 (0.65, 1.64)	0.64 (0.38, 1.09)
EIEC/*Shigella* spp.	0.60 (0.39, 0.94)	0.90 (0.62, 1.29)	0.91 (0.65, 1.27)	0.66 (0.42, 1.05)	0.99 (0.67, 1.46)
STEC	1.27 (0.79, 2.03)	0.95 (0.57, 1.57)	1.47 (0.97, 2.22)	0.86 (0.55, 1.36)	0.64 (0.39, 1.05)
Any virus	1.09 (0.95, 1.26)	1.03 (0.89, 1.20)	1.16 (1.02, 1.31)	0.94 (0.82, 1.09)	0.89 (0.77, 1.03)
Adenovirus	1.88 (1.41, 2.51)	1.32 (0.96, 1.81)	1.33 (0.99, 1.78)	1.42 (1.07, 1.87)	0.99 (0.73, 1.35)
Enterovirus	0.93 (0.76, 1.13)	0.92 (0.75, 1.12)	1.12 (0.95, 1.32)	0.82 (0.67, 1.01)	0.82 (0.67, 1.00)
Any protozoa	0.91 (0.68, 1.21)	1.02 (0.78, 1.34)	1.05 (0.82, 1.34)	0.86 (0.64, 1.16)	0.98 (0.74, 1.29)
Giardia	0.91 (0.67, 1.23)	1.09 (0.83, 1.43)	1.02 (0.79, 1.33)	0.89 (0.65, 1.22)	1.06 (0.80, 1.42)
Gene targets with <5% prevalence were omitted from PR analyses: *V.cholera*, astrovirus, norovirus, rotavirus, sapovirus, *Cryptosporidium*, *Entamoeba,* and all STHs (*Ascaris, Trichuris, Ancylostoma,* and *Necator*).					

Table 10: Mean difference in log$_{10}$-transformed gene copy estimates, comparing intervention arms to control and single intervention arms to combined intervention.

	Compared to control arm			Compared to combined arm	
	Nutrition-only	Sanitation-only	Combined	Nutrition-only	Sanitation-only
Bacteria					
CAMP	0.22 (-0.60 - 1.03)	-0.40 (-1.27 - 0.48)	-0.24 (-1.00 - 0.53)	0.45 (-0.37 - 1.28)	-0.16 (-1.05 - 0.73)
CDIF	0.31 (-1.14 - 1.75)	-0.37 (-1.92 - 1.19)	0.19 (-1.16 - 1.54)	0.11 (-1.29 - 1.52)	-0.56 (-2.08 - 0.96)
EAEC_aaic	-0.38 (-1.35 - 0.60)	0.35 (-0.52 - 1.22)	-0.28 (-1.08 - 0.52)	-0.10 (-1.11 - 0.92)	0.63 (-0.28 - 1.54)
EAEC_aata	-0.36 (-1.26 - 0.53)	-0.24 (-1.05 - 0.56)	0.12 (-0.69 - 0.93)	-0.48 (-1.43 - 0.47)	-0.36 (-1.24 - 0.51)
EPEC_bfpa	-1.08 (-3.47 - 1.31)	-1.13 (-3.16 - 0.89)	1.15 (-0.38 - 2.67)	-2.23 (-4.57 - 0.11)	-2.28 (-4.24 - -0.32)
EPEC_eae	-0.12 (-0.80 - 0.56)	-0.54 (-1.17 - 0.09)	0.24 (-0.33 - 0.82)	-0.37 (-1.05 - 0.32)	-0.78 (-1.41 - -0.15)
ETEC_LT	-0.62 (-1.64 - 0.40)	-0.47 (-1.56 - 0.61)	-0.03 (-1.06 - 1.00)	-0.59 (-1.65 - 0.47)	-0.44 (-1.57 - 0.68)
ETEC_stp	0.13 (-1.73 - 1.98)	1.29 (-0.61 - 3.20)	0.87 (-0.97 - 2.71)	-0.75 (-2.61 - 1.11)	0.42 (-1.49 - 2.33)
SALM	1.42 (-0.04 - 2.87)	0.27 (-0.97 - 1.52)	0.80 (-0.33 - 1.93)	0.62 (-0.93 - 2.17)	-0.53 (-1.89 - 0.83)
IPAH	0.22 (-1.33 - 1.77)	-0.28 (-1.68 - 1.13)	1.17 (0.07 - 2.26)	-0.95 (-2.43 - 0.53)	-1.44 (-2.77 - -0.11)
STEC1	-1.46 (-2.97 - 0.06)	-1.71 (-3.07 - -0.34)	-0.00 (-1.49 - 1.49)	-1.46 (-2.84 - -0.07)	-1.70 (-2.92 - -0.49)
STEC2	-0.20 (-1.45 - 1.04)	0.09 (-1.20 - 1.37)	0.72 (-0.47 - 1.91)	-0.92 (-2.06 - 0.22)	-0.63 (-1.82 - 0.56)
Viruses					
ADEV	0.50 (-0.48 - 1.49)	0.67 (-0.32 - 1.67)	0.48 (-0.42 - 1.38)	0.03 (-0.96 - 1.02)	0.20 (-0.80 - 1.20)
ENTV	-0.40 (-1.09 - 0.28)	-0.35 (-0.97 - 0.26)	-0.26 (-0.77 - 0.24)	-0.14 (-0.84 - 0.56)	-0.09 (-0.72 - 0.54)
Protozoa					
GIAR	-1.73 (-3.02 - -0.44)	0.23 (-1.16 - 1.62)	-0.14 (-1.47 - 1.18)	-1.58 (-3.06 - -0.11)	0.37 (-1.19 - 1.93)
Gene targets with <5% prevalence were omitted from PR analyses: *V.cholera*, astrovirus, norovirus, rotavirus, sapovirus, *Cryptosporidium*, *Entamoeba,* and all STHs (*Ascaris, Trichuris, Ancylostoma,* and *Necator*).					

References

1. Prendergast, A. J. & Humphrey, J. H. The stunting syndrome in developing countries. *Paediatr Int Child Health* **34**, 250–265 (2014).
2. Victora, C. G. *et al.* Maternal and child undernutrition: consequences for adult health and human capital. *The Lancet* **371**, 340–357 (2008).
3. Luby, S. P. *et al.* Effects of water quality, sanitation, handwashing, and nutritional interventions on diarrhoea and child growth in rural Bangladesh: a cluster randomised controlled trial. *Lancet Glob Health* **6**, e302–e315 (2018).
4. Null, C. *et al.* Effects of water quality, sanitation, handwashing, and nutritional interventions on diarrhoea and child growth in rural Kenya: a cluster-randomised controlled trial. *Lancet Glob Health* **6**, e316–e329 (2018).
5. Ruel, M. T. & Alderman, H. Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *Lancet* **382**, 536–551 (2013).
6. Humphrey, J. H. Child undernutrition, tropical enteropathy, toilets, and handwashing. *Lancet* **374**, 1032–1035 (2009).
7. Guerrant, R. L., Schorling, J. B., McAuliffe, J. F. & de Souza, M. A. Diarrhea as a cause and an effect of malnutrition: diarrhea prevents catch-up growth and malnutrition increases diarrhea frequency and duration. *Am J Trop Med Hyg* **47**, 28–35 (1992).
8. Ikeda, N., Irie, Y. & Shibuya, K. Determinants of reduced child stunting in Cambodia: analysis of pooled data from three demographic and health surveys. *Bull World Health Organ* **91**, 341–349 (2013).
9. Humphrey, J. H. *et al.* Independent and combined effects of improved water, sanitation, and hygiene, and improved complementary feeding, on child stunting and anaemia in rural Zimbabwe: a cluster-randomised trial. *Lancet Glob Health* **7**, e132–147 (2019).
10. Vyas, S., Kov, P., Smets, S. & Spears, D. Disease externalities and net nutrition: Evidence from changes in sanitation and child height in Cambodia, 2005–2010. *Econ Hum Biol* **23**, 235–245 (2016).
11. Harris, M., Alzua, M. L., Osbert, N. & Pickering, A. Community-Level Sanitation Coverage More Strongly Associated with Child Growth and Household Drinking Water Quality than Access to a Private Toilet in Rural Mali. *Environ Sci Technol* **51**, 7219–7227 (2017).
12. Pickering, A. J., Djebbari, H., Lopez, C., Coulibaly, M. & Alzua, M. L. Effect of a community-led sanitation intervention on child diarrhoea and child growth in rural Mali: A cluster-randomised controlled trial. *Lancet Glob Health* **3**, e701–e711 (2015).
13. Briceño, B., Coville, A., Gertler, P. & Martinez, S. Are there synergies from combining hygiene and sanitation promotion campaigns: Evidence from a large-scale cluster-randomized trial in rural Tanzania. *PLoS One* **12**, e0186228–e0186228 (2017).
14. Clasen, T. *et al.* Effectiveness of a rural sanitation programme on diarrhoea, soil-transmitted helminth infection, and child malnutrition in Odisha, India: a cluster-randomised trial. *Lancet Glob Health* **2**, e645–e653 (2014).
15. Knee, J. *et al.* Effects of an urban sanitation intervention on childhood enteric infection and diarrhea in Maputo, Mozambique: A controlled before-and-after trial. *Elife* **10**, 2020.08.20.20178608 (2021).

16. Center for Disease Control and Prevention (CDC). *National Health and Nutrition Examination Survey (NHANES): Anthropometry Procedures Manual*. https://wwwn.cdc.gov/nchs/data/nhanes/2017-2018/manuals/2017_Anthropometry_Procedures_Manual.pdf (2017).

17. Mutz, D. C., Pemantle, R. & Pham, P. The Perils of Balance Testing in Experimental Design: Messy Analyses of Clean Data. *Am Stat* **73**, 32–42 (2019).

18. Liu, J. *et al.* Use of quantitative molecular diagnostic methods to identify causes of diarrhoea in children: a reanalysis of the GEMS case-control study. *The Lancet* **388**, 1291–1301 (2016).

19. Rothman, K. J. No Adjustments Are Needed for Multiple Comparisons. *Epidemiology* **1**, (1990).

20. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)* **57**, 289–300 (1995).

21. Marschner, I. C. Optimal design of clinical trials comparing several treatments with a control. *Pharm Stat* **6**, 23–33 (2007).

22. Zanello, G., Srinivasan, C. S. & Shankar, B. What Explains Cambodia's Success in Reducing Child Stunting-2000-2014? *PLoS One* **11**, e0162668 (2016).

23. National Institute of Statistics, Directorate General for Health, and I. I. *Cambodia Demographic and Health Survey 2014*.

24. Pickering, A. J. *et al.* The WASH Benefits and SHINE trials: interpretation of WASH intervention effects on linear growth and diarrhoea. *Lancet Glob Health* **7**, e1139–e1146 (2019).

25. Prendergast, A. J. *et al.* Putting the 'A' into WaSH: a call for integrated management of water, animals, sanitation, and hygiene. *Lancet Planet Health* **3**, e336–e337 (2019).

26. Leroy, J. L., Ruel, M., Habicht, J.-P. & Frongillo, E. A. Using height-for-age differences (HAD) instead of height-for-age z-scores (HAZ) for the meaningful measurement of population-level catch-up in linear growth in children less than 5 years of age. *BMC Pediatr* **15**, 145 (2015).

27. Resendiz-Nava, C., Esquivel-Hernandez, Y., Alcaraz-Gonzalez, A., Castaneda-Serrano, P. & Nava, G. M. PCR Assays Based on invA Gene Amplification are not Reliable for Salmonella Detection. *Jundishapur J Microbiol* **In Press**, (2019).

28. Shiras, T. *et al.* Shared latrines in Maputo, Mozambique: exploring emotional well-being and psychosocial stress. *BMC Int Health Hum Rights* **18**, 30 (2018).

29. Platts-Mills, J. A. *et al.* Use of quantitative molecular diagnostic methods to assess the aetiology, burden, and clinical characteristics of diarrhoea in children in low-resource settings: a reanalysis of the MAL-ED cohort study. *Lancet Glob Health* **6**, e1309–e1318 (2018).

30. Rogawski, E. T. *et al.* Use of quantitative molecular diagnostic methods to investigate the effect of enteropathogen infections on linear growth in children in low-resource settings: longitudinal analysis of results from the MAL-ED cohort study. *Lancet Glob Health* **6**, e1319–e1328 (2018).