PUBLIC HOSPITALS PREPAREDNESS IN THE PROVISION OF BREAST AND CERVICAL CANCER SERVICES IN BUSIA AND TRANS-NZOIA COUNTIES IN KENYA.

Grace W. Mwaura,[1,2*] Obed K. Limo[1], Gerald O. Lwande[1], Kenneth Too [1], Richard Mugo[1], Ann W. Mwangi[1,2], Nicholas Kirui [1,2,4], Philip Tonui,[1,2], Peter Itsura,[1,2], Jemimah Kamano[1,2,4]

1. Academic Model Providing Access to Healthcare (AMPATH), Eldoret-Kenya.
2. Moi University School of Medicine, Eldoret-Kenya.
3. Moi Teaching and Referral Hospital, Eldoret-Kenya.
4. Ministry of Health, Division of Non-Communicable Diseases - Kenya

Funding: This study was funded by Access Accelerated and World Bank (Grant Number: 0356).

* **Corresponding author:**

Grace Wandia Mwaura

Academic Model Providing Access to Healthcare

P.O. Box 4606-30100; Eldoret-Kenya.

Tel: +254722960519

E-mail: *wandiahgrace@gmail.com*

Running head: *Facilities preparedness to provide cervical and breast cancer services.*

Abstract

Purpose: To assess the preparedness of public health care facilities in the provision of breast and cervical cancer services. Specifically, healthcare provider's knowledge on risk factors, screening, symptoms, diagnosis and treatment as well as availability of medical equipment required for breast and cervical cancer management.

Methods: A cross-sectional service provision assessment (SPA) survey conducted in Busia and Trans-Nzoia counties of Western Kenya between October and December 2018. Interviewer assisted questionnaires were used to collect data from healthcare workers while a structured facility questionnaire was used to assess the level of preparedness of the selected public healthcare facilities stratified by their level of care. Statistical analysis was done using STATA version 15.

Results: We enrolled 73 healthcare workers 37 (50.6%) of whom were nurses, followed by clinical officers and medical officers. The highest proportion of knowledge on risk factors and screening of breast and cervical cancer was reported among medical officers or consultant physicians, followed by clinical officers. Nurses scored highly on the symptoms of breast and cervical cancer. The medical equipment required for breast and cervical cancer screening and diagnosis were found in most facilities; however, there were no core-biopsy needles or mammograms found. A single LEEP equipment was found in a health center within Trans Nzoia while two LEEP equipment were stationed at the Busia county hospital.

Conclusion: A below average level of knowledge on breast and cervical cancer among the healthcare workers attending to patients in public healthcare facilities was found in both Busia and Trans Nzoia counties. Furthermore, there was a disparity in the distribution and quantity of priority medical equipment for the screening, diagnosis and treatment of breast and cervical cancer in the two county hospitals.

Keywords: *Cervical Cancer, Breast Cancer, Preparedness, Service Provision Assessment (SPA), Public Healthcare Facilities, Western Kenya, Busia, Trans Nzoia.*

Introduction

Cancer is a major cause of morbidity and mortality across the globe. According to the International Agency for Research on Cancer global cancer observatory report (1), approximately 1 in 5 people develop cancer during their lifetime. Of these, 1 in 8 men and 1 in 11 women die from the disease. With a rising proportion of the ageing populations globally, coupled with other socio-economic risk factors, the prevalence of cancer is on the rise. The most common forms of cancer among women globally are breast, colorectal, lung, cervical, and thyroid cancers (2). The likelihood of breast cancer is 1 in 4 cancers diagnosed among women globally (1,2). In Africa the incidence for breast cancer in 2020 was estimated at 16.8% followed by cervical cancer at 10.6% (3).   In Kenya, the incidence of breast cancer in 2020 was estimated at 16.1% followed by second placed cervical cancer at 12.4% of all women diagnosed with cancer (2). With a greater transition to a more modernized lifestyle in Kenya and many other sub-Saharan African countries, the incidence of breast and cervical cancer have been steadily rising (4). These low- and medium-income countries (LMIC) have been shown to have a great burden of various forms of cancer despite advancements in screening, diagnosis and management options (5). To adequately address the disease burden attributed to these two cancers, there is need to have effective and adequately prepared health service delivery systems. The level of preparedness and effectiveness of these health systems can be assessed using multiple dimensions to assess the progress in addressing the disease burden. Although various tools could be used in this assessment, a Service Provision Assessment (SPA) survey could be used to provide a comprehensive understanding of the health service delivery system in a particular country (6). This is because they support the strengthening of health systems in developing countries by collecting information on the availability of different resources based in

the health facilities within a country. The SPA survey describes the facility-types with the ability

to offer specific health services. It also describes the infrastructural and human resource

preparedness to offer the said services in the health charters (6). Furthermore, it assesses

adherence to the set quality standards for healthcare service provision and whether the clients

and service providers are content with the service delivery environment.

The World Health Organization's Global action plan for the prevention and control of

noncommunicable diseases between 2013-2020 targeted a 25% reduction in risk of premature

death attributed to non-communicable diseases including cancer (7). Furthermore, the United

Nations Agenda for Sustainable Development recommends a one-third reduction in the

proportion of premature death associated with non-communicable diseases such as cancer by the

year 2030 using multiple approaches such as prevention, treatment and promotion of mental

health and well-being (8). The public and private health care service delivery institutions should

have an 80% availability of the affordable basic technologies and essential medicines for the

management of these noncommunicable diseases. Furthermore, the report (7) recommends

periodic needs assessment and evaluation of resource need such as workforce, institutional and

research capacity to address the non-communicable diseases. The eight key indicators include a

comprehensive range of health services provided to the target population, the services should be

directly and permanently accessible without barriers of entry, they should adequately cover the

defined target population and provide continuity of care across the network of services, health

conditions, levels of care, and over the life cycle. The quality of these services should be high as

evidenced by their effectiveness and safety as well as being organized around the person and not

the disease or the financing mechanism. They should be properly coordinated at the local level

and across the different types of providers, types of care and levels of service delivery with

adequate accountability and efficiency (7).

In Kenya, the healthcare service delivery system is stratified into four tiers as (9): Tier 1

(Community Health Services), Tier 2 (primary healthcare facilities such as dispensaries and

health centers), Tier 3 (secondary referral facilities such as sub county and county hospitals) and

Tier 4 (Tertiary facilities such as national referral hospitals). The primary care for cancer

involves prevention of risk factors and screening, enhancing health promotion through public

awareness, social mobilization and community engagement which are carried out in all the four

tiers of healthcare service provision. Specifically, screening services begin from dispensaries and

health centers towards higher tier facilities (10). On the hand, cancer diagnosis and treatment is

predominantly offered in secondary and tertiary referral facilities (10). According Kenya's

National Cancer Policy for 2019-2030, it is argued that nearly half (40%) of the cancers can be

prevented by mitigating the risk factors and implementing already established evidence-based

prevention strategies (9). The common risk factors include being overweight or obese (27.9%),

inadequate consumption of fruits and vegetables (94%), tobacco and alcohol use among 13.3%

and 19.3% of the newly diagnosed individuals respectively (9). It is estimated that many women

harbor human papilloma virus (HPV) types 16 (9.1%) and 18 (63.1%) strains that is attributed to

invasive cervical cancer (11). However, it has been previously reported that there is a low

awareness of cancer and its risk factors in Kenya (12) both in the general population and among

health care service providers (13,14). There are also gaps in the implementation of some of the

policies and guidelines for prevention of cancer. This necessitates a local study to determine

healthcare workers knowledge on screening, diagnosis and treatment of breast and cervical

cancer.

The Kenya Government set medium term plans for the implementation of vision 2030. The

current third medium-term plan (2018-2022) set the universal healthcare coverage (UHC) policy

to ensure the provision of high quality and affordable healthcare for all its citizens (15). To

achieve this, the government noted an insufficiency of medical equipment in both county and

national hospitals. The government rolled-out the managed equipment scheme (MES) as a

private-public partnership (PPP) between the government and private medical equipment service

providers (16). For the diagnosis and management of cancer, the government supplied medical

imaging equipment such as X-rays, Ultrasounds, CT-Scans and MRI machines (9). Although this

project was noble, it was noted that the managed equipment scheme was not properly

implemented due to inadequate personnel (17) to operate them and the backbone supporting

infrastructure such as availability of electricity (16). Because of this challenge, this study aimed

to assess the preparedness of health care facilities in provision of Breast and Cervical Cancer

services in Trans Nzoia and Busia Counties in Kenya. It established the level of healthcare

workers knowledge on the screening, diagnosis and treatment of breast and cervical cancers as

well the availability of medical equipment and supplies for screening, diagnosis and management

of the two diseases in selected two counties of Western Kenya.

Methods

This cross-sectional study was conducted in Busia and Trans Nzoia counties in Western Kenya,

whose breast and cervical cancer oncology programmes are supported by the Academic Model

Providing Access To Healthcare – AMPATH (18,19). Trans Nzoia County is in the former Rift

Valley Province of Kenya. The county borders Bungoma to the west, Uasin Gishu and

Kakamega to the south, Elgeyo Marakwet to the east and the republic of Uganda to the

Northwest. Trans Nzoia covers an area of 2495.5 Km^2 with a population of 956,559 of whom

49.7% are male. Approximately 50% of Trans-Nzoia population is aged 18 to 64 years and 2.8%

aged above 65 years. The life expectancy in this County is 60.5years which is lower than the

national average at 63.4 years. Trans-Nzoia County has a high poverty rate of 50.1%. Trans

Nzoia County has 198 community units served by 38 dispensaries, 8 health centers, 6 sub-

County health facilities and 1 County level hospital a total 53. Busia County borders Kakamega

county to the east, Bungoma to the north, Lake Victoria and Siaya county to the South and

Uganda to the west. It is 1628.4 Kilometer Squared (Km^2) with a population of 953,337 people

of whom 52% are above 18 years, and 47.8% are males. Life expectancy is low at 47years.

Poverty levels are high at 64.2% compared to the national levels at 45.9%. Busia has 184

community units served by 65 healthcare facilities namely: 47 dispensaries, 12 health centers, 5

sub-county and 1 County hospital. The two counties have been reported to have a significantly

high prevalence of breast and cervical cancer.

The study adopted service provision assessment (SPA) survey technique to assess the healthcare

providers level of knowledge on breast and cervical cancer as well as determine the availability

of medical equipment as per the WHO priority list between September to December 2018. An

interviewer assisted questionnaire was used to collect socio-demographic, knowledge of breast

and cervical cancer risk factors, screening strategies, symptoms, diagnosis and treatment of the

disease among the healthcare workers enrolled. A structured facility questionnaire was used to

explore the selected institutions' preparedness to screen, diagnose and treat breast and cervical

cancer. Specifically, the facility questionnaire assessed level of care provided and availability of

specific equipment as per the WHO priority list. Data collection was done by trained research

assistants who administered questionnaires which had been   pre-programmed into the Research

Electronic Data capture (REDCap) database as well as a mobile application which allowed for

direct data entry, daily cleaning and validation by the team. Healthcare workers responses from

the questionnaire were scored to assess knowledge on breast and cervical cancer and weighted

means computed with their corresponding standard deviations. Descriptive statistical analysis of

frequency with corresponding proportions were used to compare the availability of medical

equipment in the two counties under review. All analysis were computed using standard

statistical (STATA) version 15. This study received ethical approval from the Institutional

Research and Ethics Committee (IREC) of MTRH/Moi University (approval number 0002090).

This was followed by a research permit from the National Commission for Science technology

and Innovation (NACOSTI) of Kenya (approval number NACOSTI/P/18/74238/24329) and the

two County Health Management Teams (CHMTs). A written informed consent was obtained

from all healthcare workers prior to enrollment. Data confidentiality was ensured by

deidentifying participants responses while privacy was enhanced through used of password

protected database and limited access to the datasets.

Results

Healthcare workers' knowledge on breast and cervical cancer

This study enrolled 73 healthcare service providers in two counties (Busia and Trans Nzoia) in

Western Kenya. They were stratified based on the level of training as either clinical officers

(m=29), nurses (n=37), medical officers or consultants (n=7) and assessed on their knowledge of

risk factors, symptoms, screening, diagnosis and treatment of breast and cervical cancer. The

mean number of correct answers were scored based on the probable total correct responses.

On Breast cancer there were varying responses on the probable risk factors for breast cancer.

Most clinical officers thought that a history of alcohol consumption was a major risk factor

followed by using contraceptive pills. On the other hand, most of the nurses also thought that

alcohol consumption and having a first child at an older age were the major risk factors for breast

cancer. Similarly, more than half of medical officers and consulting clinicians considered alcohol

consumption as a risk factor for breast cancer. All the healthcare workers enrolled described

mastectomy alone as the treatment for breast cancer. There were higher levels of knowledge on

breast cancer risk factors and screening among medical officers or consultant physicians

compared to clinical officers and nurses. However, nurses appeared to have a higher mean

knowledge score on the symptoms for breast cancer.

Regarding cervical cancer, the overall mean knowledge score was highest among medical

officers or consultant and lowest among nurses across all domains assessed namely: risk factors,

screening, diagnosis and treatment) Notably nurses had more knowledge on symptoms of both

cervical and breast cancers compared to other cadres (Table 1).

Medical equipment distribution

During the initial baseline survey conducted in both counties (Busia and Trans Nzoia), it was

noted that Busia county only had 6 ultrasound machines available for breast cancer screening and

with half of them (3) in sub-county hospitals, one in a health center while the county referral

hospital had two available. It was striking that 35 (87.5%) of the 40 healthcare facilities in the

county did not have a single ultrasound machine. Furthermore, x-ray equipment that would

support in further screening was only available in 5 (12.5%) healthcare facilities within Busia

County, with the county hospital and one-subcounty having two x-ray equipment each. For

cervical cancer screening, speculum availability was nearly evenly distributed in the healthcare

facilities in Busia County with 25% (n=10) of the facilities having between 6-10 speculums

while 22.5% (n=9) having more than 21. Gynecological beds were only available in 12 (30%) of

all the healthcare facilities assessed in Busia county with most of the gynecological beds being

concentrated in County and sub-County hospitals. No public healthcare facility in Busia county

had a biopsy needle for cervical cancer diagnosis while sterilization equipment were found in

90% of all the public healthcare facilities. There were seven (17.5%) had cryotherapy equipment

available in the public healthcare facilities in Busia County, however nearly all (97.5%) of the

healthcare facilities did not have a LEEP equipment for the management of cervical lesions with

the only available ones (n=2) being in Busia County referral hospital (Table 2).

In Trans Nzoia County, there were 7 ultrasound machines serving 33 public healthcare facilities

within the county for the screening of breast cancer. There was only one out of the six sub-

county hospitals with an ultrasound machine, while 2 of the 15 health centers had 2 ultrasound

machines each and the county referral hospital equally had 2 machines. This study noted that

there were 9 x-ray machines that could support in cancer screening within Trans Nzoia county

where a county and sub-county hospital having 2 each while a health center having three x-ray

machines. Most (36.4%; n=12) of the facilities assessed had a 1-5 speculum for cervical cancer

screening followed by those with 6-10 speculum in 11 (33.3%) facilities. However, a single

dispensary did not have any speculum. Public healthcare facilities in Trans Nzoia county had 44

gynecological beds distributed among 27 of the 33 facilities with most of the gynecological beds

being concentrated in County (4) and sub-County (12) hospitals. Similar to Busia County, no

public healthcare facility in Trans Nzoia county had a biopsy needle for cervical cancer

diagnosis. However, only 4 (12.1%) of the facilities surveyed in Trans Nzoia county lacked a

sterilization equipment. There were 13 (39.4%) cryotherapy equipment available in the public

healthcare facilities within Trans Nzoia County, where majority (8) of these were in health

centers. This study reports the availability of a single LEEP equipment stationed at a health

center in Trans Nzoia county (Table 3).

Discussion

Knowledge on breast and cervical cancer

In this study there was a low level of knowledge among health care providers on the risk factors,

symptoms, screening, diagnosis and treatment of both breast and cervical cancer. This finding is

consistent with a previous study conducted in Kenya (20) which found low level of knowledge

among healthcare workers on breast and cervical cancer. This finding was across various cadres

and level of training of the healthcare workers. This low level of knowledge could directly

influence the kind of care patients with both cancers could receive in public healthcare facilities

located in the devolved governance units of Kenya. In a community study conducted in the

coastal region of Kenya (12), the key barriers affecting access breast cancer care were lack of

awareness on breast and cervical cancer risk factors, signs and symptoms among the community

members enrolled. In addition to stigma towards patients seeking cervical cancer treatment and

the overall negative attitudes towards patients with cancer, cultural beliefs and personal

discomfort were some of the other reasons for given low access to cancer care. Among the health

care workers who were interviewed, it was noted that they too had limited knowledge on breast

cancer (12). The Kenya Cancer Policy of 2019 to 2030 further acknowledges that there exist a

knowledge-gap in the level of knowledge on various forms of cancer among human resources for

health (9). The Ministry of Health in Kenya has instituted various mechanisms to train more

healthcare workers on by conducting capacity building activities and mentorship to primary care

providers in counties by coordinating specialist outreach programs in collaboration with tertiary

facilities. Furthermore, Kenya has initiated diploma, masters and oncology fellowship programs

in addition to ongoing pre-service and in service training for primary care workers to strengthen

cancer care service delivery (9,10).

Low levels on knowledge in cancer are not limited to healthcare workers practicing in public

hospitals in Kenya. In Rwanda (21), it was reported that cancer care is mainly offered by

physicians who work in collaboration with colleagues in USA and Canada through virtual tumor

boards. The authors (21) further noted that in Rwanda, there are gaps in prevention and early

detection of breast and cervical cancer despite introduction of mitigation measures such as breast

surgery fellowship at the national military hospital.

Medical Equipment

This study based its assessment on the availability and distribution of medical equipment for the

management of cancer on the World Health Organization's priority list of medical devices for

cancer management of 2017 (22). This priority list provides an implementation reference to be

adopted based on the needs of the healthcare system in the specific country and serves as a

standard by which we can assess preparedness to provide cancer care. The guideline stipulates

the minimum requirements for prevention, diagnosis, treatment, follow up and palliative care for

all cancers. For breast cancer prevention, physical examination through bimanual palpation of

breasts and locoregional lymph nodes as well as mammography is recommended in low resource

healthcare settings, while an MRI breast may be used in combination with mammography in

certain high-risk patients. Among the 73 facilities assessed, none of them had a mammogram or

MRI to support breast cancer screening. This is because most of these facilities are available in

national teaching hospitals as well as a few tier – four hospitals in some counties in Kenya. This

finding is consistent to that reported in Rwanda (21) where MRI machines were available in one

national public and one private hospital. The diagnosis of breast cancer involves thorough

bimanual palpation of the breasts and locoregional nodes, blood testing that includes complete

blood count and blood chemistries (liver and renal function tests, alkaline phosphatase and

calcium), breast imaging initially bilateral mammography followed by an ultrasound of the

breast and locoregional nodes. Following imaging, an ultrasound-guided core needle biopsy, a

stereo-static guided or manual biopsy sampling is done for histopathological diagnosis and

biomarker analysis of the sample. In the facilities assessed in this study, ultrasound services were

available in 5 of the 40 public healthcare facilities in Busia County while in Trans Nzoia only 4

out of the 33 facilities in the county. However, none of these counties reported having biopsy

needles to support ultrasound-guided core needle biopsy for breast cancer diagnosis. This is

because only a few county hospitals in Kenya have the capacity to offer histopathology services

contributing to diagnostic delays (9). Computerized tomography scanning of the chest as well as

a chest x-Ray in low resource settings could be used in the staging of breast cancer. In this study,

x-ray services were not available in many facilities in both Trans Nzoia and Busia counties with

only 9 and 5 machines available respectively. The current cancer Policy in Kenya (9) highlights

that a major challenge in the adequate distribution of diagnostic equipment to healthcare

institutions in Kenya is a mismatch between human resource capacity to operationalize the

equipment and supplies. Furthermore, the policy notes a lack of clear policy guideline on

distribution, placement, monitoring and evaluation of medical equipment. That is why this study

noted a lack of consistency in the distribution of ultrasound and x-ray machines across the

various tiers of the healthcare facilities in the counties reviewed. All these equipment availability

challenges in both counties could be a major contributor to the late diagnosis of breast cancer

currently being witnessed in Kenya (2,9,10).

The current cervical cancer screening and prevention strategies (11,23) involve preventive

immunization against human papilloma virus (HPV) for girls, screening of pre-cancerous lesions

using a speculum or vaginal examinations and inspection with acetic acid (VIA), conducting of a

HPV test and treating of pre-cancerous lesions using cryotherapy and large loop excision of

transformation (LEEP). In this study, most facilities assessed in Trans Nzoia had a minimum of

1-5 speculums to support screening and a single gynecological bed available, however only a

single sub-county facility had LEEP and a cryotherapy equipment. In Busia county, most

facilities had 11-15 specula available, with two LEEP equipment in the county hospital and five

cryotherapy equipment. A biopsy needle is needle is needed in the definitive diagnosis of

invasive cervical cancer in the second care level (22). However, no facility in both counties had

a core biopsy needle. An abdominal ultrasound is often used in the staging of cervical cancer as

well as chest x-ray. However, there were limited ultrasound and x-ray facilities in the studies

sampled.

312   **Conclusions and Recommendations**

A below average level of knowledge on breast and cervical cancer among the healthcare workers

attending to patients in public healthcare facilities was found in both Busia and Trans Nzoia

counties. Furthermore, there was a disparity in the distribution and quantity of priority medical

equipment for the screening, diagnosis and treatment of breast and cervical cancer in the two

county hospitals. From the findings of this study, there is need for targeted in-service training

among nurses on breast and cervical cancer risk factors and screening modalities as they are the

first contacts with suspected cancer patients and are also involved patient medical education.

Advanced cancer diagnostic equipment such as core-biopsy needles and mammograms should be

provided in these public healthcare facilities to support breast and cervical cancer staging and

risk assessment. Specialized training of healthcare workers is needed to reduce the mismatch

between human resource capacity to operationalize the medical equipment and their supply.

Finally, there is need for proper guidelines to guide the distribution of medical equipment by
facilities level of care and human resource availability.

REFERENCES

1. IARC. Global Cancer Observatory [Internet]. World Health Organization. 2021 [cited
2021 May 11]. Available from: https://gco.iarc.fr/

2. Globocan. Kenya Cancer Report. 2020.

3. Tiruneh FN, Chuang KY, Ntenda PAM, Chuang YC. Individual-level and community-
level determinants of cervical cancer screening among Kenyan women: A multilevel
analysis of a Nationwide survey. BMC Womens Health [Internet]. 2017 Nov 15 [cited
2021 May 3];17(1):1–14. Available from:
https://link.springer.com/articles/10.1186/s12905-017-0469-9

4. de Sanjose S, Tsu VD. Prevention of cervical and breast cancer mortality in low-and
middle-income countries: A window of opportunity. Int J Womens Health. 2019;11:381–
6.

5. Kangethe JM, Gachuno OW, Komu JG, Mutai KK, Lwembe RM. Reasons for Low
Uptake of Cervical Cancer Preventive Services in Kenya and other Low-Middle Income
Countries: A Review Article. 2018;06.

6. USAID. The DHS Program - Service Provision Assessments (SPA) [Internet]. 2021 [cited
2021 May 12]. Available from: https://dhsprogram.com/methodology/Survey-
Types/SPA.cfm

7. WHO. Global action plan for the prevention and control of noncommunicable diseases

2013-2020. Geneva; 2013.

8.   UNASD. Transforming our World: The 2030 Agenda for sustainable development. United

Nations Agenda for Sustainable Development. 2016.

9.   MOH-Kenya. Kenya Cancer Policy 2019-2030. Nairobi; 2019.

10.   MOH-Kenya. Kenya National Cancer Screening Guidelines [Internet]. Ministry of Health

- Cancer Care Guidelines. 2018. Available from: www.health.go.ke

11.   ICO/IARC. Human Papillomavirus and Related Cancers - Fact Sheet 2018 (Kenya)

[Internet]. 2019 [cited 2021 May 12]. Available from: www.hpvcentre.net

12.   Sayed S, Ngugi AK, Mahoney MR, Kurji J, Talib ZM, MacFarlane SB, et al. Breast

Cancer knowledge, perceptions and practices in a rural Community in Coastal Kenya.

BMC Public Health [Internet]. 2019 Feb 12 [cited 2021 May 12];19(1):1–13. Available

from: https://doi.org/10.1186/s12889-019-6464-3

13.   Makau-Barasa LK, Greene SB, Othieno-Abinya NA, Wheeler S, Skinner A, Bennett A V.

Improving access to cancer testing and treatment in Kenya. J Glob Oncol. 2017;4:1–8.

14.   Njuguna F, Martijn H, Langat S, Musimbi J, Muliro H, Skiles J, et al. Factors influencing

time to diagnosis and treatment among pediatric oncology patients in Kenya. Pediatr

Hematol Oncol [Internet]. 2016 [cited 2021 May 12];33(3):186–99. Available from:

https://doi.org/10.3109/08880018.2016.1169566

15.   GOK. Kenya Vision 2030: Third Medium Term Plan (2018-2022). Transforming Lives:

Advancing socio-economic development through the "Big Four." 2018.

16.   Mutua J, Wamalwa N. Leasing of Medical Equipment Project in Kenya: Value for Money
366   Assessment   [Internet].   2020.   Available   from:
367   https://www.africaportal.org/publications/leasing-medical-equipment-project-kenya-
368   value-money-assessment/

369   17.   Atieno OM, Opanga S, Martin A, Kurdi A, Godman B. Pilot study assessing the direct
370   medical cost of treating patients with cancer in Kenya; findings and implications for the
371   future.   J   Med   Econ   [Internet].   2018;21(9):878–87.   Available   from:
372   https://doi.org/10.1080/13696998.2018.1484372

373   18.   Strother RM, Asirwa FC, Busakhala NB, Njiru E, Orang'o E, Njuguna F, et al. The
374   evolution of comprehensive cancer care in Western Kenya. J Cancer Policy [Internet].
375   2013;1(1–2):e25–30. Available from: http://dx.doi.org/10.1016/j.jcpo.2013.04.001

376   19.   Rosen B, Itsura P, Tonui P, Covens A, van Lonkhuijzen L, Orang'o EO. Development of
377   a comprehensive and sustainable gynecologic oncology training program in western
378   Kenya, a low resource setting. Gynecol Oncol Reports [Internet]. 2017;21(June):122–7.
379   Available from: http://dx.doi.org/10.1016/j.gore.2017.06.014

380   20.   Makau-Barasa LK, Greene SB, Othieno-Abinya NA, Wheeler S, Skinner A, Bennett A V.
381   Improving access to cancer testing and treatment in Kenya. J Glob Oncol. 2018 Aug
382   4;2018(4).

383   21.   Rubagumya F, Costas-Chavarri A, Manirakiza A, Murenzi G, Uwinkindi F, Ntizimira C,
384   et al. State of Cancer Control in Rwanda: Past, Present, and Future Opportunities. JCO
385   Glob Oncol. 2020;(6):1171–7.

22.   World Health Organization (WHO). WHO list of priority medical devices for cancer

management. Geneva: World Health Organization; 2017. [Internet]. Vol. 18, WHO press.

2017.   856   p.   Available   from:

https://apps.who.int/iris/bitstream/handle/10665/255262/9789241565462-

eng.pdf?sequence=1

23.   Khozaim K, Orang'O E, Christoffersen-Deb A, Itsura P, Oguda J, Muliro H, et al.

Successes and challenges of establishing a cervical cancer screening and treatment

program in western Kenya. Int J Gynecol Obstet [Internet]. 2014;124(1):12–8. Available

from: http://dx.doi.org/10.1016/j.ijgo.2013.06.035

Table 1: Level of Provider Knowledge

Knowledge Indicators	Mean score Knowledge on breast cancer			Expected Score	Mean score Knowledge on cervical cancer			Expected Score
	Provider Training level				Provider Training level			
	CO N= 29	Medic/ Consultant N =7	Nurse N= 37		CO N= 29	Medic/ Consultant N =7	Nurse N= 37	
Risk Factors	3.48(1.81)	3.71(1.89)	2.27(1.35)	**11**	3.79(1.29)	4(1))	1.05(1.05)	**10**
Symptoms	1.57(1.57)	1.5(1.5)	1.74(1.74)	**10**	1.25(1.25)	1.35(1.35)	2.65(0.79)	**7**
Screening	2.62(0.86)	2.71(0.95)	1.97(0.76)	**4**	2.93(0.84)	3(0.58)	0.89(0.61)	**4**
Diagnosis	0.93(0.53)	0.86(0.69)	0.51(0.56))	**2**	1.07(0.53)	1.71(0.76)	1.05(1.05)	**3**
Treatment	1	1	1	**6**	2.72(1.03)	3.29(0.95)	2.46(0.96)	**8**

397 **Table 2: Medical equipment availability by healthcare facility's level of care in Busia County**

Variable name	Value	Total (N=40)	Sub County n=7 (18%)	Health Centre n=10 (25%)	Dispensary n=22 (55%)	County Referral n=1 (2%)
Ultrasounds machines						
	0	35 (87.5%)	4 (57.1%)	9 (90%)	22 (100%)	0 (0%)
	1	4 (10%)	3 (42.9%)	1 (10%)	0 (0%)	0 (0%)
	2	1 (2.5%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)
X-ray machine						
	0	35 (87.5%)	3 (42.9%)	10 (100%)	22 (100%)	0 (0%)
	1	3 (7.5%)	3 (42.9%)	0 (0%)	0 (0%)	0 (0%)
Speculum available						
	0	1 (2.5%)	0 (0%)	0 (0%)	1 (4.5%)	0 (0%)
	1-5	10 (25%)	0 (0%)	2 (20%)	8 (36.4%)	0 (0%)
	6-10	10 (25%)	1 (14.3%)	2 (20%)	7 (31.8%)	0 (0%)
	11-15	7 (17.5%)	4 (57.1%)	2 (20%)	1 (4.5%)	0 (0%)
	16-20	3 (7.5%)	1 (14.3%)	0 (0%)	1 (4.5%)	1 (100%)
	Above 21	9 (22.5%)	1 (14.3%)	4 (40%)	4 (18.2%)	0 (0%)
Gynecological beds						
	0	28 (70%)	4 (57.1%)	6 (60%)	18 (81.8%)	0 (0%)
	1	6 (15%)	0 (0%)	3 (30%)	3 (13.6%)	0 (0%)
	2	6 (15%)	3 (42.9%)	1 (10%)	1 (4.5%)	1 (100%)
Biopsy needles						
	0	40 (100%)	7 (100%)	10 (100%)	22 (100%)	1 (100%)
Sterilization Equipment						
	0	4 (10%)	0 (0%)	0 (0%)	4 (18.2%)	0 (0%)
	1	23 (57.5%)	3 (42.9%)	6 (60%)	14 (63.6%)	0 (0%)
	2	10 (25%)	3 (42.9%)	4 (40%)	2 (9.1%)	1 (100%)
	3	2 (5%)	0 (0%)	0 (0%)	2 (9.1%)	0 (0%)
	4	1 (2.5%)	1 (14.3%)	0 (0%)	0 (0%)	0 (0%)
Cryotherapy equipment						
	0	33 (82.5%)	5 (71.4%)	9 (90%)	19 (86.4%)	0 (0%)
	1	6 (15%)	2 (28.6%)	1 (10%)	3 (13.6%)	0 (0%)
	4	1 (2.5%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)
LEEP equipment						
	0	39 (97.5%)	7 (100%)	10 (100%)	22 (100%)	0 (0%)
	2	1 (2.5%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)

Table 3: Medical equipment availability by healthcare facility's level of care in Trans Nzoia
County

Variable name	Value	Total (N=40)	Sub County n=7 (18%)	Health Centre n=10 (25%)	Dispensary n=22 (55%)	County Referral n=1 (2%)
Ultrasounds machines						
	0	29 (87.9%)	5 (83.3%)	13 86.7%)	11 (100%)	0 (0%)
	1	1 (3%)	1 (16.7%)	0 (0%)	0 (0%)	0 (0%)
	2	3 (9.1%)	0 (0%)	2 (13.3%)	0 (0%)	1 (100%)
X-ray machine						
	0	28 (84.8%)	4 (66.7%)	14 (93.3%)	10 (90.9%)	0 (0%)
	1	2 (6.1%)	1 (16.7%)	1 (6.7%)	0 (0%)	0 (0%)
	2	2 (6.1%)	1 (16.7%)	0 (0%)	0 (0%)	1 (100%)
	3	1 (3%)	0 (0%)	0 (0%)	1 (9.1%)	0 (0%)
Speculum available						
	0	1 (3%)	0 (0%)	0 (0%)	1 (9.1%)	0 (0%)
	1-5	12 (36.4%)	0 (0%)	5 (33.3%)	7 (63.6%)	0 (0%)
	11-15	4 (12.1%)	1 (16.7%)	3 (20%)	0 (0%)	0 (0%)
	16-20	2 (6.1%)	0 (0%)	1 (6.7%)	1 (9.1%)	0 (0%)
	6-10	11 (33.3%)	3 (50%)	6 (40%)	2 (18.2%)	0 (0%)
	> 21	3 (9.1%)	2 (33.3%)	0 (0%)	0 (0%)	1 (100%)
Gynecological beds						
	0	6 (18.2%)	0 (0%)	1 (6.7%)	5 (45.5%)	0 (0%)
	1	14 (42.4%)	2 (33.3%)	8 (53.3%)	4 (36.4%)	0 (0%)
	2	11 (33.3%)	3 (50%)	6 (40%)	2 (18.2%)	0 (0%)
	4	2 (6.1%)	1 (16.7%)	0 (0%)	0 (0%)	1 (100%)
Biopsy needles						
	0	33 (100%)	6 (100%)	15 (100%)	11 (100%)	1 (100%)
Sterilization Equipment						
	0	4 (12.1%)	0 (0%)	0 (0%)	4 (36.4%)	0 (0%)
	1	20 (60.6%)	5 (83.3%)	10 66.7%)	5 (45.5%)	0 (0%)
	2	6 (18.2%)	1 (16.7%)	3 (20%)	2 (18.2%)	0 (0%)
	3	2 (6.1%)	0 (0%)	2 (13.3%)	0 (0%)	0 (0%)
	5	1 (3%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)
Cryotherapy equipment						
	0	20 (60.6%)	4 (66.7%)	7 (46.7%)	9 (81.8%)	0 (0%)
	1	13 (39.4%)	2 (33.3%)	8 (53.3%)	2 (18.2%)	1 (100%)
LEEP equipment						
	0	32 (97%)	6 (100%)	14 93.3%)	11 (100%)	1 (100%)
	1	1 (3%)	0 (0%)	1 (6.7%)	0 (0%)	0 (0%)

Acknowledgements

This research was supported with funds from Access Accelerated and by cooperative agreement between World Bank and Ministry of Health Kenya NCDs Division. The management of Moi Teaching and Referral Hospital (MTRH), Moi University and the Counties of Busia and Trans Nzoia for an implementation agreement that enabled the data collection in the selected public hospitals. We further acknowledge the study participants as well as AMPATH for providing technical support for this study.