

25

Abstract

26 Cancer is a major cause of morbidity and mortality globally. This study aimed to determine the
27 preparedness of public health care facilities in the provision of breast and cervical cancer
28 services by assessing healthcare provider's knowledge on risk factors, screening, symptoms,
29 diagnosis and treatment of the diseases as well as availability of medical equipment required in
30 their management in Western Kenya. This was achieved by using a cross-sectional service
31 provision assessment (SPA) baseline survey technique in Busia and Trans-Nzoia counties of
32 Western Kenya between October and December 2018. Data was collected using an interviewer-
33 assisted questionnaires from healthcare workers, while a structured facility questionnaire was
34 used to assess the level of preparedness of the selected public healthcare facilities. We enrolled
35 73 healthcare workers 37 (50.6%) of whom were nurses, followed by clinical officers and
36 medical officers. The highest proportion of knowledge on risk factors and screening of breast
37 and cervical cancer was reported among medical officers and consultant physicians while nurses
38 scored highly on the symptoms of breast and cervical cancer. The medical equipment required
39 for breast and cervical cancer screening and diagnosis were found in most healthcare facilities;
40 however, none of them had core-biopsy needles. A single LEEP equipment was found in a health
41 center within Trans Nzoia County while two LEEP equipment were stationed at the Busia county
42 hospital. We report a below average level of knowledge on breast and cervical cancer among the
43 healthcare workers at public healthcare facilities in both Busia and Trans Nzoia counties. There
44 was a disparity in the distribution and quantity of priority medical equipment for the screening,
45 diagnosis and treatment of breast and cervical cancer in the two county hospitals. There is need
46 for capacity building among healthcare workers as well as standardization in the distribution and
47 utilization of these medical equipment.

48 **Keywords:** *Cervical Cancer, Breast Cancer, Preparedness, Service Provision Assessment*
49 *(SPA), Public Healthcare Facilities, Western Kenya, Busia, Trans Nzoia.*

50

51

Introduction

52 Cancer is a major cause of morbidity and mortality across the globe. According to the
53 International Agency for Research on Cancer global cancer observatory report [1],
54 approximately 1 in 5 people develop cancer during their lifetime. Of these, 1 in 8 men and 1 in
55 11 women die from the disease. With a rising proportion of the ageing populations globally,
56 coupled with lifestyle changes, the prevalence of cancer is on the rise. The most common forms
57 of cancer among women globally are breast, colorectal, lung, cervical, and thyroid cancers [2].
58 The likelihood of breast cancer is 1 in 4 cancers diagnosed among women globally [1,2]. In
59 Africa the incidence for breast cancer in 2020 was estimated at 16.8% followed by cervical
60 cancer at 10.6% [3]. In Kenya, the incidence of breast cancer in 2020 was estimated at 16.1%
61 followed by second placed cervical cancer at 12.4% of all women diagnosed with cancer [2].
62 With a greater adoption of a more Westernized lifestyle (especially in diet, reproductive health
63 and reduced physical activity) in Kenya and many other sub-Saharan African countries, the
64 incidence of breast and cervical cancer has seen a steady rise [4]. These low- and medium-
65 income countries (LMIC) have been shown to bear a greater burden of various forms of cancer
66 despite advancements in screening, diagnosis and management options [5]. To adequately
67 address the disease burden attributed to these two cancers, there is need to have effective and
68 adequately prepared health service delivery systems. The level of preparedness and effectiveness
69 of these health systems can be assessed using multiple dimensions to assess the progress in
70 addressing the disease burden. Although various tools could be used in this assessment, a Service
71 Provision Assessment (SPA) survey could be used to provide a comprehensive understanding of
72 the health service delivery system in a particular country [6]. This is because they support the
73 strengthening of health systems in developing countries by collecting information on the

74 availability of different resources based in the health facilities within a country. The SPA survey
75 describes the facility-types with the ability to offer specific health services. It defines the
76 infrastructural and human resource preparedness to offer the said services in the health facilities
77 [6]. Furthermore, it assesses adherence to the set quality standards for healthcare service
78 provision and whether the clients and service providers are content with the service delivery
79 environment.

80 The World Health Organization's Global action plan for the prevention and control of
81 noncommunicable diseases 2013-2020 targeted a 25% reduction in risk of premature death
82 attributed to non-communicable diseases including cancer by 2025 [7]. Furthermore, the United
83 Nations Agenda for Sustainable Development recommends a one-third reduction in the
84 proportion of premature death associated with non-communicable diseases such as cancer by the
85 year 2030 using multiple approaches such as prevention, treatment and promotion of mental
86 health and well-being [8]. The public and private health care service delivery institutions should
87 have an 80% availability of the affordable basic technologies and essential medicines for the
88 management of these noncommunicable diseases. The report [7] recommends periodic needs
89 assessment and evaluation of resource need such as workforce, institutional and research
90 capacity to address the non-communicable diseases. The eight key indicators include a
91 comprehensive range of health services provided to the target population, the services should be
92 directly and permanently accessible without barriers of entry, they should adequately cover the
93 defined target population and provide continuity of care across the network of services, health
94 conditions, levels of care, and over the life cycle. The quality of these services should be high as
95 evidenced by their effectiveness and safety as well as being organized around the person and not
96 the disease or the financing mechanism. They should be properly coordinated at the local level

97 and across the different types of providers, types of care and levels of service delivery with
98 adequate accountability and efficiency [7].

99 In Kenya, the healthcare service delivery system is stratified into four tiers as [9]: Tier 1
100 (Community Health Services), Tier 2 (primary healthcare facilities such as dispensaries and
101 health centers), Tier 3 (secondary referral facilities such as sub county and county hospitals) and
102 Tier 4 (Tertiary facilities such as national referral hospitals). The primary care for cancer
103 involves prevention of risk factors and screening, enhancing health promotion through public
104 awareness, social mobilization and community engagement which are carried out in all the four
105 tiers of healthcare service provision. Specifically, screening services begin from dispensaries and
106 health centers towards higher tier facilities [10]. On the hand, cancer diagnosis and treatment is
107 predominantly offered in secondary and tertiary referral facilities [10]. According Kenya's
108 Cancer Policy for 2019-2030, it is argued that nearly half (40%) of the cancers can be prevented
109 by mitigating the risk factors and implementing already established evidence-based prevention
110 strategies [9]. The common risk factors include being overweight or obesity (27.9%), inadequate
111 consumption of fruits and vegetables (94%), tobacco and alcohol use at 13.3% and 19.3%
112 respectively [9]. It is estimated that many women harbor human papilloma virus (HPV) types 18
113 (63.1%) followed by 16 (9.1%) strains that is attributed to invasive cervical cancer [11].
114 However, it has been previously reported that there is a low awareness of cancer and its risk
115 factors in Kenya [12] both in the general population and among health care service providers
116 [13,14]. Although the National Cancer Screening Guidelines have been disseminated and outline
117 the key cancers for screening as well as the modalities for screening by level of care, there are
118 gaps in the implementation of these guidelines for prevention and screening of cancer. The
119 guidelines recommend screening for women 25-49 years with Human Papillomavirus (HPV)

120 testing as the gold standard method for cervical cancer screening although Visual Inspection with
121 Acetic Acid and Cytology are also recommended where HPV is not available. For breast cancer,
122 mammography is the recommended screening method for women from 40-74 years of age. This
123 necessitated a local study to determine healthcare workers knowledge on screening, diagnosis
124 and treatment of breast and cervical cancer.

125 The Kenya Government set medium term plans for the implementation of vision 2030. The
126 current third medium-term plan (2018-2022) set the universal healthcare coverage (UHC) policy
127 to ensure the provision of high quality and affordable healthcare for all its citizens [15]. To
128 achieve this, the government noted an insufficiency of medical equipment in both county and
129 national hospitals. The government rolled-out the managed equipment scheme (MES) as a
130 private-public partnership (PPP) between the government and private medical equipment service
131 providers [16]. For the diagnosis and management of cancer, the Kenyan government supplied
132 medical imaging equipment such as X-rays, Ultrasounds, CT-Scans, digital mammography and
133 MRI machines through MES in all the 47 counties [9]. Both Busia and Kitale County referral
134 hospitals received a digital mammography machine each for screening and early diagnosis of
135 breast cancer. Although this project was noble, it was noted that the managed equipment scheme
136 was not properly implemented due to inadequate personnel [17] to operate them and the
137 backbone supporting infrastructure such as availability of electricity [16]. Because of this
138 challenge, this study aimed to assess the preparedness of health care facilities in provision of
139 Breast and Cervical Cancer services in Trans Nzoia and Busia Counties in Kenya. It established
140 the level of healthcare workers knowledge on the screening, diagnosis and treatment of breast
141 and cervical cancers as well the availability of medical equipment and supplies for screening,
142 diagnosis and management of the two diseases in selected two counties of Western Kenya.

143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165

Methods

This cross-sectional study was conducted in Busia and Trans Nzoia counties in Western Kenya, whose breast and cervical cancer oncology programmes are supported by the Academic Model Providing Access To Healthcare – AMPATH [18,19]. Trans Nzoia county is in the former Rift Valley Province of Kenya, it borders Bungoma county to the west, Uasin Gishu and Kakamega counties to the south, Elgeyo Marakwet county to the east and the Republic of Uganda to the Northwest. It covers a geographical area of 2495.5 km² with a population of 956,559 of whom 49.7% are male. Approximately half of the population in Trans Nzoia county is aged 18 to 64 years with 2.8% aged more than 65 years and an average life expectancy of 60.5 years which is lower than the national average at 63.4 years. Busia county borders Kakamega county to the east, Bungoma county to the north, Siaya county to the south and Uganda to the west. It has a geographical area of 1628.4 Kilometer Squared (Km²) with a population of 953,337 of whom 52% are above 18 years of age and a low life expectancy of 47 years. The two counties have been reported to have a significantly high prevalence of breast and cervical cancer in Kenya.

The study adopted service provision assessment (SPA) survey technique to assess the healthcare providers level of knowledge on breast and cervical cancer as well as determine the availability of medical equipment as per the WHO priority list between September to December 2018. Healthcare workers heading oncology programs in the selected healthcare facilities were purposively sampled to saturation prior to obtaining written informed consents and being enrolled into the study. An interviewer assisted questionnaire was used to collect socio-demographic, knowledge of breast and cervical cancer risk factors, screening strategies, symptoms, diagnosis and treatment of the disease among the healthcare workers enrolled. Healthcare facilities were selected using a census approach and stratified by level of care offered.

166 A structured facility questionnaire was used to explore the selected institutions' preparedness to
167 screen, diagnose and treat breast and cervical cancer. Specifically, the facility questionnaire
168 assessed level of care provided and availability of specific equipment as per the WHO priority
169 list. Data collection was done by trained research assistants who administered questionnaires
170 which had been pre-programmed into the Research Electronic Data capture (REDCap) database
171 mobile application which allowed for direct data entry, daily cleaning and validation by the team.
172 Healthcare workers responses from the questionnaire were scored (with a point for every correct
173 response) to assess knowledge on breast and cervical cancer and weighted means computed with
174 their corresponding standard deviations. Descriptive statistical analysis of frequency with
175 corresponding proportions were used to compare the availability of medical equipment in the
176 two counties under review. All analysis were computed using standard statistical (STATA)
177 version 15. This study received ethical approval from the Institutional Research and Ethics
178 Committee (IREC) of MTRH/Moi University (approval number 0002090). This was followed by
179 a research permit from the National Commission for Science technology and Innovation
180 (NACOSTI) of Kenya (approval number NACOSTI/P/18/74238/24329) and the two County
181 Health Management Teams (CHMTs). Data confidentiality was ensured by deidentifying
182 participants responses while privacy was enhanced through used of password protected database
183 and limited access to the datasets.

184 **Results**

185 **Healthcare workers' knowledge on breast and cervical cancer**

186 This study enrolled 73 healthcare service providers from two counties (Busia and Trans Nzoia)
187 in Western Kenya. They were stratified based on the level of training as either clinical officers
188 (m=29), nurses (n=37), medical officers or consultants (n=7) and assessed on their knowledge of

189 risk factors, symptoms, screening, diagnosis and treatment of breast and cervical cancer. The
 190 mean number of correct answers were scored based on the probable total correct responses.

191 On Breast cancer, there were varying responses on the probable risk factors for breast cancer.
 192 Nearly half (41%) of the clinical officers thought that a history of alcohol consumption was a
 193 major risk factor followed by using contraceptive pills (31%). However, 35% of the nurses
 194 described having a first child at an older age followed by alcohol consumption (24%) as the
 195 major risk factors for breast cancer. Similarly, more than half of medical officers and consulting
 196 clinicians considered alcohol consumption as a risk factor for breast cancer. All the healthcare
 197 workers enrolled described mastectomy alone as the treatment for breast cancer. There were
 198 higher levels of knowledge on breast cancer risk factors and screening among medical officers or
 199 consultant physicians compared to clinical officers and nurses. However, nurses appeared to
 200 have a higher mean knowledge score on the symptoms for breast cancer.

201 Regarding cervical cancer, the overall mean knowledge score was highest among medical
 202 officers or consultant and lowest among nurses across all domains assessed namely: risk factors,
 203 screening, diagnosis and treatment). Notably, nurses had more knowledge on symptoms of both
 204 cervical and breast cancers compared to other cadres (Table 1).

205 **Table 1: Level of Provider Knowledge**

Knowledge Indicators	Mean score Knowledge on breast cancer				Mean score Knowledge on cervical cancer			
	Provider Training level			Expected Score	Provider Training level			Expected Score
	Clinical Officers	Medical Officers/ Consultants	Nurses		Clinical Officers	Medical Officers/ Consultants	Nurses	
Risk Factors	3.48(1.81)	3.71(1.89)	2.27(1.35)	11	3.79(1.29)	4(1)	1.05(1.05)	10
Symptoms	1.57(1.57)	1.5(1.5)	1.74(1.74)	10	1.25(1.25)	1.35(1.35)	2.65(0.79)	7
Screening	2.62(0.86)	2.71(0.95)	1.97(0.76)	4	2.93(0.84)	3(0.58)	0.89(0.61)	4
Diagnosis	0.93(0.53)	0.86(0.69)	0.51(0.56)	2	1.07(0.53)	1.71(0.76)	1.05(1.05)	3
Treatment	1	1	1	6	2.72(1.03)	3.29(0.95)	2.46(0.96)	8

206 **Medical equipment distribution**

207 During the initial baseline survey conducted in both counties (Busia and Trans Nzoia), it was
208 noted that Busia county only had 6 ultrasound machines available for breast cancer screening and
209 with half of them (3) in sub-county hospitals, one in a health center while the county referral
210 hospital had two available. It was striking that 35 (87.5%) of the 40 healthcare facilities in the
211 county did not have a single ultrasound machine. Furthermore, x-ray equipment that would
212 support in further screening was only available in 5 (12.5%) healthcare facilities within Busia
213 County, with the county hospital and one-subcounty having one and two x-ray equipment
214 respectively. For cervical cancer screening, speculum availability was nearly evenly distributed
215 in the healthcare facilities in Busia County with 25% (n=10) of the facilities having between 6-
216 10 speculums while 22.5% (n=9) having more than 21. Gynecological beds were only available
217 in 12 (30%) of all the healthcare facilities assessed in Busia county with most of the
218 gynecological beds being concentrated in County and sub-County hospitals. No public healthcare
219 facility in Busia county had a biopsy needle for breast and cancer diagnosis while sterilization
220 equipment were found in 90% of all the public healthcare facilities. There were seven (17.5%)
221 had cryotherapy equipment available in the public healthcare facilities in Busia County, however
222 nearly all (97.5%) of the healthcare facilities did not have a LEEP equipment for the
223 management of cervical lesions with the only available ones (n=2) being in Busia County referral
224 hospital (Table 2).

225

226 **Table 2: Medical equipment availability by healthcare facility's level of care in Busia County**

Variable name	Value	Total (N=40)	Sub County n=7 (18%)	Health Centre n=10 (25%)	Dispensary n=22 (55%)	County Referral n=1 (2%)
Ultrasounds machines						
	0	35 (87.5%)	4 (57.1%)	9 (90%)	22 (100%)	0 (0%)
	1	4 (10%)	3 (42.9%)	1 (10%)	0 (0%)	0 (0%)
	2	1 (2.5%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)
X-ray machine						
	0	35 (87.5%)	3 (42.9%)	10 (100%)	22 (100%)	0 (0%)
	1	3 (7.5%)	3 (42.9%)	0 (0%)	0 (0%)	0 (0%)
	2	1 (5%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)
Speculum available						
	0	1 (2.5%)	0 (0%)	0 (0%)	1 (4.5%)	0 (0%)
	1-5	10 (25%)	0 (0%)	2 (20%)	8 (36.4%)	0 (0%)
	6-10	10 (25%)	1 (14.3%)	2 (20%)	7 (31.8%)	0 (0%)
	11-15	7 (17.5%)	4 (57.1%)	2 (20%)	1 (4.5%)	0 (0%)
	16-20	3 (7.5%)	1 (14.3%)	0 (0%)	1 (4.5%)	1 (100%)
	Above 21	9 (22.5%)	1 (14.3%)	4 (40%)	4 (18.2%)	0 (0%)
Gynecological beds						
	0	28 (70%)	4 (57.1%)	6 (60%)	18 (81.8%)	0 (0%)
	1	6 (15%)	0 (0%)	3 (30%)	3 (13.6%)	0 (0%)
	2	6 (15%)	3 (42.9%)	1 (10%)	1 (4.5%)	1 (100%)
Biopsy needles						
	0	40 (100%)	7 (100%)	10 (100%)	22 (100%)	1 (100%)
Sterilization Equipment						
	0	4 (10%)	0 (0%)	0 (0%)	4 (18.2%)	0 (0%)
	1	23 (57.5%)	3 (42.9%)	6 (60%)	14 (63.6%)	0 (0%)
	2	10 (25%)	3 (42.9%)	4 (40%)	2 (9.1%)	1 (100%)
	3	2 (5%)	0 (0%)	0 (0%)	2 (9.1%)	0 (0%)
	4	1 (2.5%)	1 (14.3%)	0 (0%)	0 (0%)	0 (0%)
Cryotherapy equipment						
	0	33 (82.5%)	5 (71.4%)	9 (90%)	19 (86.4%)	0 (0%)
	1	6 (15%)	2 (28.6%)	1 (10%)	3 (13.6%)	0 (0%)
	4	1 (2.5%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)
LEEP equipment						
	0	39 (97.5%)	7 (100%)	10 (100%)	22 (100%)	0 (0%)
	2	1 (2.5%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)

228 In Trans Nzoia County, there were 7 ultrasound machines serving 33 public healthcare facilities
229 within the county for the screening of breast cancer. There was only one out of the six sub-
230 county hospitals with an ultrasound machine, while 2 of the 15 health centers had 2 ultrasound
231 machines each and the county referral hospital equally had 2 machines. This study noted that
232 there were 9 x-ray machines that could support in cancer screening within Trans Nzoia County
233 where a county and sub-county hospital having 2 each while a health center having three x-ray
234 machines. One of the X-ray machines reported in the county hospital of Trans Nzoia was a
235 digital mammogram. Most (36.4%; n=12) of the facilities assessed had a 1-5 speculum for
236 cervical cancer screening followed by those with 6-10 speculum in 11 (33.3%) facilities.
237 However, a single dispensary did not have any speculum. Public healthcare facilities in Trans
238 Nzoia County had 44 gynecological beds distributed among 27 of the 33 facilities with most of
239 the gynecological beds being concentrated in County (4) and sub-County (12) hospitals. Similar
240 to Busia County, no public healthcare facility in Trans Nzoia County had a biopsy needle for
241 cervical cancer diagnosis. However, only 4 (12.1%) of the facilities surveyed in Trans Nzoia
242 County lacked a sterilization equipment. There were 13 (39.4%) cryotherapy equipment
243 available in the public healthcare facilities within Trans Nzoia County, where majority (8) of
244 these were in health centers. This study reports the availability of a single LEEP equipment
245 stationed at a health center in Trans Nzoia County (Table 3).

246

247 **Table 3: Medical equipment availability by healthcare facility's level of care in Trans Nzoia**
 248 **County**

Variable name	Value	Total (N=40)	Sub County n=7 (18%)	Health Centre n=10 (25%)	Dispensary n=22 (55%)	County Referral n=1 (2%)
Ultrasounds machines						
	0	29 (87.9%)	5 (83.3%)	13 (86.7%)	11 (100%)	0 (0%)
	1	1 (3%)	1 (16.7%)	0 (0%)	0 (0%)	0 (0%)
	2	3 (9.1%)	0 (0%)	2 (13.3%)	0 (0%)	1 (100%)
X-ray machine						
	0	28 (84.8%)	4 (66.7%)	14 (93.3%)	10 (90.9%)	0 (0%)
	1	2 (6.1%)	1 (16.7%)	1 (6.7%)	0 (0%)	0 (0%)
	2	2 (6.1%)	1 (16.7%)	0 (0%)	0 (0%)	1 (100%)
	3	1 (3%)	0 (0%)	0 (0%)	1 (9.1%)	0 (0%)
Speculum available						
	0	1 (3%)	0 (0%)	0 (0%)	1 (9.1%)	0 (0%)
	1-5	12 (36.4%)	0 (0%)	5 (33.3%)	7 (63.6%)	0 (0%)
	11-15	4 (12.1%)	1 (16.7%)	3 (20%)	0 (0%)	0 (0%)
	16-20	2 (6.1%)	0 (0%)	1 (6.7%)	1 (9.1%)	0 (0%)
	6-10	11 (33.3%)	3 (50%)	6 (40%)	2 (18.2%)	0 (0%)
	> 21	3 (9.1%)	2 (33.3%)	0 (0%)	0 (0%)	1 (100%)
Gynecological beds						
	0	6 (18.2%)	0 (0%)	1 (6.7%)	5 (45.5%)	0 (0%)
	1	14 (42.4%)	2 (33.3%)	8 (53.3%)	4 (36.4%)	0 (0%)
	2	11 (33.3%)	3 (50%)	6 (40%)	2 (18.2%)	0 (0%)
	4	2 (6.1%)	1 (16.7%)	0 (0%)	0 (0%)	1 (100%)
Biopsy needles						
	0	33 (100%)	6 (100%)	15 (100%)	11 (100%)	1 (100%)
Sterilization Equipment						
	0	4 (12.1%)	0 (0%)	0 (0%)	4 (36.4%)	0 (0%)
	1	20 (60.6%)	5 (83.3%)	10 (66.7%)	5 (45.5%)	0 (0%)
	2	6 (18.2%)	1 (16.7%)	3 (20%)	2 (18.2%)	0 (0%)
	3	2 (6.1%)	0 (0%)	2 (13.3%)	0 (0%)	0 (0%)
	5	1 (3%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)
Cryotherapy equipment						
	0	20 (60.6%)	4 (66.7%)	7 (46.7%)	9 (81.8%)	0 (0%)
	1	13 (39.4%)	2 (33.3%)	8 (53.3%)	2 (18.2%)	1 (100%)
LEEP equipment						
	0	32 (97%)	6 (100%)	14 (93.3%)	11 (100%)	1 (100%)
	1	1 (3%)	0 (0%)	1 (6.7%)	0 (0%)	0 (0%)

250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272

Discussion

Knowledge on breast and cervical cancer

In this study there was a low level of knowledge among health care providers on the risk factors, symptoms, screening, diagnosis and treatment of both breast and cervical cancer. This finding is consistent with a previous study conducted in Kenya [20] which found low level of knowledge among healthcare workers on breast and cervical cancer. This finding was across various cadres and level of training of the healthcare workers. This low level of knowledge could directly influence the kind of care patients with both cancers could receive in public healthcare facilities located in the devolved governance units of Kenya. In a community study conducted in the coastal region of Kenya [12], the key barriers affecting access breast cancer care were lack of awareness on breast and cervical cancer risk factors, signs and symptoms among the community members enrolled. In addition to stigma towards patients seeking cervical cancer treatment and the overall negative attitudes towards patients with cancer, cultural beliefs and personal discomfort were some of the other reasons for given low access to cancer care. Among the health care workers who were interviewed, it was noted that they too had limited knowledge on breast cancer [12]. The Kenya Cancer Policy of 2019 to 2030 further acknowledges that there exist a knowledge-gap in the level of knowledge on various forms of cancer among human resources for health [9]. The Ministry of Health in Kenya has instituted various mechanisms to train more healthcare workers on by conducting capacity building activities and mentorship to primary care providers in counties by coordinating specialist outreach programs in collaboration with tertiary facilities. Furthermore, Kenya has initiated diploma, masters and oncology fellowship programs in addition to ongoing pre-service and in service training for primary care workers to strengthen cancer care service delivery [9,10].

273 Low levels on knowledge in cancer are not limited to healthcare workers practicing in public
274 hospitals in Kenya. In Rwanda [21], it was reported that cancer care is mainly offered by
275 physicians who work in collaboration with colleagues in USA and Canada through virtual tumor
276 boards. The authors [21] further noted that in Rwanda, there are gaps in prevention and early
277 detection of breast and cervical cancer despite introduction of mitigation measures such as breast
278 surgery fellowship at the national military hospital.

279 **Medical Equipment**

280 This study based its assessment on the availability and distribution of medical equipment for the
281 management of cancer on the World Health Organization's priority list of medical devices for
282 cancer management of 2017 [22]. This priority list provides an implementation reference to be
283 adopted based on the needs of the healthcare system in the specific country and serves as a
284 standard by which we can assess preparedness to provide cancer care. The guideline stipulates
285 the minimum requirements for prevention, diagnosis, treatment, follow up and palliative care for
286 all cancers. For breast cancer prevention, physical examination through bimanual palpation of
287 breasts and locoregional lymph nodes as well as mammography is recommended in low resource
288 healthcare settings, while an MRI breast may be used in combination with mammography in
289 certain high-risk patients. Among the 73 facilities assessed, none of them had an MRI machine to
290 support breast cancer screening, however each county referral hospital had a digital
291 mammogram. This finding is consistent to that reported in Rwanda [21] where MRI machines
292 were available in one national public and one private hospital. The diagnosis of breast cancer
293 involves triple assessment through a clinical breast examination by a trained health provider,
294 breast biopsy and breast imaging using mammography, ultrasound or MRI as appropriate [23].
295 Biopsy sampling is usually performed for histopathological diagnosis and biomarker analysis of

296 the sample. In the facilities assessed in this study, ultrasound services were available in 5 of the
297 40 public healthcare facilities in Busia County while in Trans Nzoia only 4 out of the 33
298 facilities in the county. However, none of these counties reported having biopsy needles to
299 support histopathological diagnosis of breast cancer. This is because only a few county hospitals
300 in Kenya have the capacity to offer histopathology services contributing to diagnostic delays [9].
301 Computerized tomography scanning of the chest as well as a chest x-Ray in low resource settings
302 could be used in the staging of breast cancer. In this study, x-ray services were not available in
303 many facilities in both Trans Nzoia and Busia counties with only 9 and 5 machines available
304 respectively. The current cancer Policy in Kenya [9] as well as the Kenya Harmonized Health
305 Facility Assessment 2018/2019 [24] highlight that the major challenge in adequate distribution
306 of diagnostic equipment to healthcare institutions in Kenya is a mismatch between human
307 resource capacity to operationalize the equipment and supplies. This could be a major reason for
308 the low utilization of the already existing medical equipment. Furthermore, the policy notes a
309 lack of clear policy guideline on distribution, placement, monitoring and evaluation of medical
310 equipment. That is why this study noted a lack of consistency in the distribution of ultrasound
311 and x-ray machines across the various tiers of the healthcare facilities in the counties reviewed.
312 All these equipment availability challenges in both counties could be a major contributor to the
313 late diagnosis of breast cancer currently being witnessed in Kenya [2,9,10].

314 The current cervical cancer screening and prevention strategies [11,25] involve preventive
315 immunization against human papilloma virus (HPV) for girls, screening of pre-cancerous lesions
316 using a speculum or vaginal examinations and inspection with acetic acid (VIA), conducting of a
317 HPV test, cytology and treating of pre-cancerous lesions using cryotherapy and large loop
318 excision of transformation (LEEP). In this study, most facilities assessed in Trans Nzoia had a

319 minimum of 1-5 speculums to support screening and a single gynecological bed available,
320 however only a single sub-county facility had LEEP and a cryotherapy equipment. In Busia
321 county, most facilities had 11-15 specula available, with two LEEP equipment in the county
322 hospital and five cryotherapy equipment. A biopsy needle is needed in the definitive
323 diagnosis of invasive breast cancer in the second care level [22]. However, no facility in both
324 counties had a core biopsy needle. An abdominal ultrasound is often used in the staging of
325 cervical cancer as well as chest x-ray. However, there were limited ultrasound and x-ray facilities
326 in the studies sampled.

327 **Conclusions and Recommendations**

328 We report a below average level of knowledge on breast and cervical cancer among the
329 healthcare workers attending to patients in public healthcare facilities within both Busia and
330 Trans Nzoia counties. Furthermore, there was a disparity in the distribution and quantity of
331 priority medical equipment for the screening, diagnosis and treatment of breast and cervical
332 cancer in the two county hospitals. From the findings of this study, there is need for targeted in-
333 service training among nurses on breast and cervical cancer risk factors and screening modalities
334 as they are the first contacts with suspected cancer patients and are also involved patient medical
335 education. Advanced cancer diagnostic equipment such as core-biopsy needles should be
336 provided in these public healthcare facilities to support breast cancer staging and risk assessment.
337 There is need for specialized training of healthcare workers to reduce the mismatch between
338 human resource capacity to operationalize the medical equipment and their supply. Finally,
339 proper guidelines should be drafted in the distribution of medical equipment by the facility's
340 level of care and human resource availability.

341 **Acknowledgements**

342 This research was supported with funds from Access Accelerated and by cooperative agreement
343 between World Bank and Ministry of Health Kenya NCDs Division. The management of Moi
344 Teaching and Referral Hospital (MTRH), Moi University and the Counties of Busia and Trans
345 Nzoia for an implementation agreement that enabled the data collection in the selected public
346 hospitals. We further acknowledge the study participants as well as AMPATH for providing
347 technical support for this study.

348

349 REFERENCES

- 350 1. IARC. Global Cancer Observatory. In: World Health Organization [Internet]. 2021 [cited
351 11 May 2021]. Available: <https://gco.iarc.fr/>
- 352 2. Globocan. Kenya Cancer Report. 2020.
- 353 3. Tiruneh FN, Chuang KY, Ntenda PAM, Chuang YC. Individual-level and community-
354 level determinants of cervical cancer screening among Kenyan women: A multilevel
355 analysis of a Nationwide survey. *BMC Womens Health*. 2017;17: 1–14.
356 doi:10.1186/s12905-017-0469-9
- 357 4. de Sanjose S, Tsu VD. Prevention of cervical and breast cancer mortality in low-and
358 middle-income countries: A window of opportunity. *Int J Womens Health*. 2019;11: 381–
359 386. doi:10.2147/IJWH.S197115
- 360 5. Kangethe JM, Gachuno OW, Komu JG, Mutai KK, Lwembe RM. Reasons for Low
361 Uptake of Cervical Cancer Preventive Services in Kenya and other Low-Middle Income
362 Countries: A Review Article. 2018;06. doi:10.18535/jmscr/v6i12.169

- 363 6. USAID. The DHS Program - Service Provision Assessments (SPA). 2021 [cited 12 May
364 2021]. Available: <https://dhsprogram.com/methodology/Survey-Types/SPA.cfm>
- 365 7. WHO. Global action plan for the prevention and control of noncommunicable diseases
366 2013-2020. Geneva; 2013.
- 367 8. UNASD. Transforming our World: The 2030 Agenda for sustainable development. United
368 Nations Agenda Sustain Dev. 2016. doi:10.1201/b20466-7
- 369 9. MOH-Kenya. Kenya Cancer Policy 2019-2030. Nairobi; 2019.
- 370 10. MOH-Kenya. Kenya National Cancer Screening Guidelines. Minist Heal - Cancer Care
371 Guidel. 2018. Available: www.health.go.ke
- 372 11. ICO/IARC. Human Papillomavirus and Related Cancers - Fact Sheet 2018 (Kenya). 2019.
373 Available: www.hpvcentre.net
- 374 12. Sayed S, Ngugi AK, Mahoney MR, Kurji J, Talib ZM, MacFarlane SB, et al. Breast
375 Cancer knowledge, perceptions and practices in a rural Community in Coastal Kenya.
376 BMC Public Health. 2019;19: 1–13. doi:10.1186/s12889-019-6464-3
- 377 13. Makau-Barasa LK, Greene SB, Othieno-Abinya NA, Wheeler S, Skinner A, Bennett A V.
378 Improving access to cancer testing and treatment in Kenya. J Glob Oncol. 2017;4: 1–8.
- 379 14. Njuguna F, Martijn H, Langat S, Musimbi J, Muliro H, Skiles J, et al. Factors influencing
380 time to diagnosis and treatment among pediatric oncology patients in Kenya. Pediatr
381 Hematol Oncol. 2016;33: 186–199. doi:10.3109/08880018.2016.1169566
- 382 15. GOK. Kenya Vision 2030: Third Medium Term Plan (2018-2022). Transform Lives Adv

- 383 socio-economic Dev through “Big Four.” 2018.
- 384 16. Mutua J, Wamalwa N. Leasing of Medical Equipment Project in Kenya: Value for Money
385 Assessment. 2020. Available: [https://www.africaportal.org/publications/leasing-medical-](https://www.africaportal.org/publications/leasing-medical-equipment-project-kenya-value-money-assessment/)
386 [equipment-project-kenya-value-money-assessment/](https://www.africaportal.org/publications/leasing-medical-equipment-project-kenya-value-money-assessment/)
- 387 17. Atieno OM, Opanga S, Martin A, Kurdi A, Godman B. Pilot study assessing the direct
388 medical cost of treating patients with cancer in Kenya; findings and implications for the
389 future. *J Med Econ.* 2018;21: 878–887. doi:10.1080/13696998.2018.1484372
- 390 18. Strother RM, Asirwa FC, Busakhala NB, Njiru E, Orang’o E, Njuguna F, et al. The
391 evolution of comprehensive cancer care in Western Kenya. *J Cancer Policy.* 2013;1: e25–
392 e30. doi:10.1016/j.jcpo.2013.04.001
- 393 19. Rosen B, Itsura P, Tonui P, Covens A, van Lonkhuijzen L, Orang’o EO. Development of
394 a comprehensive and sustainable gynecologic oncology training program in western
395 Kenya, a low resource setting. *Gynecol Oncol Reports.* 2017;21: 122–127.
396 doi:10.1016/j.gore.2017.06.014
- 397 20. Makau-Barasa LK, Greene SB, Othieno-Abinya NA, Wheeler S, Skinner A, Bennett A V.
398 Improving access to cancer testing and treatment in Kenya. *J Glob Oncol.* 2018;2018.
399 doi:10.1200/JGO.2017.010124
- 400 21. Rubagumya F, Costas-Chavarri A, Manirakiza A, Murenzi G, Uwinkindi F, Ntizimira C,
401 et al. State of Cancer Control in Rwanda: Past, Present, and Future Opportunities. *JCO*
402 *Glob Oncol.* 2020; 1171–1177. doi:10.1200/go.20.00281
- 403 22. World Health Organization (WHO). WHO list of priority medical devices for cancer

404 management. Geneva: World Health Organization; 2017. WHO press. 2017. Available:
405 [https://apps.who.int/iris/bitstream/handle/10665/255262/9789241565462-](https://apps.who.int/iris/bitstream/handle/10665/255262/9789241565462-eng.pdf?sequence=1)
406 [eng.pdf?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/255262/9789241565462-eng.pdf?sequence=1)

407 23. MOH-Kenya. Kenya national cancer treatment protocols. Natl Cancer Control Progr.
408 2019.

409 24. MOH-Kenya. Kenya Harmonized Health Facility Assessment. Encycl Public Heal. 2018.
410 doi:10.1007/978-1-4020-5614-7_1413

411 25. Khozaim K, Orang'O E, Christoffersen-Deb A, Itsura P, Oguda J, Muliro H, et al.
412 Successes and challenges of establishing a cervical cancer screening and treatment
413 program in western Kenya. Int J Gynecol Obstet. 2014;124: 12–18.
414 doi:10.1016/j.ijgo.2013.06.035

415

416 **Authors' contribution**

417 All the authors equally contributed to the writing and final approval prior to submission.