

1 Title:
2 Cardiovascular Disease in the Peruvian Highlands: Local Perceptions, Barriers, and Paths to Preventing
3 Chronic Diseases in Andean Adults

4
5 Giuliana Sanchez Samaniego 1,2,3
6 Stella Hartinger Peña 1,2,3
7 Paula Skye Tallman 4, 5
8 Daniel Mäusezahl 1,2

9
10 Affiliations:

11 1 Swiss Tropical and Public Health Institute, Basel, Switzerland
12 2 University of Basel, Basel, Switzerland
13 3 Universidad Peruana Cayetano Heredia, Lima Peru.
14 4 The Field Museum of Natural History, Chicago, IL, USA
15 5 Loyola University in Chicago, Chicago, IL, USA

16
17 Abstract

18
19 Objectives: Public health interventions can be improved by understanding peoples' explanatory models of
20 disease. We explore awareness and perceptions of cardiovascular diseases (CVD) and options for
21 preventive actions in young rural adults highlanders.

22 Methods: 46 men and women from communities in Cajamarca, Peru were purposively selected to
23 participate in eight focus groups, where participants discussed their understanding and perceived causes
24 of CVD as well as barriers and pathways to healthy lifestyles.

25 Results: Fresh foods, physical activity, unpleasant emotions, and healthcare access were cited as
26 important determinants of healthy lifestyles. Barriers to healthy diets included lacking nutritional
27 knowledge, fluctuating food prices, and limited access to foodstuffs. Women felt particularly vulnerable to
28 CVD and identified gendered barriers to manage stress and engage in leisure sports. Low health literacy,
29 distance and poor doctor-patient relationships prevented participants from fully accessing healthcare.

30
31 Conclusions: CVD prevention should consider local knowledge of these diseases and of healthy lifestyles
32 and harness ongoing programmes that have successfully promoted good nutrition in children and
33 pregnant women. In concert with public-private partnerships, governments should include health
34 prevention for the entire family.

35
36 Keywords:

37 Cardiovascular disease, Andes, nutrition, prevention, health system

38

39 **Introduction**

40 Cardiovascular diseases (CVD) accounted for approximately 18.6 million deaths in 2019 and are
41 considered the leading cause of death globally [1]. CVDs are a special challenge for low and middle-
42 income countries, where they coexist with a high prevalence of infectious diseases and maternal health
43 conditions [2], and burden health systems that are plagued by inequitable access to care [3].

44 Peru is a case in point. The country faces the double challenge of malnutrition and infectious
45 diseases and increases in CVD [4]. The number of studies of CVD in Peru is growing [5-8] and prior
46 research shows that highland populations have a lower CVD risk in comparison to low-altitude sites [9].
47 However, current research predominantly utilises an epidemiological perspective to understand who is at
48 greatest risk and often misses individual perceptions of CVD and locally relevant factors that limit the
49 adoption of healthy behaviours.

50 To address this gap, this study documents local understandings of CVD and healthy lifestyles
51 among Andean adults in Cajamarca, Peru. We explore participants' personal awareness of CVD and the
52 barriers to adopting healthy behaviours using the Health Belief Model (HBM). The HMB framework
53 focuses on understanding personal attitudes, beliefs and practices, which is crucial for the development
54 of intervention approaches that are culturally acceptable, people-driven and converge people closer to the
55 health system [10-12]. We see this as a first step to determining locally relevant behavioural programmes
56 that can help slow the enormous wave of chronic disease risk that threatens the health of Andean adults.

57

58 **Methods**

59 *Study Setting*

60 This study was carried out in the provinces of San Marcos and Cajabamba in the Cajamarca
61 region, in the northern highlands of Peru, 1800-3900 MASL [13]. These provinces include semi-urban and
62 rural communities. The majority of the population lives in houses made of clay-sand walls and earthen
63 floors. Farming and livestock are the principal source of income [13]. Several governmental programmes
64 are active in the area, including the nutritional interventions “*Vaso de leche*” (for pregnant women and
65 children under 6-years-old) and “*Qaliwarmá*” (for school age children), and two conditional cash transfer
66 programmes, “*Juntos*” (for pregnant women and families with children under 15 years old), and “*Pension*
67 *65*” (for adults over 65 years old that live in extreme poverty) [14]. Different ministries, including the
68 Ministry of Development and Social Inclusion, the Ministry of Health and the Ministry of Economy and
69 Finance, implement these programmes [14].

70

71 *Study Design and Participants*

72 We conducted focus group discussions with adults in the Province of San Marcos and
73 Cajabamba. Participants were parents of children enrolled in a previous study in the area [13].
74 Additionally, at least one parent had undergone a metabolic syndrome examination [15].

75

76 *Recruitment*

77 We used purposive sampling to select participants. Subjects from eight out of 68 local
78 communities were invited to participate. The field supervisor and a fieldworker visited the participants'
79 household and invited them to take part in the focus groups 3-4 days before the scheduled session.
80 Between four to 10 male and female individuals participated in eight mixed focus groups [16].

81

82 *Focus Group Discussions (FG)*

83 Focus groups were held at a central location such as a school or community centre, or at a
84 participant's house. During the focus group, the first author acted as the moderator, and two assistants
85 worked as observers. The moderator facilitated the discussion and encouraged all participants to discuss
86 the questions included in the FG discussion guide. The FG discussion guide was based on the six
87 constructs of the Health Belief Model [10, 11]; perceived susceptibility, perceived severity, perceived
88 benefits, perceived barriers, cues to action, and self-efficacy. We complemented this framework by
89 adding three additional sections on patterns of distress, perceived causes, and gendered differences. The
90 latter was added due to the higher prevalence of metabolic syndrome among women than men [15].

91

92 *Data Analysis*

93 We recorded all FG discussions and two assistants took notes on verbal communication.
94 Following each session, project staff transcribed the audio recordings, identified the participant speaking,
95 and re-listened to the tapes to ensure accuracy. An inductive approach was used for data analysis. Data
96 was organised using the Dedoose qualitative analysis software [17].

97 The codebook consisted of topics identified in the FG guideline and emergent themes from the
98 data. The moderator and the two observers coded the data. If differences in coding appeared, consensus
99 was reached after a team discussion. Similar responses between participants of the same or different
100 focus group were identified from coded extracts. Extracts were organised into tables, with identifiers for
101 the focus group, the participant number, and the theme code. Additional demographic and health
102 information for the participants was available from the larger study this research was embedded in [13,
103 18].

104

105 *Ethics*

106 The Universidad Peruana Cayetano Heredia (UPCH) Ethical Review Board approved the study.
107 Informed consent was obtained from all individuals included in the study. They all consented to the
108 recording of the session.

109

110 **Results**

111 *Characteristics of participants*

112 A total of 46 participants (37 women and 9 men) partook in the eight FGs. Each FG lasted
113 between 30 minutes and one hour. Demographic and health information are shown in Table 1.

114

115 Insert Table 1. Here

116

117 *Main Findings*

118 *Understandings of CVD*

119 In all FGs at least one participant knew a person who suffered from a CVD, but there was
120 dissatisfaction with the knowledge that was available and accessible about CVD. One man in his mid-30s
121 stated, “Yes, we have heard about these diseases, we even know heart transplants exist. All of that. But
122 we don’t have information to prevent these diseases, nor what they do, nor what causes them”.

123 Following up on why there is a lack of information, a woman in her mid-20s expanded that she
124 had not received information on CVDs from the health services, “Because the health staff in this health
125 post only cares about the main reason of your visit. For example, child growth control. They just examine
126 that and it’s done”. Another man in his mid-30s expressed a similar sentiment, “About nutrition, about
127 other diseases, for example, tuberculosis, HIV, cancer. For all those... yes, there is prevention... different
128 pamphlets, but very few about the heart”. High levels of illiteracy are an additional factor complicating
129 understandings of CVD. As a woman in her mid-30s stated, “The one who knows how to read, can read.
130 But the one who doesn’t know, only observes”.

131 While the participants expressed dissatisfaction with the level of information available for CVD
132 and cited that they have not had in-depth discussions of CVD previously, they still expressed knowledge
133 about the causes and symptoms of CVD. Specifically, they mentioned that symptoms of CVD included
134 agitation, pain, weakness and strong heart beating; and that the ultimate consequences of CVD were
135 heart attacks and death. In all focus groups the participants stated that anyone could develop a heart
136 disease, no matter their weight or age. Additionally, participants identified a number of drivers of CVD,
137 which fell into four general categories: nutrition, physical activity, emotions, and access to healthcare.

138

139 *Nutrition*

140 Study participants expressed that CVD is associated with being overweight and “having fat
141 around the heart”. Interestingly, a woman in her early 30s stated, “When we have fat in the heart too, and
142 is not necessary that we are fat” and other community members clearly articulated that being overweight
143 or underweight could be a determinant of CVD. Another woman in her mid-30s expanded, “If you are
144 overweight or you are underweight, it is equally bad.” and that “Nobody is free ... no one is free neither

145 the fat nor the skinny. I believe that not only the fat person gets sick, furthermore the skinny one may be
146 sicker”.

147 Participants not only acknowledged that both under and overnutrition were problematic, but also
148 made direct connections between nutritional health and food. A woman in her late-20s confided,
149 “Sometimes, we only eat to fill our belly, without realizing that it is not truly nourishing us”. Further,
150 participants explained how structural factors limited their ability to obtain good nutrition. A woman in her
151 early-30s stated that, “Sometimes we eat things that are bad for us and we know that they are going to be
152 bad for us. But we eat out of necessity because there is no other thing to eat”.

153 The identification of structural barriers to adopting a nutritious diet were common and included not
154 being able to access healthy foods, not having the economic resources to purchase them, and lacking
155 knowledge about how to prepare healthy foods. These barriers varied across geography and occupation.
156 More than one third of the participants were farmers, but reported that they did not regularly consume
157 products from their crops. A man in his mid-30s told us that, “50% [of all goods] you have to buy and
158 another 50% is produced in the field”.

159 Participants living in the low altitude valley (1800-2200 MASL) reported that they can grow all
160 types of food. However, community members in higher altitudes are limited in their crop variety and also
161 had issues with access to purchasing fresh food products such as vegetables, fruits and animal protein.
162 Barriers to obtaining these products included the cost to reach a market (which could be between 30min–
163 2 hours away), time spent getting there, and issues with food storage. As a woman in her late-20s pointed
164 out, “Food cannot be stored for a long time, maybe 2 or 3 days because we do not have a refrigerator, a
165 place where to keep the food”.

166 On the other hand, local *bodegas* provide constant access to non-perishable foods and drinks,
167 which are cheaper. As a man in his mid 30s quipped, “Mostly you decide what food to buy according to
168 the economy, right?” But even buying foods at the *bodegas* was complicated because food prices
169 changed every week because of the many intermediaries involved. As a woman in her mid-30s explained,
170 “You know... the intermediaries always have to take advantage, right? They live off that, so sometimes
171 they sell a product at a certain price and the next week they change it. Then, in the local store, items are
172 more expensive.” Local challenges to accessing, purchasing, and storing foods lead people to
173 preferentially buy non-perishable foods such as rice, noodles, oil, and sugar, especially those participants
174 living in distant communities.

175 176 *Physical Activity*

177 Participants indicated that physical activity was beneficial for their wellbeing and recognised it as
178 a preventive measure for developing CVD. They mentioned being highly active because of their daily
179 agricultural and farming work, which they enjoyed and kept them busy. For example, a man in his late-
180 20s stated that, “We get health benefits, and economic benefits.... to be useful (for field work)”. However,
181 participants explained that recreational physical activities such as sports were often limited due to lack of

182 time, religious and local beliefs, and gendered factors. A man in his mid-30s shed light on this saying,
183 "The word of God [local church] has forbidden us to play football. We always bet sometimes, right?
184 Sometimes we have anger at the partner because he is beating us. Sometimes we do not play well."

185 Women appear to face dual pressures. On the one hand, participation in sport appears to be
186 expected of mothers. A mother in her late-20s expanded saying, "Sometimes in the schools, in
187 kindergartens, the moms have to participate. There is no way you do not participate. If the babies see that
188 their mother does not want to, then they will not want too. So, you have to participate, no matter what."
189 Yet there is also gendered pressure to refrain from sports as a woman in her mid-30s stated that
190 sometimes women are expected to stay at home and take care of their family and not spend leisure time
191 in sports. Gendered dynamics in CVD risk factors were also evident in group discussions regarding the
192 perceived role of emotional stress in CVD.

193

194 *Negative Emotions*

195 Participants articulated that negative emotions could contribute to heart problems and also that
196 knowing they have health problems can produce negative emotions. Speaking about a family member, a
197 woman in her late-20s stated that, "My aunt... sometimes she suffers attacks. She has been told to avoid
198 being melancholic, to avoid worries. Nothing, nothing. No worries, because if she is worried or
199 melancholy she can die." Indeed, participants connected both physical and emotional stress to strain on
200 the heart. A young woman explained that, "Sometimes when you lose your patience. You feel that your
201 heart beats faster at times like this. Sometimes you have anger, let's say, it's a moment ...", while a
202 woman in her late 30s expressed a similar sentiment in relation to physical activity, saying, "When you
203 walk and feel agitated, then you feel pain and sometimes "stitches". Well then, you think it is the heart,
204 right?".

205 These quotes indicate that negative emotions or over-exertion create heart problems, but
206 participants also indicated that simply knowing about having heart problems can produce substantial
207 household distress. Indeed, a number of participants expressed that they would feel sad and worried if
208 they were diagnosed with any CVD as the costs of treatment, or losses due to death, could drive
209 economic insecurity for the entire household. Women worried about themselves and their partners
210 saying, "If my husband gets sick, what should I do? Who works? Who maintains us?" and "If my husband
211 gets sick, I have to spend money, I do worry."

212 Discussions in the FGs revealed why women appeared to be more prone to this type of worrying.
213 A woman in her late-20s explained, "It seems to me, I say it is because men go to the farm, they occupy
214 themselves there, they forget about everything. On the other hand, the women stay at home and
215 sometimes we have worries about the children... everything, that can also be it [a risk factor of CVD] or
216 maybe not". A man in his late-30s expressed a similar explanation, musing that "Maybe... [Women are at
217 higher risk] because sometimes we, men, go to work, it is a single job and the ladies stay at home, they

218 work different tasks...they worry sometimes and think "There are things that I don't have.... I do not have
219 money".

220 Finally, participants expressed that they would rather not know about a potential cardiovascular
221 condition because of the stress it might cause, saying, "It is better to not know (that you have a CVD).
222 Just stay quite during the pain, then it goes away and you forget.... if you know what you have, it is
223 worse." However, even for those who are interested in knowing about their CVD risk, there are substantial
224 barriers to accessing knowledge, affordable treatment, and care at the local health posts.

225

226 *Healthcare*

227 Similarly, to the challenges reported in accessing healthy fresh foods, participants cited costs
228 associated with transport and food as a barrier to healthcare. As a result, participants stated that they
229 typically only went to the clinic for a pressing health issue and not for regular check-ups. A woman in her
230 late-20s stated, "If it does not hurt, I don't go to the doctor. If it hurts, I will go to check what is going on.
231 On the contrary, we should have general check-ups, for example annually." Appointments at the health
232 post also needed to be scheduled in person, potentially creating two lengthy and costly trips, and creating
233 a high initial barrier to seek care.

234

235 In addition to challenges in accessing care, participants criticised the quality of patient-provider
236 relationships. They felt that they were not treated well nor receiving proper attention because health
237 professional were inexperienced and prescribed painkillers for every condition. According to a woman in
238 her mid-20s, previously, "There was a good nurse, but they relocated her to the hospital. And here, we
239 only have trainees. This is what happens here at the health post". and when she does visit the health
240 post, "Oh, this hurt, oh, paracetamol, ibuprofen. All the time it is the same medicines we receive. When
241 we go with our children, we already know what they will prescribe to us."

242 More problematically, participants were receiving conflicting medical advice that was not
243 evidence-based. A woman in her mid-30s revealed that

244

245 "I went to the health post with my results [lipids and glucose measurements]. They [health
246 professional] told me: "No, there is no medicine, you can only take warm water before eating". I
247 told them I couldn't eat in the morning. They told me that anyway I should eat a little bit.... but I
248 should eat. They told me to start eating less but not to stop eating. They told me better to buy
249 wine and drink a glass before going to sleep to get rid of the fat. But, I have not seen any results."

250

251 This type of medical advice appears to be intersecting with local explanatory models for the
252 treatment of CVD. One woman in her early 30s stated, "For example, if the heart is not well and they
253 (health professional) tell you it is because of the fat.. then you try to eat things that do not affect your
254 heart. Right? I know that you have to drink pineapple juice and because you know this is good for the

255 heart then you drink it..., this is a medicine.... right? A similar sentiment was expressed by a woman in
256 her mid-20s, “Yes, sometimes one feels bad. You can try to look for flowers, and drink flower waters, you
257 can use also lemon, orange, custard apple”

258 It is unclear whether these remedies were coming from healthcare practitioners, traditional
259 healers, or fellow community members. What is apparent is that the public health messaging for
260 preventing and treating CVD is not completely clear. Despite this, participants related CVD to nutrition
261 and exercise and many participants showed an interest in adopting healthier diets and increasing physical
262 exercise. However, they also identified a number of substantial challenges to enacting these behaviours
263 in the short and long-term, which are key issues to address in launching successful behavioural
264 interventions to prevent CVD in Andean communities in Peru.

265

266 **Discussion**

267 In this paper, we explored how Andean adults in Peru understood CVD, including the drivers of
268 CVD risk, and the barriers to adopting healthy lifestyles that can reduce risk. While the understandings of
269 CVD were basic, study participants stated that nutrition, physical activity, negative emotions, and
270 healthcare were important drivers of CVD risk. Here we expand on these findings with particular attention
271 to how they can inform the National Guideline of Prevention and Control of Non-Communicable Diseases
272 and their strategies for vulnerable high-altitude populations [19].

273 Many of the participants in this study knew that CVD was potentially life-threatening and
274 expressed a desire to be better informed. However, participants made it clear that health communication
275 strategies via text or print-outs would not reach illiterate community members. Instead, they suggested
276 that nutritional workshops with demonstrations of healthy cooking and nutritional discussion groups could
277 effectively engage community members.

278 Such nutritional workshops, if designed holistically, could address many of the barriers to the
279 adoption of healthy eating behaviours [20, 21]. Participants understood that healthy diets mean eating
280 various types of food in proper amounts but felt their ability to adopt these diets was limited because they
281 did not know how to prepare healthy meals and could not afford and access readily diverse food items.
282 While there are existing programmes providing food to children and pregnant women in these areas,
283 including *Vaso de Leche* and *Qaliwarma*, adults without children are notably excluded. As main
284 programme targets, women have to comply with regular maternal and children health checkups and
285 ensure children's school attendance [14], making them central stakeholders for such programmes and a
286 regular point of contact with the health system. Women are the key health information channels into the
287 family. As primary knowledge-bearers and with their expressed desire for nutritional skill-building they are
288 the ideal partners for people-centred engagement in CVD prevention in remote Andean communities.

289 We suggest that additional studies are conducted in close collaboration with Andean women to
290 design nutritional interventions that can improve health and wellbeing at all life stages and are appropriate
291 for this unique context. There are existing governmental programmes such as Haku Wiñay (“My

292 enterprising farm” in Quechua language), which support the raising and trading of domestic animals and
293 entrepreneurship in small-scale agriculture[14]. Is it possible to expand these programmes to support
294 women in small-scale subsistence activities that can provide healthy foods that can dually reduce CVD
295 risk in adults and undernutrition in children? Identifying nutrient-rich and culturally-valued crops, suitable
296 for both, home consumption and trading can enhance food security and strengthen income generating
297 capacities. Additionally, opportunities exist to build on public-private partnerships for CVD prevention. For
298 example, the national dairy company collects milk from the remotest areas of the Andes at least weekly.
299 Considering that the company maintains this regular supply chain with remote outreach, building public-
300 private partnership could help providing healthy foods to highlanders who otherwise travel only monthly or
301 bi-weekly to local markets

302 However, to be feasible, such initiatives need to be linked to system-wide changes and
303 interventions. Peru has successfully executed multi-level actions to address childhood undernutrition and
304 anemia using a combination of programmes in the areas of health, education, cash transfer, water and
305 sanitation, housing, and agriculture [22, 23]. CVD prevention action would need the same approach and
306 could piggy back off of ongoing programmes with already established contacts within the health system.
307 Specifically, CVD risk needs to be assessed at the peripheral primary care level measuring by
308 anthropometrics and blood pressure, and short risk questionnaires [24, 25]. Health data should then be
309 saved in a national repository of chronic conditions similar to the “Wawared” system for pregnant women
310 and children[26]. With surveillance systems in place, monitoring, evaluation and access to treatment can
311 become feasible in distant populations. Task-shifting [27] is another option that has contributed to
312 maternal and child health programmes in Peru in the past and could be utilised for CVD interventions.
313 Task-shifting entails working with community health workers (CHW) to bring preventative strategies to the
314 periphery and is potentially helpful in settings with health workforce shortages [28]. Task-shifting requires
315 the training of CHW in new skills, the use of standardised protocols, and the provision of adequate
316 equipment for CVD prevention and communication. In this way, health professionals can expand from a
317 child-centred health approach to include adult health. The above recommendations are inherently people-
318 centred and community-based, however they are the first approach into creating more comprehensive
319 national guidelines for CVD prevention [19].

320 In 2015, non-communicable diseases were included in a global development agenda [8]. This
321 target requires multi-level action beyond the health sector, as showed by the success of child nutrition
322 interventions in Peru [22, 23]. Thus, system-wide health interventions are possible in the country and can
323 dually address CVD risk in adults and continue supporting the nutritional health of children. Such
324 approaches designed in partnership with local women, could also potentially address emotional distress
325 as a notable locally perceived CVD risk factor, by promoting health and wellbeing and improving food
326 security and access to care.

327

328 The study was conducted in young adults in their 30's and perceptions of CVD may differ from
329 older individuals. Villagers participating in the FGs knew each other, potentially influencing their
330 participation. However, they collectively reflected a community view, which was particularly valuable when
331 discussing locally feasible and acceptable intervention approaches.

332 Most of the FG participants were women but as household managers and family-health
333 knowledge-bearers with the most regular contact with health facilities we believe they and their
334 participating husbands closely represent the broader community perspective on NCD health matters.

335

336 This research revealed that there are many feasible entry points of intervention that can prevent
337 Andean community members from succumbing to the wave of CVD risk that is currently engulfing the
338 globe. Specifically, local women are key stakeholders who can influence their own health and the health
339 of all family members. Therefore, they must be empowered with the knowledge and resources to enact
340 healthy lifestyles. We see substantial opportunities for this to occur by expanding existing government
341 programmes to make healthy foods and healthcare accessible, promoting healthy lifestyles and by
342 building on public-private partnerships that can support access healthy food for all community members.

343

344

345 Acknowledgements

346 We thank the study participants for their valuable time and kind participation. We appreciate and thank
347 the local authorities of the RED IV de Salud San Marcos, for their continuous support. We also express
348 our gratitude to Mrs Angelica Fernandez for helping and organising community meetings and Karen Meza
349 and Forlly Chavez for supporting the coding of the transcripts and organization of extracts

350

351 References

352

- 353 1. Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM et al. Global burden
354 of Cardiovascular Diseases and risk factors, 1990-2019: Update from the GBD 2019 study. *J Am Coll*
355 *Cardiol* (2020) 76:2982-3021. doi:10.1016/j.jacc.2020.11.010
- 356 2. Roth GA, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N et al. Global, regional, and national
357 age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a
358 systematic analysis for the Global Burden of Disease Study 2017. *Lancet* (2018) 392:1736-88.
359 doi:10.1016/s0140-6736(18)32203-7
- 360 3. Agyemang C, van den Born B-J. Limited access to CVD medicines in low-income and middle-
361 income countries: poverty is at the heart of the matter. *Lancet GlobHealth* (2018) 6:e234-e35.
362 doi:10.1016/s2214-109x(18)30048-2

- 363 4. Huicho L, Trelles M, Gonzales F, Mendoza W, Miranda J. Mortality profiles in a country facing
364 epidemiological transition: an analysis of registered data. *BMC Public Health* (2009) 9:47.
365 doi:10.1186/1471-2458-9-47
- 366 5. Benziger CP, Bernabé-Ortiz A, Gilman RH, Checkley W, Smeeth L, Málaga G et al. Metabolic
367 abnormalities are common among South American hispanics subjects with normal weight or excess body
368 weight: The CRONICAS cohort study. *Plos One* (2015) 10. doi:10.1371/journal.pone.0138968
- 369 6. Miranda JJ, Bernabé-Ortiz A, Diez-Canseco F, Málaga G, Cárdenas MK, Carrillo-Larco RM et al.
370 Towards sustainable partnerships in global health: the case of the CRONICAS Centre of Excellence in
371 Chronic Diseases in Peru. *Global Health* (2016) 12. doi:10.1186/s12992-016-0170-z
- 372 7. Bernabe-Ortiz A, Carrillo-Larco RM, Gilman RH, Checkley W, Smeeth L, Miranda JJ et al.
373 Contribution of modifiable risk factors for hypertension and type-2 diabetes in Peruvian resource-limited
374 settings. *J Epidemiol Community Health* (2016) 70:49-55. doi:10.1136/jech-2015-205988
- 375 8. Carrillo-Larco RM, Bennett JE, Di Cesare M, Gregg EW, Bernabe-Ortiz A. The contribution of
376 specific non-communicable diseases to the achievement of the Sustainable Development Goal 3.4 in
377 Peru. *PLoS One* (2020) 15:e0240494. doi:10.1371/journal.pone.0240494
- 378 9. Bernabé-Ortiz A, Carrillo-Larco RM, Gilman RH, Checkley W, Smeeth L, Miranda JJ. Impact of
379 urbanisation and altitude on the incidence of, and risk factors for, hypertension. *Heart* (2017) 103:827-33.
380 doi:10.1136/heartjnl-2016-310347
- 381 10. Krummel DA, Humphries D, Tessaro I. Focus groups on cardiovascular health in rural women:
382 implications for practice. *J Nutr Educ Behav* (2002) 34:38-46. doi:10.1016/s1499-4046(06)60223-6
- 383 11. K Glanz BKR, K. Viswanath. *Health Behaviour and Health education*. 4th ed. San Francisco, CA:
384 Jossey-Bass (2008).
- 385 12. Collins TE, Nugent R, Webb D, Placella E, Evans T, Akinnawo A. Time to align: development
386 cooperation for the prevention and control of non-communicable diseases. *BMJ* (2019) 366:l4499.
387 doi:10.1136/bmj.l4499
- 388 13. Hartinger SM, Nuno N, Hattendorf J, Verastegui H, Karlen W, Ortiz M et al. A factorial cluster-
389 randomised controlled trial combining home-environmental and early child development interventions to
390 improve child health and development: rationale, trial design and baseline findings. *BMC Med Res*
391 *Methodol* (2020) 20:73. doi:10.1186/s12874-020-00950-y
- 392 14. Pillaca-Medina S, Chavez-Dulanto PN. How effective and efficient are social programs on food
393 and nutritional security? *Food and Energy Security* (2017) 6:e120. doi:10.1002/fes3.120
- 394 15. Sanchez-Samaniego G, Mäusezahl D, Carcamo C, Probst-Hensch N, Verastegui H, Hartinger
395 SM. Improved cookstoves and the prevalence of metabolic syndrome in the rural Peruvian Andes: a
396 quasiexperimental study. *Research Square[Preprint]* (2020) . doi:10.21203/rs.2.14701/v1(Accessed
397 October 21, 2020)
- 398 16. Rabiee F. Focus-group interview and data analysis. *Proc Nutr Soc* (2007) 63:655-60.
399 doi:10.1079/pns2004399

- 400 17. Dedoose Version 8.0.35, web application for managing, analyzing, and presenting qualitative and
401 mixed method research data. (2018) Los Angeles, CA. Available from: www.dedoose.com.
- 402 18. Sanchez-Samaniego G, Maüsezahl D, Carcamo C, Verastegui H, Probst-Hensch N, Hartinger S.
403 Improved cookstoves and the prevalence of metabolic syndrome in the rural Peruvian Andes - a quasi-
404 experimental study. (2019).
- 405 19. Ministerio de Salud Lineamientos de política y estrategias para la prevención y control de
406 Enfermedades No Trasnmisibles (ENT) 2016-2020. (2016).
407 [https://cdn.www.gob.pe/uploads/document/file/193750/192473_RM_229-2016-MINSA.pdf20180904-](https://cdn.www.gob.pe/uploads/document/file/193750/192473_RM_229-2016-MINSA.pdf20180904-20266-1ap2x89.pdf)
408 [20266-1ap2x89.pdf](https://cdn.www.gob.pe/uploads/document/file/193750/192473_RM_229-2016-MINSA.pdf20180904-20266-1ap2x89.pdf) [Accessed January 15, 2021]
- 409 20. Sriram U, Sandreuter K, Graham M, Folta S, Pullyblank K, Paul L et al. Process evaluation of
410 Strong Hearts, healthy communities: A rural community-based cardiovascular disease prevention
411 program. *J Nutr Educ Behav* (2019) 51:138-49. doi:10.1016/j.jneb.2018.10.011
- 412 21. Seguin RA, Eldridge G, Graham ML, Folta SC, Nelson ME, Strogatz D. Strong Hearts, healthy
413 communities: a rural community-based cardiovascular disease prevention program. *BMC Public Health*
414 (2016) 16:86. doi:10.1186/s12889-016-2751-4
- 415 22. Huicho L, Segura ER, Huayanay-Espinoza CA, de Guzman JN, Restrepo-Mendez MC, Tam Y et
416 al. Child health and nutrition in Peru within an antipoverty political agenda: a Countdown to 2015 country
417 case study. *Lancet Glob Health* (2016) 4:E414-E26. doi:Doi 10.1016/S2214-109x(16)00085-1
- 418 23. United States Agency for International Development (USAID). Peru: National Anemia Profile.
419 (2016). [https://www.spring-nutrition.org/sites/default/files/publications/anemia-](https://www.spring-nutrition.org/sites/default/files/publications/anemia-profiles/spring_nap_peru.pdf)
420 [profiles/spring_nap_peru.pdf](https://www.spring-nutrition.org/sites/default/files/publications/anemia-profiles/spring_nap_peru.pdf) [Accessed January 15, 2021]
- 421 24. Bernabe-Ortiz A, Perel P, Miranda JJ, Smeeth L. Diagnostic accuracy of the Finnish Diabetes
422 Risk Score (FINDRISC) for undiagnosed T2DM in Peruvian population. *Prim Care Diabetes* (2018)
423 12:517-25. doi:10.1016/j.pcd.2018.07.015
- 424 25. World Health Organization (WHO). STEPS Surveillance Manual: The WHO STEPwise approach
425 to chronic disease risk factor surveillance. Geneva; 2005.
- 426 26. Pérez-Lu JE, Romero RI, Bayer AM, García PJ. Wawared Peru: reducing health inequities and
427 improving maternal health by improving information systems in health. *Rev Peru Med Exp Salud Publica*
428 (2015) 32.
- 429 27. Westgard C, Naraine R, Paucar Villacorta DM. Performance evaluation of community health
430 workers: Case study in the Amazon of Peru. *J Community Health* (2018) 43:908-19. doi:10.1007/s10900-
431 018-0503-3
- 432 28. Joshi R, Alim M, Kengne AP, Jan S, Maulik PK, Peiris D et al. Task shifting for non-
433 communicable disease management in low and middle income countries--a systematic review. *PLoS One*
434 (2014) 9:e103754. doi:10.1371/journal.pone.0103754

435
436 Captions

437 Tables

438 Table 1:

439 Demographic and health characteristics of young adult participants of the Provinces of San
440 Marcos and Cajabamba, Peru 2017.

441 Legend:

442 SD: standard deviation

443 MASL: metres above sea level

444

	Women		Men	
	N	Mean \pm SD or N (%)	N	Mean \pm SD or N (%)
Age (years)	37	29.1 \pm 6.0	9	29.3 \pm 4.0
Complete secondary	34	9 (27%)	6	4 (67%)
Work in the agriculture sector	34	23 (68%)	6	6 (100%)
Housewife	34	6 (18%)		
Living over 2500 MASL	37	15 (41%)	9	6 (67%)
Body mass index	34		6	
Underweight		0		0
Normal weight		15		3
Overweight		13		3
Obese		6		0
Metabolic syndrome	34	10 (29%)	6	3 (50%)

445

446