Title: Prevalence of Fetal Alcohol Spectrum Disorder (FASD) in Greater Manchester, UK: an

active case ascertainment study

Robyn McCarthy MSc[1], Raja A.S. Mukherjee PhD[1,2], Kate M. Fleming PhD[3], Jonathan Green
FRCPsych[4], Jill Clayton-Smith MD[5], Alan D. Price PhD[1], Clare S. Allely PhD[1], Penny A. Cook
PhD*[1]
*Correspondence to: Penny A. Cook

School of Health and Society,

University of Salford,

Allerton Building

Frederick Road

Salford M6 6PU

p.a.cook@salford.ac.uk

Tel: +44(0)161 295 2804

1. School of Health and Society, University of Salford , UK

2.   Fetal Alcohol Spectrum Disorder Service, Surrey & Borders NHS Trust, UK School of

Health and Society, University of Salford, UK

3. Institute of Population Health Sciences, University of Liverpool, UK Division of

4.Neuroscience & Experimental Psychology, University of Manchester, UK Royal Manchester

Children's Hospital, Manchester, UK

5.Royal Manchester Children's Hospital, Manchester, UK Division of Evolution & Genomic

Sciences, University of Manchester, UK

Funding

This project was funded by The Greater Manchester Health and Social Care Partnership. The funders had no role in study design, data collection and analysis, decision to publish, or the preparation of the manuscript.

Conflicts of interest

RASM has received honoraria from speaking for various pharmaceutical companies related to ADHD including Takeda, Flynn and Jannsen, as well as support for entry to an international conference related to ADHD. RASM is also a non-paid advisor to various ASD, ADHD and FASD charities in England and Internationally (Australia and Europe). RASM has organised conferences and obtained non educational grants from various sources including Pharma and private hospitals. AP is an non-paid member of the Professional Advisory Panel of the not-for-profit organisation, ÉNDpae. All other authors declare that they have no competing interests.

Acknowledgments

We thank the parents, guardians, children, head teachers and staff at participant schools. We also thank Róisín Reynolds and Rachael Nielsen from the Greater Manchester Health and Social Care Partnership for their support. Philip May allowed us to use the Maternal Risk Questionnaire, the University of New Mexico by the Fetal Alcohol Syndrome Epidemiology Research Team initially developed in collaboration with Sharon Wilsnack (University of North Dakota), and adapted and updated multiple times for use in prevalence studies in the United States, South Africa, and Italy. We thank those who represented families affected by

46   FASD, led by Anna Webster. We thank the members of an independent steering group,

47   including Anna Webster Jen Michaels, Joanne Higham, Mary Scanlon and Susan McGrail.

Abstract

Background: Despite high levels of prenatal alcohol exposure in the UK, evidence on the prevalence of fetal alcohol spectrum disorders (FASD) is lacking. This paper reports on FASD prevalence in a small sample of children in primary school.

Methods: A two-phase active case ascertainment study was conducted in three mainstream primary schools in Greater Manchester, UK. Schools were located in areas that ranged from relatively deprived to relatively affluent. Initial screening of children aged 8-9 years used pre-specified criteria for elevated FASD risk (small for age; special educational needs; currently/previously in care; significant social/emotional/mental health symptoms). Screen positive children were invited for detailed ascertainment of FASD using gold standard measures including medical history, facial dysmorphology, neurological impairment, executive function and behavioural difficulties.

Results: Of 220 eligible children, 50 (23%) screened positive and 12% (26/220) proceeded to phase-two assessment. Twenty had a developmental disorder, of which, four had FASD and four were assessed as possible FASD. The crude prevalence rate of FASD in these schools was 1.8% (95%CI: 1.0%,3.4%) and when including possible cases was 3.6% (2.1%,6.3%). None of these children had previously identified with a developmental diagnosis.

Conclusions: FASD was found to be common in these schools, but limitations to the sampling restrict inferences to a population prevalence. Most of these children's needs had not previously been identified.

70   Keywords (5)

71   FASD; Fetal Alcohol Syndrome (FAS); neurodevelopmental disorder; attention deficit

72   hyperactivity disorder (ADHD); Autism Spectrum Disorder (ASD)

Introduction

An estimated 10% of pregnancies globally are exposed to alcohol, a potent teratogen that can lead to physical and neurodevelopmental birth defects (Popova et al., 2017), collectively known as Fetal Alcohol Spectrum Disorders (FASD) (Cook et al., 2016). FASD is an umbrella term that includes the diagnoses of Fetal Alcohol Syndrome (FAS), pFAS (partial Fetal Alcohol Syndrome), ARBD (Alcohol-Related Birth Defects), and ARND (Alcohol-Related Neurodevelopmental Disorder)/ND-PAE (Neurobehavioral Disorder Associated with Prenatal Alcohol Exposure).

The prevalence of FASD is estimated to be 0.8% globally and highest in Europe, at 2% (Lange et al., 2017a). There are no direct estimates of prevalence in the four countries with the highest known rates of prenatal alcohol exposure (Ireland, Belarus, Denmark and UK), all of which have rates of over 40% pregnancies exposed to alcohol (Popova et al., 2017). For the UK, the modelled estimate suggests 3.2% of children and young people may have FASD (Lange et al., 2017a). A national study in the US, on populations relatively similar to those in the UK, found a weighted estimate of 3-10% for FASD in children in primary school (May et al., 2018). Where a UK cohort with high levels of exposure (79% of mothers drank during the pregnancy, with 25% at binge levels) was used, 6–17% of children screened positive for features of FASD (McQuire et al., 2019). However, the lack of direct evidence of prevalence contributes to under-investment in diagnostic, treatment and prevention services (Scholin et al., 2021).

Active case ascertainment studies, the basis of global and national prevalence estimates (Lange et al., 2017a; May et al., 2018), are considered the 'gold standard' method of estimating prevalence, and involve screening a cross section of the general population of

children (Roozen et al., 2016). Passive methods are less useful because individuals with

FASD are often not diagnosed (May & Gossage, 2001; Morleo et al., 2011) for a number of

reasons, including a lack of knowledge/training among healthcare/educational professionals

(Mukherjee et al., 2015); difficulty in differentiating features of FASD from other commonly

co-occurring disorders (e.g. Attention-Deficit/Hyperactivity Disorder, ADHD, or Autism

Spectrum Disorder, ASD) (Chasnoff, Wells & King, 2015; Mukherjee et al., 2011; Young et al.,

2016).

The aim of this study was to provide direct evidence of the prevalence of FASD in a small

sample of UK children aged 8 to 9 years.

Materials and Methods

Setting

The setting was Greater Manchester, North West England (population 2.8 million), an area

with a higher than England average level of alcohol harm (Public Health England, 2021), and

relatively high deprivation (Ministry of Housing Communities and Local Government, 2019).

This study was part of a wider initiative, the 'Preventing Alcohol Exposed Pregnancy

Programme' taking place in four of the ten Greater Manchester local authority areas. The

initiative also included increased awareness raising and interventions with women who

were pregnant or at risk of unplanned pregnancy (Reynolds et al., 2021).

Design

We report on a cross-sectional study to detect cognitive impairments and associated

conditions, including FASD, using an Active Case Ascertainment method. Based on the World

Health Organization (WHO) standard protocol for FASD prevalence studies (World Health

Organisation, 2012), children in school year three/four (8-9 years of age at enrolment) who

were able to communicate in English were invited to take part in a two-phase approach.

Children with a known risk factor for FASD went to the second phase 'Children with a known

risk factor for FASD went to the second phase (full assessment, see section below, phase 1:

initial screening, Figure 1). Child assessments took place between July 2019 and March

2020. Restrictions related to the COVID-19 pandemic prevented face to face data collection

from mid-March 2020, at which point there was one outstanding parent interview, which

was conducted by telephone in March 2020.

Sample size

129 A sample size of 170 was adequate to get a preliminary indication of prevalence in a

130 selected sample of schools (based on estimated true prevalence=0.03, precision=0.05;

131 sensitivity=0.85; specificity=0.85). We aimed to recruit three schools (assuming

132 approximately 60 pupils in the relevant age category per school).

School recruitment

In this pragmatic, small-scale study, schools were purposefully selected. Key informants (e.g.

local government officials who work with schools, Education Psychologists) were consulted

and suggested four schools, all of which agreed to take part. One school was excluded due

to disengagement from the study and deviation from the protocol (see supplementary file 1

for details of a partial dataset). We initially aimed to include a specialist school providing

Social Emotional and Mental Health (SEMH) support. All such schools within the study area

were contacted. One such school agreed to take part but withdrew after being unable to

gain consent from any parents.

In February 2020, an extension of data collection was granted with the aim to recruit further

schools to increase the baseline sample size. Four additional schools were recruited.

However, the imposition of COVID-19 related restrictions, including lockdown, meant that

the study could not be completed in the second wave of schools. Partial results are reported

in Supplementary file 1.

Phase 1: Initial Screening

Parents of all eligible children were sent a letter describing the study, with the option to opt

out, at least five days before data collection. Parents who wished to remove their child from

the study at this stage were advised to return the letter with the opt-out option selected.

Researchers were present in school to assess physical measurements. Height was measured

using a mobile stadiometer, weight using Marsden MBF-6000 scales and head

circumference (OFC) using a Sec 201 measuring tape. To become eligible for phase 2 (full

assessment), one or more of the following criteria had to be met: height or weight below

the ninth centile or OFC below the second centile; identified by the school or parents as

having difficulties with learning, maladaptive behaviour, inattention or hyperactivity issues;

have an Education Health and Care plan; be a currently or previously looked after child;

already have diagnosed difficulties with behaviour including ADHD or conduct disorder.

Exclusion criteria: disabilities or behavioural abnormalities known to be caused by well-

characterised and already identified genetic factors (e.g. Down Syndrome, Williams

Syndrome) or by post-natal brain injuries. These were excluded because the

physical/behavioural stigmata overlap with FASD in presentation, and an FASD label cannot

be attributed as the primary aetiological factor.

Phase 2a: Measures completed by parents/carers

Parents of those meeting criteria for phase 2 were sent information sheets and consent

forms to opt themselves and their child into the study either by email, letter sent home, by

telephone or in person by the Special Educational Needs Coordinator or head teacher.

Where children were currently under the care of the local authority consent was obtained

from the supervising social worker. For parent-report assessments, most took place at

school in a private room for two-hour sessions. Where requested by the parent,

assessments took place in their home.

Parent report assessments used the validated measures listed in Figure 1. The medical

history was taken using a structured questionnaire developed originally at the University of

New Mexico (May et al., 2018). The schedule included questions on pre-pregnancy,

pregnancy and current substance use (alcohol, tobacco, prescription and illicit drugs), folic

acid use in pregnancy and birth complications. General questions on current alcohol

consumption were asked first as part of wider lifestyle set of questions, followed by

questions about alcohol consumption in pregnancy. If alcohol was consumed during

pregnancy, further questions ascertained the level and timing of consumption. Birth

mothers were interviewed privately in order to encourage open reporting of alcohol use.

For looked after children, fostered or adopted children, the parent/carer with most

knowledge of the child was interviewed.

Phase 2b: Measurements on children

Measurements took place in school during school hours. Dysmorphology of fetal alcohol

syndrome (FAS) facial characteristics was assessed from photographs (using standardised

alignment of the participant's head relative to the camera lens) taken by trained

researchers. Images were analysed by FAS Facial Photographic Analysis Software (Astley,

2015) and validated by an experienced clinical geneticist. Neurological impairment was

assessed using the Developmental NEuroPSYchological Assessment (NEPSY) to assess

memory, attention, and executive functions. The NEPSY subtests (Inhibition, Narrative

Memory and Word List Interference, Animal Sorting, Clocks and Memory for Faces) were

those that had been identified in previous FASD research (Rasmussen et al., 2013). The

Wechsler Intelligence Scale for Children (WISC V) (Kaufman et al., 2015) was used to assess

the child's cognition, as used in previous FASD studies (Raldiris et al., 2018).

Phase 2c: Ruling out genetic causes and case conference

The results of the assessments were compared to the case definition (Table 1), derived from

the WHO protocol for prevalence studies (World Health Organisation, 2012) and

international guidelines (Cook et al., 2016). Where deficits met criteria in three subdomains,

and PAE was present, FASD was considered. Where a child had all three sentinel facial

features associated with FAS, FASD was considered in the absence of reported PAE. For all

those considered for FASD, a Microarray Comparative Genomic Hybridisation (array CGH)

test was used to rule out other disorders that may present with a similar behavioural

presentation of genetic origin (Douzgou et al., 2012). A saliva sample was taken using the

Oragene DNA OG-575 (DNAGenotek, 2020). The Oligo (Oxford Gene Technology, Oxford,

UK) Hx60k array was carried out by the North Western Regional Genetics Laboratory within

Manchester University NHS Foundation Trust, Manchester.

Findings for each child discussed during case conferences attended by the study team

including clinicians with expertise in the diagnosis of FASD. Cases of FASD were defined

according to internationally recognised guidelines (Cook et al., 2016). Possible cases were

where FASD was suspected but information was missing or unclear, for example regarding

alcohol use.

Data analysis

Prevalence was estimated as the total number of children with FASD and probable FASD as

the numerator, and the total number of eligible children at the same site as the

denominator. This generates a conservative or minimum estimate of prevalence as it

assumed that all children who were not examined did not have FASD. To obtain confidence

intervals we accounted for clusters (schools) by using nonparametric bootstrapping,

resampling with replacement clusters with 10000 bootstrap runs.

Ethical considerations

Parents were invited to take part and were offered a report on their child's learning and

behaviour issues, which could include a range of learning difficulties, including possible ASD,

ADHD and FASD. The text of the information sheet made it clear that the main objective was

to ascertain FASD prevalence.

A small number of participants took part in genetic testing, which can occasionally identify

markers for other health conditions. The information sheet sought to ensure that the

parents/carers understood possible outcomes of the genetic test before giving consent.

Ethical approval was granted by the University of Salford Ethics committee in May 2019

(reference: HSR1819-100).

Results

Participating schools

Schools represented a range of deprivation when measured by the index of multiple deprivation. School 1 was located in one of England's 20% most deprived areas, and school 3, 30%. School 2 was located among the 20% most affluent areas in England.

Flow of participants

A total of 220 children were invited and 203 children took part in the phase 1 physical measures (height, weight and OFC) (Figure 2). Fifty children were eligible for full assessment (phase 2) because of screening positive on the physical measures, parent/teacher concerns, already acknowledged as SEN, or currently or previously being in local authority care (Table 2).

Assessment was not possible on 24 children due largely to parents not giving consent or being uncontactable.

Full assessment took place with 26 children, four of whom had FASD and a further four were identified as possible FASD (Table 2). Other developmental disorders identified are detailed in supplementary file 2.

Of the children with full data sets, none had reported exposure to known teratogenic antiepileptic pharmaceuticals. Two had exposure to illegal drugs and over two thirds of the children assessed (68%) has some prenatal exposure to alcohol (Table 3).

Of the FASD cases, none had a prior clinical diagnosis of any neurodevelopmental disorder or an Education, Health and Care (EHC) Plan, and only one child had been identified by the school as having special educational needs. Three out of the four cases had high risk

prenatal alcohol exposure reported, while in the fourth, alcohol was not reported but the

child had severe FAS facial features, was small for age, a low full scale IQ of 66, and a normal

CGH array (a summary of cases and possible cases assessed against the FASD criteria is given

in supplementary file 2).

Of the possible cases where FASD was suspected but not confirmed, none had a clinical

diagnosis of any neurodevelopmental disorder and only one had an EHC Plan. All were on

the schools' special educational needs register, had prenatal exposure to alcohol, and had

deficits in at least three subdomains. Of the 12 children with a suspected disorder that was

deemed unlikely to be FASD, 11 had some level of prenatal exposure to alcohol (Table 3).

Prevalence Rates

262 We calculated a conservative (minimum) prevalence of FASD of 1.8% 4/220 (95% CI [1.0,

263 3.4]), and a conservative (minimum) prevalence that also included possible FASD of 3.6%

264 8/220 (95%CI [2.1, 6.3]) within our study population.

Discussion

Prevalence of FASD

This is the first FASD active case ascertainment study to be carried out in the UK, a country

with one of the highest rates of drinking in pregnancy in the world. These prevalence

estimates, though not necessarily generalisable to other communities, are in line with a

modelled population prevalence estimate for the UK of 3.2% (Lange et al., 2017b) and

consistent with a screening prevalence of 6-17% in a secondary analysis of data from a

cohort study of children born in the 1990s (McQuire et al., 2019). As per May et al. (2018),

our prevalence estimates could be considered as 'conservative' because we assumed all

those who we did not fully assess did not have FASD. Further work examining the

prevalence of FASD taking account of maternal sociodemographic factors and trends in

alcohol consumption would allow a more precise estimate of the likely burden of this

condition in the UK.

The study region (Greater Manchester) sees approximately 34,000 babies born per year

(ONS, 2020), yet diagnoses only around 36 cases of FASD per year (unpublished data from a

Freedom of Information request). This low rate of diagnosis in relation to the example local

prevalence we have demonstrated here suggests that increased recognition and diagnostic

capacity is required. This is important because early diagnosis and support for families

affected by FASD can prevent or mitigate adverse secondary outcomes, such as school

exclusions, poor job prospects and mental ill-health (Alex & Feldmann, 2012; Landgren, et

al., 2019; Rangmar, et al., 2016Streissguth, et al., 2004).

Strengths and Limitations

We demonstrated successful engagement with the three schools who made intensive

efforts on our behalf to recruit parents. The schools covered a range of communities in

different levels of deprivation, with two serving relatively deprived populations and one

relatively affluent. A larger, more definitive study would use a random sampling technique,

stratified by deprivation level, to select schools.

We were unable to obtain sufficient information for almost half (24/50) of the children who

were identified as being at higher risk of FASD (i.e. those who were identified in phase 1

screening). A further nine were actively withdrawn before phase 1 screening and selection

took place. It is not possible to know whether the proportion of cases would have been

higher or lower in these groups that did not take part. It is also possible that there were

cases of FASD in the remaining 161 children with no apparent risk factors. It is possible the

shame and stigma associated with FASD or developmental disorders in general may have

impacted on participation. Low participation has been a documented issue for other

prevalence studies (Caccanti et al,. 2014; May et al., 2011; Okuliez-Kozaryn, et al., 2017) in

the European region.

Extensive effort was made to contact each parent, and for those not taking part, the reasons

were documented (see reasons given in Figure 2). The opportunity to have children's needs

assessed was the major incentive for schools to take part in the study. For children in the

care of foster parents, consent was required from the local authority and this was

particularly difficult to obtain.

We had initially hoped to include a further four schools in the prevalence calculations,

however, disruptions due to COVID-19 prevented completion in these schools. Partial

findings are presented in supplementary file 1.

We were not successful in collecting data in a Social Emotional and Mental Health (SEMH)

school despite recruiting an enthusiastic head teacher at such a school. Anecdotally, parents

felt they had nothing to gain by taking part as their child was already benefiting from

specialist support. Similarly, pupil referral units were not included in this study. Together

with SEMH schools, this is another setting where FASD prevalence is anticipated to be

higher than the mainstream school setting.

It is likely that prenatal alcohol exposure was underestimated in this study, as dose and

duration of alcohol consumption during pregnancy was calculated from retrospective self-

reporting, relying on recall of behaviours taking place up to 10 years prior. Social desirability

bias also affects reporting of alcohol consumption during pregnancy (Caccanti et al,. 2014,

Smith et al., 2014) and we do not know how much the potential for perceived shame

associated with alcohol consumption in pregnancy may have affected reporting.

It may be beneficial for future research to consider how methods could be adjusted for

European populations to optimise participation and accuracy of report of alcohol

consumption in these populations.

Other general limitation in studies that measure the prevalence of FASD, it that whilst there

are a wide range of physical and neurological features associated with FASD, which when

taken together increase the likelihood of FASD, the only features that are widely accepted

as discriminating in the absence of confirmed alcohol exposure are the distinctive facial

features. However, significant facial features are thought to only occur in a minority of cases

of FASD, and, likewise, in this study only two out of the eight cases and possible cases

showed these distinctive features. Previous prevalence studies have used a higher threshold

for the physical measurements at phase 1 (e.g. a threshold of 10th percentile for OFC: May

et al., 2011), which would have made the screening stage more sensitive and may have led

to more cases being identified. Instead, we used the Canadian diagnostic approach, where

the detailed evaluation stage was more focused towards the neurocognitive deficits. We

ruled out obvious genetic causes through chromosome microarray analysis and clinical

geneticist review. Whilst this helped to improve confidence about the aetiological basis of

the presentation, it is not possible to prove that the alcohol exposure caused the deficits.

The converse also true: it is not possible to rule out alcohol as an important causal factor.

Every effort was taken to reduce bias that may give rise to false positives. The case

definition used validated developmental assessments and was described in advance in the

study protocol and followed internationally recognised guidance (Cook et al., 2016).

An improvement on our methodology would have been to include a random sample of

children for full assessment. This would have enabled us to obtain characteristics of the

birth mothers in a group of children without any risk factors for FASD (i.e. those who did not

meet the phase 1 screening criteria), which would have allowed more sophisticated

modelling (and extrapolation) of the likely prevalence in the entire population of 8-9 year

olds (May et al., 2018). In this small-scale study, we determined that this would have been

difficult. Parents were prepared to take part if they thought it might benefit their child, for

example by getting more information to inform their special educational needs or to obtain

a diagnosis. The fact that our assessment also identified possible ASD, ADHD and other

neurodevelopmental conditions was a significant 'pull factor' for schools and parents. It was

also notable that the only children actively opted into the study because of parent concerns

(and no other risk factor) were from the school in the most affluent setting (school 2). Thus,

motivations to take part may differ depending on socioeconomic status. We would conclude

that further research would be needed into how to incentivise parents of children with

apparently typical development to take part.

Conclusions

This is the first study in the UK to directly assess FASD in a systematically ascertained sample

of children. It found FASD in 1.8% (1.0%-3.4%) of the population studied, or 3.6%

(2.1%,6.3%) when possible cases were also included. Due to the small sampling frame of

schools included and limitations of baseline information obtained on contacted families, we

can only conclude this represents local prevalence data in typical mainstream schools rather

than being able confidently to infer a 'population prevalence' of FASD. There are

uncertainties too about this prevalence found, since half the children screened positive

were lost to full ascertainment, and case identification may have been higher if all cases had

been seen. Further research is needed to identify how to improve participation and

accuracy of PAE in European populations.

It was already suspected from modelling and screening studies that FASD is highly prevalent

in the UK (Lange et al., 2017a; McQuire et al., 2019). Confirmation that this is the case in a

sample of schools should be used to increase the awareness of FASD, and invest in

diagnostic services, treatment and support for those affected by FASD.

Abbreviations

ADHD	Attention-Deficit/Hyperactivity Disorder
ARBD	Alcohol-Related Birth Defects
ARND	Alcohol-Related Neurodevelopmental Disorder)/
ASD	Autism Spectrum Disorder
CCC	Children's communication Checklist
FAS	Fetal Alcohol Syndrome
FASD	Fetal Alcohol Spectrum disorders
FSIQ	Full Scale Intelligence Quotient
IDACI	Income Deprivation Affecting Children Index
IMD	Index of Multiple Deprivation
ND-PAE	Neurobehavioral Disorder Associated with Prenatal Alcohol Exposure
NEPSY II	NEuroPSYchological Assessment 2nd Edition
NHS	National Health Service
OFC	Occipital Frontal Circumference
PAE	Prenatal Alcohol Exposure
pFAS	partial Fetal Alcohol Syndrome
SCQ	Social Communication Questionnaire
SDQ	Strengths and Difficulties Questionnaire
SEMH	Social, Emotional And Mental Health School
SEN	Special Educational Needs
SENCO	Special Educational Needs Coordinator
SFA	Small for age
SSP	Short sensory profile
WISC V	Weschler Intelligence scale for Children 5th Edition

References

Alex, K. & Feldmann, R. 2012. Children and adolescents with fetal alcohol syndrome (FAS): better social and emotional integration after early diagnosis. *Klin Padiatr,* 224**,** 66-71.

Astley, S. J. 2015. Palpebral fissure length measurement: accuracy of the FAS Facial Photographic Analysis Software and inaccuracy of the ruler. *Journal of Population Therapeutics and Clinical Pharmacology, 22(1).*

Ceccanti, M., Fiorentino, D., Coriale, G., Kalberg, W. O., Buckley, D., Hoyme, H. E., ... & May, P. A. (2014). Maternal risk factors for fetal alcohol spectrum disorders in a province in Italy. *Drug and Alcohol Dependence*, 145, 201-208.

Chasnoff, I J., Wells, A.M., King, L. 2015. "Misdiagnosis and missed diagnoses in foster and adopted children with prenatal alcohol exposure." *Pediatrics* 135.2, 264-270.

Cook, J. L., Green, C. R., Lilley, C. M., Anderson, S. M., Baldwin, M. E., Chudley, A. E., Conry, J. L., Leblanc, N., Loock, C. A., Lutke, J., Mallon, B. F., Mcfarlane, A. A., Temple, V. K. & Rosales, T. 2016. Fetal alcohol spectrum disorder: a guideline for diagnosis across the lifespan. *Can. Med. Assoc. J.,* 188**,** 191-197.

Dnagenotek 2020. Oragene 500-series. *Genotek.*

Douzgou, S., Breen, C., Crow, Y. J., Chandler, K., Metcalfe, K., Jones, E., Kerr, B. & Clayton-Smith, J. 2012. Diagnosing fetal alcohol syndrome: new insights from newer genetic technologies. *Arch. Dis. Child.,* 97**,** 812-817.

Kaufman, A. S., Raiford, S. E. & Coalson, D. L. 2015. *Intelligent testing with the WISC-V,* USA, John Wiley & Sons.

Kelly, Y., Sacker, A., Gray, R., Kelly, J., Wolke, D., & Quigley, M. A. (2009). Light drinking in

pregnancy, a risk for behavioural problems and cognitive deficits at 3 years of age?.

International journal of epidemiology, 38(1), 129-140.

Landgren, V., Svensson, L., Gyllencreutz, E., Aring, E., Grönlund, M. A., & Landgren, M.

(2019). Fetal alcohol spectrum disorders from childhood to adulthood: a Swedish

population-based naturalistic cohort study of adoptees from Eastern Europe. BMJ

open, 9(10), e032407.

Lange, S., Probst, C., Gmel, G., Rehm, J., Burd, L. & Popova, S. 2017a. Global Prevalence of

Fetal Alcohol Spectrum Disorder Among Children and Youth A Systematic Review

and Meta-analysis. *JAMA Pediatr,* 171**,** 948-956.

Lange, S., Probst, C., Gmel, G., Rehm, J., Burd, L. & Popova, S. 2017b. Global Prevalence of

Fetal Alcohol Spectrum Disorder Among Children and Youth: A Systematic Review

and Meta-analysis. *JAMA Pediatr,* 171**,** 948-956.

May, P. A., Chambers, C. D., Kalberg, W. O., Zellner, J., Feldman, H., Buckley, D., Kopald, D.,

Hasken, J. M., Xu, R., Honerkamp-Smith, G., Taras, H., Manning, M. A., Robinson, L.

K., Adam, M. P., Abdul-Rahman, O., Vaux, K., Jewett, T., Elliott, A. J., Kable, J. A.,

Akshoomoff, N., Falk, D., Arroyo, J. A., Hereld, D., Riley, E. P., Charness, M. E., Coles,

C. D., Warren, K. R., Jones, K. L. & Hoyme, H. E. 2018. Prevalence of Fetal Alcohol

Spectrum Disorders in 4 US Communities. *JAMA,* 319, 474-482.

May, P. A., Fiorentino, D., Coriale, G., Kalberg, W. O., Hoyme, H. E., Aragon, A. S., Buckley,

D., Stellavato, C., Gossage, J. P., Robinson, L. K., Jones, K. L., Manning, M. & Ceccanti,

M. 2011. Prevalence of children with severe fetal alcohol spectrum disorders in

communities near Rome, Italy: new estimated rates are higher than previous

estimates. *Int. J. Environ. Res. Public. Health,* 8**,** 2331-51.

May, P. A. & Gossage, J. P. 2001. Estimating the prevalence of fetal alcohol syndrome - A

summary. *Alcohol Research & Health,* 25**,** 159-167.

Mcquire, C., Mukherjee, R., Hurt, L., Higgins, A., Greene, G., Farewell, D., Kemp, A. &

Paranjothy, S. 2019. Screening prevalence of fetal alcohol spectrum disorders in a

region of the United Kingdom: A population-based birth-cohort study. *Prev. Med.,*

118**,** 344-351.

Ministry of Housing Communities and Local Government 2019. The English Indices of

Deprivation. https://www.gov.uk/government/statistics/english-indices-of-

deprivation-2019.

Morleo, M., Woolfall, K., Dedman, D., Mukherjee, R., Bellis, M. A. & Cook, P. A. 2011. Under-

reporting of foetal alcohol spectrum disorders: an analysis of hospital episode

statistics. *BMC Pediatr.,* 11.

Mukherjee, R., Layton, M., Yacoub, E. & Turk, J. 2011. Autism and autistic traits in people

exposed to heavy prenatal alcohol: data from a clinical series of 21 individuals and

nested case control study. *Advances in Mental Health and Intellectual Disabilities*.

Mukherjee, R., Wray, E., Curfs, L. & Hollins, S. 2015. Knowledge and opinions of professional

groups concerning FASD in the UK. *Adoption and Fostering,* 39**,** 212-224.

Okulicz-Kozaryn, K., Borkowska, M., & Brzózka, K. (2017). FASD prevalence among

schoolchildren in Poland. *Journal of Applied Research in Intellectual Disabilities*,

30(1), 61-70.

ONS 2020. Births in England and Wales: summary tables.

Popova, S., Lange, S., Probst, C., Gmel, G. & Rehm, J. 2017. Estimation of national, regional,

and global prevalence of alcohol use during pregnancy and fetal alcohol syndrome: a

systematic review and meta-analysis. *Lancet Global Health,* 5**,** E290-E299.

Public Health England 2021. Public Health Profiles.

https://fingertips.phe.org.uk/search/alcohol%20specific%20death.

Raldiris, T. L., Bowers, T. G. & Towsey, C. 2018. Comparisons of intelligence and behavior in

children with fetal alcohol spectrum disorder and ADHD. *Journal of Attention*

Disorders, 22 (10), 959-970.

Rangmar, J., Hjern, A., Vinnerljung, B., Strömland, K., Aronson, M., & Fahlke, C. (2015).

Psychosocial outcomes of fetal alcohol syndrome in adulthood. *Pediatrics*, 135(1),

e52-e58.

Rasmussen, C., Tamana, S., Baugh, L., Andrew, G., Tough, S. & Zwaigenbaum, L. 2013.

Neuropsychological impairments on the NEPSY-II among children with FASD. *Child*

Neuropsychology, 19**,** 337-349.

Reynolds, R., Mccarthy, R. & Cook, P. A. 2021. We do things differently here: the Greater

Manchester approach to preventing alcohol-exposed pregnancy. *Perspectives in*

Public Health. September 2021.

Roozen, S., Peters, G. J. Y., Kok, G., Townend, D., Nijhuis, J. & Curfs, L. 2016. Worldwide

Prevalence of Fetal Alcohol Spectrum Disorders: A Systematic Literature Review

Including Meta-Analysis. *Alcoholism-Clinical and Experimental Research,* 40**,** 18-32.

Sayal, K., Heron, J., Draper, E., Alati, R., Lewis, S. J., Fraser, R., ... & Gray, R. (2014). Prenatal

exposure to binge pattern of alcohol consumption: mental health and learning

outcomes at age 11. *European Child & Adolescent Psychiatry*, 23(10), 891-899.

Scholin, L., Mukherjee, R. a. S., Aiton, N., Blackburn, C., Brown, S., Flemming, K. M., Gard, P.

R., Howlett, H., Plant, M., Price, A. D., Shields, J., Smith, L. A., Suttie, M., Zammitt, D.

C., Cook, P. A. & Collaboration, U. F. R. 2021. Fetal alcohol spectrum disorders: an

overview of current evidence and activities in the UK. *Arch. Dis. Child.*

Smith, L., Savory, J., Couves, J. & Burns, E. 2014. Alcohol consumption during pregnancy:

Cross-sectional survey. *Midwifery,* 30**,** 1173-1178.

Streissguth, A. P., Bookstein, F. L., Barr, H. M., Sampson, P. D., O'malley, K. & Young, J. K.

2004. Risk factors for adverse life outcomes in fetal alcohol syndrome and fetal

alcohol effects. *J. Dev. Behav. Pediatr.,* 25**,** 228-38.

World Health Organisation 2012. WHO Global Prevalence Study on FASD. WHO Research

Initiative on Alcohol, Health and Development, WHO International Collaborative

Research Project on Fetal Alcohol Spectrum Disorders (FASD).

Young, S., Absoud, M., Blackburn, C., Branney, P., Colley, B., Farrag, E., Fleisher, S., Gregory,

G., Gudjonsson, G. H., Kim, K., O'malley, K. D., Plant, M., Rodriguez, A., Ozer, S.,

Takon, I., Woodhouse, E. & Mukherjee, R. 2016. Guidelines for identification and

treatment of individuals with attention deficit/hyperactivity disorder and associated

fetal alcohol spectrum disorders based upon expert consensus. *BMC Psychiatry,* 16**,**

324.

Figure legends

487 Figure 1. Study design

488 Figure 2. Flow of participants through the study.

489 SFA: small for age; SEN: on school educational needs register; LAC: 'looked after child' i.e. in

490 the care of the local authority; PrevLAC: previously looked after, i.e. adopted.

Table 1

Table 1: Measures used and Fetal Alcohol Spectrum Disorder (FASD) domains of

impairment

Domain	Sub-domain	Measure
Growth		Height below 9th percentile and Weight below 9th percentile, medical history
Face		Photographic measurements 4 digit FAS Facial Measurement software.
Central Nervous System	Brain	OFC below 2nd centile, medical history or brain scan report
	Hard/ Soft Neurological signs	SSP Defined differences in 2 or more domains History of formally diagnosed motor disorder e.g. dyspraxia History of epilepsy or other neurological issues (from medical history)
	Communication	CCC-2, Vineland communication domain. Defined by abnormality >1.5 standard deviations below the mean
	Cognition	WISC V: Full Scale IQ 2 standard deviations from norm or 2 standard deviations between different subdomains
	Executive Function	BRIEF overall score: Score above clinical cut off range NEPSY executive function domain Score 1.5 standard deviations from norm or 2 standard deviations between subdomains
	Memory	WISC V, NEPSY Memory domain
	Attention	SDQ attention domain scores above clinical cut offs

		NEPSY attention domain
	Adaptive behaviour, social skills	SCQ (score above 15), SDQ (score above clinical cut off), Vineland Social domain, Vineland Daily Living Skills domain (both scores 1.5 standard deviations from mean)
	Academic	Teacher report, below average level, has EHC Plan
Alcohol consumption of birth mother		History information sheet

498 BRIEF: Behaviour Rating Inventory of Executive Function

499 CCC-2: Communication Checklist for Children

500 EHC Plan: Education Health and Care Plan

501 FAS: Fetal Alcohol Syndrome

502 IQ: Intelligence Quota

503 NEPSY: NEuroPSYchological Assessment

504 OFC: Occipitofrontal Circumference

505 SSP: Short Sensory Profile

506 SCQ: Social Communication Questionnaire

507 SDQ: Strengths and Difficulties Questionnaire

508 WISC-V: Wechsler Intelligence Scale for Children Fifth UK Edition

511 **Table 2: Screening and assessment data by school**

	School1 (baseline N=44)	School2 (baseline N=118)	School3 (baseline N=59)	Total (N=220)
Screened at stage one n (% of baseline)	38 (86)	109 (92)	56 (95)	203 (92)
Sex, n(% of baseline)				
Male	18 (40)	55 (46)	30 (51)	103 (47)
Female	20 (45)	54 (46)	26 (44)	100 (45)
Age, Years (y) & months (m)				
Median	8y11m	8y10m	8y6m	8y10m
[Interquartile range]	[8y9m, 9y5m]	[8y5m, 9y6m]	[8y3m, 8y11m]	[8y 5m, 9y4m]
(range)	(8y3m, 9y9m)	(8y0m, 9y10m)	(8y0m, 9y8y)	(8y0m, 9y10m)
Physical screen positive n (% of baseline)				
Height <9[th] percentile	4 (9)	7 (6)	2 (3)	13 (6)
Weight <9[th] percentile	2 (5)	3 (3)	2 (3)	7 (3)
OFC <2th percentile	1 (2)	1(1)	1 (2)	3 (1)
Reason for recruitment n (% of invited)				
SEN	14(66)	7 (43)	5 (50)	26 (13)
SFA	3(14)	5 (31)	4 (40)	12 (6)
LAC	2(10)	3 (18)	1 (10)	6 (3)
SFA+SEN	2(10)	0	0	2 (1)
SFA + prevLAC	0	1 (6)	0	1 (0.5)
Parent opt-in	0	3 (16)	0	3 (1)
Invited to second phase	21	19	10	50
Phase two assessments	**N=10**	**N=9**	**N=7**	**N=26**
Referral reason (% of phase two participants)				
SEN	5 (50)	3 (33)	4 (57)	12 (46)
SFA	3 (30)	0	2 (29)	5 (19)
LAC	0	2 (22)	1 (14)	3 (12)
SFA and SEN	2 (20)	0	0	2 (8)
SFA and PrevLAC	0	1 (11)	0	1 (4)
parent opt in	0	3 (33)	0	3 (11)
Maternal Ethnicity, n(%)				

White UK	10 (100)	8 (89)	6 (86)	24 (92)
Non white	0	1 (11)	0	1 (4)
Not reported	0	0	1 (14)	1 (4)
Maternal educational qualifications, highest received n(%)				
<4 GCSEs	2 (20)		1 (14)	3 (12)
4+GCSEs	4 (40)		2 (29)	6 (23)
2+A levels	2 (20)	3 (33)	1 (14)	6 (23)
Degree	2 (20)	4 (44)	2 (29)	8 (30)
Missing data	0	2 (22)	1 (14)	3 (12)
Full assessment				
Deficits in three subdomains met, n(% of assessed)	8 (80)	7 (78)	6 (86)	21 (80)
FASD, n (% of assessed)				
FASD	1 (10)	3 (33)	0 (0)	4 (15)
FASD and possible FASD	3 (30)	4 (44)	1 (10)	8 (30)
Other outcomes				
ASD	0 (0)	2 (22)	3 (43)	5 (19)
ADHD	2 (20)	0 (0)	1 (14)	3 (12)
DLD	2 (20)	1 (10)	0 (0)	3 (12)
General learning disability	0 (0)	0 (0)	2 (28)	2 (8)
No disorder	3 (30)	2 (20)	0 (0)	5 (19)
FASD prevalence per 100 [95% CI]				
FASD	2.3 [0.3,14.8]	2.5 [0.8, 7.4]	No observations	1.8 [1.0,3.4]
FASD and possible FASD	7.0 [2.2,19.8]	3.4 [1.3, 8.7]	1.7 [0.2,11.1]	3.6 [2.1,6.3]

SEN: on school educational needs register; SFA: small for age; LAC: 'looked after child', i.e.

under care of local authority; PrevLAC: previously looked after child, i.e. adopted; OFC:

occipital frontal circumference; GCSE: General Certificate of Education, qualification usually

taken at age 16 years; A level: Advanced Level, qualification usually taken at age 18 years;

FASD: fetal alcohol spectrum disorder; ASD: autism spectrum disorder; ADHD: attention

deficit hyperactivity disorder; DLD: developmental language disorder.

Table 3

520   **Table 3: prenatal exposure data from cases with a complete dataset**

Reported Prenatal exposures	FASD N=4	Possible FASD n=4	Other N= 18	TOTAL N=26
Anticonvulsants	0	0	0	0
Illegal drugs	1(25%)	0	1 (6%)	2 (8%)
Alcohol				
Significant risk[a]	3 (75%)	3(75%)	4 (22%)	10 (38%)
Low risk[b]	0	1 (25%)	7 (40%)	8 (30%)
No alcohol reported	1(25%)	0	7 (40%)	8 (30%)

521   [a] Cut off for significant risk defined as four or more alcoholic drinks on four or more

522   occasions or greater than seven units a week throughout the term of the pregnancy

523   [b] Low risk: alcohol consumed, but not meeting threshold above (Kelly, et al., 2009; Sayal et

524   al., 2013).

Table legends

Table 2: Measures used and Fetal Alcohol Spectrum Disorder (FASD) domains of impairment

BRIEF: Behaviour Rating Inventory of Executive Function

CCC-2: Communication Checklist for Children

EHC Plan: Education Health and Care Plan

NEPSY: NEuroPSYchological Assessment

OFC: Occipitofrontal Circumference

SSP: Short Sensory Profile

SCQ: Social Communication Questionnaire

SDQ: Strengths and Difficulties Questionnaire

WISC-V: Wechsler Intelligence Scale for Children Fifth UK Edition

Table 2: Screening and assessment data by school

SEN: on school educational needs register; SFA: small for age; LAC: 'looked after child', i.e.

under care of local authority; PrevLAC: previously looked after child, i.e. adopted; OFC:

occipital frontal circumference; GCSE: General Certificate of Education, qualification usually

taken at age 16 years; A level: Advanced Level, qualification usually taken at age 18 years;

FASD: fetal alcohol spectrum disorder; ASD: autism spectrum disorder; ADHD: attention

deficit hyperactivity disorder; DLD: developmental language disorder.

Table 3: prenatal exposure data from cases with a complete dataset

[a]Cut off for significant risk defined as four or more alcoholic drinks on four or more

occasions or greater than seven units a week throughout the term of the pregnancy.

[b]Low risk: alcohol consumed, but less than four or more occasions or less than seven units a

week

Figure 1

INVITATION

3 schools, all children aged 8 or 9 years in Years 3 and 4

PHASE 1

Initial screening with children (participation on the basis of attendence at school on the measurement days and no parental opt out)

Height, Weight and Occipitofrontal measurement

All children screened positive at step 1 invited to step two and either:

small / small head (occipitofrontal) circumference (OFC)

and/or identfied as SEN

and/or teacher-identified /parent-identified behaviour problem or learning difficulty

and/or looked after child

PHASE 2a: Parent Assessment

Parent opt-in required, assessed with parent/teacher-completed measurement tools:

Social Communication Questionnaire (SCQ)

Strengths and Difficulties Questionnaire (SDQ)

Short Sensory Profile (SSP)

Vineland Adaptive Behavior Scales II (VABS-II)

Communication checklist for children (CCC-2)

Medical History sheet

PHASE 2b: Child Assessment

child assessed for (child-completed)

Cognitive (WISC V and NEPSY)

face morphology (photography)

PHASE 2c

genetic testing

(rule out genetic causes)

case conference

children fully meet criteria for FASD

or possible cases

Figure 2