Pulsed electromagnetic fields can prevent bone loss and improve balance function

in older adults with primary osteoporosis: a systematic review and meta-analysis

Siyi Zhu[a,b], Yi Li[a,b], Liqiong Wang[a,b], Jinming Huang[a,b], Kangping Song[a,b], Xinling Gan[a,b],

Xiaona Xiang[a,b], Chengqi He[a,b*], Lin Yang[a,b*]

[a]Rehabilitation Medical Center, West China Hospital, Sichuan University, Chengdu, Sichuan,

China.

[b]Rehabilitation Medicine key laboratory of Sichuan Province, West China Hospital, Sichuan

University, Chengdu, Sichuan, China.

*Corresponding author: Dr. Chengqi He or Dr. Qiang Gao or Dr. Lin Yang at the Rehabilitation

Medical Center, West China Hospital, Sichuan University, No. 37 Guoxue Alley, Wuhou

District, 610041, Chengdu, China. Email: hxkfhcq@126.com (CH) or 5648197@qq.com (LY).

Abbreviated title: Pulsed electromagnetic fields for primary osteoporosis

Key words: Osteoporosis, Electromagnetic Fields; Bone Density; Physical Therapy

Modalities; Disability Evaluation

Word Count: 250 words (Abstract)

3730 words (Main text)

ABSTRACT

Question(s): What is the effectiveness of pulsed electromagnetic fields (PEMFs) for the management of primary osteoporosis in older adults?

Design: Systematic review and meta-analysis of randomised or quasi-randomised controlled trial.

Participants: Older adults aged over 50 years with primary osteoporosis.

Intervention: PEMFs at extremely low frequencies.

Outcome measures: Bone mass, number of incident fractures, functional assessments, and adverse events.

Results: Eight trials including 396 participants met the inclusion criteria. Low certainty evidence showed that PEMFs was non-inferior to conventional pharmacological agents in preventing the decline of Bone Mineral Density (BMD) at the lumbar (MD 0.01; CI -0.04 to 0.06) and femur neck (MD 0.01; CI -0.02 to 0.04), and improving balance function measured by Berg Balance Scale (BBS) (MD 0.01; CI -0.09 to 0.11) and Timed Up and Go test (MD -0.04; CI -0.80 to 0.72), directly after intervention. The similar effects were observed in BMD and BBS at 12- and 24-weeks follow-up from baseline with moderate certainty evidence. Very low certainty evidence showed that PEMFs (versus exercise) had small but significant effect on BMD at the femur neck (MD 0.10; CI 0.01 to 0.20), and no effect on BMD at the lumbar (MD 0.15; CI -0.04 to 0.35).

Conclusion: PEMFs had positive effects non-inferior to first-line treatment on BMD and balance function in older adults with primary osteoporosis, but with low to very low certainty evidence and short-term follow-ups. There is a need for high-quality randomised controlled trials evaluating PEMFs for the management of primary osteoporosis.

Registration: PROSPERO CRD42018099518.

INTRODUCTION

Osteoporosis is a systemic and multifactorial skeletal disorder characterized by low bone mineral density and skeletal fragility that occur with aging, with a consequent increase of susceptibility to low-trauma fractures [1]. The most prevalent symptoms of osteoporosis are fractures at vertebrae, proximal femur (hip), and wrist affecting patients' physical function and quality of life. It is estimated that over 200 million people worldwide are affected by osteoporosis, accounting for 8.9 million fractures annually [2]. The possibility of osteoporotic fractures exceeds 40% and the probability of hip fracture alone could target 20% in white female population over 50 years old [3]. In China, a higher incidence of hip fractures in men than in women was reported [4]. Each year, osteoporotic fractures account for over 432,000 hospitalizations and 2.5 million medical visits in the USA [5]. The treatment costs for fractures were recorded at nearly $17 billion in the USA in 2005 [6], and €31.7 billion in Europe in 2000 [7]. Therefore, osteoporosis has been identified as a major health burden globally by WHO, due to its high prevalence, disability rate, related mortality and poor quality of life [8].

Rehabilitation interventions given its important roles in modifying risk factors related to fractures, restoring function and improving quality of life are frequently recommended as an option in the nonpharmacological management of osteoporosis [9,10]. Pulsed electromagnetic fields (PEMFs) at a specific intensity and frequency have been proved effective in attenuating bone loss and the relief of pain and discomfort after osteoporosis. As we reviewed elsewhere [11], PEMFs were found to be positive at promoting bone formation by stimulating the formation and differentiation of osteoblasts, and negative at inhibiting the function of osteoclasts in bone resorption. Experimental studies suggest that PEMFs may exert effects on Ca^{2+}-related receptors on the bone cell membrane which play a regulatory role in the maintenance of bone remodelling [12]. Further, the exposure of PEMFs could influence the physiopathology of osteoporosis by targeting inflammation and potentially relieving pain via these regulatory processes and improvements in bone remodeling [13].

PEMFs have been widely used as an clinical option for the management of pain and discomfort related to osteoporosis since the introduce of its usage for non-union fractures was approved by FDA in 1979 [14]. However, clinical trials evaluating the effectiveness of PEMFs have been conducted with inconsistent results, to which parameters of PEMFs used in studies, follow-up time points and clinical settings differ across studies may lead [11]. In order to expand upon the

76 current knowledge on whether PEMFs is an effective physical agent for osteoporosis clinically, a
77 systematic review and meta-analysis of clinical trials was performed to compare PEMFs with
78 placebo or sham or other agents for the management of primary osteoporosis in older adults.
79

METHODS

81 This systematic review and meta-analysis was conducted in accordance with the Cochrane
82 Handbook for Systematic Reviews of Interventions [15] and reported based on Preferred Reporting
83 Items for Systematic Reviews and Meta-analyses guidelines (PRISMA) [16]. The protocol of this
84 study is available in PROSPERO (CRD42018099518) [17].
85

Identification and selection of studies

We searched the MEDLINE (via Ovid), EMBASE (via Ovid), Web of Science, CENTRAL and
CCTR (via The Cochrane Library), Physiotherapy Evidence Database (via PEDro website),
CNKI, VIP, Wan Fang, ClinicalTrials.gov (https://www.clinicaltrials.gov/) and Current
controlled trials (www.controlled-trials.com) from the inception dates to December 9, 2018,
using the keywords *pulsed electromagnetic fields* and *osteoporosis*. The Open Grey
(http://www.opengrey.eu/) was searched for the Grey Literature research. The detailed electronic
search strategies are provided in Supplementary Appendix I. An additional search was performed
under a mechanism of living systematic review [18] to identify recently published randomized
clinical trials (RCTs) from December 10, 2018 to April 30, 2021 using the databases and
keywords described above. The whole procedure was assisted by a librarian from Sichuan
University.
Randomised controlled trials or quasi-randomised trials examining the effects of PEMFs
compared to placebo or sham or other agents for the treatment of primary osteoporosis (including
those with previous fractures) were included if they met the inclusion criteria listed in Box 1.
Studies were excluded if the study population had a diagnosis of corticosteroid-induced
osteoporosis or other secondary osteoporosis (e.g., rheumatoid arthritis), studies where
participants had a history of hip replacement or osteoporosis fractures related surgery, the study
type was observational studies, review articles, abstracts, conference reports and book chapters.
A three-stage screening methodology was performed to select relevant RCTs for this review.
Primarily, all titles were screened by one reviewer (SYZ) for eligibility and irrelevant papers

were excluded accordingly. Secondary, two reviewers (YL and LQW or KPS) independently
reviewed each study title and abstract. Thirdly, two independent reviewers (XNX and JMH or
XLG) accessed the full text to assess against the eligibility criteria for each potentially eligible
study. A third reviewer (CQH or LY) was involved for any disagreement.
111

Assessment of characteristics of studies

Quality assessment

The risk of bias was assessed by using the Cochrane Collaboration's 'Risk of bias' tool [15]. Seven
key domains were assessed by two reviewers (SYZ and LQW): 1) the randomization sequence
generation, 2) allocation concealment, 3) blinding of participants and personnel, 4) blinding of
outcome assessment, 5) incomplete outcome data, 6) selective reporting, and 7) other bias. The
included studies were graded as low, unclear, or a high risk of bias. Methodological quality was
assessed with the use of Physiotherapy Evidence Database (PEDro) tool [19], which was proved
reliable [20] and valid [21]. Each criterion in the PEDro scale with a range of 0-10 was scored 1
("yes") or 0 ("no, don't know/unclear"). Generally, trials with a PEDro summary score of over
five. were considered to have adequate methodological quality [22]. Finally, we used Grading of
Recommendations, Assessment, Development and Evaluations (GRADE) [23] to describe the
overall quality of the body of evidence.
125

126 *Participants*

127 To be included, studies involved participants were healthy older adults (including those with
128 previous fractures) aged over 50 years with primary osteoporosis [24], recognized by two distinct
129 types [25]: 1) type I occurred in postmenopausal women; 2) type II, known as senile osteoporosis,
130 occurred in both men and women.
131

132 *Interventions*

133 All RCTs applying electromagnetic fields with pulsed signal and extremely low frequencies
134 (between 5 and 300 Hz) for the management of primary osteoporosis were included. The
135 parameters (frequency and intensity) of PEMFs, sessions per week and total duration of the
136 treatment period were recorded to describe the interventions.
137

Outcome measures

All outcomes were continuous data and recorded as the percent change from baseline to post-

intervention and different follow-up timepoints. To be included, trials had to provide original

data or sufficient information about at least one of outcomes on bone mass, number of incident

fractures, self-reported data on the changes in balance and quality of life, physical activity and

function, and adverse events. Primary outcomes were Bone mass (e.g., Bone Marrow Density or

Bone Mineral Content) immediately post-intervention and at follow-ups. Secondary outcomes

were functional assessments (e.g., Berg Balance Scale, Timed Up and Go test), quality of life

(e.g., EuroQoL (EQ 5D)) and adverse events (e.g., falls, number of incident fractures, death).

Data extraction and analysis

Two independent reviewers (SYZ and YL) extracted the following information from eligible

studies: lead author; year of publication; original country; subject characteristics; study design;

treatment information; intervention protocol; outcome measures; raw outcome data; follow-up

period and other relevant information. Disagreements were resolved by consensus.

All meta-analyses were performed using analysis was performed using Review Manager

(RevMan) software (The Cochrane Collaboration, version 5.4). For each included study, the

mean difference (MD) of percentage change with 95% confidential intervals (CIs) was calculated

when the outcome measures were consistent across studies or else the standard mean difference

(SMD) was calculated instead for continuous outcomes (reporting mean and standard deviation

(SD) or standard error (SE) of the mean). If the MD was not reported, it was calculated as the

change between values of the baseline and post-intervention. In the case that the value of SD

(SD_{diff}) was not reported, it was obtained 1) by multiplying SEs of means by the square root of

the sample size when standard errors (SEs) of the means were reported, or 2) with SDs at the

baseline ($SD_{baseline}$) and post-intervention SD (SD_{post}) in addition to the within-groups bivariate

correlation coefficient (r) [26]:

$$SD_{diff} = \sqrt{SD_{baseline}^2 + SD_{post}^2 - (2 \times r \times SD_{baseline} \times SD_{post})}$$

The I^2 statistic was employed for evaluating heterogeneity and a standard Chi^2 test was employed

for detecting whether significant heterogeneity existed. Heterogeneity was statistically significant

at $P < 0.10$ after due consideration of I^2 statistic, of which a value greater than 50% was

considered substantial heterogeneity [27]. The random-effects model was applied where the
evidence of heterogeneity was found.

The comparison was established between PEMFs and placebo control or exercise in the meta-
analysis. The subgroup analysis was conducted to detect the effectiveness relative to different
follow-up timepoints (postintervention; follow-up at 12, 24 weeks from baseline). To evaluate the
quality and consistency of pooled results, the sensitivity analysis was conducted by deleting each
included study. Where the data allowed, assessment of publication bias was performed. All tests
were two-tailed, and $P < 0.05$ was considered statistically significant.

RESULTS

Flow of studies through the review

In total, 806 articles were identified by our search, of which 124 duplicate articles were removed.
Based on title and abstract screening, 632 of these articles were excluded. Full texts of 50 articles
were read, a further 42 articles were excluded, remaining 8 articles included in the data extraction
and analysis of the review (Figure 1) [28-35].

Characteristics of studies

Quality

All included studies achieved PEDro scores over 5, among which five studies achieved the score
over 8 (Table 1)[28,29,31-33]. Of the 8 included studies, 2 studies were rated as 'low risk of bias' in all

domains[31,32], and other studies were classified as 'unclear risk of bias' for at least 1 aspect or

'high risk of bias' for at least 2 aspects. In results of GRADE, the quality of the evidence for the
comparison between PEMFs versus placebo control was low or moderate, and that for the
comparison between PEMFs versus exercise was very low. The results of the risk of bias and
GRADE are presented in the Supplementary Tables and Figures.

Participants

In total, data were extracted for 396 participants, comprising 183 participants in PEMFs group
and 213 participants in placebo control (alendronate/pharmacological therapy) or exercise group.
Mean age ranged from 56.3 to 70 years, with a gender ratio of 49 to 347. Participant

characteristics are detailed in Table 2.

Intervention

The frequency and intensity of PEMFs exposure varied at 8-100 Hz and 1.2-5 mT separately. A

range of 30-36 PEMFs sessions, with 30-60 min/session, were prescribed for participants, and 4

to 72 weeks follow-up were conducted across all studies. Control intervention types included

first-line pharmacological agents (e.g., Alendronate, intake of Vitamin D and Calcium) and

exercise (e.g., whole body vibration and resistance training). The intervention characteristics of

the included studies are detailed in the Table 2.

Outcome measures

The outcome measures in each study with the categories of bone mass and functional assessments

are detailed in Table 2.

Effect of intervention I: PEMFs versus control group

Bone Mineral Density (BMD)

Five studies (study population, N=248) [28,30-33] and three studies (study population, N=124)

[30,32,33] reported data on percentage change in BMD at the lumbar and femur neck respectively

after intervention directly (Figure 2). Low certainty evidence showed that PEMFs has no effect

on BMD at the lumbar (MD 0.01; CI -0.04 to 0.06) and femur neck (MD 0.01; CI -0.02 to 0.04)

with no statistically significant heterogeneity (lumbar: I^2= 0%, P =0.48; femur neck: I² = 0%, P =

0.79). Two studies (study population, N=125) [31,32] performed follow-ups at 12 and 24 weeks from

baseline on percentage change in BMD at the lumbar (Figure 2). Moderate certainty evidence

showed that there is no statistically significant effect on BMD at the lumbar at the 12 weeks

follow-up (MD -0.01; CI -0.28 to 0.25) and the 24 weeks follow-up (MD -0.02; CI -0.30 to 0.26)

with no statistically significant heterogeneity (12 weeks: I² = 0%, P = 1.00; 24 weeks: I² = 0%, P

= 0.98). No significant changes in heterogeneity and overall effect were observed in the

sensitivity analysis.

Berg Balance Scale (BBS)

Three studies (study population, N=168) [31,32,35] conducted the assessment of BBS after

intervention directly (Figure 3). Low certainty evidence showed that there is no statistically

significant effect on percentage change in BBS (MD 0.01; CI -0.09 to 0.11) with no statistically

significant heterogeneity ($I^2 = 0\%$, P = 1.00). Two studies (study population, N=125) [31,32] carried

out follow-ups at 12 and 24 weeks from baseline on percentage change in BBS (Figure 3).

Moderate certainty evidence showed that no statistically significant effect is detected on BBS at

the 12 weeks (MD 0.00; CI -0.13 to 0.14) and 24 weeks follow-up (MD -0.00; CI -0.15 to 0.14)

with evidence of no statistically significant heterogeneity (12 weeks: $I^2 = 0\%$, P = 0.99; 24 weeks:

$I^2 = 0\%$, P = 0.99). No significant changes in heterogeneity and overall effect were observed in

the sensitivity analysis.

Timed Up and Go (TUG) test

Two studies (study population, N=127) [31,35] assessed the percentage change in TUG test after

intervention directly (Figure 3). Low certainty evidence showed that there is no statistically

significant effect on TUG (MD -0.04; CI -0.80 to 0.72) with no statistically significant

heterogeneity ($I^2 = 0\%$, P = 0.97). No significant changes in heterogeneity and overall effect were

observed in the sensitivity analysis.

Effect of intervention II: PEMFs versus exercise group

Three studies (study population, N=110) [29,33,34] investigated the effect of PEMFs on percentage

change in BMD at the lumbar and femur neck respectively after intervention directly (Figure 4).

Very low certainty evidence showed that PEMFs has small but significant effect on BMD at the

femur neck (MD 0.10; CI 0.01 to 0.20), and no effect on BMD at the lumbar (MD 0.15; CI -0.04

to 0.35), both with no statistically significant heterogeneity respectively (femur neck: $I^2 = 28\%$, P

= 0.25; $I^2 = 47\%$, P = 0.15). In the sensitivity analysis, one study [29] was found to be a contributor

to results of non-significant heterogeneity and the effect on BMD at the femur neck. After

excluding the study, the heterogeneity was reduced to 0% (femur neck: P = 0.96; P = 0.86), and

the small but significant effect on BMD at the femur neck was eliminated (MD 0.00; CI -0.15 to

0.16).

DISCUSSION

This systematic review and meta-analysis of 8 studies involving 396 participants demonstrated

PEMFs as a physical therapy was non-inferior to conventional pharmacological agents in

preventing the decline of BMD and balance function for the management of primary osteoporosis

in older adults. Meanwhile, we also found that PEMFs might be slightly more effective in

increasing BMD than exercise. According to our knowledge, no systematic review and meta-

analysis was initiated before, except one network meta-analysis exploring effects of

nonpharmacological interventions including PEMFs on balance function only [36], and several

narrative reviews including clinical studies were retrieved [11,37,38]. Our results are in consistent

with findings from previous reviews[11,36,37] that PEMFs achieved positive effects on BMD and

balance function for older adults with primary osteoporosis, implicating that PEMFs may

potentially become a promising treatment option.

Bisphosphonates and exercise were both identified as the first-line interventions for the

management of osteoporosis in the latest evidence-based guideline [10]. Our study established

comparisons between PEMFs and active placebo or exercise based on groups set by included

studies, and sub-group analysis was stratified by different intervals between the baseline, post-

intervention, and follow-ups. BMD, as a surrogate measure for therapeutic effectiveness, can be

assessed by various methods, among which dual x-ray absorptiometry (DXA) was proved to be

reliable in the diagnosis of osteoporosis and have relatively good responsiveness in RCT [39,40].

Our study demonstrated that there was no difference between PEMFs and active placebo in

improving BMD at the lumbar and femur neck in all sub-group meta-analysis, which suggests

PEMFs is nearly effective as pharmacological agents for osteoporosis, and the effect could last

for at least 24 weeks. Moreover, the effects of PEMFs versus exercise on BMD at the lumbar was

not significant, while that was detected as small but significant on BMD at the femur neck. Our

results were considered fairly stable as the sensitivity analysis only detected one study [29]affecting

the heterogeneity and effect of PEMFs versus exercise on BMD at the femur neck. This is likely

due to variations in exercise characteristics (e.g., duration, training load and training volume).

Therefore, this part of results should be interpreted with caution.

Impaired balance function is an important risk factor increasing the incidence of falling and

fracture, which is modifiable by balance-improving interventions [41]. In line with results on BMD,

no statistically significant difference was observed for balance function measured by BBS and

TUG after the intervention of PEMFs versus active placebo. In the sub-group analysis, the effect

of PEMFs on BBS at 12, 24 weeks follow-up was also not statistically significant, confirming

that PEMFs is as effective as conventional pharmacological agents in improving balance function

which could last for 24 weeks. In consistent with a previous systematic review [36], it reported that

PEMFs exert positive effect on BBS and TUG tests reflecting balance function. However, the

network meta-analysis was conducted to further compare the effects of five interventions on

balance function with the conclusion that balance and strength training was better than other

interventions. Only one study using data on PEMFs was included in the analysis, compared to

that we included 3 studies, and no study was ever conducted to directly compare the effect of

PEMFs versus other non-pharmacological interventions on balance function, combining these

two may explain, in part, the conflicting results. Furthermore, our sensitivity analysis detecting

no changes in levels of heterogeneity and effect confirmed our results as relatively robust.

To overcome the shortcomings of the "statistically significant difference", the minimum

clinically important difference (MCID) defined as "the smallest change that is important to

patients" is employed to generate a threshold value for such change [42]. Any patient whose

responses help them reach the MCID threshold is considered as responders. Thus, a certain

proportion of responders to the total participants involved in a trialed intervention indicates the

likelihood of patients under the same condition also responding favorably to the same

intervention [43]. However, no definite consensus reached on the MCID of BMD and balance

function. Some evidence showed that changes by 2-5% at the lumbar and 8% at the proximal

femur [44], a point-drop in BBS associated with a 3-4% increase in risk of falling [45], and an

improvement of 2-3 seconds in TUG test were considered as MCID for the older population [46]. In

our study, no study included used the MCID and responder rate to evaluate the effect of PEMFs,

thus, it is hard to determine the clinical importance of improvements achieved by PEMFs on

BMD and balance function compared with placebo and exercise. Furthermore, the successful

treatment of osteoporosis is prevention of fractures, while no treatment can completely eliminate

fracture risk [10]. Although a certain increase in BMD and the improvement in balance function for

osteoporosis may result in a reduction in fracture risk, no risk of fracture was assessed and

reported in studies included in our review. Therefore, studies with reporting measures of MCID

and fracture risk in the future are required to further confirm that the effect of PEMFs in treating

osteoporosis is of clinical importance to clinicians and patients.

This review has several strengths. To date, we are the first to conduct a systematic review and

meta-analysis to explore the effect of PEMFs on BMD and balance function for primary

osteoporosis. Secondary, the results of our study were fairly stable due to studies included were

of appropriate methodological quality, and no significant heterogeneity was found in the analysis.

Furthermore, a key finding of this review was that PEMFs as an intervention alone achieved an

effect non-inferior to first-line treatment (pharmacological agents and exercise) on improvements

in BMD and balance function for older adults with primary osteoporosis. At last, a librarian

familiar with the development of searches and a mechanism of living systematic review were

involved over a course of 3 years to ensure no study was missed in compliance with the study

protocol.

There are several limitations to this review that deserve consideration. The primary limitation of

our review was the limited number of included studies with only 8 studies comprising of 396

participants for the analysis, from which the level of evidence generated were moderate to very

low. In some sub-group analyses, only 2-3 studies were included from which some uncertainty in

the results interpretation may exist based on data extracted. In addition, as the maximum PEMF

treatment session lasted for 16 weeks and the longest follow-up was 24 weeks for the analysis,

the error among the minimum percentage change in BMD and balance function may stay

undetectable and further deepened the uncertainty [44]. Furthermore, the clinical relevance of the

findings was limited, combined with no study reported data on MCID, responder rate and fracture

risk and only two studies [31,32] included used intention-to-treat (ITT) analysis. According to

accumulated evidence [11,47], the parameters of non-pharmacological interventions, including

intensity, frequency, and duration, were critical to impact changes in outcome measures for

osteoporosis, while our study could not conduct the sub-group analysis based on different

parameters due to limited information retrieved. Another potential limitation was the systematic

search was limited to English and Chinese manuscripts available in full text, and some relevant

trials may be missed.

CONCLUSION

In summary, moderate to very low certainty evidence showed that PEMFs as an intervention

alone has positive effects non-inferior to first-line treatment (pharmacological agents and exercise)

on BMD and balance function in older adults with primary osteoporosis and should be considered

as a promising option in the management of osteoporosis. Although uncertainty about responses

to the intervention and changes in outcome measures may be existed but undetectable, our

findings may still be fairly stable as we consistently found similar results in the primary and
sensitivity analyses. For further deeply confirming the effect of PEMFs for osteoporosis, adding
endpoints like fracture risk and events, and outcome measures like MCID and responder rate to
the core collection are necessary. In the future, researchers planning a PEMFs study should
optimize the study design, with taking factors not limited to undetectable errors under outcomes,
parameters of interventions, longer follow-up period, larger sample size and ITT analysis into
consideration, to generate high certainty evidence.

DECLARATIONS

Acknowledgements
We are grateful for the partnership and support from the library of Sichuan University for the
development of search strategy, also and the clinical research and statistics office of West China
Hospital for data processing and analysis.

Funding
This study was supported by the National Natural Science Foundation (81972146 to Chengqi He,
and 82002393 to Siyi Zhu), the Department of Science and Technology of Sichuan Province
(2020YJ0210 to Chengqi He, 2021YFS0004 and 2021YJ0424 to Siyi Zhu), China Postdoctoral
Science Foundation (2020M673251), Health Commission of Sichuan Province (20PJ034), and
West China Hospital of Sichuan University (2019HXBH058 to Siyi Zhu). The funders played no
role in the design, conduct, or reporting of this study.

Availability of Data and Materials
Not applicable

Authors' contributions
All authors have made substantial contributions to the study design. SYZ and YL have
contributed to the ongoing data collection. All listed authors have contributed to the drafting of
the manuscript.

Competing interests

The authors declare that they have no competing interests.

REFERENCES

1. Eastell R, O'Neill TW, Hofbauer LC, et al. Postmenopausal osteoporosis. *Nature reviews Disease primers.* 2016;2:16069.
2. International Osteoporosis Foundation. EPIDEMIOLOGY OF OSTEOPOROSIS AND FRAGILITY FRACTURES. https://www.osteoporosis.foundation/facts-statistics/epidemiology-of-osteoporosis-and-fragility-fractures. Published 2021. Accessed May 12, 2021.
3. Bessette L, Ste-Marie L-G, Jean S, et al. The care gap in diagnosis and treatment of women with a fragility fracture. *Osteoporosis International.* 2008;19(1):79-86.
4. International Osteoporosis Foundation. KEY STATISTICS FOR ASIA. https://www.osteoporosis.foundation/facts-statistics/key-statistic-for-asia. Published 2021. Accessed May 12, 2021.
5. US D. Bone health and osteoporosis: A Report of the Surgeon General. *http://www surgeongeneral gov/library/bonehealth/content html.* 2004.
6. Burge R, Dawson☐Hughes B, Solomon DH, Wong JB, King A, Tosteson A. Incidence and economic burden of osteoporosis☐related fractures in the United States, 2005–2025. *Journal of bone and mineral research.* 2007;22(3):465-475.
7. Kanis J, Johnell O. Requirements for DXA for the management of osteoporosis in Europe. *Osteoporosis international.* 2005;16(3):229-238.
8. Kanis J, Group WS. Assessment of Osteoporosis at the Primary Health Care Level. Geneva. *World Health Organization.* 2008.
9. Tu KN, Lie JD, Wan CKV, et al. Osteoporosis: a review of treatment options. *Pharmacy and Therapeutics.* 2018;43(2):92.
10. Camacho PM, Petak SM, Binkley N, et al. American Association of Clinical Endocrinologists/American College of Endocrinology clinical practice guidelines for the diagnosis and treatment of postmenopausal osteoporosis—2020 update. *Endocrine Practice.* 2020;26:1-46.
11. Zhu S, He H, Zhang C, et al. Effects of pulsed electromagnetic fields on postmenopausal osteoporosis. *Bioelectromagnetics.* 2017;38(6):406-424.
12. Petecchia L, Sbrana F, Utzeri R, et al. Electro-magnetic field promotes osteogenic differentiation of BM-hMSCs through a selective action on Ca 2+-related mechanisms. *Scientific reports.* 2015;5:13856.
13. Vincenzi F, Targa M, Corciulo C, et al. Pulsed electromagnetic fields increased the anti-inflammatory effect of A2A and A3 adenosine receptors in human T/C-28a2 chondrocytes and hFOB 1.19 osteoblasts. *PloS one.* 2013;8(5):e65561.
14. Andrew C, Bassett L, Pawluk RJ, Pilla AA. Augmentation of bone repair by inductively coupled electromagnetic fields. *Science.* 1974;184(4136):575-577.
15. Higgins JP, Thomas J, Chandler J, et al. *Cochrane handbook for systematic reviews of interventions.* John Wiley & Sons; 2019.
16. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71.
17. Siyi Z, Wei X, Hongchen H, et al. Pulsed electromagnetic fields (PEMFs) for preventing

428		and treating primary osteoporosis: systematic review and meta-analysis of randomized
429		controlled trials. PROSPERO 2018 CRD42018099518.
430		https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42018099518.
431		Published 2021. Accessed May 16, 2021.
432	18.	Cochrane Library. Living systematic reviews. https://community.cochrane.org/review-
433		production/production-resources/living-systematic-
434		reviews#:~:text=Living%20Evidence%20Network-
435		,What%20is%20a%20living%20systematic%20review%3F,the%20evidence%20(i.e.%20
436		monthly%20searches). Published 2021. Accessed May 16, 2021.
437	19.	Sherrington C, Herbert R, Maher C, Moseley A. PEDro. A database of randomized trials
438		and systematic reviews in physiotherapy. *Manual therapy.* 2000;5(4):223-226.
439	20.	Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro
440		scale for rating quality of randomized controlled trials. *Physical therapy.* 2003;83(8):713-
441		721.
442	21.	de Morton NA. The PEDro scale is a valid measure of the methodological quality of
443		clinical trials: a demographic study. *Australian Journal of Physiotherapy.*
444		2009;55(2):129-133.
445	22.	da Costa BR, Hilfiker R, Egger M. PEDro's bias: summary quality scores should not be
446		used in meta-analysis. *Journal of clinical epidemiology.* 2013;66(1):75-77.
447	23.	Group GW. Grading quality of evidence and strength of recommendations. *BMJ: British*
448		*Medical Journal.* 2004;328(7454):1490.
449	24.	Soen S, Fukunaga M, Sugimoto T, et al. Diagnostic criteria for primary osteoporosis: year
450		2012 revision. *Journal of bone and mineral metabolism.* 2013;31(3):247-257.
451	25.	Glaser DL, Kaplan FS. Osteoporosis: definition and clinical presentation. *Spine.*
452		1997;22(24):12S-16S.
453	26.	Higgins JP, Li T, Deeks JJ. Choosing effect measures and computing estimates of effect.
454		*Cochrane handbook for systematic reviews of interventions.* 2019:143-176.
455	27.	Sedgwick P. Meta-analyses: what is heterogeneity? *Bmj.* 2015;350:h1435.
456	28.	Abdelaal AA, Taha MM, Amin DI, Draz AH. Effect of pulsed electromagnetic therapy
457		versus low-level laser therapy on bone mineral density in the elderly with primary
458		osteoporosis: a randomized, controlled trial. *Bulletin of Faculty of Physical Therapy.*
459		2017;22(1):34-39.
460	29.	Elsisi HFEM, Mousa GSM, ELdesoky MTM. Electromagnetic field versus circuit weight
461		training on bone mineral density in elderly women. *Clinical interventions in aging.*
462		2015;10:539.
463	30.	Giordano N, Battisti E, Geraci S, et al. Effect of electromagnetic fields on bone mineral
464		density and biochemical markers of bone turnover in osteoporosis: a single-blind,
465		randomized pilot study. *Current Therapeutic Research.* 2001;62(3):187-193.
466	31.	Liu H, Liu Y, Yang L, Wang C, Wu Y, He C. Curative effects of pulsed electromagnetic
467		fields on postmenopausal osteoporosis. *Sheng wu yi xue gong cheng xue za zhi= Journal*
468		*of biomedical engineering= Shengwu yixue gongchengxue zazhi.* 2014;31(1):48-52.
469	32.	Liu HF, Yang L, He HC, et al. Pulsed electromagnetic fields on postmenopausal
470		osteoporosis in southwest China: A randomized, active☐controlled clinical trial.
471		*Bioelectromagnetics.* 2013;34(4):323-332.
472	33.	Shanb AA, Youssef EF, Muaidi QI, Alothman AA. Whole body vibration versus
473		magnetic therapy on bone mineral density in elderly osteoporotic individuals. *Journal of*
474		*back and musculoskeletal rehabilitation.* 2017;30(4):903-912.

475   34.   Shanb A-SA, Youssef EF, El-Barkouky MG, Kamal RM, Tawfick AM. The effect of
476   magnetic therapy and active exercise on bone mineral density in elderly women with
477   osteoporosis. *Journal of Musculoskeletal Research.* 2012;15(03):1250016.

478   35.   Wu Y-C, Xu J, Yang L, Xiong E-F, Xie W, He C-Q. The effect of low-frequency pulsed
479   electromagnetic fields on balance ability of patients with post-menopausal osteoporosis.
480   *Sichuan da xue xue bao Yi xue ban= Journal of Sichuan University Medical science*
481   *edition.* 2014;45(1):116-119.

482   36.   Zhu L, Wu W, Chen M, et al. Effects of Nonpharmacological Interventions on Balance
483   Function in Patients with Osteoporosis or Osteopenia: A Network Meta-Analysis of
484   Randomized Controlled Trials. *Evidence-Based Complementary and Alternative*
485   *Medicine.* 2021;2021.

486   37.   Wang R, Wu H, Yang Y, Song M. Effects of electromagnetic fields on osteoporosis: a
487   systematic literature review. *Electromagnetic biology and medicine.* 2016;35(4):384-390.

488   38.   Wang T, Yang L, Jiang J, et al. Pulsed electromagnetic fields: promising treatment for
489   osteoporosis. *Osteoporosis International.* 2019;30(2):267-276.

490   39.   Blake GM, Fogelman I. The role of DXA bone density scans in the diagnosis and
491   treatment of osteoporosis. *Postgraduate medical journal.* 2007;83(982):509-517.

492   40.   Kanis JA, Melton III LJ, Christiansen C, Johnston CC, Khaltaev N. The diagnosis of
493   osteoporosis. *Journal of bone and mineral research.* 1994;9(8):1137-1141.

494   41.   Choi M, Hector M. Effectiveness of intervention programs in preventing falls: a
495   systematic review of recent 10 years and meta-analysis. *Journal of the American Medical*
496   *Directors Association.* 2012;13(2):188. e113-188. e121.

497   42.   Stratford PW, Binkley JM, Riddle DL, Guyatt GH. Sensitivity to change of the Roland-
498   Morris back pain questionnaire: part 1. *Physical therapy.* 1998;78(11):1186-1196.

499   43.   Farrar JT, Portenoy RK, Berlin JA, Kinman JL, Strom BL. Defining the clinically
500   important difference in pain outcome measures. *Pain.* 2000;88(3):287-294.

501   44.   Cranney A, Welch V, Wells G, et al. Discrimination of changes in osteoporosis outcomes.
502   *The Journal of rheumatology.* 2001;28(2):413-421.

503   45.   Shumway-Cook A, Baldwin M, Polissar NL, Gruber W. Predicting the probability for
504   falls in community-dwelling older adults. *Physical therapy.* 1997;77(8):812-819.

505   46.   Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for
506   frail elderly persons. *Journal of the American geriatrics Society.* 1991;39(2):142-148.

507   47.   Bonaiuti D, Shea B, Iovine R, et al. Exercise for preventing and treating osteoporosis in
508   postmenopausal women. *Cochrane Database of Systematic Reviews.* 2002(2).

509
510

511 Box of inclusion criteria for eligible studies
512
513 **Box 1.** Inclusion criteria
514

Design
- Randomised or quasi-randomised controlled trial
- Published in a peer-reviewed journal
- Full text available in Chinese and English

Participants
- Healthy older adults (including those with previous fractures) aged over 50 years with primary osteoporosis

Intervention
- Electromagnetic fields with pulsed signal and extremely low frequencies (between 5 and 300 Hz)

Outcome measures
- Primary outcomes: bone mass, number of incident fractures
- Secondary outcomes: Functional assessments, quality of life, adverse events

Comparisons
- PEMFs versus sham/nothing
- PEMFs versus placebo/pharmacological agents
- PEMFs versus exercise/other physical agent or intervention
- PEMFs plus other intervention versus other intervention

515
516
517

Table 1
PEDro scores of included studies.

Study	Eligibility criteria and source	Random allocation	Concealed allocation	Baseline comparability	Blind subjects	Blind therapists Blind	Blind assessors	Adequate follow-up	Intention-to-treat analysis	Between-group comparisons	Point estimates and variability	Total score (0 to 10)
Liu Hf et al, 2013[32]	Y	Y	Y	Y	N	N	Y	Y	Y	Y	Y	9
Liu Hf et al, 2014[31]	Y	Y	Y	Y	N	N	N	Y	Y	Y	Y	8
Shanb et al, 2017[33]	Y	Y	Y	Y	N	N	Y	Y	N	Y	Y	8
Shanb et al, 2012[34]	Y	Y	N	N	N	N	N	Y	N	Y	Y	5
Elsisi et al, 2015[29]	Y	Y	Y	N	Y	Y	Y	Y	N	Y	Y	9
Giordano et al, 2001[30]	Y	Y	N	Y	N	N	Y	Y	N	Y	Y	7
Wu Yc et al, 2014[35]	Y	Y	N	Y	N	N	N	Y	N	Y	Y	6
Ashraf et al, 2017[28]	Y	Y	Y	Y	N	N	Y	Y	N	Y	Y	8

Table 2
Characteristics of the included studies.

Study	Participant				Intervention		Outcome measures			Follow-up
	Sample size	Study arms	Age (y)	Male/Female	Type of intervetion	Dose per protocol[a]	Primary	Secondary	Efficacy/Effect	
Liu Hf et al, 2013[32]	41	1. PEMFs (n=20) 2. AL (n=21)	61.70 (1.34) 63.14 (4.32)	0/41	PEMFs	3.82 mT, 8 Hz, 40min/session for a total of 30 sessions.	BMDL	BMDF; 25(OH)D; LE MMT; BBS	Effective as alendronate.	24 weeks
Liu Hf et al, 2014[31]	84	1. PEMFs (n=44) 2. AL (n=40)	62.41 (5.47) 62.63 (4.54)	0/84	PEMFs	3.8 mT, 8-12 Hz, 40min/session for a total of 30 sessions.	BMDL; BMDF	VAS; TUG; BBS	Effective as alendronate.	72 weeks
Shanb et al, 2017[33]	68	1. PEMFs (n=25) 2. WBV (n=25) 3. AL (n=18)	63.9 (3.9) 64.1 (4.4) 64.5 (4.03)	23/45	PEMFs	5 mT, 33 Hz, 50 min/session, 2 sessions/week for 16 weeks.	BMDL; BMDF	Serum calcium and vitamin D	Effective in conjunction with pharmacological treatment.	N/A
Shanb et al, 2012[34]	30	1. PEMFs (n=15) 2. Exercise (n=15)	60-70	0/30	PEMFs	5 mT, 33 Hz, 50 min/session, 3 sessions/week for 12 weeks.	BMDL; BMDF	N/A	Effective as exercise.	N/A
Elsisi et al, 2015[29]	30	1. PEMFs (n=15) 2. CRT-Exercise (n=15)	64.73 (3.08) 65.13 (2.44)	0/30	PEMFs	5 mT, 33 Hz, 30 min/session, 3 sessions/week for 12 weeks.	BMDL; BMDF	N/A	More effective than CRT exercise.	N/A
Giordano et al, 2001[30]	40	1. PEMFs (n=20) 2. Placebo (n=20)	56.3 (4.0) 55.9 (3.1)	0/40	PEMFs	1.2 mT, 100 Hz, 60 min/session, 3 sessions/week for 12 weeks.	BMDL; BMDF	Serum and urinary calcium, phosphate; Serum ALP, osteocaltin and PICP; Urinary hydroxyproline	Not effective in increasing BMD, but effective in stimulating osteogenesis.	1 month
Wu Yc et al, 2014[35]	43	1. PEMFs (n=24) 2. AL (n=19)	59.08 (4.65) 59.53 (5.40)	0/43	PEMFs	3.8 mT, 8-12 Hz, 40min/session for a total of 30 sessions.	TUG; BBS; Balance function	N/A	Effective as alendronate.	N/A

Ashraf et al, 2017[28]	60	1. PEMFs (n=20)	59.85 (2.35)	26/34	PEMFs	4 mT, 33 Hz, 30 min/session, 3 sessions/week for 12 weeks.	BMDL	N/A	More effective than LLLT and AL group.	N/A
		2. LLLT (n=20)	60.20 (2.17)							
		3. Med (n=20)	60.00 (2.62)							

[a]Waveforms; Intensity; Frequency; Duration (weeks).

[b]Abbreviations. PEMFs=Pulsed electromagnetic fields, AL=alendronate, WBV=Whole body vibration, CRT=Circuit weight training, LLLT=low-level laser therapy, mT=millitesla, Hz=Hertz, BMDL=lumbar bone mineral density, BMDF=femur bone mineral density, 25(OH)D=serum 25OH vitamin, LE MMT=total lower-extremity manual muscle test, BBS=Berg Balance Scale, VAS=Visual Analogue Scale, N/A=not applicable, TUG=Timed Up and Go, ALP=Alkaline Phosphatase, PICP=Procollagen type I carboxy-terminal propeptide, Med=Medication.

Figure 1. PRISMA flow of studies through the review.

Figure 2. Forest plot analysis of the effects of PEMFs on BMD at the lumbar (A) and femur neck (B) compared with placebo control. Data are presented as mean difference (MD) between treatment and control groups with a 95% confidence interval (CI). SD = Standard Deviation; BMD = Bone Mineral Density; PEMFs = Pulsed Electromagnetic Fields.

Figure 3. Forest plot analysis of the effects of PEMFs on balance function measured by BBS (A) and TUG test (B) compared with placebo control. Data are presented as mean difference (MD) between treatment and control groups with a 95% confidence interval (CI). SD = Standard Deviation; BBS = Berg Balance Scale; TUG = Timed Up and Go; PEMFs = Pulsed Electromagnetic Fields.

Figure 4. Forest plot analysis of the effects of PEMFs on BMD at the lumbar (A) and femur neck (B) compared with exercise group. Data are presented as mean difference (MD) between treatment and control groups with a 95% confidence interval (CI). SD = Standard Deviation; BMD = Bone Mineral Density; PEMFs = Pulsed Electromagnetic Fields.