Virologic features of SARS-CoV-2 infection in children

Lael M. Yonker[1,2,3*], Julie Boucau[4*], James Regan[5], Manish C. Choudhary[3,5], Madeleine D.

Burns[1], Nicola Young[1], Eva J. Farkas[1], Jameson P. Davis[1], Peter P. Moschovis[2,3], T. Bernard

Kinane[2,3], Alessio Fasano[1,2,3], Anne M. Neilan[2,3,6], Jonathan Z. Li[3,5*], Amy K. Barczak[3,4,6*]

1. Massachusetts General Hospital, Mucosal Immunology and Biology Research Center,

Boston, MA, USA

2. Massachusetts General Hospital, Department of Pediatrics, Boston, MA, USA

3. Harvard Medical School, Boston, MA, USA

4. Ragon Institute of MGH, MIT and Harvard, Cambridge, MA, USA

5. Brigham and Women's Hospital, Department of Medicine, Boston, MA, USA

6. Massachusetts General Hospital, Department of Medicine, Boston, MA, USA

*Authors contributed equally

Correspondence:

Lael Yonker: Massachusetts General Hospital, 55 Fruit St, Jackson 14, Boston, MA 02114.

617-724-2890. Lyonker@mgh.harvard.edu

Word count abstract: 200

Word count main text: 2894

Abstract

Background: Data on pediatric COVID-19 has lagged behind adults throughout the pandemic.

An understanding of SARS-CoV-2 viral dynamics in children would enable data-driven public

health guidance.

Methods: Respiratory swabs were collected from children with COVID-19. Viral load was

quantified by RT-PCR; viral culture was assessed by direct observation of cytopathic effects and

semiquantitative viral titers. Correlations with age, symptom duration, and disease severity were

analyzed. SARS-CoV-2 whole genome sequences were compared with contemporaneous

sequences.

Results: 110 children with COVID-19 (median age 10 years, range 2 weeks-21 years) were

included in this study. Age did not impact SARS-CoV-2 viral load. Children were most infectious

within the first five days of illness, and severe disease did not correlate with increased viral

loads. Pediatric SARS-CoV-2 sequences were representative of those in the community and

novel variants were identified.

Conclusions: Symptomatic and asymptomatic children can carry high quantities of live,

replicating SARS-CoV-2, creating a potential reservoir for transmission and evolution of genetic

variants. As guidance around social distancing and masking evolves following vaccine uptake in

older populations, a clear understanding of SARS-CoV-2 infection dynamics in children is critical

for rational development of public health policies and vaccination strategies to mitigate the

impact of COVID-19.

Keywords: SARS-CoV-2, Pediatric COVID-19, Viral dynamics

Background

Since the SARS-CoV-2 virus ignited the COVID-19 global pandemic, the impact of the virus on children and the role that children play in this pandemic has been understudied. Initially, epidemiology reports suggested that children may have been relatively spared from infection, however, as COVID-19 testing became more available, it has been increasingly recognized that children can be infected with SARS-CoV-2 at rates comparable to adults [1, 2]. To date, over 4.1 million children in the Unites States have been reported as testing positive for COVID-19 [3]. Since the winter of 2020-2021, children under 19 years of age have represented one of the age groups with the highest rates of infection [4], which likely reflects a combination of increased number of infections among children plus increased vaccination rates amongst adults. Most children generally have milder symptoms when infected with SARS-CoV-2 [5], although a small subset of individuals develop severe disease. In the US, over 16,000 children have been hospitalized for acute COVID-19 with over 300 deaths reported[3]. A baseline understanding of the viral characteristics of SARS-CoV-2 infection in children is a necessary prerequisite to understanding the pathogenesis of severe presentations of COVID-19 [6].

At a population perspective, the role that children play in viral transmission remains poorly understood. Epidemiologic studies suggest that children exhibit lower transmission rates than adults [7], however, these findings are potentially confounded by higher rates of asymptomatic or pauci-symptomatic infection in children, increased social isolation by children early in the pandemic, and reduced COVID-19 testing in children. To date, one small study demonstrated that live virus can be cultured from children [8]. However, the types of systematic studies that have informed our understanding of the viral dynamics of SARS-CoV-2 in adult populations [9-11] have not similarly been carried out in children. As vaccination has become available for adults and adolescents in many places in the world and our understanding of transmission dynamics have evolved, masking and distancing policies are being relaxed[12]. Policy changes

have necessarily been made despite the paucity of data providing insight into the role that

pediatric disease might play in ongoing transmission. As viral variants that enhance the potential

for transmission and/or reduce vaccine efficacy emerge [13-15], the importance of identifying

potential reservoirs of viral replication and transmission has been brought into the spotlight.

Defining the virologic features of SARS-CoV-2 infection in children and the potential for children

to transmit virus will facilitate rational public health decision-making for pediatric populations.

In this work, we sought to define fundamental virologic features of SARS-CoV-2 in a pediatric

population across a range of disease severity. We analyzed respiratory swabs from children

presenting to urgent care clinics or the hospital with symptomatic and asymptomatic COVID-19

infection. Clinical factors, such as age, COVID-19 risk factors, and disease severity were

compared with viral features including SARS-CoV-2 viral load, isolation of replication-competent

virus, and whole viral sequencing. Our data indicate that age, from infancy through adulthood, is

not a predictor of viral infection dynamics, and that children of all ages can have high SARS-

CoV-2 viral loads of replication-competent virus, including variants, displaying comparable

dynamics to those seen in adults.

Methods

Sample collection

Infants, children and adolescents $\leq$21 years of age presenting to Massachusetts General

Hospital urgent care clinics or the hospital with either symptoms concerning for or known

exposure to COVID-19 (4/2020-4/2021) were prospectively offered enrollment in the Institutional

Review Board-approved MGH Pediatric COVID-19 Biorepository (IRB # 2020P000955) [16].

After informed consent, and assent when appropriate, was obtained verbally, a research-

designated swab of the nasopharynx, oropharynx and/or anterior nares was obtained and

placed in phosphate buffered saline. Samples were aliquoted and stored at -80°C. Samples

from patients who tested positive for COVID-19 on clinical SARS-CoV-2 RT-PCR testing were

analyzed. Nasal samples from adults hospitalized with acute COVID-19 [10] (4/2020-8/2020;

enrolled in Institutional Review Board-approved MGH COVID-19 Biorepository, IRB #

2020P000804) with duration of symptoms equal to the hospitalized pediatric cohort were

selected for comparative studies.

Clinical data collection

Demographic and clinical factors were recorded through a combination of manual chart reviews

and data extraction from electronic health records (EHR), then collected in a REDCap database

[17] through the Partners Electronic Health Record Registry of Pediatric COVID-19 Disease

(IRB # 2020P003588). Trained reviewers collected demographics, SARS-CoV-2 risk factors,

comorbid conditions, medications, COVID-19 related symptoms, and laboratory tests. Outcome

of initial presentation to care, admission status, and complications of COVID-19 disease were

also extracted by manual review.

SARS-CoV-2 viral load quantification

Virions were pelleted from anterior nasal, oropharyngeal, and nasopharyngeal swab fluids by

centrifugation at approximately 21,000 x g for 2 hours at 4°C. RNA was extracted using Trizol-

LS (Thermofisher) according to the manufacturer's instructions. RNA was then concentrated by

isopropanol precipitation, and SARS-CoV-2 RNA was quantified using the CDC N1 primers and

probe [https://www.cdc.gov/coronavirus/2019-ncov/lab/rt-pcr-panel-primer-probes.html] as

previously described [10]. As there was no significant difference in viral load from respiratory

secretions obtained from the anterior nares, nasopharynx or oropharynx of participants

(**Supplemental Figure 1**), samples were analyzed together, regardless of collection site.

Viral Culture

Vero-E6 cells (ATCC) were maintained in D10+ media [Dulbecco's modified Eagle's media

(DMEM) (Corning) supplemented with HEPES (Corning), 1X Penicillin/Streptomycin (Corning),

1X Glutamine (Glutamax, ThermoFisher Scientific) and 10% Fetal Bovine serum (FBS) (Sigma)]

in a humidified incubator at 37°C in 5% CO2. Vero-E6 cells were passaged every 3-4 days,

detached using Trypsin-EDTA (Fisher Scientific) and seeded at 150,000 cells per wells in 24

well plates for culture experiments and 20,000 cells per well in 96w plates the day before

inoculation for median tissue culture infectious dose (TCID50) experiments.

After thawing, each specimen was filtered through a Spin-X 0.45µm filter (Corning) at 10,000 x g

for 5min. 50µL of the supernatant was then diluted in 450µL of D^+ media [DMEM supplemented

with HEPES, 1X Penicillin/Streptomycin and 1X Glutamine]. The viral culture experiments were

performed as previously reported [18] with the following modifications: 100µL of the solution was

used to inoculate wells in a 24 well plate and 1mL of D_2^+ media [D+ media with 2% FBS] was

added to each well after 1 hour of incubation. The plates were then placed in a 5% CO2

incubator at 37°C. For TCID50 measurements conducted in parallel, 25µL of the Spin-X flow-

through was used to inoculate Vero-E6 cells in a 96 well plate in the presence of 5µg/mL of

polybrene (Santa Cruz Biotechnology) using 5-fold dilutions (5^{-1} to 5^{-6}) and 4 repeats for each

sample. The plates were centrifuged for 1 hour at 2,000 x g at 37°C before being placed in a 5%

CO2 incubator at 37°C. The SARS-CoV-2 isolate USA-WA1/2020 strain (BEI Resources) was

used as a positive control for CPE in both culture and TCID50 experiments.

Viral culture and TCID50 plates were observed at 3- and 6-days post-infection with a light

microscope and wells showing CPE were counted. The TCID50 titers were calculated using the

Spearman-Karber method. For the culture plates, the supernatant of the wells displaying CPE

was harvested 10-14 days post-infection and RNA was isolated using a QIAamp Viral RNA Mini

kit (QIAGEN) for confirmation of the viral sequence.

SARS-CoV-2 sequencing

cDNA synthesis was performed using Superscript IV reverse transcriptase (Invitrogen). Whole

viral amplification was performed with the Artic protocol using multiplexed primer pools designed

with Primal Scheme generating 400-bp tiling amplicons [19, 20]. PCR products were pooled and

Illumina library construction was performed using the Nextera XT Library Prep Kit (Illumina). The

comparison dataset included 183 representative contemporaneous SARS-CoV-2 genomes from

Massachusetts present in GISAID to assess for local clustering. Nucleotide sequence alignment

was performed with MAFFT (multiple alignment using fast Fourier transform) [21]. Best-fit

nucleotide substitution GTR+G+I was used for the datasets using model selection in IQ-Tree

followed by maximum likelihood phylogenetic tree construction using IQ-Tree web server with

1000-bootstrap replicates [22].

Analysis

All statistical analyses were performed using parametric comparisons in GraphPad Prism

(Version 9.1.1), including Pearson correlation, ANOVA with multiple comparisons, and unpaired

t test.

Results

Clinical cohort

One-hundred-ten children diagnosed with COVID-19 with a mean age of 10 years (range 0-21

years) were included in the study (**Table 1**). There were slightly more boys (56%) than girls

(44%) with SARS-CoV-2 infection included in our analyses. One third of the participants were

White (33%), 10% African American/Black, and 4% were Asian; one third (38%) reported their

ethnicity as Hispanic. Past medical history in children is reported in **Supplemental Table 1**.

Thirty children were asymptomatic but were identified as having COVID-19: twenty-six children

(27%) presented to urgent care/COVID-19 testing sites because of a COVID-19 exposure, while

four children (4%) were identified on routine screening during hospital admission. Eight (7%)

presented with COVID-19 symptoms but had no known COVID-19 contact. The majority of

participants with COVID-19 did not require hospitalization (72 children, 65%). Thirty-six children

(33%) were hospitalized with COVID-19, although only 18 children (16%) required supplemental

oxygen and/or invasive or non-invasive respiratory support (referred to as "moderate/severe

COVID-19").

Age did not impact SARS-CoV-2 viral load or recovery of replication competent virus

Age is a well-established risk factor for developing severe COVID-19. Accordingly,

asymptomatic patients were significantly younger than patients with mild disease, and pediatric

patients who were hospitalized with hypoxemia were significantly older than asymptomatic

children or children with mild disease (**Figure 1A**). However, viral load was not increased in

more severe disease: asymptomatic children and children with mild disease displayed

significantly higher viral loads than adults hospitalized with COVID-19 with comparable duration

of symptoms (less than 10 days) (**Figure 1B**). However, there were no differences in viral load

between pediatric patients hospitalized with moderate/severe disease and hospitalized adults of

similar duration of illness (**Figure 1B**) (Adult demographics are detailed in **Supplemental Table**

2).

The age of each infected child was analyzed to determine whether age impacted viral load.

There was no significant correlation of age with viral load (**Figure 1C**), nor were there significant

differences between ages when grouped by school levels: 0-4 years (infant through pre-school),

5-10 years (elementary school), 11-16 years (middle school), 17 and older (high school and

higher education) (ANOVA, P = 0.12) (**Figure 1D**). Thus, a child's age did not appear to impact

viral load: all children, from 2 weeks through 21 years of age, were equally capable of carrying a

high viral load.

As SARS-CoV-2 RNA detection by RT-PCR does not specify whether replication-competent

virus is being shed, we next sought to ascertain risk factors for shedding live virus by performing

viral culture assays for recoverable SARS-CoV-2 in parallel with viral load testing. From the 110

participants, we collected 126 samples; live virus was cultured from 33 samples coming from 32

participants. Of note, eight of these children with culturable SARS-CoV-2 were asymptomatic.

Higher viral load was significantly predictive of shedding of live virus (t test, P < 0.0001) (**Figure**

2A). Consistent with the results for viral load, age was not correlated with viral culture results;

virus was recovered from children ages 1 month through 21 years (**Figure 2B**). Semi-

quantitative assessment of the amount of virus shed by an individual participant was assessed

by median tissue culture infectious dose (TCID50). TCID50 for culture-positive specimens

correlated strongly with viral load (Pearson correlation r = 0.7, P < 0.0001) (**Figure 2C**) but did

not correlate with age across all pediatric participants (**Figure 2D**).

Children with COVID-19 were most infectious within first five days of illness

To define the likely period of infectiousness in our pediatric population, we analyzed viral load,

culturability, and TCID50 in comparison with duration of symptoms. Of note, duration of

symptoms does not necessarily indicate duration of infection, as time infection was acquired

cannot be confirmed. Consistent with prior reports in adults [9], viral loads in children were the

highest earliest in the course of illness and declined over time after symptom onset (Pearson, r

= -0.4, P <0.001) (**Figure 3A**). Viral load was highest in the first two days of symptoms, with

significant decrease after 5 days of symptoms and further decline after 10 days of symptoms (P

< 0.0001) (**Figure 3B**). Analysis of pediatric viral culture results demonstrated that children

tested early after symptom onset were more likely to shed replication competent virus (P =

0.004) (**Figure 3C)**. Correspondingly, semi-quantitative assessment of infectious viral shedding

in children showed that the TCID50 was higher early after symptom onset and decreased over

time. When grouped by days of symptoms, children in days 0-2 of their symptoms had the

highest infectivity, while children with greater than six days of illness shed less virus (P = 0.004)

(**Figure 3D**).

We then sought to assess whether COVID-19 severity impacted the relationship between viral

load and age in pediatric cohorts of varying severity: asymptomatic, mildly symptomatic, and

moderate/severe pediatric COVID-19 patients. None of these COVID-19 severity groups

revealed any correlation of age with viral load (**Figure 4A**). Further, there were no differences in

viral clearance over the duration of illness, not only when comparing mild pediatric COVID-19

with moderate/severe COVID-19, but also when comparing pediatric COVID-19 with adults

hospitalized with COVID-19 (**Figure 4B**). Duration of symptoms did not affect the finding that

viral load does not correlate with age of the infected child (**Supplemental Figure 2**).

Pediatric SARS-CoV-2 sequences were representative of those found in the community

We successfully performed whole-viral sequencing of 57 respiratory samples from 54 children.

Phylogenetic analysis of these pediatric sequences with contemporaneous Massachusetts

sequences from GISAID showed that they were representative of the spectrum of sequences

found in the community (**Figure 5**). Notable variants identified in the pediatric samples included

four Alpha (B.1.1.7) and three Iota (B.1.526.2) variants. To validate our culture results on a

subset of culture-positive samples, we sequenced virus isolated from the supernatant from 8

positive samples. Sequences from the supernatant and respiratory specimens were identical in

7 cases and demonstrated only 1 nucleotide change in the last case.

Discussion

As the global COVID-19 pandemic took hold, infected older adults suffered high rates of hospitalization and death while infected children typically experienced paucisymptomatic or asymptomatic infection. While it is now clear that children can become infected with and transmit SARS-CoV-2, viral dynamics in children have been understudied, and a full understanding of the dynamics of infection in children is needed to inform public health policies specific to the pediatric population. Here, we show that pediatric patients of all ages, from infancy to young adulthood, can carry a high SARS-CoV-2 viral load in their upper airways, particularly early in the course of infection, and an elevated viral load corresponds with high levels of viable, replicating virus. Pediatric sequences were largely reflective of those found in the general community and the presence of novel variants was identified.

Our findings have significant implications for both public health policy and the potential role of universal vaccination of pediatric populations in fully curbing the COVID-19 pandemic. As vaccination has rolled out in adult populations, public health policies are being adjusted to account for changes in risk that result from vaccination. Our results emphasize the importance of considering and clarifying how these policy changes relate to children. As adult populations have been vaccinated, pediatric cases have represented a growing proportion of infections, currently accounting for up to 25% of all COVID-19 cases across different regions of the United States[3]. Our results suggest that the low rates of transmission in settings such as schools and daycares cannot be attributed to low viral loads, low rates of viral shedding, or rapid clearance of virus in younger patient populations. As changes in masking and distancing policies are implemented for vaccinated adults, consideration of how and whether policies changes will be applied for children will be critical for ongoing reduction of new COVID-19 cases.

Our results additionally suggest that pediatric populations have the potential to serve as a community reservoir of actively replicating virus, with implications for both new waves of

infection and the evolution of viral variants. The duration of natural and vaccine-induced

immunity for each vaccine in clinical use are not yet known. If a community reservoir of actively

replicating virus is maintained and transmitted within unvaccinated pediatric populations, that

population could then serve as a source of new infections as vaccine-induced immunity wanes

in vaccinated adult populations. In addition, viral genomic variants were readily identified in the

pediatric samples and these variants have the potential to impact viral transmission [23-25],

disease severity [26, 27], and vaccine efficacy [28]. Ongoing viral replication within pediatric

populations has the potential to serve as a source of existing and new viral variants that

interfere with eradication efforts.

Our study has several limitations. First, the data collected here represent a single medical

center and affiliated pediatric urgent care/COVID-19 screening clinics. However, these were

amongst the few pediatric testing centers encompassing a large catchment area during the

duration of this study, and patients enrolled spanned a wide range of symptoms. Additionally,

many of these samples were collected early in the pandemic and SARS-CoV-2 variants of

interest have shifted over time. Ongoing studies analyzing shifts in virologic features of SARS-

CoV-2 infection in children alongside studies of infection in adults are needed to better

understand the full reach of the COVID-19 pandemic.

Ultimately, our data suggest that although age is generally protective against severe disease,

children, especially early in the infection course, carry high viral loads of SARS-CoV-2, which

can include viral variants. Our results underline the importance of defining public health policy

with viral dynamics in children in mind and of including pediatric populations in vaccine efforts

aimed at eradication.

Acknowledgements:

The viral culture work was performed in the Ragon Institute BSL3 core, which is supported in

part by the NIH-funded Harvard University Center for AIDS Research (P30 AI060354). This

research was supported by the National Heart, Lung, and Blood Institute (5K08HL143183 to

LMY), the Department of Pediatrics at Massachusetts General Hospital *for* Children (to LMY),

the Massachusetts Consortium for Pathogen Readiness and a gift from Mark and Lisa Schwartz

(to JZL).

Conflicts of Interest

The authors do not report any conflicts of interest.

1. Mehta NS, Mytton OT, Mullins EWS, et al. SARS-CoV-2 (COVID-19): What Do We Know About
Children? A Systematic Review. Clin Infect Dis **2020**; 71:2469-79.
2. Heald-Sargent T, Muller WJ, Zheng X, Rippe J, Patel AB, Kociolek LK. Age-Related Differences
in Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Levels in
Patients With Mild to Moderate Coronavirus Disease 2019 (COVID-19). JAMA Pediatr **2020**;
174:902-3.
3. https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/children-and-
covid-19-state-level-data-report/.
4. https://www.mass.gov/info-details/covid-19-response-reporting.
5. Yonker LM, Neilan AM, Bartsch Y, et al. Pediatric Severe Acute Respiratory Syndrome
Coronavirus 2 (SARS-CoV-2): Clinical Presentation, Infectivity, and Immune Responses. J Pediatr
2020; 227:45-52 e5.
6. Fialkowski A, Gernez Y, Arya P, Weinacht KG, Kinane TB, Yonker LM. Insight into the pediatric
and adult dichotomy of COVID-19: Age-related differences in the immune response to SARS-
CoV-2 infection. Pediatr Pulmonol **2020**.
7. Soriano-Arandes A, Gatell A, Serrano P, et al. Household SARS-CoV-2 transmission and
children: a network prospective study. Clin Infect Dis **2021**.
8. L'Huillier AG, Torriani G, Pigny F, Kaiser L, Eckerle I. Culture-Competent SARS-CoV-2 in
Nasopharynx of Symptomatic Neonates, Children, and Adolescents. Emerg Infect Dis **2020**;
26:2494-7.
9. Weiss A, Jellingso M, Sommer MOA. Spatial and temporal dynamics of SARS-CoV-2 in COVID-
19 patients: A systematic review and meta-analysis. EBioMedicine **2020**; 58:102916.
10. Fajnzylber J, Regan J, Coxen K, et al. SARS-CoV-2 viral load is associated with increased
disease severity and mortality. Nat Commun **2020**; 11:5493.
11. Wolfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients
with COVID-2019. Nature **2020**; 581:465-9.
12. https://www.cdc.gov/coronavirus/2019-ncov/index.html.
13. Madhi SA, Baillie V, Cutland CL, et al. Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine
against the B.1.351 Variant. N Engl J Med **2021**.
14. Liu Y, Liu J, Xia H, et al. BNT162b2-Elicited Neutralization against New SARS-CoV-2 Spike
Variants. N Engl J Med **2021**.
15. Shinde V, Bhikha S, Hoosain Z, et al. Efficacy of NVX-CoV2373 Covid-19 Vaccine against the
B.1.351 Variant. N Engl J Med **2021**.
16. Lima R, Gootkind EF, De la Flor D, et al. Establishment of a pediatric COVID-19 biorepository:
unique considerations and opportunities for studying the impact of the COVID-19 pandemic on
children. BMC Med Res Methodol **2020**; 20:228.
17. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data
capture (REDCap)--a metadata-driven methodology and workflow process for providing
translational research informatics support. J Biomed Inform **2009**; 42:377-81.
18. Choi B, Choudhary MC, Regan J, et al. Persistence and Evolution of SARS-CoV-2 in an
Immunocompromised Host. N Engl J Med **2020**; 383:2291-3.

19. Gonzalez-Reiche AS, Hernandez MM, Sullivan MJ, et al. Introductions and early spread of
SARS-CoV-2 in the New York City area. Science **2020**; 369:297-301.
20. Quick J. nCoV-2019 sequencing protocol v1. Available at:
https://www.protocols.io/view/ncov-2019-sequencing-protocol-bbmuik6w.
21. Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence alignment,
interactive sequence choice and visualization. Brief Bioinform **2019**; 20:1160-6.
22. Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ. W-IQ-TREE: a fast online phylogenetic
tool for maximum likelihood analysis. Nucleic Acids Res **2016**; 44:W232-5.
23. Zhou B, Thao TTN, Hoffmann D, et al. SARS-CoV-2 spike D614G change enhances replication
and transmission. Nature **2021**; 592:122-7.
24. Deng X, Garcia-Knight MA, Khalid MM, et al. Transmission, infectivity, and neutralization of
a spike L452R SARS-CoV-2 variant. Cell **2021**.
25. Davies NG, Abbott S, Barnard RC, et al. Estimated transmissibility and impact of SARS-CoV-2
lineage B.1.1.7 in England. Science **2021**; 372.
26. Davies NG, Jarvis CI, Group CC-W, et al. Increased mortality in community-tested cases of
SARS-CoV-2 lineage B.1.1.7. Nature **2021**; 593:270-4.
27. Funk T, Pharris A, Spiteri G, et al. Characteristics of SARS-CoV-2 variants of concern B.1.1.7,
B.1.351 or P.1: data from seven EU/EEA countries, weeks 38/2020 to 10/2021. Euro Surveill
2021; 26.
28. Garcia-Beltran WF, Lam EC, St Denis K, et al. Multiple SARS-CoV-2 variants escape
neutralization by vaccine-induced humoral immunity. Cell **2021**; 184:2372-83 e9.
379
380

381   **Table 1**: Participant Demographics, past medical history, reason for presenting for SARS-CoV-2

382   RT-PCR testing, and disease severity of children with COVID-19 (n=110).

	COVID+ children (N=110)
Age, average (max, min)	10 (0,21)
Sex, n (%)	
Male	62 (56)
Female	48 (44)
Race, n (%)	
White	36 (33)
Black	11 (10)
Asian	4 (4)
Other	45 (41)
Ethnicity, n (%)	
Hispanic	42 (38)
Past Medical History, n (%)	
Obesity	28 (25)
Asthma	11 (10)
Other	39 (35)
Reason for presenting for COVID-19 testing, n (%)	
Asymptomatic, exposure	30 (27)
Symptomatic, exposure	72 (65)
Symptomatic, no exposure	8 (7)
COVID-19 Severity, n (%)	
Hospitalized	36 (33)
Supplemental oxygen	18 (16)
Outpatient	75 (68)

Figure 1: COVID-19 disease severity and SARS-CoV-2 viral load across age groups. **A.** Age of

pediatric patients with SARS-CoV-2 infection, stratified by disease severity: asymptomatic

(n=27), mild disease, outpatient (n=48), moderate/severe COVID-19, hospitalized (n=18).

Analyzed by ordinary one-way ANOVA. **B.** SARS-CoV-2 viral load was quantified across a

range of disease severities. Patients presenting with $\leq$10 days of symptoms were compared,

including asymptomatic pediatric outpatients (n=27), mildly symptomatic pediatric outpatients

(n=44), moderate/severe pediatric hospitalized patients with oxygen requirement (n=18), and

moderate/severe adult hospitalized patients (n=29). Analyzed by ordinary one-way ANOVA. **C.**

Viral load for each specimen (n=110) was determined by qPCR, plotted against participant age

and analyzed by Pearson correlation. **D**. Viral load levels reported by school age group: 0-4

years old (yo)– infant through pre-school (n=32), 5-10yo – elementary school (n=23), 11-16yo –

middle school (n=33), 17yo and over – high school and higher education (n=22). Analyzed by

ordinary one-way ANOVA. Dotted lines depict limit of detection. * P<0.05, *** P<0.001, **** P<

0.0001, ns = not significant

Figure 2: SARS-CoV-2 culture results across age groups and viral load. **A-B**. Samples with

observable CPE (culture +) (n=31) or without observable CPE (culture -) (n=95) plotted against

viral load (**A**) and participant age (**B**) and compared using t test. **C-D.** Semiquantitative viral titer

expressed as TCID50/mL for culture positive samples plotted against corresponding viral load

(**C**) or participant age (**D**), Analyzed using Pearson correlation. Dotted line depicts limit of

detection. **** P < 0.0001

Figure 3: Culture positivity and duration of symptoms. **A**. Viral load for each specimen was

determined by qPCR and plotted against the duration of symptoms (in days). Analysis by

Pearson correlation. **B**. Viral load reported by binned duration of symptoms. Ordinary one-way

ANOVA used for analysis. **C**. Duration of symptoms for samples with observable CPE (culture

+, n=29) and without observable CPE (culture -, n=85). Analysis by t test. **D**. Semiquantitative

viral titer reported by binned duration of symptoms, analyzed by ordinary one-way ANOVA.

Dotted lines depict limit of detection. ** P < 0.01, **** P < 0.0001

Figure 4: Correlation of viral load with age and duration of illness based on disease severity.

A. Correlation of viral load and age, stratified by asymptomatic (n=30), mild outpatient (n=48)

and moderate/severe hospitalized (n=18) cohorts. **B.** Viral load of hospitalized adult (n=29),

hospitalized pediatric participants requiring respiratory support (n=18) and pediatric outpatients

with mild disease (n=48), plotted against duration of symptoms. Dotted lines depict limit of

detection.

Figure 5. Phylogenetic analysis of pediatric and community SARS-CoV-2 sequences. Maximum

likelihood tree generated from pediatric sequences (red) and 183 contemporaneous

Massachusetts sequences from GISAID.

Supplemental Table 1: Past medical history of children infected with SARS-CoV-2.
433

Past Medical History - Pediatric Patients
Acute recurrent pancreatitis
ADHD
Alopecia
Asthma
Autism
Cardiac conduction disorder
Chronic kidney disease
Coats Plus Syndrome
Crohn's disease
Cystic Fibrosis
Eczema
Epilepsy
G6PD Deficiency
Glutaric acidemia, type 1
Liver transplant
Mitochondrial complex 1 deficiency
Obesity
Obstructive sleep apnea
Prematurity
Sickle cell trait
Speech delay
Subglottic stenosis
Tracheobronchomalacia

436 **Supplemental Table 2**: Demographics, past medical history and disease severity of adults with

437 COVID-19 (n=29) included in analysis of SARS-CoV-2 viral load.

	COVID+ adults (n=29)
Age, average (max, min)	62 (26,93)
Sex, number (%)	
Male	13 (45)
Female	16 (55)
Race, number (%)	
White	20 (69)
Black	5 (17)
Asian	0 (0)
Other	4 (14)
Ethnicity, number (%)	
Hispanic	4 (14)
COVID-19 Severity	
Hospitalized, number (%)	21 (100)
Respiratory support	22 (78)
Outpatient, number (%)	0 (0)

Supplemental Figure 1: Viral load by sample collection location. Viral load of samples collected

from nasopharynx (n=60), anterior nares (n=47), or oropharynx (n=19) were compared and

analyzed by ordinary one-way ANOVA. ns = non-significant.

Supplemental Figure 2: Viral load plotted against pediatric patient age, for patients with 0-2

days symptoms (n=67), 3-5 days symptoms (n=30), and ≥6 days of symptoms (n=29). No

significant correlation in any grouping, when analyzed by Pearson correlation. Dotted lines

depict limit of detection. ns = not significant.

451