

1 **Incidence of delirium in an acute geriatric community hospital: an exploratory analysis of an**
2 **observational cohort study controlled with a meta-analysis of incidences from literature.**

3
4 Marthe E. Ribbink MD^{1*}, Emma Stornebrink, MD^{1*}, Remco Franssen MD, PhD¹, Annemarieke de
5 Jonghe MD, PhD², Janet L. MacNeil Vroomen PhD¹, Bianca M. Buurman RN, PhD^{1,3} on behalf of the
6 AGCH-study group.

7
8 ¹Amsterdam University Medical Centre, University of Amsterdam, Department of Internal Medicine,
9 Section of Geriatric Medicine, Amsterdam Public Health Research Institute, Amsterdam, The
10 Netherlands

11 ²Tergooi hospitals, Department of Geriatrics, The Netherlands

12 ³ACHIEVE – Centre of Applied Research, Faculty of Health, Amsterdam University of Applied
13 Sciences, Amsterdam, The Netherlands

14 *MER and ES share first authorship.

15
16 **Corresponding author**

17 M.E. Ribbink, MD

18 Amsterdam UMC, University of Amsterdam, Department of Internal Medicine, Section of Geriatric
19 Medicine, D3-335

20 PO Box 22600, 1100 DD Amsterdam, The Netherlands

21 Tel: 0031-20 5661647, email: m.e.ribbink@amsterdamumc.nl

22
23 **Keywords:** delirium, older adults, hospitalisation, community hospital, intermediate care.

24

25

26

27

28

29

30

31

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

32 **Declaration of Sources of Funding**

33 This work was supported by ZonMw, the Netherlands Organization for Health Research and
34 Development [grant number 808393598041] and the PVE- fund. These funders played no role in the
35 design or execution of this research. Moreover, the care provided at the AGCH is provided in a
36 partnership between Cordaan, a community and home-care organization and the Amsterdam
37 University Medical Centre, location Academic Medical Centre. The AGCH is financially supported by
38 Zilveren Kruis, a health insurance company. Zilveren Kruis played a role in the initiation and design,
39 but no role in the execution, analysis and writing of this research.

40

41 **Conflicts of interest**

42 All authors declare no conflicts of interest.

43

44 **Acknowledgements**

45 The authors thank all professionals who have worked on the development of the AGCH. Also, we
46 would like to thank the members of the AGCH study group, these are the clinicians who work at the
47 Geriatrics Department of the Amsterdam University Medical Centres and who support the data
48 collection at the AGCH. We would also like to thank Marije Wolvers for her advice on the statistical
49 analysis.

50 **Collaborators**

51 The AGCH study group: R H A van den Broek; W J Frenkel; M J Henstra; M A van Maanen; C J M
52 Melkert, J L Parlevliet; E P van Poelgeest; M N Resodikromo; K J Kaland; I Oudejans; P M A van Rijn; N
53 van der Velde; H C Willems; D Wyatt

54

55

56

57

58

59

60

61

62

63

64

65 **Abstract**

66 **Background** Delirium in hospitalised older adults is associated with negative health outcomes.

67 Admission to an alternative care setting may lower the incidence of delirium. The Acute Geriatric
68 Community Hospital (AGCH) was recently opened in the Netherlands and uses a multi-component
69 non-pharmacological intervention strategy to prevent delirium.

70 **Objective** To describe the incidence of delirium found in the AGCH and compare this incidence to a
71 hospital control group found in literature.

72 **Design** Prospective cohort study; exploratory meta-analysis of proportions.

73 **Setting** The AGCH is an acute geriatric unit in an intermediate care facility.

74 **Participants** Patients aged >65 years with acute medical conditions admitted to the AGCH.

75 **Methods** Delirium assessment using the Confusion Assessment Method (CAM) upon admission and
76 on day one, two and three or until delirium had resolved. Patients' charts were reviewed if CAM was
77 missing. In an logistic mixed-effects model, the delirium incidence rate in AGCH was compared to
78 pooled delirium incidence rates from six studies found in a high-quality review.

79 **Results** 214 patients from the AGCH (mean age 81.9 years, 47% male, 12% with a history of
80 dementia) were included in the analysis. Delirium developed in 8% (18/214) (95% CI [confidence
81 interval] 5-13%) of patients during AGCH admission compared to 16% (95% CI 12-21%) in hospitals.
82 Admission to the AGCH was associated with a decreased delirium incidence rate compared to the
83 hospital control group (OR [odds ratio]= 0.49, 95% CI 0.24-0.98, p-value=.044).

84 **Conclusions** The delirium incidence in the AGCH was relatively low compared to those incidences
85 found in general hospitals.

86 **Introduction**

87 A common complication of hospitalisation in older adults is the development of delirium, an acute
88 disturbance in attention and several cognitive functions (1). The aetiology of delirium is considered
89 multifactorial (2). Delirium is associated with negative health outcomes, including functional and
90 cognitive decline, institutionalisation, and mortality (3, 4). The prevalence and incidence of delirium
91 is known to vary between settings and populations, with new-onset delirium during hospitalisation
92 ranging from 10% to 56% (5).

93 An alternative to conventional hospitalisation is admission to an acute geriatric unit outside
94 of a general hospital. This unit may be better adapted to the needs of older adults (6). In the
95 Netherlands, the Acute Geriatric Community Hospital (AGCH) was introduced in 2018 (7). This
96 geriatrician-led unit within an intermediate care facility integrates specialised medical treatment with
97 geriatric nursing care. A non-pharmacological multi-component delirium prevention strategy has
98 been implemented, consisting of nurses encouraging early mobilisation, preventing overstimulation
99 (single rooms, noise reduction), management of delirium-inducing drugs and improving orientation
100 through e.g. family involvement (6, 7). It is unknown what the effect of these interventions is on the
101 incidence of delirium in this new care setting. This is the first unit of its kind in the Netherlands but
102 there are other examples internationally (8).

103 We hypothesize that the non-pharmacological interventions at the AGCH may reduce the
104 incidence of delirium compared to incidence rates found in general hospitals. Therefore, the
105 objective of this study in the AGCH was to determine the incidence of delirium compared to those
106 incidence rates found in literature from general hospitals.

107 As secondary aims, we determined the duration of delirium and we quantified the use of
108 pharmacological delirium treatment. The duration of delirium is relevant as it can also be shortened
109 by a multi-component non-pharmacological intervention (9). Moreover, it is clinically relevant to
110 know if patients (with or without delirium) were prescribed antipsychotics and/or benzodiazepines

111 for the pharmacological treatment or prevention of delirium. According to the Dutch delirium
112 guideline, prescription of antipsychotics and/or benzodiazepines is not indicated for the prevention
113 of delirium, but may be indicated when delirium is present in a patient (10).

114

115 **Methods**

116 *Design and setting*

117 Data from a prospective cohort study were used. The study protocol was published
118 elsewhere (7). Data collection started in February 2019 and was temporarily ceased in March 2020
119 during the Covid-19 pandemic. The aim of the study was to measure the readmission to hospital rate
120 90 days post-discharge and to compare this rate to the rate found in two historical cohort groups of
121 older adults admitted to general hospitals. This exploratory analysis focuses on one of the secondary
122 outcomes, incident delirium.

123 Patients seen at the emergency department (ED) of the Amsterdam University Medical
124 Centres were assessed by an on-call geriatrician. Patients admitted to the AGCH were 65 years or
125 older, presenting with an acute medical problem requiring hospitalisation and one or more geriatric
126 conditions, such as a fall, functional impairment or polypharmacy (11). Patients who did not require
127 hospitalisation, but needed short-term care in an intermediate care facility, i.e. nursing home, were
128 excluded from admission to the AGCH. See the study protocol (7) and appendix 1 for complete
129 admission eligibility criteria.

130 *Ethical considerations*

131 The local Ethics Committee of the Amsterdam UMC, location AMC waived the obligation for
132 the study to undergo formal ethical approval as described under Dutch law. We included patients
133 who, or whose legal representative, could provide written informed consent.

134

135 *Control population*

136 We wanted to compare the delirium incidence in our newly opened acute geriatric
137 community hospital in the Netherlands to those incidences found in older adults in acute hospital
138 medical or geriatric units. However we did not recruit a control group during the study period. Two
139 historical control groups are planned to be used to compare the study's primary outcome to, but
140 these studies did not measure incident delirium (7, 12, 13). We therefore searched for sources of
141 aggregated data on the incidence rate of delirium in medical or geriatric (non-surgical) inpatients
142 with a mean age of about 80 years (search strategy- appendix 2). Furthermore, we excluded sources
143 with case-control studies or with studies reporting the incidence of patients participating in an
144 interventional study (excluded studies- appendix 3). Finally, we selected six studies from a review by
145 Inouye et al. as a control group (5). One of the aims of this review was to provide an overview of
146 delirium incidences in various hospital departments found in high quality studies— with a sample size
147 of 100 or more, a prospective sampling method and meeting the Strengthening the Reporting of
148 Observational Studies in Epidemiology [STROBE] criteria (14)., that were published between Jan 1st
149 2004 and August 31st 2012. The six studies that we selected reported delirium incidence on general
150 medical and geriatric wards. We checked in each paper if multi-component delirium prevention
151 strategies were implemented at the participating ward.

152 *Primary Outcome*

153 Incident delirium was the primary outcome of this study and was defined as '*the number of*
154 *new cases that develop within a cohort over a defined period*' based on the National Institute for
155 Health and Care Excellence (NICE) guideline on delirium (15). As we wanted to evaluate if patients
156 developed delirium during admission to the AGCH, patients were excluded from our analysis if
157 delirium was already present at the ED. The diagnosis of delirium was made by the geriatrician or
158 geriatric nurse specialist by clinical assessment and using the Confusion Assessment Method (CAM)
159 (16). The CAM was filled out upon presentation to the ED and during the first three days of admission

160 or until delirium had resolved. Nurses screened for signs of possible delirium, three times a day,
161 during the first three days of admission using the Delirium Observation Screening Scale (DOSS) (17).
162 Patients were assessed by the same geriatrician/nurse specialist for several consecutive days to
163 recognize changes in mental status. On the weekend an on-call geriatrician assessed delirium status
164 if delirium was clinically suspected. The DOSS and nursing chart covering the previous 24 hours were
165 also considered in the delirium assessment. If there was a possible delirium after day three of
166 admission, CAM assessments were continued until delirium had resolved.

167 *Duration of delirium*

168 The duration of delirium was counted from the day the diagnosis was made until the CAM
169 was permanently negative and/or the treating physician stated the delirium had resolved. In patients
170 with an unresolved delirium at the time of discharge, we defined the first day of delirium until
171 discharge as the duration of delirium. We reported the median duration of delirium as this measure
172 is adequate for reporting skewed data and is least influenced when in some patients delirium had not
173 resolved upon discharge.

174 *Use of antipsychotics and/or benzodiazepines*

175 The administration of haloperidol, other antipsychotics, and benzodiazepines was collected
176 from patients' charts. The researcher also checked if patients categorized as not delirious had
177 received antipsychotics. This was 1) a check to see if no patient with a delirium diagnosis was missed
178 by the researchers and 2) a measure to quantify the use of antipsychotics and/or benzodiazepines as
179 a preventive measure for delirium, although prescribing medication to prevent delirium is not
180 recommended by the Dutch guideline on delirium (10).

181 *Statistical analysis*

182 Descriptive statistics, chi-square, t-test, and Mann-Whitney U test were used to compare
183 patients with and without delirium upon admission. To compare incidence rates from literature we

184 pooled studies in a meta-analysis of proportions, using a random-effects model (18). We tested if the
185 difference in delirium incidence was statistically significant by creating a logistic mixed-effects meta-
186 regression model with the location of the study (hospital versus AGCH) as a moderator (19). We
187 found significant heterogeneity in our analysis, meaning there were large variations in estimates
188 between studies. We did not perform meta-regression of other covariates because the number of
189 included studies was limited (<10) (18). All analyses were performed using SPSS version 26.00 (IBM
190 SPSS Statistics, IBM Corporation, Armonk, NY) and R version 3.6.1. We used the metaphor
191 (Viechtbauer, 2010) and meta (Schwarzer et al., 2015) packages in R.

192 **Results**

193 Between January 31, 2019 and March 13, 2020, a total of 466 consecutive patients were admitted to
194 the AGCH (figure 1). 34 of these admissions were readmissions of study participants and 125 patients
195 were not approached based on the exclusion criteria. Of the remaining patients, 46 patients did not
196 consent to participation. Three patients were excluded because delirium assessments at the ED were
197 missing and 44 because of a prevalent delirium. The sample for this study therefore consisted of 214
198 patients (figure 1). Mean (SD) age was 81.9 (8.1) years, 47.2% was male, 12.1% had a diagnosis of
199 dementia, and 47.2% of the patients was frail (table 1). Development of delirium during admission
200 occurred in 18 out of 214 patients, which is an incidence rate of 8.4% (95% CI [confidence interval] 5-
201 13%). The median (IQR [interquartile range]) duration of delirium in the AGCH was 2.5 days (1.0-5.3)
202 (table 1). Incomplete resolution of the delirium at the time of discharge occurred in two patients.
203 Mean length of stay (SD) was 9.6 (7.3) days in all patients, 9.4 (7.4) days in patients with no delirium
204 and 11.9 (6.4) days in patients with delirium. Median length of stay (IQR) was 7.0 (5.0-11.00) days in
205 patients with no delirium and 10 (7.5-16.8) days in patients with delirium.

206 *Pharmacological treatment for delirium*

207 Eleven out of 18 patients (61.1%) with a diagnosis of delirium were administered medication for the
208 treatment of delirium. Haloperidol was administered most frequently (n=11), followed by additional

209 benzodiazepines (n=4), and any other type of antipsychotics (n=1). The regular prescription of
210 haloperidol was 0.5-2.0mg per dose, typically given once a day, or twice in case of severe delirium,
211 with a maximum of three dosages. Five (5 out of 196, 2.6%) patients without delirium were
212 administered haloperidol, either as prevention due to a high risk of delirium or as treatment for pre-
213 existing symptoms unrelated to delirium (table 1).

214 *Delirium incidence in comparison to reference group from literature*

215 The reference group was based on six studies that were included in the 2014 review by Inouye et al.
216 (5): in total 1546 study participants with an overall mean age of 80 years (figure 2 and table 2).

217 None of the studies, except for Friedman et al. (20) reported to have multi-component delirium
218 prevention strategies in place at the ward, we therefore assumed usual care was delivered. The
219 pooled delirium incidence rate of these six studies was 16% (95% CI random effects model 12-21%)
220 (figure 2). The meta-analysis showed a high heterogeneity ($I^2=84\%$). In a separate logistic mixed-
221 effects model comparing general hospitals (reference category) versus the AGCH, we found that
222 admission to the AGCH was associated with a decrease in delirium incidence (OR [odds ratio]= 0.49,
223 95% CI 0.24-0.98, $p=.044$).

224 *Adherence to CAM evaluations and missing data*

225 Based on the CAM evaluation and daily delirium assessment by the attending physician/nurse
226 specialist we could ascertain the presence delirium in the first three days of admission in all patients.
227 In patients with delirium 27.8% of total CAM evaluations and 46.3% of total DOSS scores were
228 missing during the first three days of admission. For patients without delirium 46.9% and 66.7% were
229 missing, respectively. In 15% of all cases all three CAM evaluations were missing, here we only used
230 physicians' notes and discharge letters to ascertain whether delirium had occurred. Discharge letters
231 mentioned whether a patient had experienced a delirium or not in 88.9% of patients with delirium
232 and in 84.8% of patients without delirium.

233 **Discussion**

234 We implemented a non-pharmacological multi-component delirium prevention strategy at the AGCH
235 and found an incidence rate of delirium of 8.4%. This incidence is significantly lower compared to
236 rates found in hospital medical or geriatric wards. We would expect to find a higher incidence in our
237 study sample, because delirium risk factors such as higher age, previous delirium, dementia,
238 comorbidities, poor ADL-function and polypharmacy were highly prevalent, moreover patients were
239 admitted for acute medical events requiring hospitalisation which increases the risk on developing
240 delirium compared to non-hospitalized patients (10, 21). A 2016 Cochrane review on interventions in
241 preventing delirium in hospitalised patients reports moderate quality evidence that multi-component
242 interventions in medical, non-surgical, patients lower delirium incidence, with a risk ratio of 0.63
243 (95% CI 0.43-0.92) for patients who receive a multi-component intervention versus those who did
244 not (6). Our findings are in line with this finding because the incidence that we found is on the lower
245 spectrum of those incidences reported in literature (5). Moreover, the median duration of delirium of
246 2.5 days is comparable to the duration that is found in literature on non-pharmacological
247 interventions (22). The prescription rate of medication (61.1%) may be lower in the AGCH compared
248 to other studies, which report rates of 74-86% (23, 24). The Dutch guideline on delirium, and
249 international guidelines alike, recommends to take a cautious approach to the prescription of
250 medication for the treatment of delirium (10, 25). Also, only a few patients received medication, in
251 this case haloperidol, for the prevention of delirium, meaning that there were not many non-
252 delirious patients receiving haloperidol. This is relevant because, administration of anti-psychotics
253 such as haloperidol could lower delirium incidence rates in high incidence groups (10). Furthermore,
254 in the Netherlands physicians prescribe relatively low dosages of haloperidol, 0.5-2.0 mg per dose.

255 *Limitations*

256 This study has multiple limitations. Firstly, the incidence rate of delirium could have been influenced
257 by selection bias: legal representatives of patients could not always be contacted to obtain consent.

258 Secondly, several CAM and DOSS assessments were not reported during the first three days of
259 admission. However, daily clinical delirium assessment by a geriatrician/trained nurse specialist and
260 nursing charts from every nursing shift (three per day) were available. Therefore, we do not have
261 missing data in the outcome incident delirium. Also, as this was an exploratory analysis of a
262 secondary outcome, no sample size was calculated, and our analysis may therefore be
263 underpowered. We also did not have insight into all of the baseline characteristics of the control
264 group from literature; the mean age was similar, but we did not or could not review information on
265 ADL (activities of daily living)-impairments or cognitive status available for each of these studies. This
266 makes it more difficult assess the comparability of this control group to our study sample. Moreover,
267 we could not definitively ascertain that ‘usual care’ was delivered in each unit or what this was
268 composed of. Especially in the geriatric units, such as the Acute Care for the Elderly Unit described in
269 Friedman et al. 2008 (20), an adapted environment may have been in place which is similar to the
270 environment of the AGCH and this could lower the risk of developing delirium in hospital. Finally, we
271 did not collect data on illness severity, which can be associated with delirium (26). Even so, patients
272 admitted to the AGCH suffer from acute medical illnesses for which they would otherwise have been
273 admitted to a general hospital. Also, they did have many other known delirium risk factors (21). In
274 addition patients with solely social or care indications, who require admission to intermediate care
275 but do not require hospital treatment are, by protocol, excluded from admission to the AGCH.

276 *Clinical and research implications*

277 Our exploratory analysis shows that the incidence rate of delirium in the AGCH is significantly lower
278 than in studies performed in general hospitals. This may indicate that this acute geriatric unit
279 situated in intermediate care could prevent delirium in older adults, at risk for delirium and requiring
280 hospital admission. However, as this study is only exploratory, a randomized controlled study or a
281 two-armed observational study using e.g. inversely weighted propensity scores (27) would be needed
282 to ascertain whether there is indeed a lower incidence of delirium, as was found in this study.

283 **Conclusion**

284 Delirium is a common complication in older persons during hospitalisation. Preliminary evidence on
285 the implementation of a multi-component, non-pharmacological intervention at the newly opened
286 Acute Geriatric Community Hospital resulted in a relatively low delirium incidence rate compared to
287 incidences reported in general hospitals.

288 References

- 289 1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fifth*
290 *Edition*. Arlington, VA, 2013.
- 291 2. Inouye SK, Charpentier PA; Precipitating factors for delirium in hospitalized elderly persons.
292 Predictive model and interrelationship with baseline vulnerability. *Jama* 1996;**275**(11):852-7.
- 293 3. Inouye SK, Rushing JT, Foreman MD, et al.; Does delirium contribute to poor hospital
294 outcomes? A three-site epidemiologic study. *J Gen Intern Med* 1998;**13**(4):234-42. doi:
295 10.1046/j.1525-1497.1998.00073.x.
- 296 4. Fick DM, Agostini JV, Inouye SK; Delirium superimposed on dementia: a systematic review. *J*
297 *Am Geriatr Soc* 2002;**50**(10):1723-32. doi: 10.1046/j.1532-5415.2002.50468.x.
- 298 5. Inouye SK, Westendorp RG, Saczynski JS; Delirium in elderly people. *Lancet*
299 2014;**383**(9920):911-22. doi: 10.1016/s0140-6736(13)60688-1.
- 300 6. Siddiqi N, Harrison JK, Clegg A, et al.; Interventions for preventing delirium in hospitalised
301 non-ICU patients. *Cochrane Database Syst Rev* 2016;**3**:Cd005563. doi:
302 10.1002/14651858.CD005563.pub3.
- 303 7. Ribbink ME, Macneil-Vroomen JL, van Seben R, et al.; Investigating the effectiveness of care
304 delivery at an acute geriatric community hospital for older adults in the Netherlands: a protocol for a
305 prospective controlled observational study. *BMJ Open* 2020;**10**(3):e033802. doi: 10.1136/bmjopen-
306 2019-033802.
- 307 8. Colprim D, Martin R, Parer M, et al.; Direct admission to intermediate care for older adults
308 with reactivated chronic diseases as an alternative to conventional hospitalization. *J Am Med Dir*
309 *Assoc* 2013;**14**(4):300-2. doi: 10.1016/j.jamda.2012.12.003.
- 310 9. Inouye SK, Bogardus ST, Jr., Charpentier PA, et al.; A multicomponent intervention to prevent
311 delirium in hospitalized older patients. *N Engl J Med* 1999;**340**(9):669-76. doi:
312 10.1056/nejm199903043400901.
- 313 10. Nederlandse Vereniging voor Klinische Geriatrie; Richtlijn Delier Volwassenen. 2013.
- 314 11. Buurman BM, Hoogerduijn JG, de Haan RJ, et al.; Geriatric conditions in acutely hospitalized
315 older patients: prevalence and one-year survival and functional decline. *PLoS One*
316 2011;**6**(11):e26951. doi: 10.1371/journal.pone.0026951.
- 317 12. Buurman BM, Parlevliet JL, van Deelen BA, et al.; A randomised clinical trial on a
318 comprehensive geriatric assessment and intensive home follow-up after hospital discharge: the
319 Transitional Care Bridge. *BMC Health Serv Res* 2010;**10**:296. doi: 10.1186/1472-6963-10-296.
- 320 13. Reichardt LA, Aarden JJ, van Seben R, et al.; Unravelling the potential mechanisms behind
321 hospitalization-associated disability in older patients; the Hospital-Associated Disability and impact
322 on daily Life (Hospital-ADL) cohort study protocol. *BMC Geriatr* 2016;**16**:59. doi: 10.1186/s12877-
323 016-0232-3.
- 324 14. von Elm E, Altman DG, Egger M, et al.; [The Strengthening the Reporting of Observational
325 Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies]. *Rev Esp*
326 *Salud Publica* 2008;**82**(3):251-9. doi: 10.1590/s1135-57272008000300002.
- 327 15. National Institute for Health and Care Excellence; Delirium: Diagnosis, Prevention and
328 Management. London, 2010.
- 329 16. Inouye SK, van Dyck CH, Alessi CA, et al.; Clarifying confusion: the confusion assessment
330 method. A new method for detection of delirium. *Ann Intern Med* 1990;**113**(12):941-8. doi:
331 10.7326/0003-4819-113-12-941.
- 332 17. Schuurmans MJ, Shortridge-Baggett LM, Duursma SA; The Delirium Observation Screening
333 Scale: a screening instrument for delirium. *Res Theory Nurs Pract* 2003;**17**(1):31-50. doi:
334 10.1891/rtnp.17.1.31.53169.
- 335 18. Wang N. *How to Conduct a Meta-Analysis of Proportions in R: A Comprehensive Tutorial*,
336 2018.

- 337 19. Viechtbauer W. *Comparing Estimates of Independent Meta-Analyses or Subgroups*.
338 https://www.metafor-project.org/doku.php/tips:comp_two_independent_estimates (Date Accessed
339 2020 Accessed, date last accessed)
- 340 20. Friedman SM, Mendelson DA, Bingham KW, et al.; Hazards of hospitalization: residence prior
341 to admission predicts outcomes. *Gerontologist* 2008;**48**(4):537-41. doi: 10.1093/geront/48.4.537.
- 342 21. Ahmed S, Leurent B, Sampson EL; Risk factors for incident delirium among older people in
343 acute hospital medical units: a systematic review and meta-analysis. *Age Ageing* 2014;**43**(3):326-33.
344 doi: 10.1093/ageing/afu022.
- 345 22. Martinez FT, Tobar C, Beddings CI, et al.; Preventing delirium in an acute hospital using a non-
346 pharmacological intervention. *Age Ageing* 2012;**41**(5):629-34. doi: 10.1093/ageing/afs060.
- 347 23. van Velthuisen EL, Zwakhalen SMG, Mulder WJ, et al.; Detection and management of
348 hyperactive and hypoactive delirium in older patients during hospitalization: a retrospective cohort
349 study evaluating daily practice. *Int J Geriatr Psychiatry* 2018;**33**(11):1521-1529. doi:
350 10.1002/gps.4690.
- 351 24. Nguyen PV-Q, Malachane A, Minh TTV; Antipsychotic prescription patterns in the
352 management of delirium symptoms in hospitalized elderly patients. *Proceedings of Singapore*
353 *Healthcare* 2017;**26**(4):230-234. doi: 10.1177/2010105817697733.
- 354 25. Excellence NifHaC. *Delirium: prevention, diagnosis and management*.
355 <https://www.nice.org.uk/guidance/cg103/chapter/1-Guidance#interventions-to-prevent-delirium>
356 (Date Accessed 2010 Accessed, date last accessed)
- 357 26. Inouye SK, Viscoli CM, Horwitz RI, et al.; A predictive model for delirium in hospitalized
358 elderly medical patients based on admission characteristics. *Ann Intern Med* 1993;**119**(6):474-81. doi:
359 10.7326/0003-4819-119-6-199309150-00005.
- 360 27. Austin PC, Stuart EA; Moving towards best practice when using inverse probability of
361 treatment weighting (IPTW) using the propensity score to estimate causal treatment effects in
362 observational studies. *Stat Med* 2015;**34**(28):3661-79. doi: 10.1002/sim.6607.
- 363 28. Katz S, Ford AB, Moskowitz RW, et al.; Studies of illness in the aged. The index of ADL: A
364 standardized measure of biological and psychosocial function. *Jama* 1963;**185**:914-9. doi:
365 10.1001/jama.1963.03060120024016.
- 366 29. Fried LP, Ferrucci L, Darer J, et al.; Untangling the concepts of disability, frailty, and
367 comorbidity: implications for improved targeting and care. *J Gerontol A Biol Sci Med Sci*
368 2004;**59**(3):255-63. doi: 10.1093/gerona/59.3.m255.
- 369 30. Folstein MF, Folstein SE, McHugh PR; "Mini-mental state". A practical method for grading the
370 cognitive state of patients for the clinician. *J Psychiatr Res* 1975;**12**(3):189-98. doi: 10.1016/0022-
371 3956(75)90026-6.
- 372 31. Charlson ME, Pompei P, Ales KL, et al.; A new method of classifying prognostic comorbidity in
373 longitudinal studies: development and validation. *J Chronic Dis* 1987;**40**(5):373-83. doi:
374 10.1016/0021-9681(87)90171-8.
- 375 32. Franco JG, Valencia C, Bernal C, et al.; Relationship between cognitive status at admission
376 and incident delirium in older medical inpatients. *J Neuropsychiatry Clin Neurosci* 2010;**22**(3):329-37.
377 doi: 10.1176/appi.neuropsych.22.3.329
- 378 10.1176/jnp.2010.22.3.329.
- 379 33. White S, Calver BL, Newsway V, et al.; Enzymes of drug metabolism during delirium. *Age*
380 *Ageing* 2005;**34**(6):603-8. doi: 10.1093/ageing/afi189.
- 381 34. Adamis D, Treloar A, Darwiche FZ, et al.; Associations of delirium with in-hospital and in 6-
382 months mortality in elderly medical inpatients. *Age Ageing* 2007;**36**(6):644-9. doi:
383 10.1093/ageing/afm094.
- 384 35. Eeles EM, Hubbard RE, White SV, et al.; Hospital use, institutionalisation and mortality
385 associated with delirium. *Age Ageing* 2010;**39**(4):470-5. doi: 10.1093/ageing/afq052.

- 386 36. Inouye SK, Zhang Y, Han L, et al.; Recoverable cognitive dysfunction at hospital admission in
387 older persons during acute illness. *J Gen Intern Med* 2006;**21**(12):1276-81. doi: 10.1111/j.1525-
388 1497.2006.00613.x.
- 389 37. Leslie DL, Marcantonio ER, Zhang Y, et al.; One-year health care costs associated with
390 delirium in the elderly population. *Arch Intern Med* 2008;**168**(1):27-32. doi:
391 10.1001/archinternmed.2007.4.
- 392 38. Siddiqi N, House AO, Holmes JD; Occurrence and outcome of delirium in medical in-patients:
393 a systematic literature review. *Age Ageing* 2006;**35**(4):350-64. doi: 10.1093/ageing/afl005.
- 394

Figure 1 – Participant flow-chart Acute Geriatric Community Hospital (AGCH) study.




Table 1. Baseline characteristics of the total study population grouped by patients with and without delirium.

	Total (n = 214)	No delirium (n = 196) 91.6%	Incident delirium (n = 18) 8.4%	p value^a
Age (years), mean (SD)	81.9 (8.1)	81.6 (8.0)	85.2 (8.8)	.08
Male, n (%)	101 (47.2)	93 (47.4)	8 (44.4)	.81
Born in the Netherlands, n (%)	160 (74.8)	146 (74.5)	14 (77.8)	.99
Marital status, n (%)				.58
Married/living together	69 (32.2)	65 (33.2)	4 (22.2)	
Single/Divorced	45 (21.0)	40 (20.4)	5 (27.8)	
Widow(er)	99 (46.3)	90 (45.9)	9 (50.0)	
Living arrangement before admission, n (%)				.80
Independent	174 (81.3)	158 (80.6)	16 (88.9)	
Nursing home	5 (2.3)	5 (2.6)	-	
Senior residence	33 (15.4)	31 (15.8)	2 (11.1)	
Other	2 (0.9)	2 (1.0)	-	
Level of education, n (%)				.44
Primary school	37 (17.3)	32 (16.3)	5 (27.8)	
Elementary technical/domestic science school	45 (21.0)	43 (21.9)	2 (11.1)	
Secondary vocational education	63 (29.4)	58 (29.6)	5 (27.8)	
Higher-level high school/third-level education	49 (22.9)	43 (21.9)	6 (33.3)	
Polypharmacy (≥ 5 medications), n (%)	160 (74.8)	147 (75.0)	13 (72.2)	.80
Primary admission diagnosis, n (%)				.44
Pneumonia	40 (18.7)	38 (19.4)	2 (11.1)	
Urinary tract infection (UTI)	27 (12.6)	25 (12.8)	2 (11.1)	
Other infections (excl. pneumonia/UTI)	21 (9.8)	8 (4.1)	3 (16.7)	
Congestive heart failure	20 (9.3)	18 (9.2)	2 (11.1)	

Neurologic disorders	19 (8.9)	17 (8.7)	2 (11.1)	
COPD exacerbation	15 (7.0)	15 (7.7)	-	
Fall(s)	13 (6.1)	12 (6.1)	1 (5.6)	
Gastrointestinal disease	10 (4.7)	10 (5.1)	-	
Electrolyte disturbance	6 (2.8)	4 (2.0)	2 (11.1)	
Other	43 (20.1)	39 (19.9)	4 (22.2)	
Katz-ADL^b score two weeks before admission, median (IQR)	1.0 (0.0-2.0)	1.0 (0.0-2.0)	2.0 (0.8-3.0)	.054
Katz-ADL^b score upon admission, median (IQR)	2.5 (1.0-4.0)	2.0 (1.0-4.0)	3.5 (0.8-5.3)	.34
Frailty^c, n (%)	101 (47.2)	92 (46.9)	9 (50.0)	.94
Unknown	59 (27.6)	55 (28.1)	4 (22.2)	
MMSE^d score, median (IQR)	25.0 (22.0-28.0)	25.0 (23.0-28.0)	23.0 (20.0-24.8)	.035
- Unknown or not done, n (%)	56 (26.2)	50 (25.5)	6 (33.3)	
History of dementia, n (%)	26 (12.1)	22 (11.2)	4 (22.2)	.17
Cognitive impairment^e, n (%)	88 (41.1)	76 (38.8)	12 (66.7)	.022
Charlson comorbidity index score^f, median (IQR)	3.0 (1.0-4.0)	2.5 (1.0-4.0)	3.0 (1.0-4.0)	.99
History of delirium/confusion during sickness, n (%)	55 (25.7)	48 (24.5)	7 (38.9)	.23
Duration of delirium, in median days (IQR) NA= not applicable	NA	NA	2.5 (1.0-5.3)	
Pharmacological treatment for delirium, n NA= not applicable	NA	NA	11	
- Haloperidol	16	5	11	
- Other antipsychotics	NA	NA	1	
- Benzodiazepines	NA	NA	4	

^a Incident delirium compared to no delirium

^b Katz Index score range 0-6, with a higher score indicating more dependence in activities of daily living (ADL) (28)

^c Based on Fried criteria for frailty range 0-5 with a score of 3 and higher indicating presence of physical frailty (29)

^d Mini Mental State Exam score ranging 0-30, MMSE score \leq 23 indicating cognitive impairment (30)

^e All patients with a diagnosis of dementia, a MMSE score \leq 23, or, in case of missing MMSE score, subjective cognitive problems

^f Range of 0-31, with a higher score indicating more or more severe comorbidity (31)




Figure 2 – Meta-analysis of proportions of delirium incidences in older hospitalised medical patients found in literature (5). The pooled incidence rate of these six studies was 16% (95% CI [confidence interval]random effects model 12-21%).

Table 2 –Studies included in meta-analysis of proportions of delirium incidence, studies from a 2014 review on delirium incidence by Inouye et al. CAM, Confusion Assessment Method; DRS, Delirium Rating Scale; DSM–IV, Diagnostic and Statistical Manual of Mental Disorders (4rd Ed). * original sample included prevalent delirium

Study (authors, publication year, country)	Study setting	Study design	Patients with delirium (n)	Patients without delirium (n)	Total sample (n)	Incidence rate (%)	Age, years	Mean age	Sex: male/female	Criteria for delirium	Assessment frequency (h)
Franco <i>et al.</i> , 2010, Colombia (32)	Medical	Case control	34	257	291	12%	≥59	74	105/186	CAM,DRS	24
White 2005, United Kingdom (33)	Medical	Cohort	29	178	207*	14%	≥75	82	Only number of males in total group	DAM,DSMIV	24
Adamis 2007, United Kingdom (34)	Geriatric	Cohort	47	117	164	29%	≥70	85	54/110	CAM,DRS	72
Friedman 2008, United States (20)	Geriatric	Cohort	42	171	212	20%	≥65	79	85/127	CAM	24
Eeles 2010, United Kingdom (35)	Medical	Cohort	23	175	198*	11%	≥74	83	117 males in total group	DSMIV	48
Inouye et al. 2006, United States (36)	Medical	Cohort	60	400	460	13%	≥70	80	183/277	CAM	24

Appendix 1 – Admission criteria Acute Geriatric Community Hospital (AGCH)

Criteria upon assessment at the emergency department:

- 1) Acute medical problems in older patients that require hospitalization, e.g., acute events such as a pneumonia or an exacerbation of chronic conditions such as heart failure or minor acute events in very frail patients.
- 2) Hemodynamic stability.
- 3) No need for complex diagnostic testing such as CT- or MRI scans during admission.
- 4) Expecting to return to previous living situation in 14 days.
- 5) Geriatric conditions e.g., delirium, cognitive impairment, falls and/or functional impairment.

Appendix 2 – Search strategy studies reporting incident delirium in medical inpatients.

Methods

We looked for sources of aggregated data (guidelines) or reviews reporting the incidence of delirium in older adults admitted to acute medical units and extracted the data from the individual studies that reported in the guideline.

Secondly, we performed a search on PubMed looking for (systematic) reviews or meta-analyses reporting incidence of delirium in older adults admitted to acute medical units.

Search strategy in PubMed

((delirium[MeSH Terms]) AND ((internal medicine[MeSH Terms])OR(medical[Title/Abstract])) NOT (surgical[Title/Abstract]) NOT (orthopaedic[Title/Abstract]) NOT (hip fracture[Title]) NOT (fracture[Title]) NOT (pediatric[Title/Abstract]) NOT (postoperative[Title/Abstract]))

Filters applied: Meta-Analysis, Systematic Review

Criteria for including studies were:

Inclusion criteria

- *Older adults aged 60 years and older*
- *Admitted to medial or geriatric wards of acute or general hospitals*
- *Prospective cohort and cross-sectional studies*
- *Control arms of clinical trials*

Exclusion criteria

- *Mean age study population <65 years*
- *Including data from intervention arm of clinical trials*
- *Case-control studies*
- *Retrospective studies*
- *Data from surgical wards or intensive care units*

Appendix 3 – found and excluded studies.

Clinical guideline on delirium and or review:

1) **Excluded: NICE guideline delirium 2010 (15)** We found a guideline reporting the incidence of delirium in general medicine wards; the NICE guideline for delirium which reports an incidence of 15.2% (range 12.5-17.9) based on two studies.(15) One of these studies included the intervention arm of a clinical trial(37) and therefore we decided not to use the NICE guideline report on incident delirium as a control group.

2) **Included: Inouye et al. 2014 (5)**- This is the high-quality review from which we used included studies as a control group.

Results from PubMed search looking for systematic reviews or meta-analyses:

Our search yielded 21 hits (1,124 without filters) in which we found two relevant systematic reviews based on screening title and abstract:

1) **Excluded: Siddiqi et al. 2006 (38)** We found a systematic review by Siddiqi et al.(38) that reported on the incidence of delirium in 14 studies, due to the smaller sample sizes and studies being published before 2005 we decided not to use this systematic review.

2) **Excluded: Ahmed et al. 2014 (21)** We found a systematic review by Ahmed et al.,(21) including 11 studies that report delirium incidence in older medical patients, but in this study there was a risk of bias because it aimed to describe risk factors for delirium and not solely the incidence of delirium. This meant case-control studies were used and possible relevant studies were excluded. Therefore, we decided not to use this review.

Flow-chart of selection of studies (aggregated data and (systematic reviews) for hospital control group :

