Title: LITE SABR M1: a phase I trial of Lattice SBRT for large tumors

Authors: Sai Duriseti MD, PhD, James Kavanaugh, PhD, Jeff Szymanski MD, PhD, Yi Huang PhD, Franco Basarabescu, BS, Aadel Chaudhuri MD, PhD, Pamela Samson, MD, MS, Alexander Lin, MD, PhD, Clifford Robinson MD, Matthew B Spraker MD, PhD

Affiliation: Department of Radiation Oncology, Washington University in St. Louis

DISCLAIMER: This pre-print manuscript has not been peer reviewed and is therefore incomplete. During the peer review process, referees assess the manuscript, identify weaknesses, and suggest edits to the authors. This could include additional analyses or adding new interpretations of the results. Typically, medical journals will not publish an article until the editors are satisfied with the authors' edits according to referee comments. If you have any questions, please contact the corresponding author.

Corresponding Author:

Matthew B. Spraker, MD, PhD

Email: mspraker@wustl.edu

Twitter: @SprakerMDPhD

Abstract

Purpose: Stereotactic body radiotherapy (SBRT) is an attractive treatment option for patients with metastatic and/or unresectable tumors, however its use is limited to smaller tumors. Lattice is a form of spatially fractionated radiotherapy that may offer safe delivery of SBRT for bulky tumors. We previously described Lattice SBRT, which delivers 20 Gy in 5 fractions with a simultaneous Lattice boost to 66.7 Gy. The goal of this study was to prospectively evaluate the acute toxicity and quality of life (QoL) of patients with large tumors (> 5 cm) treated with Lattice SBRT.

Methods: This was a single-arm phase I trial conducted between October 2019 and August 2020. Patients with tumors >4.5 cm were eligible. Lattice SBRT was delivered every other day then patients were evaluated for 90 days. The primary outcome was the rate of treatment-associated grade 3+ acute toxicity by Common Terminology Criteria for Adverse Events (CTCAE) version 5.0 criteria. Other outcomes included patient reported toxicity and QoL inventories, and GTV changes following Lattice SBRT.

Results: Twenty patients (22 tumors) were enrolled. Median gross tumor volume (GTV) was 783 cc (range: 64-3713 cc). Fifty percent of tumors were in the thorax, 45% abdomen/pelvis, and 5% extremity. There was no grade 3+ toxicity probably or definitely attributable to Lattice SBRT. There was one case of grade 4 toxicity possibly associated with Lattice SBRT.

Conclusions: This phase I study met its primary endpoint of physician reported safety, confirmed with secondary PRO-CTCAE outcomes. Early efficacy, as demonstrated by PROMIS QoL and imaging, is encouraging. An ongoing phase II clinical trial of Lattice SBRT is evaluating the safety and efficacy of this novel technique.

Introduction

Hypofractionated radiotherapy is an appealing treatment for patients with metastatic and/or unresectable tumors in need of palliative radiotherapy.[1] Stereotactic body radiotherapy (SBRT) has been increasingly used in this setting as it may improve local control and better palliate symptoms as compared with conventional fractionation.[2,3] However, SBRT faces limitations when used for large tumors. Locoregional control and toxicity are both thought to be worse for larger tumors treated with SBRT compared with smaller tumors.[4,5] Toxicity has also been reported to be worse for larger tumors treated with SBRT.[6–8] As a consequence, consensus guidelines support use of SBRT for tumors 5 cm or smaller.[9]

Spatially-fractionated radiotherapy is hypothesized to allow for safe, hypofractionated, dose escalated radiotherapy for large tumors.[10] Spatially-fractionated radiotherapy can be delivered in a grid-like pattern (i.e. GRID) using 3D conformal planning techniques. It can also be delivered using an intensity modulated technique (i.e. Lattice radiotherapy) that simultaneously delivers a conformal lower dose of radiation to a planning target volume (PTV) with a simultaneous integrated boost to a geometric lattice of spherical sub-volumes confined to gross tumor.[11] Both GRID and Lattice radiotherapy have been used as definitive treatment, as a boost after conventional radiation, and de novo or for reirradiation. While the literature has been limited to retrospective studies, GRID and Lattice have associated with significant clinical responses and limited toxicities in large and/or radioresistant tumors, including non-small cell lung cancer and sarcomas.[12,13,14,15]

We previously described a novel spatially fractionated SBRT technique – Lattice SBRT – that delivers 20 Gy in 5 fractions with a 66.7 Gy simultaneous integrated Lattice boost using volumetric modulated arc therapy (VMAT).[16] Here we present the results of the first prospective study of Lattice SBRT, LITE SABR M1: a single-arm phase I study of Lattice SBRT for large tumors (NCT04133415).

Materials/Methods

Study Design and Patients

This study was a prospective, single-arm, single-institution phase 1 trial approved by the Siteman Cancer Center Protocol Review and Monitoring Committee and the Human Research Protection Office at Washington University in St. Louis. All eligible patients provided signed consent for voluntary participation in this study.

Eligible patients were at least 18 years of age, had a histologically confirmed cancer diagnosis, had a tumor of at least 4.5 cm in largest axial diameter, and Eastern Cooperative Group (ECOG) performance status of 2 or less. All patients received a multidisciplinary recommendation for radiotherapy for palliation of symptoms associated with the tumor target, asymptomatic progression, or as definitive treatment for locally advanced unresectable and/or metastatic tumors. Patients who received or planned to receive systemic therapy before or after Lattice SBRT were eligible, although a 2-week washout was recommended before or after Lattice SBRT. Concurrent systemic therapy was not allowed. Patients were ineligible if they previously received radiotherapy to the same area, if the tumor needed emergent surgical intervention, if they were pregnant, or had uncontrolled HIV with CD4 below 350 $^{cells}/_{\mu l}$ or on certain anti-retroviral therapies.

Treatment Planning and Delivery

Our treatment planning and quality assurance (QA) process has been detailed previously, and a guide of the treatment planning process is available.[16] Computed tomography (CT) simulation was completed with 3 mm slices and four-dimensional (4D) CT was used to assess tumor motion for all patients with thoracic and abdominal tumors above the iliac crest. Gated radiotherapy was used if tumor motion was more than 5 mm in a single direction. The planning process is fully described in the companion paper, but it is briefly summarized as follows. The treating radiation oncologist created a gross tumor volume (GTV) and fusion of prior diagnostic imaging was allowed. They then created a 5–10 mm isotropic expansion of the GTV to form the PTV_2000. Using the GTV, a dosimetrist and/or physicist created a PTV_6670 (i.e. the Lattice boost volume) and other Lattice planning volumes (i.e. PTV_Avoid). Volumetric-modulated arc therapy (VMAT) Lattice SBRT plans were generated in the Eclipse Treatment Planning System (Varian, Palo Alto, CA). The protocol required that

all plans meet consensus 5-fraction SBRT organ at risk (OAR) constraints published by the American Association of Medical Physicists.[17] If a plan could not meet the OAR constraints, the PTV_6670 was modified.

Lattice SBRT was delivered every other day for a total of 5 fractions. Patients could receive Lattice SBRT to multiple sites on alternating days if the radiotherapy plans did not overlap. Patients could also receive other conventional palliative radiotherapy courses to other lesions during the study period as directed by the treating physician. On-board imaging was required to verify positioning for all patients. Cone-beam CT (CBCT) and orthogonal kV films or fluoroscopic imaging were obtained immediately prior to treatment for tumor position verification. Structures were created and displayed on the CBCT from critical dose lines to assist with OAR position verification at the time of treatment.

Follow Up, Physician, and Patient Reported Outcomes

Patients were assessed for acute toxicity through 90 days after completion of therapy and were evaluated in clinic at baseline, once during therapy, and at 14, 30, and 90 days. Physician-reported and patient-reported outcomes (PROs) were evaluated at each visit using CTCAE version 5.0 and PRO-CTCAE, respectively. PROMIS anxiety (Item Bank v1.0 – Emotional Distress – Anxiety), physical global health (v1.2, physical 2a), depression (v1.0, short form 4a), pain interference (v1.0, short form 4a), and physical function (v2.0, short form 10a) were assessed at each visit.[18,19]

Statistical Analysis

The primary endpoint of the study was the rate of treatment-associated grade 3 or greater CTCAE v5.0 toxicity within 90 days of completion of Lattice SBRT. Early stopping of the trial was based on a Pocock-type stopping boundary. We assumed the expected treatment emergent severe adverse event rate of 30% with an unacceptable rate of higher than 40%. The trial opened on 10/31/2019 with an enrollment goal of 10 patients. The trial was expanded since accrual was robust and this would allow for better detection of possible severe acute toxicities prior to opening a planned phase II trial. It was expanded to 15 patients on 4/23/2020 and then to 20 patients on 6/17/2020. Trial accrual by month is exhibited in Supplementary Figure 1. The trial completed enrollment on 08/11/2020.

Secondary objectives included assessment of patient-reported toxicity and quality of life (QOL) at baseline and in follow up. Toxicity was assessed using PRO-CTCAE according to site-specific inventories. A composite score was calculated for each pertinent symptom for thoracic tumors and abdominal/pelvic tumors at all time points. The composite score combines the severity, frequency, and interference elements of each toxicity into a single score for efficient reporting.[20] PROMIS anxiety (Item Bank v1.0 – Emotional Distress – Anxiety), global health – physical (2a, v1.2), pain interference (Short Form 4a), and physical function (Short Form 10a) was used to asses PROMIS QOL scores.[21] PROMIS inventories were scored using the HealthMeasures Scoring Service (https://www.assessmentcenter.net/ac_scoringservice) and the median T score was calculated for each patient for each available time point.

An exploratory analysis of tumor imaging response after Lattice SBRT was completed. Follow up images at least 30 days from completion of treatment for all patients were imported into the Eclipse TPS (Varian, Palo Alto) and fused with each patient's CT simulation series. Using the originally contoured GTV_2000 as a benchmark, the treated lesions were contoured on each available diagnostic CT scan and assessed for percent change from baseline by volume.

Results

Patient and Disease Characteristics and Disease-specific Outcomes

Between October 2019 and August 2020, 20 patients underwent Lattice SBRT. Two patients were treated to two sites each for a total of 22 tumors treated. The CONSORT diagram for the study is shown in Figure 1. Patient and tumor characteristics are shown in Table 1. The population was 50% female, median age was 67 years (range: 31 to 86), and median ECOG status was 1 (range 0 to 2). Thirteen patients (65%) identified as White, 6 patients (30%) identified as Black, and 1 patient (5%) identified as Asian. Nine patients had soft tissue sarcomas (45%), 7 non-small cell lung cancers (35%), 1 thymic carcinoma (5%), 1 malignant mesothelioma (5%), 1 endometrial adenocarcinoma (5%), and 1 colonic adenocarcinoma (5%). Median GTV was 579.2 cc (range: 54.2–3713.5 cc) and median largest diameter of the tumor was 11.1 cm (range: 5.6–21.4 cm). Fifteen patients (75%) had undergone at least 1 course of chemotherapy and 3 patients (15%) received radiotherapy to other sites prior to receiving Lattice SBRT. Eight patients (40%) received chemotherapy after

completion of Lattice SBRT. The median time from CT simulation to first fraction was 12 days (range: 6–21 days). The median length of the course of treatment was 11 days (range 9–23 days). Three patients (15%) underwent Lattice SBRT as their only form of therapy. At 90-day follow up, 80% of treated patients were alive for evaluation.

Toxicity

All patients successfully completed the prescribed Lattice SBRT course. Cumulative acute toxicities within 90 days of therapy are shown in Table 2. Only one patient (Patient 11) had grade 3 or greater toxicity possibly related to Lattice SBRT. This patient underwent Lattice SBRT to a 3713.5 cc primary retroperitoneal leiomyosarcoma and a 1426.3 cc right thigh metastasis. The patient had known ureteral involvement from retroperitoneal tumor and had percutaneous nephrostomy tubes placed before starting treatment. After 2 fractions of Lattice SBRT to the retroperitoneal tumor and 1 fraction to the extremity metastasis, the patient was admitted for grade 4 urosepsis requiring pressor support, antibiotics, and nephrostomy tube exchange. The patient resumed treatment to the retroperitoneal and extremity tumors 12 and 14 days after the last fraction to each site, respectively. There was also grade 3 transaminitis concurrent with the sepsis and grade 2 diarrhea that resolved after treatment. On plan review, a segment of the intra-renal portion of the tube was inside of the 2000 cGy isodose line, and the soft tissue traversed by the tube was within the 1000 cGy isodose line.

Patient 2 experienced grade 2 pneumonitis possibly associated with Lattice SBRT. The patient was treated with Lattice SBRT to a 616.9 cc right, anterior mediastinal metastasis from solitary fibrous tumor. Sixty days after completion of therapy, the patient reported increased dyspnea on exertion consistent with radiation pneumonitis. No imaging was obtained. This was treated with prednisone and the symptoms resolved. There were no deaths associated with Lattice SBRT.

Patient reported outcomes

Prior to treatment, 95% of the total cohort (19 of 20 eligible) completed baseline PRO evaluation. At 14 days post-treatment, 84% of eligible patients (16 of 19 eligible) completed post-treatment evaluation, with 3 patients not completing PRO follow up at this time point, and 1 patient unavailable due to enrollment on hospice. At 30 days post-treatment, 56% (11/18) completed post-treatment evaluation, with 7 patients not

completing PRO follow up at this time point, 1 unavailable due to enrollment on hospice, and 1 deceased within 30 days follow up. At 90 days post-treatment, 62% (8/13) completed post-treatment evaluation, with 5 lost to PRO follow up at this time point, 3 unavailable due to enrollment on hospice, and 4 deceased within 90 days follow up.

The proportions of patients thoracic and abdominal/pelvic tumors with grade 0-1 and grade 2+ composite PRO-CTCAE scores at baseline and 14 days after therapy are shown in Figure 2. In patients with thoracic targets, the proportion of patients with grade 2+ shortness of breath decreased from baseline (33%) to 14 days (11%) after Lattice SBRT. The proportion of patients with grade 2+ slightly increased for vomiting (11% to 22%) and was unchanged for pain and fatigue from baseline to 14 days. In patients with abdominal or pelvic tumors, there was a decrease in the proportion of patients with grade 2+ pain (86% to 57%), GI/GU (71%-43%), and anxiety (43% to 14%) while fatigue remained constant (86%) from baseline to 14 days. PRO-CTCAE data for all patients is shown in Supplementary Table 1.

Median PROMIS T scores are shown in Table 3. These demonstrate that, at baseline, patients had more anxiety (median=55.7), depression (median=55.4), pain interference (median=61.3), and less physical global health (median=32.4) and physical function (median=34.5) than healthy individuals. At 14 days after treatment, patients improved on all measures. While data from only seven patients were available at 90 days, median anxiety (40.5), physical global health (39.2), depression (41.0), pain interference (41.6), and physical function (43.3) were all improved. PRO-CTCAE and PROMIS responses at 30- and 90-day follow up are available in the supplementary materials.

Imaging Response

Thirteen patients with 14 tumors had diagnostic imaging meeting criteria for analysis following Lattice SBRT. The tumor volume at each imaging time point, normalized by the original GTV volume, for all evaluable tumors is depicted in Supplementary Figure 2. The first follow up scan was a median of 1.8 months following Lattice SBRT, and all evaluable GTVs had decreased in size (median 24.4% decrease, range: 2.4–91.7%). Eleven patients (12 tumors) had a second follow up scan at a median of 4.5 months, and all but one tumor continued to shrink, with one tumor growing slightly (median 47.4% decrease, range: -5.3–96.1%).

Discussion

This prospective phase I study found Lattice SBRT is safe to deliver as palliative radiotherapy for patients with very large tumors, including those who receive systemic therapy before or after radiation. We did not observe acute grade 3+ toxicities probably or definitely attributable to Lattice SBRT, and PROs suggest that the acute effects of Lattice SBRT did not add significant toxicity or harm QoL in the acute setting. Of note, there was only one instance of grade 2 radiation pneumonitis in patients with thoracic tumors (N=10). This patient had a large solitary fibrous tumor of the right upper lobe of the lung, abutting the mediastinum. While the Lattice SBRT plan met protocol constraints, it is of interest to note that the patient received immune checkpoint inhibitor therapy alone approximately 8 months prior to commencement of radiation and the patient experienced significant toxicity associated with the drug. It is possible that this patient had higher propensity to experience radiation pneumonitis than other patients.[22]

Few studies have evaluated SBRT for tumors larger than 5 cm and all are retrospective.[6–8] However, on average, the patients treated in these series had much smaller tumors than those treated in the current study with a reported median largest dimension 5.6 cm (versus 11.1 cm in this study) or median GTV of 52.9 cc (versus 579.2 cc) with a reported max 597.8 cc (versus 3713 cc). Despite this difference, this study was associated with similar or lower toxicity rates compared with the prior work. For example, the largest study completed by Verma and colleagues reported grade 3+ toxicities in 5% of patients with nine cases (10%) of grade 2+ radiation pneumonitis.[8] Two other retrospective series show a similar grade 3+ toxicity rates of 10%[6] and 13%[7] with grade 2+ pneumonitis rates of 12%[6] and 7%[7]. Therefore, it appears that our spatially fractionated SBRT approach may offer a reasonable acute toxicity profile consistent with conventional SBRT, even for tumors significantly larger than those reported in the literature.

The secondary and exploratory analyses of PRO data and GTV change suggest that Lattice SBRT is efficacious as a palliative treatment for patients with large tumors. PRO-CTCAE data show that patients did not experience increased toxicity with Lattice SBRT except one patient each who experienced worse anxiety and nausea. PROMIS inventories demonstrated that patients had improved physical function, pain, global health, anxiety, and depression at 14 days after treatment and this improvement was preserved at 90 days. Analysis of

GTVs demonstrated clinically significant shrinkage (median 24.4% by volume) within a median of 81 days following Lattice SBRT despite the fact that most patients had cancers that are traditionally considered radioresistant (i.e. sarcomas, NSCLC). Further, 36% of evaluable tumors (N=11) had greater than 80% reduction of volume by the second imaging time point 3-6 months after treatment.

This study has important limitations. First, toxicity was only prospectively evaluated for 90 days. As SBRT can be associated with significant late toxicity, it is vital to evaluate for late toxicity associated with Lattice SBRT. Also, the study evaluated a highly heterogeneous cohort in order to assess the safety of a promising novel radiotherapy option for patients that are otherwise treated with a range of radiotherapy doses and fractionations, other local therapy options, or supportive care. While Lattice SBRT appears very efficacious, it is possible that established simpler radiotherapy options that can be planned and delivered within 1 day, such as 20 Gy in 5 fractions using 3D conformal radiotherapy could offer similar symptom relief. The ongoing prospective phase II study of Lattice SBRT at our center will address some of these limitations (NCT 04553471). It is enrolling patients into pre-specified thoracic, abdomen, pelvis, and sarcoma sub-groups, and will observe patients for local control and toxicity for 1 year.

References

1. Spencer K, Parrish R, Barton R, Henry A. Palliative radiotherapy. *BMJ*. 2018;360:k821. doi:10.1136/bmj.k821

2. Shiue K, Cerra-Franco A, Shapiro R, et al. Histology, Tumor Volume, and Radiation Dose Predict Outcomes in NSCLC Patients After Stereotactic Ablative Radiotherapy. *J Thorac Oncol*. 2018;13(10):1549-1559. doi:10.1016/j.jtho.2018.06.007

3. Nguyen Q-N, Chun SG, Chow E, et al. Single-Fraction Stereotactic vs Conventional Multifraction Radiotherapy for Pain Relief in Patients With Predominantly Nonspine Bone Metastases: A Randomized Phase 2 Trial. *JAMA Oncol*. 2019;5(6):872-878. doi:10.1001/jamaoncol.2019.0192

4. Ricco A, Davis J, Rate W, et al. Lung metastases treated with stereotactic body radiotherapy: the RSSearch® patient Registry's experience. *Radiat Oncol*. 2017;12(1):35. doi:10.1186/s13014-017-0773-4

5. Allibhai Z, Taremi M, Bezjak A, et al. The Impact of Tumor Size on Outcomes After Stereotactic Body Radiation Therapy for Medically Inoperable Early-Stage Non-Small Cell Lung Cancer. *Int J Radiat Oncol*. 2013;87(5):1064-1070. doi:10.1016/j.ijrobp.2013.08.020

6. Peterson J, Niles C, Patel A, et al. Stereotactic Body Radiotherapy for Large (> 5 cm) Non–Small-Cell Lung Cancer. *Clin Lung Cancer*. 2017;18(4):396-400. doi:10.1016/j.cllc.2016.11.020

7. Woody NM, Stephans KL, Marwaha G, Djemil T, Videtic GMM. Stereotactic Body Radiation Therapy for Non-Small Cell Lung Cancer Tumors Greater Than 5 cm: Safety and Efficacy. *Int J Radiat Oncol*. 2015;92(2):325-331. doi:10.1016/j.ijrobp.2015.01.045

8. Verma V, Shostrom VK, Kumar SS, et al. Multi-institutional experience of stereotactic body radiotherapy for large (≥5 centimeters) non–small cell lung tumors. *Cancer*. 2017;123(4):688-696. doi:https://doi.org/10.1002/cncr.30375

9. Videtic GMM, Donington J, Giuliani M, et al. Stereotactic body radiation therapy for early-stage non-small cell lung cancer: Executive Summary of an ASTRO Evidence-Based Guideline. *Pract Radiat Oncol*. 2017;7(5):295-301. doi:10.1016/j.prro.2017.04.014

10. Billena C, Khan AJ. A Current Review of Spatial Fractionation: Back to the Future? *Int J Radiat Oncol • Biol • Phys*. 2019;104(1):177-187. doi:10.1016/j.ijrobp.2019.01.073

11. Wu X, Perez NC, Zheng Y, et al. The Technical and Clinical Implementation of LATTICE Radiation Therapy (LRT). *Radiat Res*. 2020;194(6):737-746. doi:10.1667/RADE-20-00066.1

12. Huhn JL, Regine WF, Valentino JP, Meigooni AS, Kudrimoti M, Mohiuddin M. Spatially Fractionated GRID Radiation Treatment of Advanced Neck Disease Associated with Head and Neck Cancer. *Technol Cancer Res Treat*. 2006;5(6):607-612. doi:10.1177/153303460600500608

13. Mohiuddin M, Miller T, Ronjon P, Malik U. Spatially Fractionated Grid Radiation (SFGRT): A Novel Approach in the Management of Recurrent and Unresectable Soft Tissue Sarcoma. *Int J Radiat Oncol*. 2009;75(3, Supplement):S526. doi:10.1016/j.ijrobp.2009.07.1200

14. Blanco Suarez JM, Amendola BE, Perez N, Amendola M, Wu X. The Use of Lattice Radiation Therapy (LRT) in the Treatment of Bulky Tumors: A Case Report of a Large Metastatic Mixed Mullerian Ovarian Tumor. *Cureus*. 7(11). doi:10.7759/cureus.389

15. Amendola BE, Perez NC, Wu X, Amendola MA, Qureshi IZ. Safety and Efficacy of Lattice Radiotherapy in Voluminous Non-small Cell Lung Cancer. *Cureus*. 11(3). doi:10.7759/cureus.4263

16. Duriseti S, Kavanaugh J, Goddu S, et al. Spatially fractionated stereotactic body radiotherapy (Lattice SBRT) for large tumors. *Adv Radiat Oncol*. 2021;0(0). doi:10.1016/j.adro.2020.100639

17. Benedict SH, Yenice KM, Followill D, et al. Stereotactic body radiation therapy: The report of AAPM Task Group 101: Stereotactic body radiation therapy: The report of TG101. *Med Phys*. 2010;37(8):4078-4101. doi:10.1118/1.3438081

18. PROMIS. Accessed December 26, 2020. https://www.healthmeasures.net/explore-measurement-systems/promis

19. Dueck AC, Scher HI, Bennett AV, et al. Assessment of Adverse Events From the Patient Perspective in a Phase 3 Metastatic Castration-Resistant Prostate Cancer Clinical Trial. *JAMA Oncol*. 2020;6(2). doi:10.1001/jamaoncol.2019.3332

20. Basch E, Becker C, Rogak LJ, et al. Composite grading algorithm for the National Cancer Institute's Patient-Reported Outcomes version of the Common Terminology Criteria for Adverse Events (PRO-CTCAE). *Clin Trials*. 2021;18(1):104-114. doi:10.1177/1740774520975120

21. Jensen RE, Potosky AL, Moinpour CM, et al. United States Population-Based Estimates of Patient-Reported Outcomes Measurement Information System Symptom and Functional Status Reference Values for Individuals With Cancer. *J Clin Oncol*. 2017;35(17):1913-1920. doi:10.1200/JCO.2016.71.4410

22. Shaverdian N, Beattie J, Thor M, et al. Safety of thoracic radiotherapy in patients with prior immune-related adverse events from immune checkpoint inhibitors. *Ann Oncol*. 2020;31(12):1719-1724. doi:10.1016/j.annonc.2020.09.016

Table 1: Patient clinical information, disease information, gross tumor volume (GTV), time from last day of systemic therapy prior to Lattice SBRT (days), time from completion of Lattice SBRT to initiation of subsequent systemic therapy (days)

Participant	ECOG	Indication	Sex	Site	Histology	GTV (cc)	Time prior to lattice	Time after lattice
1[*]	2	Palliation	F	Thorax	NSCLC, poorly-differentiated	1784.2	--	--
2[*]	1	Palliation	M	Thorax	Solitary fibrous tumor	616.9	281	--
3[*]	1	Palliation	F	Thorax	NSCLC, adenocarcinoma	551.4	--	6
4[*]	1	Oligoprogression	M	Thorax	Leiomyosarcoma	64.3	49	--
5[*]	0	Oligoprogression	M	Thorax	NSCLC, squamous cell carcinoma	318.4	126	143
6	2	Definitive	M	Abdomen	Undifferentiated pleomorphic sarcoma	644.8	52	39
7[*]	2	Palliation	M	Thorax	Neuroendocrine tumor	606.9	396	--
8	1	Palliation	F	Thorax	NSCLC, squamous cell carcinoma	251.1	--	--
9[*]	1	Palliation	M	Abdomen	Round cell sarcoma	67.0	14	--
10[*]	0	Oligoprogression	M	Thorax	Rhabdomyosarcoma, pleomorphic type	54.2	58	1
11[*+]	1	Definitive	F	Abdomen	Leiomyosarcoma	3713.5	--	--
				Extremity		1426.3		
12[*▯]	2	Palliation	F	Thorax	NSCLC, adenocarcinoma	206.9	--	24
				Abdomen		316.7		
13[*]	2	Palliation	F	Pelvis	Rhabdomyosarcoma, embryonal type	1666.6	21	--
14[*]	1	Oligoprogression	F	Pelvis	NSCLC, poorly-differentiated	208.3	37	--
15	1	Definitive	F	Thorax	Thymoma	815.1	77	--
16[*]	1	Palliation	F	Thorax	Mesothelioma	1486.4	48	4
17[*]	1	Palliation	M	Abdomen	Endometrial, adenocarcinoma	131.7	342	11
18[*]	2	Palliation	F	Pelvis	Colon adenocarcinoma	3616.2	29	--
19[*]	2	Palliation	F	Pelvis	Desmoplastic small round cell tumor	462.7	15	16
20[*]	2	Oligoprogression	M	Abdomen	Leiomyosarcoma	1573.2	47	--

NSCLC = Non-small cell lung cancer

[*]   Patients with metastatic disease

[+]   Patient 11 had 2 sites of disease, the primary retroperitoneal tumor and a single site of metastatic disease in the extremity.

[▯]   Patient 12 had 2 sites of disease, the primary tumor in the right lung and a single site of metastatic disease in the right adrenal gland.

Table 2: <u>Attribution of toxicity for patients treated with Lattice SBRT.</u> The total number of toxicities graded per CTCAE version 5.0 criteria are shown.

	Unrelated			Unlikely			Possibly		
Grade	2	3	4	2	3	4	2	3	4
Total Events	53	14	1	6	2	1	1[*+]	2[+]	1[+]

*　　Patient 2 had increased dyspnea and work of breathing about 60 days after completion of therapy, clinically consistent with radiation pneumonitis. The patient was treated with prednisone and had resolution of his symptoms.

[+]　　Patient 11 required hospitalization for urosepsis after 2 fractions of Lattice SBRT to the abdominal tumor. The patient had Grade 3 transaminitis, attributed to sepsis physiology. The patient also had Grade 2 diarrhea that self-resolved, however given Lattice SBRT delivery to the abdomen, an acute radiotherapy toxicity could not be excluded.

Table 3: Median (range), of PROMIS-QOL inventory T scores were assessed at baseline and each follow up time point, as available.

	Baseline (N=19)	*14-day (N=15)*	*30-day (N=11)*	*90-day (N=7)*
Anxiety	55.7 (40.5–72.5)*	40.5 (40.5–67.6)	48.4 (40.5–62.4)	40.5 (40.5–52.2)
Physical global health	32.4 (25.9–57.8)	39.2 (25.9–57.8)	39.2 (25.9–57.8)	39.2 (32.4–57.8)
Depression	55.4 (41–73.3)	41 (41–71.6)	41 (41–60.5), 11	41 (41–55.3)◻
Pain interference	61.3 (41.6–75.6)	57.5 (41.6–75.6)	54.8 (41.6–66.7)+	41.6 (41.6–59.7)◻
Physical function	34.5 (23.5–61.5)	37.5 (23.5–61.5)	39.8 (27–61.5)	43.3 (30.1–61.5)

*   One patient did not complete the inventory (N=18)
+   One patient did not complete the inventory (N=17)
◻   Two patients did not complete the inventory (N=5)
◻   One patient did not complete the inventory (N=6)

Figure 1: LITE SABR M1 Consort diagram

24 patients screened for eligibility

24 patients eligible for Lattice SBRT
    4 patients completed simulation but did not start treatment
        1 patient underwent simulation, but elected to pursue alternative therapy prior to starting SBRT
        1 patient elected to pursue hospice prior to starting SBRT
        1 patient died from respiratory failure prior to starting SBRT
        1 patient died from tumor cachexia prior to starting SBRT

20 patients completed 5 fractions Lattice SBRT

12 patients completed 90-day follow-up
    4 patients died within 90 days of completing Lattice SBRT (death unrelated to therapy)
    3 patients enrolled on hospice and withdrew from study
    1 patient progressed and withdrew from study

Figure 2 <u>Proportion of patients reporting PRO-CTCAE grade 0-1 and grade 2+ at baseline and 14 days.</u> A site-specific (thorax, gastrointestinal/pelvic) PRO-CTCAE inventory was assessed at each time point per protocol. 3A shows composite scores for pain, shortness of breath (SOB), nausea, vomiting, Anxiety, and fatigue for patients with thoracic tumors. 3B shows composite scores for pain, gastrointestinal (GI) and genitourinary symptoms (GU), anxiety, and fatigue.

Supplementary Materials

Supplementary Figure 1

Supplementary Figure 2

Change in GTV on follow up imaging. Diagnostic images were fused with CT Sim data in the TPS to contour tumors and measure volumetric change after completion of Lattice SBRT. Change relative to baseline GTV_2000 is shown against time (in months).

Supplementary Table 1
PRO-CTCAE data for all patients at all follow-up time points. Patients are grouped by treatment area (thoracic vs abdominal/pelvic). An entry of -- indicates that the PRO was not collected at that time point due to death, loss to follow-up, or withdrawal from study.

Thoracic

	Days after completion	Composite Score						
		Pain	SOB	Cough	Nausea	Vomiting	Anxiety	Fatigue
Patient 1	Baseline	1	3	2	0	0	3	2
	14	--	--	--	--	--	--	--
	30	0	1	1	0	0	2	3
	90	--	--	--	--	--	--	--
Patient 2	Baseline	2	1	--	--	--	0	2
	14	--	--	--	--	--	--	--
	30	1	0	--	--	--	0	0
	90	1	0	--	--	--	0	0
Patient 3	Baseline	3	2	--	1	0	2	2
	14	2	0	--	1	0	--	2
	30	--	--	--	--	--	--	--
	90	--	--	--	--	--	--	--
Patient 4	Baseline	0	0	--	0	0	0	0
	14	0	0	--	0	0	0	0
	30	0	0	--	0	0	0	0
	90	0	0	--	0	0	0	0
Patient 5	Baseline	0	0	--	0	0	0	0
	14	0	0	--	0	0	0	0
	30	0	0	--	0	0	0	0
	90	0	0	--	0	0	0	0
Patient 8	Baseline	0	0	--	1	0	2	0
	14	2	2	--	2	2	2	2
	30	0	1	--	0	0	1	1
	90	1	1	--	2	2	1	1
Patient 10	Baseline	--	--	--	--	--	--	--
	14	--	--	--	--	--	--	--
	30	--	--	--	--	--	--	--
	90	--	--	--	--	--	--	--
Patient 12	Baseline	2	0	--	0	0	1	3
	14	3	1	--	0	0	3	3
	30	--	--	--	--	--	--	--
	90	--	--	--	--	--	--	--
Patient 15	Baseline	3	3	--	2	2	3	2
	14	2	1	--	2	2	3	1

	30	--	--	--	--	--	--	--
	90	--	--	--	--	--	--	--
Patient 16	Baseline	--	--	--	--	--	--	--
	14	--	--	--	--	--	--	--
	30	--	--	--	--	--	--	--
	90	--	--	--	--	--	--	--

Abdominal/Pelvic

		Composite Score			
	Days after completion	Pain	GI/GU	Anxiety	Fatigue
Patient 6	Baseline	0	--	0	2
	14	0	--	1	2
	30	1	--	2	2
	90	0	--	1	2
Patient 7	Baseline	--	--	--	--
	14	--	--	--	--
	30	--	--	--	--
	90	--	--	--	--
Patient 9	Baseline	3	2	3	3
	14	3	1	2	3
	30	0	3	2	3
	90	--	--	--	--
Patient 11	Baseline	3	1	0	0
	14	2	2	0	2
	30	2	1	0	2
	90	--	--	--	--
Patient 13	Baseline	4	2	3	3
	14	2	0	1	2
	30	--	--	--	--
	90	--	--	--	--
Patient 14	Baseline	4	1	2	1
	14	3	0	2	2
	30	--	--	--	--
	90	3	3	2	0
Patient 17	Baseline	2	3	1	2
	14	2	3	1	1
	30	1	2	1	0
	90	--	--	--	--
Patient 18	Baseline	--	--	--	--
	14	--	--	--	--
	30	--	--	--	--

	90	--	--	--	--
Patient 19	Baseline	2	2	0	3
	14	2	0	0	2
	30	--	--	--	--
	90	--	--	--	--
Patient 20	Baseline	4	4	2	4
	14	2	3	1	3
	30	3	--	2	2
	90	1	1	2	2