Effects of immunosuppressive therapy reduction and early post-infection graft function in kidney transplant recipients with COVID-19

Gaetano Alfano[1,2*], Francesca Damiano[1*], Francesco Fontana[1], Camilla Ferri[3], Francesco Giaroni[3], Andrea Melluso[3], Martina Montani[3], Niccolò Morisi[3], Tei Lorenzo[3], Jessica Plessi[3], Silvia Giovanella[2,3], Giulia Ligabue[3], Giacomo Mori[2], Giovanni Guaraldi[4], Riccardo Magistroni[1,2], Gianni Cappelli[1,2]

1 Nephrology, Dialysis and Transplant Unit, University Hospital of Modena, Modena, Italy

2 Clinical and Experimental Medicine Ph.D. Program, University of Modena and Reggio Emilia, via del Pozzo 71, 41124 Modena, Italy

 3 Surgical, Medical and Dental Department of Morphological Sciences, Section of Nephrology, University of Modena and Reggio Emilia, via del Pozzo 71, 41124 Modena, Italy

4 Clinic of Infectious Diseases, University Hospital of Modena, Modena, Italy

* both authors contributed equally as first author

Corresponding author:

Gaetano Alfano, MD

Surgical, Medical and Dental Department of Morphological Sciences, Section of Nephrology, University Hospital of Modena and Reggio Emilia, via del Pozzo, 71

41124 - Modena (Italy)

E-mail:gaetano.alfano@unimore.it

ORCID: 0000-0003-0591-8622

Short title

Kidney transplantation & COVID-19

Authorship credit

Conception: Gaetano Alfano and Francesca Damiano

Collection of data Camilla Ferri, Francesco Giaroni, Andrea Melluso, Martina Montani, Niccolò Morisi, Tei Lorenzo, Jessica Plessi

Analysis and interpretation of data: Gaetano Alfano, Francesco Giaroni, Francesca Damiano

Drafting the article: Gaetano Alfano, Francesco Fontana, Silvia Giovanella, Giulia Ligabue, Giacomo Mori

Intellectual Contribution: Francesco Fontana Gianni Cappelli, Giovanni Guaraldi

Revising the article: Gianni Cappelli, Giovanni Guaraldi

Approval of the version to be published: all authors

Acknowledgments

Special thanks are due to Dr. Roberto Pulizzi, Francesc Facchini and Marco Ballestri, the nephrologists involved in the "Kideny Transplant Program" and

Laura Bonaretti and all nurses of the "Kidney Transplantation Outpatient Clinic" at the University Hospital of Modena for their precious support in managing KT recipients.

Text word count: 2189

Abstract word count: 247

Number of references: 34

Number of tables: 3

Abstract

Kidney transplant (KT) recipients with COVID-19 are at high risk of poor outcomes because of comorbidities and long-term immunosuppressive therapy (IST). There are little data on the effect of IST reduction and early graft function after COVID-19.

We conducted a retrospective study on 45 consecutive KT recipients followed at the University Hospital of Modena who tested positive for COVID-19 by RT-PCR analysis. We detailed clinical management and outcomes of these patients.

Median age of patients was 56.1 (interquartile range, [IQR] 47.3-61.1) years with a predominance of male (64.4%) and patients of Caucasian origin (91.1%). Kidney transplantation vintage was 10.1 (2.7-16) years, and more than half (55.6%) was on triple IST. Therapeutic management included antimetabolite (62.8%) and calcineurin inhibitor withdrawal (22.2%), and suspension of IST in severely ill patients.

Of the 45 patients, 88.9% became symptomatic and 40% required hospitalization. Overall mortality accounted for 17.8% (n=8). There were no differences in outcomes between full- and reduced-dose IST at the end of follow-up. Overall, early graft function after COVID-19 showed a stable and unmodified kidney function in 95% of survivors. Risk factors for death were age (odds ratio [OR]: 1.19; 95% CI: 1.01-1.39), and years spent on immunosuppression (OR: 1.96; 95% CI: 0.38-10.03-4.9). One patient experienced symptomatic reinfection with COVID-19 after primary infection and anti-SARS-CoV-2 mRNA vaccine.

COVID-19 impacted the graft and general survival of KT recipients. Short-term graft outcome after COVID-19 was favorable in most survivors. Age and transplantation vintage are independent predictors of death in our patients.

Keywords: kidney transplantation; immunosuppression; mortality; graft function; SARS-CoV-2

Introduction

Since SARS CoV-2 was identified in December 2019, the pandemic spreads quickly all around the world with a disruptively impact on social and economic life. This virus yielded several new challenges to our healthcare systems that had to copes with an increase of morbidity and mortality among the most vulnerable populations.[1] Kidney transplant (KT) recipients is a subset of the population at high risk of severe COVID-19 relate consequences due to comorbidities, consequences of chronic kidney disease (CKD) and the burden of immunosuppressive therapy (IST), which in some subjects include also therapies administered before kidney transplantation to treat glomerulonephritis or underlying autoimmune diseases. [2]

Data collected so far reported that transplant recipients were at higher risk of morbidity and mortality compared to the general population [3,4]. Despite the great emphasis on IST reduction to face the potentially lethal consequences of COVID-19, no confirming data supports its beneficial effect in terms of survival or clinical and laboratory manifestations. Additional uncertainty arises from recent literature when the tempered immune response is suggested to prevent COVID-19– induced systemic inflammatory syndrome leading acute respiratory distress syndrome and sepsis. Furthermore, few data are available regarding early graft outcomes after COVID-19 in survivors.[5]. Multiple causes, including kidney tropism of SARS-CoV-2 [6] may be responsible for non-reversible episodes of AKI[7] that can severely impair graft survival. Lastly, a concerning issue is the hyporesponsiveness to anti-SARS-CoV-2 vaccination[8]. Numerous studies have confirmed that KT recipients have a blunted immune response to mRNA vaccine[9]. Only 48% of patients was able to mount a protective serologic response to SARS-CoV-2[10]. Caillard et al [11]reported that about one-third of kidney transplant patients had severe manifestations, including a fatal outcome, after completing vaccination with mRNA. This group of patients is therefore expected to remain vulnerable to the severe complications of COVID-19 until new strategies will be implemented to reduce the susceptibility of these subjects.

Considering all the uncertainties in the management of KT recipients and the still current risk of infection in this cohort of patients we report our experience in managing KT recipients with

COVID-19. In particular, we will focus on the impact of early IST reduction and early graft function after the resolution of the infection.

Material and methods

Kidney transplant outpatient clinic

This kidney transplant outpatient clinic follows more than 500 KT recipients, including combined liver and pancreas-kidney transplantation. Outpatient service was delivered by a senior nephrologist with experience in kidney transplantation, one fellow and three nurses. A 24-h, 7/7 days per week service was available for KT recipients in case of kidney-related pathologic process (anuria, fluid overload) or infections. This service was also offered to the subjects transplanted in our Center but living far from it.

During COVID-19 all the patients were instructed to call the clinic in case of COVID-19 symptoms. Despite the reduction of non-essential healthcare services, our outpatient clinic continued to deliver care for KT recipients adopting all the containment measures (triage at entry, masking, social distance and cleaning hands) to prevent COVID-19 diffusion. A telephonic triage was performed for all patients before reaching the hospital to intercept some paucisymptomatic patients.

Patients with symptoms were invited to perform RT-PCR on nasal swab and were visited in a dedicated room to assessed vital parameters and clinical conditions. According to the severity of symptoms, patients were sent home or to the emergency room. To reduce the workload to the emergency room, patients were managed as an outpatient unless they developed severe symptoms requiring hospital admission. Remote patient monitoring was performed principally with phone calls and email.

According to our internal protocol and taking into account European expert opinions[12,13], immunosuppressive was modulated as follow:

- for asymptomatic or mild COVID-19 (e.g., mild upper respiratory and/or gastrointestinal symptoms, temperature < 38°C without dyspnea) in patients on triple therapy (calcineurin-inhibitors [CNI] + mycophenolate acid [MPA]/ azathioprine [AZA] + steroids), MPA or AZA was

withdrawn, and a dual therapy (CNI + steroid) was continued. If patient the patient was on dual therapy (CNI+ mammalian target of rapamycin inhibitor [MPA/mTOR-i]), MPA/mTOR was withdrawal and replaced with a low dose of steroids.

-for moderate and severe COVID-19 all immunosuppressors but steroid were stopped

COVID-19 population

The study population was comprised of kidney transplant recipients with COVID-19 followed in the nephrology outpatient clinic at the University Hospital of Modena. This outpatients clinic delivers care to all patients after three months from kidney transplantation.

We retrospectively reviewed the electronic charts of all KT recipients with COVID-19 from March 7, 2019, to June 25, 2021. The diagnosis of COVID-19 was performed through reverse transcriptase-polymerase chain reaction (RT-PCR) assay on a nasal swab. We excluded patients aged<18 years. Kidney function was estimated by glomerular fraction rate (eGFR) using CKD-EPI equation. Missing occurred for patients admitted to a hospital located far from our Center.

This study has been authorized by the local Ethical Committee of Emilia Romagna (n. 839/2020). The study protocol complies with the guidelines for human studies and includes evidence that the research was conducted ethically in accordance with the World Medical Association Declaration of Helsinki.

Statistical analysis

Baseline characteristics were described using median (interquartile range [IQR]), mean and standard deviation (SD), or frequencies, as appropriate. The chi-square or Fisher's test, and student's t-test were used to compare categorical and continuous variables between groups, respectively. Univariate and multivariate logistic regression were performed to test the association between mortality and baseline patient characteristics. Variables that were significant on univariate analysis (P =<0.1) were entered into the multivariate model to identify independent predictors (P= < 0.05). Results were expressed as odds ratios (OR) and 95% confidence intervals (CI).

Univariate and multivariate logistic regression analysis determined risk factors for death. A P value of <0.05 was considered statistically significant. All statistical analyses were performed using SPSS® statistical software.

Results

Population

From the spread of COVID-in Italy, 45 KT recipients followed in at our Center contracted COVID-19. The demographic and clinical characteristics of the patients were detailed in Table 1. This group of patients included two (4.4%) combined **liver**-**kidney** and one (2.2%) **heart**-**kidney** transplant recipient.

Age of patients ranged from 19.2 to 83.5 years and median was 56.1 (IQR, 47.3-61.1) years. COVID-19 was prevalent in males (64.4%) and occurred after a median of 10,1 (2,7-16,01) years from transplantation.

Before COVID-19 infection serum creatine sCr measured 1.45 (IQR 1.1-1.8) mg/dl corresponding a median eGFR of 48.4 ml/min. At the time of COVID-19 diagnosis, 55.6% of the patients was in triple standard IST. Reduction of IST was practiced in more than half (62.8) of subjects. MPA (88.8%), CNI or mTOR-i (22.2%) were the most frequent discontinued agents.

Hospitalization and death rate in patients who underwent IST reduction were 51.8% and 14.8%, respectively. However, despite IST reduction, hospitalization (P=0.71) and death (P=1) rates were not different compared to the full-dose IST group. Overall, forty patients (88.9%) developed symptoms of COVID-19 and 18 of them (45%) required hospitalization. Overall, eight patients (17.8%) died for COVID-19.

Univariate and multivariate logistic regression was performed to detect predictors of mortality (Table 2). Multivariate analysis found that age (OR=1.19 [95%CI 1.01-1.39]; P=0.034) and years spent on immunosuppressive therapy (OR=1.17 [95%CI 1.01-1.35]; P=0.040) were associated with mortality in this group of patients.

Among survivors, one patient with a CKD stage 4 (GFR=20 ml/min) before SARS-CoV-2 infection, developed irreversible graft failure requiring HD. In one patient presenting with de-novo proteinuria (4100 mg/die) after resolution of COVID-19, graft biopsy revealed IgA glomerulonephritis (available data to do allow to classify these histological findings as de-novo or recurrent IgA glomerulonephritis). Lastly, patients experienced symptomatic reinfection with COVID-19 after primary infection and anti-SARS-CoV-2 mRNA vaccine. Early post-COVID-19 follow-up of 25 (78.3%) patients showed that pre- and post-COVID variations of sCr, eGFR and 24-hour proteinuria were not statistically significant in outpatients after resolution of COVID-19. A significantly lower sCr level (P=0.019) and eGFR (P=0.028) were measured after dismission in hospitalized patients. No differences were noted in the level of proteinuria. (Table 3)

Discussion

Numerous reports have alerted the scientific community regarding the unfavorable outcome of COVID-19 in patients with a reduced immune response. [1,14] The results of this study confirmed that COVID-19 poses KT recipients at high risk of severe consequences. In our cohort of KT recipients, COVID-19 carried a higher rate of symptoms, hospitalization and mortality compared to the general population[15,16].

We found that in a cohort of 45 KT recipients of 56.1 years with COVID-19, 40% of them developed severe symptoms requiring hospitalization. Overall mortality was 17.8%, higher than the national mortality reported in the general population, which ranges between 0.1-9.2% among countries worldwide and account for about 2% globally[17].

Since IST portends poor outcomes in patients with infections, a new therapeutic strategy was set to reduce the burden of IST in COVID-patients. In the attempt to reconstitute the immune system against SAR-CoV-2 infection, we systematically minimized anti-rejection therapy. All KT recipients, who communicated their COVID-19 positivity, were advised to withdraw at least antimetabolite agents (MFA or AZA). In hospitalized patients, IST was reduced or suspended according to the clinical

conditions of the patient. Nevertheless, hospitalization and death rate in the reduced-dose IST group was not dissimilar from full-dose IST group. A potential explanation for this result is that reduction of immunosuppressive therapy was not able to reconstitute an efficient immune response after the long-term burden of IST.

Although IST reduction did not lead to a favorable outcome, it is worth mentioning that the overall mortality estimated in our cohort was tendentially lower than that reported in several monocentric and multicentric studies where this approaching up to 32.5%[18–25]. Our results are in line with the population-based data on 1013 KT recipients affected by COVID-19 collected by French and Spain national registries, which reported a 28-day mortality of 20%[26]. In Brescia (Italy), Bossini et al.[23] reported a higher overall mortality rate among KT recipients (28%) during the first wave of the infection. Similar to our therapeutic strategy, they practiced immunosuppression cessation in all hospitalized patients and introduced or increased the glucocorticoid dose. At first glance, the causes of this different mortality rate are unknown. The different timing of enrollment made the two cohorts not perfectly comparable. The enrollment of all patients of the Brescia cohort during the first wave has probably magnified the risk of side effects of therapeutic regimens administered during the first wave[27,28] and the challenges to delivering the standard of care in an overwhelmed and unprepared hospital. Lastly, a lower age (56.1 vs 60 years) in our cohort of patients has probably contributed to lead to a better prognosis.

Multivariate analysis showed that the predictors of death were age and time elapsed on IST, in line with previous studies.

Age is widely associate with COVID-19 severity and death in KT recipients[29,30] as well as the general population [31]. Centers for Disease Control (CDC) claims that 8 out 10 COVID-19 death in the U.S. has been reported in adults aged more 65 years and the risk of hospitalization and death increases enormously with age.[32]

The effect of immunosuppression is still controversial in KT recipients[33]. Immunosuppression is known to dysregulate innate and adaptive immunity, exposing the patients to severe infections. On

the other hand, severe COVID-19 infection has been associated with a dysregulated inflammatory response (IL-6, IL-1, and chemokines) leading to ARDS and sepsis. The new insights support a promising role of immunosuppressants in the most severe case on COVID-19(i.e., tocilizumab, steroid)[34]. In our opinion, the interplay between immunosuppression and pathogenesis of COVID-19 should be contextualized with the stage of SARS-CoV-2 infection. The high rate of symptomatic disease is probably secondary to the host susceptibility due to immunodepression and the burden of CKD. If the IST is somewhat beneficial to reduce the pro-inflammatory release of cytokines in the second phase of infection, more studies are required to elucidate which molecular pathway is involved.

Lastly, we report a short-term good graft function in survival after COVID-19. These data indicate a stable early graft function (sCr and 24-hour proteinuria) in outpatients COVID-19 who were not hospitalized. Conversely, hospitalized KT recipients had a statistically significant improvement in renal function. As supported by Dacina et al[5] we speculate that lower sCr after the COVID-19 episode was due to the minimization or withdrawn of CNI, a 'drug holiday' without apparently dire consequences in terms of graft rejection for 95% of survivors.

Some limitations of the study should be enunciated. It is a retrospective study, with small sample size and a short follow-up after COVID-19. The small number of patients in full- and reduced-dose of IST group may have reduced the probability to observe an underlying difference between these two groups. Furthermore, we cannot exclude that in some cases, reduction of the IST occurred after a short delay from the diagnosis of COVID-19. However, all patients with symptoms or at risk of contagious underwent nasal swab as fast as possible in an ambulatory setting.

Conclusion

Reduction of immunosuppression did not reduce the risk of severe COVID-19 or death. Age and time spend on kidney transplantation are independent predictors of death in our patients. Short-term

follow-up after COVID-19 showed an excellent graft function in most survivors. Primary infection or vaccination do not exclude the risk of SARS-CoV-2infection in KT recipients.

Conflicts of interest

All the authors have declared no competing interest

Funding

This study was not funded

Bibliography

[1]   Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. Nature. 2020;584:430–436.

[2]   Khairallah P, Aggarwal N, Awan AA, et al. The impact of COVID-19 on kidney transplantation and the kidney transplant recipient - One year into the pandemic. Transpl Int Off J Eur Soc Organ Transplant. 2021;34:612–621.

[3]   Caillard S, Chavarot N, Francois H, et al. Is COVID-19 infection more severe in kidney transplant recipients? Am J Transplant. 2021;21:1295–1303.

[4]   Fisher AM, Schlauch D, Mulloy M, et al. Outcomes of COVID-19 in hospitalized solid organ transplant recipients compared to a matched cohort of non-transplant patients at a national healthcare system in the United States. Clin Transplant. 2021;35.

[5]   Elec AD, Oltean M, Goldis P, et al. COVID-19 after kidney transplantation: Early outcomes and renal function following antiviral treatment. Int J Infect Dis. 2021;104:426–432.

[6]   Cravedi P, Mothi SS, Azzi Y, et al. COVID-19 and kidney transplantation: Results from the TANGO International Transplant Consortium. Am J Transplant Off J Am Soc Transplant Am Soc Transpl Surg. 2020;20:3140–3148.

[7]   Bajpai D, Deb S, Bose S, et al. Recovery of kidney function after AKI due to COVID-19 in kidney transplant recipients. Transpl Int Off J Eur Soc Organ Transplant. 2021;

[8]   Benotmane I, Gautier-Vargas G, Cognard N, et al. Weak anti–SARS-CoV-2 antibody response after the first injection of an mRNA COVID-19 vaccine in kidney transplant recipients. Kidney Int. 2021;99:1487–1489.

[9]   Boyarsky BJ, Werbel WA, Avery RK, et al. Antibody Response to 2-Dose SARS-CoV-2 mRNA Vaccine Series in Solid Organ Transplant Recipients. JAMA. 2021;325:2204.

[10]   Benotmane I, Gautier-Vargas G, Cognard N, et al. Low immunization rates among kidney transplant recipients who received 2 doses of the mRNA-1273 SARS-CoV-2 vaccine. Kidney Int. 2021;99:1498–1500.

[11] Caillard S, Chavarot N, Bertrand D, et al. Occurrence of severe COVID-19 in vaccinated transplant patients. Kidney Int [Internet]. 2021 [cited 2021 Jun 5];0. Available from: https://www.kidney-international.org/article/S0085-2538(21)00509-3/abstract.

[12] Expert-opinion-on-ISD-in-Covid-19.pdf [Internet]. [cited 2021 May 26]. Available from: https://www.era-edta.org/en/wp-content/uploads/2020/04/Expert-opinion-on-ISD-in-Covid-19.pdf.

[13] Tsalouchos A, Salvadori M. La pandemia del nuovo coronavirus 2019 ed il trapianto renale. G Clin Nefrol E Dialisi. 2020;32:60–63.

[14] Myint PK, Carter B, Barlow-Pay F, et al. Routine use of immunosuppressants is associated with mortality in hospitalised patients with COVID-19. Ther Adv Drug Saf. 2021;12:2042098620985690.

[15] Cascella M, Rajnik M, Aleem A, et al. Features, Evaluation, and Treatment of Coronavirus (COVID-19). StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 [cited 2021 Jun 5]. Available from: http://www.ncbi.nlm.nih.gov/books/NBK554776/.

[16] Mappe Coronavirus [Internet]. [cited 2021 Jun 5]. Available from: https://mappe.protezionecivile.gov.it/it/mappe-emergenze/mappe-coronavirus.

[17] COVID-19 Map [Internet]. Johns Hopkins Coronavirus Resour. Cent. [cited 2021 May 27]. Available from: https://coronavirus.jhu.edu/map.html.

[18] Pérez-Sáez MJ, Blasco M, Redondo-Pachón D, et al. Use of tocilizumab in kidney transplant recipients with COVID-19. Am J Transplant Off J Am Soc Transplant Am Soc Transpl Surg. 2020;20:3182–3190.

[19] Aziz H, Lashkari N, Yoon YC, et al. Effects of Coronavirus Disease 2019 on Solid Organ Transplantation. Transplant Proc. 2020;52:2642–2653.

[20] Coll E, Fernández-Ruiz M, Sánchez-Álvarez JE, et al. COVID-19 in transplant recipients: The Spanish experience. Am J Transplant Off J Am Soc Transplant Am Soc Transpl Surg. 2021;21:1825–1837.

[21] Akalin E, Azzi Y, Bartash R, et al. Covid-19 and Kidney Transplantation. N Engl J Med [Internet]. 2020 [cited 2021 May 27]; Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7200055/.

[22] Fernández-Ruiz M, Andrés A, Loinaz C, et al. COVID-19 in solid organ transplant recipients: A single-center case series from Spain. Am J Transplant Off J Am Soc Transplant Am Soc Transpl Surg. 2020;20:1849–1858.

[23] Bossini N, Alberici F, Delbarba E, et al. Kidney transplant patients with SARS-CoV-2 infection: the brescia renal COVID task force experience. Am J Transplant [Internet]. 2020 [cited 2021 May 27]; Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7361317/.

[24] Cravedi P, Mothi SS, Azzi Y, et al. COVID-19 and kidney transplantation: Results from the TANGO International Transplant Consortium. Am J Transplant Off J Am Soc Transplant Am Soc Transpl Surg. 2020;20:3140–3148.

[25] Nair V, Jandovitz N, Hirsch JS, et al. COVID-19 in kidney transplant recipients. Am J Transplant [Internet]. 2020 [cited 2021 May 28]; Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7267603/.

[26] Jager KJ, Kramer A, Chesnaye NC, et al. Results from the ERA-EDTA Registry indicate a high mortality due to COVID-19 in dialysis patients and kidney transplant recipients across Europe. Kidney Int. 2020;98:1540–1548.

[27] Gérard A, Romani S, Fresse A, et al. "Off-label" use of hydroxychloroquine, azithromycin, lopinavir-ritonavir and chloroquine in COVID-19: A survey of cardiac adverse drug reactions by the French Network of Pharmacovigilance Centers. Therapies. 2020;75:371–379.

[28] Izcovich A, Siemieniuk RA, Bartoszko JJ, et al. Adverse effects of remdesivir, hydroxychloroquine, and lopinavir/ritonavir when used for COVID-19: systematic review and meta-analysis of randomized trials [Internet]. Infectious Diseases (except HIV/AIDS); 2020 [cited 2021 May 27]. Available from: http://medrxiv.org/lookup/doi/10.1101/2020.11.16.20232876.

[29] Coll E, Fernández-Ruiz M, Sánchez-Álvarez JE, et al. COVID-19 in transplant recipients: The Spanish experience. Am J Transplant Off J Am Soc Transplant Am Soc Transpl Surg. 2021;21:1825–1837.

[30] Oto OA, Ozturk S, Turgutalp K, et al. Predicting the outcome of COVID-19 infection in kidney transplant recipients. BMC Nephrol. 2021;22:100.

[31] Levin AT, Hanage WP, Owusu-Boaitey N, et al. Assessing the age specificity of infection fatality rates for COVID-19: systematic review, meta-analysis, and public policy implications. Eur J Epidemiol. 2020;35:1123–1138.

[32] CDC. COVID-19 and Your Health [Internet]. Cent. Dis. Control Prev. 2020 [cited 2021 May 27]. Available from: https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/older-adults.html.

[33] Chavarot N, Gueguen J, Bonnet G, et al. COVID-19 severity in kidney transplant recipients is similar to nontransplant patients with similar comorbidities. Am J Transplant. 2021;21:1285–1294.

[34] Pérez-Sáez MJ, Blasco M, Redondo-Pachón D, et al. Use of tocilizumab in kidney transplant recipients with COVID-19. Am J Transplant Off J Am Soc Transplant Am Soc Transpl Surg. 2020;20:3182–3190.

Table 1. Demographics and clinical characteristics of KT recipients

Variable	All patients*
	(n=45)
Age, years	56.1 (47.3-61.1)
Range	19.2-83.5
Males, n. (%)	29 (64.4)
Race/ethnicity	
White	41 (91.1)
Black	4 (8.9)
Transplant vintage	10.1 (2.7-16.01)
sCr pre-COVID-19	1.45 (1.18-1.84)
eGFR pre-COVID-19	48.4 (36-64)
24-h proteinuria, mg/dl	87.4 (0.52-188.5)
Immunosuppressive therapy	
CNI	39 (86.7)
mTOR-i	8 (17.8)
MPA	31 (68.9)
Steroid	36 (80)
Triple therapy	25 (55.6)
Double therapy	19 (42.2)
Monotherapy	1 (2.2)
Reduction IS therapy	27 (62.8)
MPA withdrawal	24 (88.8)
CNI or mTOR-i withdrawal	6 (22.2)
Increase steroid	9 (20)
Comorbidities	
HIV, HCV or HBV	6 (13.3)
Diabetes	5 (11.1)
Neoplasia	10 (22.2)
Graft rejection	4 (8.9)
CVD#	12 (26.7)
Autoimmune	4 (8.9)
Previous severe infection	13 (28.9)
Symptomatic COVID-19	40 (88.9)
Hospitalization	19 (42.2)
Post-COVID-19 follow-up	70.5 (51-109)
Mortality	8 (17.8)

CNI denotes calcineurin-inhibitors; CVD, cardiovascular disease; eGFR, estimated glomerular filtration renal; HCV, hepatitis C; HBV, hepatitis B; MPA, mycophenolate acid; mTOR-I, mammalian target of rapamycin inhibitor; sCr, serum creatinine

Table 2. Logistic regression to identify predictors of AKI

		Univariate			Multivariate			
	Variable	**OR**	**CI (95%)**		*P*-value	OR	CI 3(95%)		*P*-value
Sex									
Male		4,40	0,78	24,81	0,093				
Age (1-yr increase)		1,11	1,02	1,22	**0,016**	1,19	1,01	1,39	**0,034**
KT vintage (1-yr increase)		1,10	1,00	1,21	0,053	1,17	1,01	1,35	**0,040**
Steroid-based IST		1,93	0,21	18,08	0,564				
Reduction IST		1,33	0,26	6,869	0,74				
Increase of steroid		0,52	0,06	4,85	0,564				
Triple IST		0,51	0,10	2,620	0,42				
Double IST		1,96	0,38	10,026	0,42				
GFR		0,99	0,95	1,026	0,57				
GFR< 45ml/min		1,47	0,32	6,80	0,624				
GFR 45-59 ml/min		0,68	0,15	3,16	0,624				
sCr		1,33	0,26	6,87	0,736				
Graft rejection		1,52	0,14	16,91	0,732				
Autoimmune disease		0,00	0,00		0,999				
HIV/HCV/HBV		2,58	0,38	17,43	0,330				
Previous sever infection		0,73	0,13	4,19	0,722				
Diabetes		1,11	0,11	11,49	0,932				
Neoplasm		1,12	0,19	6,70	0,897				
Cardiovascular disease		1,73	0,34	8,76	0,506				

eGFR denotes estimated glomerular filtration rate; HCV, hepatitis C; HBV, hepatitis B; IST, immunosuppressive therapy; MPA, mycophenolate acid; mTOR-I, mammalian target of rapamycin inhibitor*;* sCr, serum creatinine.

Table 3. Early graft function post-COVID-19 in hospitalized and non-hospitalized KT recipients

	Non-hospitalized patients			Hospitalized patients		
	Pre-COVID-19	Post-COVID-19	P-value	Pre-COVID-19	Post-COVID-19	P-value
sCr. mg/dl	1.31 (1.2-1.76)	1.33 (1.08- 1.7)	0.85	1.44 (1.1-1.8)	1.32 (1-1.7)	**0.019**
eGFR. ml/min	48.8 (40.5-62.1)	56.7 (41.5-67)	0.25	48. 4 (36-64)	56.7 (41.5-67)	**0.028**
24-h proteinuria	102 (6.2-205)	89.4 (37.2-246.4)	0.08	102 (6.2-205)	89.4 (37.2-246.4)	0.29