Modern contraceptive use among adolescent girls and young women in Benin: a mixed-methods study.

AUTHORS :

Noudéhouénou Crédo Adelphe AHISSOU [1,2]

Lenka BENOVA [3]

Thérèse DELVAUX [3]

Charlotte GRYSEELS [3]

Jean-Paul DOSSOU [2,3]

Sourou GOUFOUDJI [2]

Lydie KANHONOU [2]

Christelle BOYI [2]

Armelle VIGAN [2]

Koen PEETERS [1,3]

Miho SATO [1]

Mitsuaki MATSUI [1]

AFFILIATIONS:

1 Department of Global Health, Nagasaki University School of Tropical Medicine and Global Health
Nagasaki 852-8523, Japan

2 Centre de Recherche en Reproduction Humaine et en Démographie
Cotonou, Benin

3 Department of Public Health, Institute of Tropical Medicine Antwerp
Antwerp, Belgium

CORRESPONDING AUTHOR

Mitsuaki MATSUI

e-mail: mmatsui@nagasaki-u.ac.jp

c/o Nagasaki University School of Tropical Medicine and Global Health
Sakamoto 1-12-4, Nagasaki 852-8523, Japan

Word count – 5082 words

ABSTRACT

Objectives

The study aimed to assess the determinants of modern contraceptive method use among young women in Benin.

Design

A mixed-methods design.

Setting and participants

We used the Benin 2017-18 Demographic and Health Survey datasets for quantitative analysis. Data collection was conducted using multiple-cluster sampling method and through household survey. Qualitative part was conducted in the city of Allada, one of the Fon cultural capitals. The participants were purposively selected.

Outcomes

Contraceptive prevalence rate, unmet need for modern method, and percentage of demand satisfied by a modern method for currently married and sexually active unmarried women were measured in the quantitative part. Access barriers and utilization of modern methods were assessed in the qualitative part.

Results

Overall, 8.5% (95%CI: 7.7-9.5%) among young women ages 15 to 24 were using modern contraceptives and 13% (95%CI: 12.1-14.0%) among women ages 25 or more. Women 15-24 had a higher unmet need, and a lower demand satisfied by modern contraceptive methods compared to women aged 25 or more. 60.8% (56.9-64.7%) of all unmarried young women had unmet need for modern contraceptives. Young women were more likely to use male condoms which they

obtain mainly from for-profit outlets, pharmacies, and relatives. The factors associated with demand satisfied by a modern method were literacy, being unmarried, knowing a greater number of modern contraceptive methods, and experiencing barriers in access to health services. On the other hand, the qualitative study found that barriers to using modern methods include community norms about pre-marital sexual intercourse, perceptions about young women's fertility, spousal consent, and the use of non-modern contraceptives.

Conclusion

Contraceptive use is low among young women in Benin. The use of modern contraceptives is influenced by socio-demographic factors and social norms. Appropriate interventions might promote comprehensive sexuality education, increase community engagement, provide youth-friendly services, and address gender inequalities.

KEYWORDS

Contraception, modern method, demand satisfied, unmet need, adolescent, Benin

Strengths and limitations of this study

- This study used a nationally representative survey data which allows generalizability at the national level.
- We used a mixed methods design which enables us to compare and discuss the large-scale quantitative data with qualitative information from the ground.
- The quantitative information could have been affected by recall errors and biases.

INTRODUCTION

Despite efforts toward making access and use of contraceptive services a basic reproductive right for all women, many countries still face high rates of unintended and unwanted pregnancies, [1, 2]. In sub-Saharan Africa alone, about 14 million unintended (unwanted or mistimed) pregnancies occur every year,[3]; and adolescent girls and young women 15 to 24 years old are the most vulnerable group,[4]. The reasons behind this situation include the high prevalence (about 70%) of sexually active young women with low utilization of effective contraceptive methods (less than 10%),[4]. Additionally, unmarried sexually active adolescents are likely to have a high unmet need for contraception, which increases their risk of unintended pregnancies,[5, 6]. In developing countries, nearly half (49%) of pregnancies are unintended among adolescent girls of 15 to 19 years old,[7]. Adolescence is viewed as the starting point in the continuum of care for reproductive, maternal, neonatal, and child health; and is a phase when poor access and utilization of contraception are likely to result in poor health outcomes across the continuum of care,[8, 9]. Early and unintended pregnancies result in increased risks of maternal mortality and morbidity, premature births, low birth weight, unsafe abortions, and social consequences such as stigmatization, school dropout, and poverty,[10, 11].

In Benin, although the use of modern contraceptives has been slowly increasing since 2006, it is still relatively low. The modern contraceptive prevalence among all women reached 12% in 2018 compared to 6% in 2006,[12]. At the same time, almost half (48%) of all adolescent girls age 15 to 19 are sexually active, and one in five girls has already had a child or is pregnant,[12]. According to the government of Benin (2017), only 5.4% of women ages 15 to 24 were using modern contraceptive methods in 2017,[13]. Recent data

showed a total fertility rate of 5.7 among all women of reproductive ages 15 to 49 years old, and the modern contraceptive prevalence rate (mCPR) was estimated at 12%,[12]. Of all pregnancies in the country, 19% were unintended,[14]; and in 2017, both the maternal mortality ratio and infant mortality rate remained high at 397 per 100,000 live births, and 30 per 1,000 live births, respectively,[6, 12].

Existing studies on the use of modern contraceptive methods in Benin largely reported on women of reproductive age as a whole, rather than focusing on specific age groups, [15-20]. MacQuarrie suggested that young women should be studied separately, as they do not have the same needs for or access to contraception as adult women,[21]. However, for Benin, the literature is sparse on the use of reproductive health services among adolescent girls and young women. The only study which focused on young women in Benin was quantitative,[22], and did not explore in-depth factors that affect decision-making such as adolescent girls and young women's perceptions, preferences, and interactions with peers and providers.

Therefore, this study aimed to assess contraceptive use, perceptions, and practices among adolescent girls and young women in Benin. Specifically, we used secondary quantitative data to examine the rates of modern method use, unmet needs for modern contraception, and to evaluate the association between various factors and the demand satisfied for a modern contraceptive among sexually active young women. We then used qualitative methods to further explore the socio-cultural factors influencing the use of modern contraceptive methods among young women of ages 18 to 24 years.

METHODS

Study design

The study was conducted using a sequential explanatory mixed-methods design where the collection and analysis of quantitative data were followed by the collection and analysis of qualitative data. The qualitative results thus aim to explain and interpret the findings of the quantitative study,[23].

Quantitative strand

Data source

We used the most recent Benin Demographic and Health Survey (BDHS), conducted between November 2017 and February 2018,[12]. The BDHS2017-18 was a nationally representative cross-sectional survey conducted by the National Institute of Statistics and Economic Analysis and funded by the USAID. The BDHS 2017-18 employed a two-stage stratified cluster sampling during which data were collected with standardized questionnaires.

Study population

Included in the study were women ages 15 to 49 years at the time of the survey, residing in sampled households, and who agreed to respond to the BDHS 2017-18. We defined adolescent girls as those 15 to 19 years old and young women as those 20 to 24 years. In descriptive analyses, we compared indicators of contraceptive use in each of the age groups to all women 25 years and older. In the analysis of the determinants of modern contraceptive methods use, we focused on women who reported to be sexually active (married and unmarried) because they are exposed to

119 the risk of pregnancy, and we used the combined age groups 15 to 24 years due to
120 sample size considerations.

121 Definitions

122 Some definitions of indicators from the BDHS 2017-18 were adopted, while others
123 were generated or recategorized as necessary for our study-specific objectives.

124 *Modern method*: Respondents were asked which method of contraception they were
125 using at the time of the survey. If they selected more than one method (for example,
126 implant and condom), only the most effective method was noted in the dataset.
127 Based on the definition of Hubacher and Trussell (2015), we considered modern
128 contraceptive methods to include: oral contraceptive pills, intrauterine devices
129 (IUD), injectables, diaphragm, male/female condoms, female/male sterilization,
130 implants/Norplant, foam, or jelly, and emergency contraception,[24]. All other
131 methods were grouped as traditional methods.

132 *Current modern method users:* Women who reported using any modern
133 contraceptive method at the time of the survey were considered modern method
134 users.

135 *Contraceptive prevalence rate*: The proportion of women who reported using a
136 method at the time of the survey. This was estimated for modern contraceptive
137 methods among different groups of women based on their marital status.

138 *Sexual activity:* Women who reported sexual intercourse 30 days preceding the survey were
139 categorized as sexually active.

140 *Source of modern method:* The source of the modern method is the most recent
141 source of provision among current modern methods users. We grouped sources of

142 modern methods into public sources (government hospital, government health
143 center, mobile clinic, government community health worker, and other public
144 sectors), non-public sources (private hospital/clinic, private clinic or doctor, and
145 private mobile clinic), for-profit outlets (pharmacy, family planning clinic, and shop),
146 relatives and other sources (friends/relatives). A second categorization in the
147 analysis was done based on the theoretical capacity model of providers proposed by
148 Radovich et al.,[22]. It divides the providers into two groups, 'comprehensive' or
149 'limited'. Comprehensive providers are public or private providers that can provide
150 both short- and long-acting methods. Limited providers include service locations,
151 for-profit outlets, partners, and friends, which do not generally provide long-acting
152 methods.

153 *Unmet need for modern methods:* Unmet need for contraception describes women
154 who are at risk of pregnancy (married or sexually active), do not wish to be pregnant,
155 but are not using a modern contraceptive method. We applied an algorithm
156 proposed by Bradley et al. (2012) to identify 'unmet need for limiting' and 'unmet
157 need for spacing' among married women and sexually active unmarried women,[25].
158 Also, we adopted the definition of FP2020 (2021) to compute the 'unmet need for
159 modern methods',[4]. Because women using a traditional method are less protected
160 against pregnancy, they were included with the non-users and considered to have an
161 unmet need for modern contraception,[26]. Unmet need was estimated among the
162 denominator of all women regardless of whether they were in need of family
163 planning or not.

164 *Percentage of demand satisfied by a modern method:* This is the percentage of women at
165 risk of pregnancy and who do not wish to become pregnant, who are using modern
166 methods. The numerator is current modern method users, and the denominator is all
167 women in need of family planning (current users of any contraceptive methods and women
168 with unmet need for family planning).

169 *Existence of barriers to access to health services:* Women were asked whether getting
170 permission to go, getting money to pay for services, the distance to travel, or going alone
171 to a health center were barriers in their access to health services. We created a binary
172 variable categorizing women who said that they had none of these issues versus those who
173 had one or more.

174 *Core indicators:* The core indicators include 'contraceptive prevalence rate', 'unmet
175 need for modern method', and 'percentage of demand satisfied by a modern method'
176 for currently married and sexually active unmarried women. Demand satisfied by a
177 modern method among women ages 15 to 24 in need of family planning was the main
178 outcome variable in the logistic regression.

179 *Explanatory variables:* We included in the logistic regression the respondents'
180 socio-demographic characteristics (literacy, ethnicity, religion, marital status
181 [married/cohabiting at the time of the survey or no], parity, and household wealth
182 quintile); knowledge of modern contraceptive methods; and the existence of
183 barriers to access health services. Definitions and categories of these independent
184 variables are shown in Supplementary Materials.

185 Data analysis

186 Data analysis was carried out using Stata (version 13, Stata Corp., Texas, USA). We
187 conducted a descriptive analysis to describe the socio-demographic characteristics
188 of the study population within three age groups (15 to 19, 20 to 24, 25 or older). We
189 used logistic regression to conduct bivariate and multivariable analyses to assess the
190 associations between the explanatory variables and modern method use among
191 sexually active women of ages 15 to 24 years in need of contraception. We retained
192 all explanatory variables in multivariable logistic regression. We used the *svyset*
193 command to adjust for clustering, stratification, and the sampling weights of
194 individual women in all analyses.

195 Missing data

196 There was no missing data in any variables used except for the source of the modern
197 method. Of condom users 15 to 49 years old, 26 out of 1805 (1.4%) had a missing
198 source of method; and 19 of them were 15 to 24 years old. We grouped them within
199 the "pharmacy" category, which is the most common source for condom users.

200

201 **Qualitative strand**

202 Study site and population

203 Data collection was conducted in the city of Allada, one of the Fon cultural capitals –
204 the largest ethnic group in Benin - and located in southern Benin about 50 km from
205 Cotonou, the largest city of Benin. The region offers the advantage of being
206 semi-rural with diverse religions and care practices such as spiritual care,
207 alternative care, and biomedicine,[27]. Spiritual healing involves using mystical
208 practices, advice prescribed by traditional healers which are viewed as effective in

209 treating health problems. Such practices could include consulting deities,
210 performing rituals, using leaves, or consuming herbal teas. Alternative care refers to
211 the use of herbal infusions and powders, as well as the use of certain leaves and
212 barks to address health issues.

213 We restricted the target population of the qualitative study to women ages 18 to 24
214 years old due to consent issues with minors. We collected data from two main
215 groups: young pregnant women from whom we wanted to learn about access barriers
216 and utilization of modern methods based on their experiences, particularly in cases
217 where pregnancy might be unintended. They were recruited after pre or postnatal
218 visits, or consultation about family planning services in the main public health
219 center of Allada.

220 Healthcare providers helped us identify women that met our inclusion criteria
221 (pregnant young women). We later approached them face-to-face outside the health
222 facilities after consultations. Non-pregnant adolescent girls and young women were
223 recruited in schools through purposive sampling, and in the community using
224 convenience and snowball sampling techniques. In the community, we visited public
225 places (markets, school yards, parks) and approached young women face-to-face and
226 were selected if they met our inclusion criteria and were willing to participate in the
227 study. New potential participants were then referred by previous informants.

228 Since the use of modern contraceptive methods may also be affected by other actors
229 in decision-making and service provision, we collected data from young men ages 18
230 to 24, health care providers in public and private local health centers, relatives of
231 adolescent girls, and the leaders of a non-governmental organization (NGO) which

232 promotes contraceptive methods in the region. Those informants were all recruited
233 by purposive sampling techniques.

234 Data collection

235 Data were collected through in-depth interviews and focus group discussions with
236 young people, females, and males separately. The group discussions were conducted
237 in isolated public places (parks, youth gathering places) to ensure quietness and that
238 participants speak with ease and without fear of being seen by relatives. We
239 conducted the discussions mainly in the Fon language. The number and
240 characteristics of the participants are shown in Table 1.

241 **Table 1. Numbers and characteristics of participants in informant interviews and focus**
242 **group discussions.**

	Informant interview	Focus group discussion
Pregnant women	12	-
Non-pregnant women	13	7
Young men	3	7
Relatives of adolescent girls	2	-
Service providers, NGO leaders	6	-
Total	36	14

243

244 Based on an adapted theoretical framework from the PASS Health Seeking Behaviour
245 model of Hausmann-Muela et al.,[28], and the first findings from the quantitative
246 analysis, we designed interview guides specific to each target group. The question guides
247 focused on perceptions of ideal family size, knowledge of contraceptive methods, and
248 decision-making for contraceptive use.

249 Data analysis

250 All recorded sessions were transcribed into French. Those that were not conducted
251 in French were translated. A content analysis of transcripts was done using the
252 software MAXQDA 11. In the inductive analysis, narrative data was parsed and
253 categorized into units of meaningful information and the categories were linked to
254 understanding young women's decision-making for contraception.
255
256 **Ethical considerations**
257 The BDHS 2017-18 received approvals from the National Statistics Council and the
258 National Ethics Committee for Health Research in Benin. As DHS data are secondary
259 data, the quantitative strand did not require ethical approvals. The qualitative strand
260 was conducted as part of the "Continuum des Soins de Santé Reproductive au Bénin"
261 study which received approvals from the Local Ethics Committee for Biomedical
262 Research of the University of Parakou in Benin (approval number:
263 0092/CLERB-UP/P/SP/R/SA), the Institute of Tropical Medicine Antwerp in Belgium
264 (approval number: IRB/AB/AC/044), and Nagasaki University School of Tropical
265 Medicine and Global Health in Japan (approval number: NU-TMGH_064). The
266 participants in the interviews and the focus group discussions were informed of the
267 research objectives, their voluntary participation in the study, and there was no
268 disadvantage in refusal or withdrawal from the study. Written consent was obtained
269 from each participant.
270
271 **RESULTS**
272 **Quantitative strand**

273 Overall, 15,928 women of reproductive age (15 to 49 years old) were surveyed on
274 the DHS; 39.4% (n=6,251) were 15 to 24 years old. The sociodemographic
275 characteristics of the respondents by three age groups (15 to 19 years old; 20 to 24
276 years old; and 25 years or older) are shown in Table 2.

277 **Table 2. Socio-demographic characteristics of the quantitative analysis sample (Benin DHS 2017-18); N=15,928 (women aged 15-49)**

278

	Age groups	15 - 19 years (n=3,335)			20 - 24 years (n=2,916)			25 years or older (n=9,677)		
Variables		n	%	95%CI	n	%	95%CI	n	%	95%CI
Residence	Urban	1468	42.2	39.4-45.1	270	41,6	39.0-44.4	4307	42.8	40.8-44.8
	Rural	1867	57.8	54.9-60.7	1646	58.4	55.7-61.0	5370	57.2	55.2-59.2
Ethnicity	Fon	1227	37.4	34.4-40.4	1004	34.6	32.0-37.3	3514	37.5	35.4-39.7
	Others	2108	62.6	59.6-65.6	1912	65.4	62.7-68.0	6163	62.5	60.3-64.6
Religion	No religion	165	4.6	3.8-5.7	140	4.4	3.6-5.4	528	5.3	4.5-6.3
	Traditional	235	6.9	5.8-8.3	224	7.8	6.5-9.3	1010	10.4	9.1-11.8
	Islam	892	26.9	24.2-29.8	932	32.7	29.8-35.8	2853	29.7	27.5-32.0
	Christianity	2018	60.8	57.9-63.6	1585	53.8	50.8-56.7	5181	53.5	51.2-55.8
	Others	25	0.8	0.5-1.3	35	1.3	0.8-2.3	105	1.1	0.89-1.5
Literacy	Can read	1899	57.0	54.5-59.6	1301	44.3	41.6-47.0	2290	23.0	21.5-24.6
	Cannot read	1436	43.0	40.1-45.5	1615	55.7	53.0-58.4	7387	77.0	75.4-78.5
Marital status and sexual activity	Not married	2712	81.5	79.7-83.3	972	32.7	30.4-35.2	1074	11.1	10.3-12.0
	Sexually inactive	2344	70.3	68.2-72.3	688	23.4	21.5-25.3	822	8.6	7.9-9.4
	Sexually active	368	11.2	10.0-12.6	284	9.3	8.1-10.8	252	2.5	2.2-2.9
	Married	623	18.5	16.7-20.3	1944	67.3	64.8-69.6	8603	88.9	88.0-89.7
Parity	0	2826	84.8	83.2-86.3	974	32.5	30.5-34.7	372	3.7	3.3-4.1
	1	426	12.8	11.5-14.3	925	32.6	30.7-34.5	690	7.2	6.6-7.8
	2	79	2.2	0.2-2.9	663	23.2	21.5-25.0	1212	12.4	11.7-13.2
	3 or more	4	0.1	0.0-0.3	354	11.7	10.4-13.1	7403	76.7	75.6-77.9
	Median [IQR]	0	[0,0]		1	[0,2]		4	[3,6]	
Wealth quintile	Poorest	538	15.7	13.2-17.6	464	15.0	12.9-17.3	1854	18.5	16.6-20.6
	Poorer	658	20.3	18.3-22.5	536	18.9	17.1-20.9	1782	18.6	17.3-20.1
	Middle	631	19.8	17.8-22.0	564	19.9	17.9-22.0	1790	19.3	18.0-20.6
	Richer	691	21.2	19.1-23.6	630	22.5	20.4-24.8	1960	21.1	19.6-22.7
	Richest	817	23.4	20.6-26.4	722	23.8	21.2-26.6	2291	22.5	20.6-24.6

279

280 Modern contraceptive use was 11.2% (95%CI: 10.6-12.0%) among all women of
281 reproductive age (15 to 49 years old) versus 8.5% (95%CI: 7.7-9.5%) among women aged 15
282 to 24, and 13.0% (95%CI: 12.1-14.0%) among 25 years old women or older.
283 Contraceptive practices and knowledge in the study population are shown in Table 3,
284 where we compared four groups: adolescents and young women (15-24 years) and
285 women 25 years or older, (separately married and unmarried sexually active). Among
286 all women, the unmet need for modern contraception was significantly higher among
287 young unmarried sexually active women (60.8%) compared to all other groups.
288 Specifically, among women in need of family planning, the percentage of demand
289 satisfied by a modern method was lowest among 15 to 24 years old married women
290 (17.3%).

Table 3. Need for, demand for, and knowledge of modern contraception among sexually active women in Benin (n=12,074)

	Age	15 – 24 years old						25 years old or more						
	Marital status	Unmarried, sexually active (n=652)			Married (n=2567)			Unmarried, sexually active (n=652)			Married (n=2567)			
Indicator		n	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI	p-value
Modern contraceptive prevalence rate	Total	152	23.2	19.8 – 26.9	202	8.1	7.0 – 9.4	58	22.5	17.6 – 29.2	1132	13.0	12.1 – 14.0	<0.001
	for limiting	1	0.19	0.0 – 1.3	8	0.3	0.2 – 0.7	18	7.7	4.7 – 12.4	498	5.7	5.1 – 6.3	
	for spacing	151	23.0	19.7 – 26.7	194	7.8	6.7 – 9.0	40	14.8	10.7 – 20.2	634	7.4	6.7 – 8.1	
Unmet need for modern contraception	Total	393	60.8	56.9 – 64.7	989	38.8	36.7 – 41.0	113	44.5	38.0 – 51.0	3027	35.1	33.9 – 36.3	<0.001
	for limiting	5	0.72	0.3 – 1.9	28	1.1	0.7 – 1.7	33	13.4	9.4 – 18.7	1186	13.6	12.7 – 14.5	
	for spacing	388	60.1	56.1 – 64.0	961	37.7	35.5 – 55.6	80	31.1	25.3 – 37.5	1841	21.5	20.5 – 22.5	
Total need for modern contraception	Total	545	84.0	80.6 – 86.9	1191	46.9	44.4 – 49.3	171	67.0	60.2 – 73.2	4159	48.1	46.8 – 49.4	<0.001
	for limiting	6	0.9	0.4 – 2.2	1155	1.5	1.0 – 2.1	51	21.1	16.2 – 27.2	1684	19.2	18.2 – 20.4	
	for spacing	539	83.1	79.5 – 86.2	1376	45.4	43.0 – 47.8	120	45.9	39.1 – 52.9	2475	28.9	27.8 – 30.0	
Existence of barriers to access health services	None (0)	274	40.2	36.0 – 44.6	1,022	40.3	37.4 – 43.1	113	45.2	38.2-52.5	3,405	39.6	37.7 – 41.5	0.44
	Any (>=1)	378	59.8	55.4 – 64.0	1,545	59.7	56.8 – 62.5	139	54.8	47.5-61.9	5,198	60.4	58.5 – 62.3	
Number of modern methods known by women	Fewer than four	87	13.8	10.9 – 17.2	470	18.0	15.9 – 20.3	13	5.8	3.2-10.2	958	10.9	9.8 – 12.1	<0.001
	Four to six	316	50.0	45.2 – 54.8	1,208	46.9	44.3 – 49.5	92	35.8	29.4-42.9	3,674	43.6	41.7 – 45.4	
	More than six	249	36.2	31.8 – 40.9	889	35.0	32.6 – 37.6	147	58.4	51.4-65.2	3,971	45.6	43.6 – 47.5	
	Median [IQR]	6 [4, 7]			5 [4, 7]			7 [5, 8]			6 [5, 8]			
Percentage of demand satisfied by a modern method (as of % of women in need)			27.6	23.8 – 31.7		17.3	15.0 – 19.8		33.6	26.1-42.1		27.1	25.4 – 28.9	<0.001

294 Table 4 shows the method mix among users of modern methods. Among unmarried
295 sexually active women ages 15 to 24 years, 63.4% of modern contraceptive users
296 were using short-term methods, and more than 60% of these being male condom
297 users. The use of modern contraceptives was lower among married young women
298 (27.1%), sexually active unmarried women ages 25 or above (25.7%), and among
299 married women 25 years old or more (19.3%). Although there was no substantial
300 difference in long-term method use among married women (72.9% vs 80.7%) of 15 to
301 24 years old and 25 or older, the use of implants and IUDs was lower among young
302 women (49.4% vs 61.0%). Regarding the source of contraceptives, unmarried
303 sexually active young women obtained their contraceptives from for-profit outlets or
304 relatives (62.1%) or comprehensive providers (36.2%). The public sector was the
305 main source of methods among young and adult married women (77.0% and 78.9%).

306 **Table 4: Modern contraceptive practices among unmarried and married sexually active modern contraceptive users, by age group (n=1544)**

307

	Age	15 – 24 years old						25 years old or more					
	Marital status and Sexual activity	Unmarried, sexually active (n=152)			Married (n=202)			Unmarried, sexually active (n=58)			Married (n=1132)		
Variables		n	%	95% CI	n	%	95% CI	n	%	95% CI	n	%	95% CI
Methods of contraception	Short-term	100	63.4	54.4-71.6	53	27.1	20.6-34.7	17	25.7	15.3-39.8	229	19.3	16.8-22.1
	Oral contraceptive pills	6	2.6	1.1-6.0	26	12.3	8.3-18.0	7	11.4	5.1-23.6	151	12.7	10.6-15.1
	Male condoms	94	60.8	51.8-69.1	27	14.8	9.93-21.4	10	14.3	7.3-26.2	78	6.6	5.2-8.4
	Long-term	52	36.6	28.4-45.6	149	72.9	65.3-79.4	41	74.3	60.2-84.7	903	80.7	77.9-83.2
	Implants & IUD	36	24.7	17.8-33.3	99	49.4	41.3-57.6	30	54.4	39.4-68.7	688	61.0	57.7-64.2
	Injectables	16	11.9	6.92-19.6	48	22.9	17.3-29.7	11	19.9	10.3-34.9	191	17.6	15.4-20.0
	Others	0	-	-	2	0.6	0.2-2.47	0	-	-	24	2.2	1.4-3.3
Source of method	Public	49	33.9	25.8-43.1	143	70.8	62.9-77.6	30	61.1	47.3-73.3	824	72.4	68.8-75.8
	Non-public medical	6	3.9	1.5-9.7	16	7.23	4.2-12.3	11	13.2	7.0-23.5	111	10.4	8.4-12.8
	For-profit outlets and relatives	97	62.1	53.0-70.5	43	22.0	16.2-29.2	17	25.7	15.3-39.8	197	17.2	14.6-20.1
Capacity of provider	Comprehensive	52	36.2	27.9-45.4	156	77.0	69.8-82.9	38	70.7	56.6-81.7	892	78.9	75.9-81.6
	Limited	100	63.8	54.6-72.1	46	23.0	17.1-30.2	20	29.3	18.3-43.4	240	21.1	18.4-24.1

308

309

310 Factors associated with demand satisfied by a modern method

311 We examined the demand satisfied by a modern method among women ages 15 to 24
312 years in need of contraception (n=1736). Table 5 shows the percentages of modern
313 method users and the crude and adjusted odds ratios of each factor on the use of
314 modern methods. The multivariable analysis showed that being unmarried, higher
315 parity, existence of barriers in access to health services, and a higher number of
316 modern methods known by a woman were positively associated with demand
317 satisfied by a modern method.

318 **Table 5. Percentages and factors associated with modern method use among 15-24-year-old women in need of contraception (n=1,736)**

Variables		Modern method use (95% CI)	Crude OR [95% CI]		P-value	Adjusted OR [95% CI]		P-value
Residence	Urban	21.9 (18.8-25.2)	Ref			Ref		
	Rural	19.5 (16.9-22.5)	0.87	[0.67-1.12]	0.283	1.00	[0.72-1.38]	0.999
Ethnicity	Others	21.6 (19.1-24.4)	Ref			Ref		
	Fon	18.1 (15.0-21.7)	0.80	[0.61-1.05]	0.111	0.76	[0.55-1.05]	0.095
Religion	Christians	20.6 (18.0-23.4)	Ref			Ref		
	Islam	21.6 (18.0-25.8)	1.07	[0.80-1.42]	0.666	1.04	[0.73-1.47]	0.828
	Traditional and others	17.1 (12.0-23.9)	0.80	[0.52-1.24]	0.312	0.83	[0.53-1.30]	0.406
Literacy	Cannot read	17.8 (15.2-20.7)	Ref			Ref		
	Can read	23.6 (20.5-27.0)	1.43	[1.10-1.85]	0.007	1.20	[0.89-1.61]	0.233
Marital status	Married	17.3 (15.1-19.8)	Ref			Ref		
	Not married	27.6 (23.8-31.7)	1.82	[1.42-2.35]	<0.001	3.33	[2.21-5.01]	<0.001
Parity	None	23.4 (19.9-27.3)	Ref			Ref		
	One	19.6 (16.3-23.3)	0.80	[0.59-1.07]	0.133	2.11	[1.39-3.22]	0.001
	Two or more	18.6 (18.5-22.4)	0.75	[0.54-1.02]	0.071	2.22	[1.39-3.55]	0.001
Wealth quintiles	Poorest	21.9 (15.6-29.8)	Ref			Ref		
	Poorer	18.2 (13.9-23.4)	0.79	[0.46-1.35]	0.388	0.78	[0.45-1.33]	0.358
	Middle	20.3 (16.0-25.5)	0.91	[0.55-1.48]	0.701	0.79	[0.49-1.28]	0.333
	Richer	18.5 (14.7-23.1)	0.81	[0.49-1.35]	0.420	0.69	[0.41-1.16]	0.161
	Richest	24.0 (20.0-28.5)	1.13	[0.70-1.81]	0.625	0.94	[0.56-1.57]	0.814
Barriers to access to the services	None	17.6 (14.6-21.1)	Ref			Ref		
	Any	22.2 (19.6-25.1)	1.34	[1.02-1.76]	0.038	1.52	[1.15-2.02]	0.003
Number of modern methods known	Fewer than four	9.7 (6.30-14.8)	Ref			Ref		
	Four to six	21.1 (18.2-24.3)	2.48	[1.48-4.15]	<0.001	2.43	[1.45-4.07]	0.001
	More than six	23.7 (20.2-27.5)	2.88	[1.70-4.85]	<0.001	3.00	[1.73-5.21]	<0.001

319

320 **Qualitative strand**

321 We identified three main barriers in access to and utilization of contraception services
322 among adolescents, which are (i) socio-cultural perceptions related to sexuality and fertility
323 and (ii) perceptions about the purpose and effectiveness of contraception. A conceptual
324 model of the barriers is shown in Figure 1.

325

326 **Figure 1. A conceptual model of the barriers to access and utilization of contraception**
327 **services from qualitative analyses**

328

329 Sociocultural norms

330 Sociocultural norms including gender norms, cultural, and religious perceptions affect
331 people's perceptions about young women's fertility and the need for contraception in
332 Allada. These norms include views about pregnancy before marriage, male partners' role in
333 women's contraceptive use, local perceptions of women's fertility, resulting in different
334 perceptions of contraception such as the misuse of biomedicine, perceived effectiveness of
335 natural methods, mistrust in modern methods, and negative views about modern method
336 users.

337 *To be pregnant before marriage*

338 There is a common negative perception about sexual intercourse among adolescents.
339 Becoming pregnant as an adolescent and before marriage is not well viewed in the
340 community as judged as promiscuity. Therefore, unmarried young women avoid such a
341 judgment through abstinence or use fewer methods that require a source including fertility

342 awareness, to avoid exposure of their sexual activity. This situation prevents unmarried
343 young women from using contraceptives.

344 *"To avoid pregnancy, one must abstain from sex. If a girl really wants to avoid shame in*
345 *front of her classmates, she must refrain from sex." (less than 20 years old/pregnant)*

346 *Perceptions regarding the ideal family size*

347 In the study area, there was cultural pressure on having a certain number of children based
348 on social norms about the ideal family size. It implies that when a young woman gets
349 married, she may not be allowed to use contraception until she has several children.

350 *"If we say that someone has enough children, it will be from seven. That is when the*
351 *person will say, 'Yes, I do have children.'" (less than 20 years old/male)*

352 None of the respondents mentioned cultural pressures that might force young unmarried
353 women to have children early.

354 *Decision-making dynamics*

355 Women do not often have the decision-making power to use contraception.

356 *"One day, some people went door-to-door to inform us of the presence of an NGO at*
357 *the health center for family planning services. I then informed my husband and did the*
358 *injection for 3 months. But after he forbade me to continue it, because of rumors about*
359 *other women's bad experiences. When your husband refuses you have no choice."*
360 *(20-24 years old pregnant)*

361 Perceptions of contraception

362 *Contraception perceived as abortion*

363 Religious beliefs, especially Catholicism, also affect the utilization of contraception.

364 *"In my opinion, 'planning' is an abortion, a deadly sin. We are killing babies without*
365 *knowing and it is against the ten commandments of God. We are also killing ourselves*
366 *because we do not know the consequences of those things in our body." (less than 20*
367 *years old/non-pregnant)*

368 *Perceptions of modern method users*

369 Respondents believed that the use of contraception facilitates sexual promiscuity.
370 Contraception was also considered to be a eugenic procedure.

371 *"When she was 12 years old, people told me to have her adopt a contraceptive method.*
372 *I thought it was asking her to prostitute herself and I did not. Here now, as you see, she*
373 *came home with pregnancy while she is still having her training." (Relative of pregnant)*

374 *Perceived effectiveness of natural methods*

375 Faith in natural methods contributes to reluctance to use modern contraceptives among
376 young women.

377 *"I do not plan to use any of those things [modern methods]. My family members told*
378 *me they will prepare an herbal tea for me and I will not get pregnant for the period I*
379 *want - they know how to do it." (less than 20 years old/pregnant)*

380 Beliefs were led by fear of side effects of modern methods, and these opinions were in part
381 based on rumor or hearsay.

382 *" It happened to one of my relatives. She chose the one that is inserted in the arm.*
383 *Three months later, she had no more periods. Then when she had menstrual periods*
384 *again, it was too much. It was when it became excessive that she had to go where she*
385 *had it inserted for removal. Finally, she died." (20-24 years old/female)*

386 *Misuse of biomedicine*

387 Some also perceived 'aspirin' to be effective in preventing pregnancy and used it as a type
388 of modern method.

389 *"I often use Sedaspir [commercial name of acetylsalicylic acid] and it works well. I buy*
390 *the box at the pharmacy for 1500F CFA [equivalent to US$ 3]." (20-24 years*
391 *old/pregnant).*

392

393 **DISCUSSION**

394 We investigated modern method use, unmet need for contraception, and the reasons
395 why modern methods were not used; focusing on adolescent girls and young adult
396 women (15 to 24 years old) using the most recent Benin Demographic and Health
397 Survey (BDHS 2017-18) combined with a qualitative study conducted in the Allada area in
398 Benin.

399 Our quantitative analysis revealed that characteristics of modern method use
400 differed between unmarried sexually active and married young women ages 15 to 24.
401 Among all women, young unmarried sexually active women had both the highest
402 modern contraceptive use and the highest unmet need for modern contraceptives.
403 Modern method users tended to adopt more short-term than long-term methods,
404 especially the male condoms which they acquired from providers with a limited
405 range of methods. Young married women had the lowest percentage of demand
406 satisfied by a modern method among women who wished to delay or limit births.
407 Factors associated with modern method use among young women in need of
408 contraception were being unmarried, having at least one child, experiencing barriers
409 in access to health services, and knowing a greater number of modern

410 contraceptives. As suggested by the qualitative analysis, the reasons underlying the
411 high unmet need for modern contraceptives in young women are related to the
412 access and use of modern methods which can be categorized as 'sociocultural norms'
413 and 'perceptions of contraception'.
414 Young unmarried women had the highest unmet need for modern contraceptives as a
415 percentage of all women, which may explain the high rate of unintended pregnancy
416 in Benin. Very few unmarried young women want to have a child due to community
417 negative perceptions about pre-marital sexual intercourse and pregnancy, therefore
418 in need of contraception. However, as unmarried young women are considered in
419 the communities not to be ready for sexual intercourse they are reluctant to request
420 contraceptives despite their wish to avoid a pregnancy,[19, 20]. Moreover, the fear
421 of adverse effects contributes to young women's non-use or discontinuation of
422 modern contraceptives,[20, 29, 30].
423 Among unmarried young women in need of family planning, those with demand
424 satisfied by a modern contraceptive tend to use male condoms, the most adopted
425 modern method in Benin (45.3%), which can be easily obtained from for-profit outlets,
426 partners, and relatives. This finding is similar to that of Ali and Cleland,[31],
427 secondary data analysis from 36 African countries which showed that male condoms
428 constitute 51.3% of modern methods among unmarried women ages 15 to 24 years.
429 Our qualitative analysis suggested that while access to male condoms has fewer
430 barriers, male partners are in most cases the ones to obtain the method for use.
431 Consequently, the decision to use a male condom is likely to be made by the male
432 partner whose failure to acquire the method or unwillingness to use could be a

433 barrier to the young woman's contraceptive use. Male condoms are obtained in
434 shops, pharmacies, or from non-governmental organizations during
435 awareness-raising campaigns. Also, young women may assume that male partners
436 obtain contraceptives mainly from pharmacies which could explain why pharmacies
437 and relatives are the most frequent sources of methods reported in the DHS2017-18.
438 Young people's choice of providers including shops and pharmacies is based on their
439 values for discretion, respect of confidentiality, and the quickness of the service
440 delivery. This is in line with the findings of Murray et al.,[4], which suggested that
441 the preference of pharmacies by young women in requesting contraceptive services
442 may be driven by their perception that the providers are youth friendly.
443 Also, young married women in need of family planning had a lower percentage of
444 demand satisfied by a modern contraceptive compared to unmarried young women
445 and older married women. Most young women want to have a child once they get
446 married which explains the low unmet need as a percentage of all women. However,
447 married young women who wished to delay the next pregnancies face several
448 challenges in decision-making. For instance, the qualitative study found that women
449 are often under pressure to have children once they get married due to traditional
450 beliefs about fertility that prevent married women from using contraceptives.
451 Women are expected to demonstrate their fertility after marriage, which may be even
452 stronger for younger women as suggested by the literature,[32-36]. Moreover, in a
453 social context such as Allada, where large families of seven children are the average, the
454 family size is often decided by male partners and women feel less empowered to decide on
455 their own to adopt contraceptives. This may explain the high total fertility rate of 5.7 in

456 Benin. To get around the barrier of spousal consent, some married women request
457 contraceptives discretely, after giving several births and as their households face
458 economic challenges to raise the children. Most of the women would request long-term
459 methods such as implants to be inserted in the leg or other unusual and less exposed parts
460 of the body.

461 Our study revealed that factors associated with modern method use among those in
462 need included literacy, not being married, having children, knowing a greater
463 number of modern methods, and experiencing barriers to access health services.
464 Although 'literacy' was not often used as a variable in previous studies, among young
465 women there was evidence of a positive association between educational attainment
466 and the demand satisfied,[33, 37]. Similarly, the number of modern methods known
467 was positively associated with use. The strongest association was found between
468 women's marital status and the demand satisfied by a modern method. The
469 multivariate analysis revealed that young unmarried women were four times more
470 likely to use modern methods. This finding strengthens the evidence that married
471 young women are being left behind in terms of demand satisfied. Also, a similar
472 positive association between being unmarried and using modern contraceptives was
473 found by Dansou et al. in a study among all women of reproductive age (15 to 49
474 years old) in Benin,[20]. Several other studies conducted in other sub-Saharan
475 African countries such as Burkina Faso, Ethiopia, Nigeria, and Ghana found that
476 modern method prevalence was higher among unmarried adolescent girls compared
477 to married,[38, 39]. Young unmarried women used contraceptives to avoid pregnancy
478 because of the social stigma associated with early pregnancy in Benin, and because

479 they have no income to support a child on their own,[10, 20]. This finding is also
480 supported by our qualitative data. Having a child was associated with modern
481 method use. Other studies revealed that women with children were more likely to
482 use modern methods,[40, 41]. Health education from care providers during
483 pregnancy and in the postpartum period might be a contributing factor because the
484 BDHS 2017-8 showed that coverage of antenatal care provided by skilled personnel
485 and delivery in a health facility were 83% and 84%, respectively. Counterintuitively,
486 we found that having barriers to access to health services was associated with a higher
487 likelihood of modern methods use. We hypothesize that women who had accessed
488 contraceptive methods were more likely to report barriers, while those who did not report
489 such barriers did not theoretically exist.
490 Finally, our qualitative analysis revealed aspirin or paracetamol was used in Allada
491 to prevent pregnancy, despite the lack of a biomedical evidence-based for such an
492 effect. Boko et al. (2015) noted the same practice in Benin,[42]. The practice may be
493 driven by the fact that requesting such medicine does not expose a young woman to
494 any social stigma as the drug is not viewed as a contraceptive method by providers.
495 This common practice may imply that the use of oral contraception use among young
496 women in the DHS2017-18 is an over-estimate. In a context such as in Allada where
497 other ineffective drugs are used as oral contraceptives, it is important to promote
498 accurate knowledge as well as practices through health education activities.
499
500 **Limitations**

501 Our study has several limitations. We used secondary data, our analysis could have
502 been affected by recall errors and biases in some variables collected on the DHS. For
503 example, several assumptions were made in the DHS to assess the need for
504 contraception among women, which were criticized to have possibly induced an
505 underestimation in unmet needs for family planning,[43]. The assumptions
506 concerned the frequency of sexual activity of women, women's capacity to conceive
507 (fecundity criteria, length of lactational amenorrhea which is 24 months in DHS
508 versus 6 months clinically accepted standard), and the validity of their reports on
509 whether recent births were desired. In the qualitative study, having a male
510 interviewer might have affected responses from young women. Meanwhile, this was
511 addressed as a young female interviewer continued interviews with female
512 participants. Moreover, we could not ascertain whether a respondent in the
513 qualitative study was sexually active whereas the analysis of determinants in the
514 quantitative study focused on sexually active young women.
515

516 **Conclusion**

517 Access to and use of modern contraceptive methods is low among women in Benin,
518 especially among adolescent girls and young women. Adolescent girls and young
519 women tend to use short-term methods such as condoms which they obtain from
520 easily accessible providers and relatives. Unmarried and married women face different
521 barriers to use modern contraceptives, which need to be carefully addressed. Most
522 importantly young married women who wish to delay births should be specifically targeted
523 in interventions to address challenges related to decision-making to use contraceptives.

524 Raising awareness in communities with strong culture towards encouraging early
525 marriage and high fertility, as well as promoting comprehensive sexuality education
526 and providing youth-friendly services may help reduce the incidence of unintended
527 pregnancies among adolescent girls and young women,[44, 45].
528

529 **Acknowledgments**

530 We thank the Demographic Health Survey Program (http://dhsprogram.com/) for
531 granting us access to the BDHS2017-18 dataset. We acknowledge TOTIN Mahoutondji
532 Gilberte for her work as a research assistant during the qualitative data collection.
533 We would like to thank the study participants for sharing their experiences as well as
534 the local administrations of Allada for facilitating our study implementation.
535

536 **Contributors**

537 NCAA, MM, MS, LB, TD, J-P D, and SG were involved in conception and study design.
538 LB and MM provided statistical expertise. NCAA, AV, and CB collected qualitative
539 data in the field. MM, MS, TD, CG, LK, and KP provided qualitative data analysis
540 expertise. NCAA, MM, and MS were involved in the drafting of the manuscript. LB, TD,
541 J-P D, and CG were involved in the critical revision of the manuscript for important
542 intellectual content. All authors were involved in the final approval of the
543 manuscript and the decision to submit the manuscript for publication.
544

545 **Funding**

This study was supported by core funding from Nagasaki University School of Tropical Medicine and Global Health to MM (grant number N/A).

Competing interests

None declared.

Patient and public involvement

Patients and/or the public were not involved in the design, conduct, reporting, or dissemination plans of this research.

Patient consent for publication

Not required.

559 **References**

560 1. United Nations Population Fund (UNFPA). Programme of Action. New York: UNFPA 1994.

561 2. Singh S, Sedgh G, Hussain R. Unintended pregnancy: worldwide levels, trends, and

562 outcomes. *Stud Fam Plann* 2010;41(4):241-50. doi:

563 10.1111/j.1728-4465.2010.00250.x [published Online First: 2011/04/06]

564 3. Hubacher D, Mavranezouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa:

565 magnitude of the problem and potential role of contraceptive implants to alleviate

566 it. *Contraception* 2008;78(1):73-8. doi: 10.1016/j.contraception.2008.03.002

567 [published Online First: 2008/06/17]

568 4. FP2020. Indicators 3-4 : Unmet Need and Demand Satisfied.

569 *http://2016-2017progressfamilyplanning2020org/en/measurement-section/unmet-n*

570 *eed-and-demand-satisfied-indicators-3-4* [accessed November 2020]

571 5. Okigbo CC, Speizer IS. Determinants of Sexual Activity and Pregnancy among Unmarried

572 Young Women in Urban Kenya: A Cross-Sectional Study. *PLoS One*

573 2015;10(6):e0129286. doi: 10.1371/journal.pone.0129286 [published Online First:

574 2015/06/06]

575 6. World Health Organization (WHO). Family planning evidence brief: family planning

576 financing. Geneva: WHO, 2018.

577 7. Darroch JE, Woog V, Bankole A, et al. Costs and Benefits of Meeting the Contraceptive

578 Needs of Adolescents. *New York: Guttmacher Institute* 2016

579 8. Fescina R, B DM, JL DR, et al. Sexual and reproductive health: guides for the HPC focused

580 continuum of care of women and newborn: guides for basic practice. *Latin*

American Center for PerinatologyWomen and Reproductive HealthCLAP/WR - PAHO Montevideo: CLAP/WP; 2009? (CLAP/WR Scientific Publication; 15622) 2009

9. World Health Organization (WHO). Health for the world's adolescents: A second chance in the second decade. Geneva: WHO,2014

10. Sedgh G, Ashford LS, Hussain R. Unmet Need for Contraception in Developing Countries: Examining Women's Reasons for Not Using a Method. *New York: Guttmacher Institute* 2016

11. McCurdy RJ, Jiang X, Schnatz PF. Long-acting reversible contraception in adolescents in Sub-Saharan Africa: evidence from demographic and health surveys. *Eur J Contracept Reprod Health Care* 2018;23(5):357-64. doi: 10.1080/13625187.2018.1519535 [published Online First: 2018/11/23]

12. Institut National de la Statistique et de l'Analyse Économique (INSAE) and ICF. Enquête Démographique et de Santé au Bénin, 2017-2018. Cotonou, Bénin and Rockville, Maryland, USA : INSAE and ICF. 2019.

13. Government of Benin. FAMILY PLANNING 2020 COMMITMENT. 2017. http://www.familyplanning2020.org/news/benin-fp2020-commitment [accessed March 2020]

14. Chae S, Woog V, Zinsou C, et al. Obstacles à la pratique contraceptive des femmes au Bénin. *En Bref, New York: Guttmacher Institute* 2015

15. USAID, Communities AP. Benin's Community-Based Access to Injectable Contraceptives Pilot Project. 2015

16. Okegbe T, Affo J, Djihoun F, et al. Introduction of Community-Based Provision of Subcutaneous Depot Medroxyprogesterone Acetate (DMPA-SC) in Benin:

604 Programmatic Results. *Glob Health Sci Pract* 2019;7(2):228-39. doi:
605 10.9745/ghsp-d-19-00002 [published Online First: 2019/06/07]
606 17. Igras S, Diakité M, Lundgren R. Moving from theory to practice: A participatory social
607 network mapping approach to address unmet need for family planning in Benin.
608 *Glob Public Health* 2017;12(7):909-26. doi: 10.1080/17441692.2016.1147589
609 [published Online First: 2016/03/08]
610 18. Ahovey EC. Besoins non satisfaits en planification familiale au sein du couple : :
611 caractéristiques socio-démographiques et cadre de vie au Benin. 2002
612 19. Sedgh G, Hussain R. Reasons for contraceptive nonuse among women having unmet
613 need for contraception in developing countries. *Stud Fam Plann* 2014;45(2):151-69.
614 doi: 10.1111/j.1728-4465.2014.00382.x [published Online First: 2014/06/17]
615 20. Dansou J, Adekunle AO, Arowojolu AO. Factors behind the preference in contraceptives
616 use among non-pregnant and sexually active women in Benin. *Central African*
617 *Journal of Public Health* 2017;3(5):80-89.
618 21. MacQuarrie KLD. Unmet need for family planning among young women: levels and
619 trends. DHS Comparative Reports No 34. Rockville, Maryland, USA: ICF International,
620 2014.
621 22. Radovich E, Dennis ML, Wong KLM, et al. Who Meets the Contraceptive Needs of Young
622 Women in Sub-Saharan Africa? *J Adolesc Health* 2018;62(3):273-80. doi:
623 10.1016/j.jadohealth.2017.09.013 [published Online First: 2017/12/19]
624 23. Maxwell J. Designing a Qualitative Study. *Qualitative Research Design An Interactive*
625 *Approach (Third Edition)* 2012

626 24. Hubacher D, Trussell J. A definition of modern contraceptive methods. *Contraception*
627 2015;92(5):420-1. doi: 10.1016/j.contraception.2015.08.008 [published Online First:
628 2015/08/16]
629 25. Bradley SEK, Croft T, Fishel J, et al. Revising Unmet Need for Family Planning. *DHS*
630 *Analytical Studies No 25 Calverton, Maryland, USA: ICF International* 2012
631 26. Festin MP, Kiarie J, Solo J, et al. Moving towards the goals of FP2020 - classifying
632 contraceptives. *Contraception* 2016;94(4):289-94. doi:
633 10.1016/j.contraception.2016.05.015 [published Online First: 2016/06/12]
634 27. Vigan A, Boyi C, Ahissou C, et al. Déterminants de l'utilisation des services le long du
635 continuum des Soins de Santé Reproductive au Bénin: une exploration qualitative.
636 Rapport d'étude. . *Ministère de la Santé, Centre de Recherche en Reproduction*
637 *Humaine et Démographie (CERRHUD), and Institut de Médecine Tropicale (IMT)* 2020
638 28. Hausmann-Muela S, Muela Ribera J, Toomer E, et al. The PASS-model: a model for
639 guiding health-seeking behavior and access to care research, 2012.
640 29. Ochako R, Mbondo M, Aloo S, et al. Barriers to modern contraceptive methods uptake
641 among young women in Kenya: a qualitative study. *BMC Public Health* 2015;15:118.
642 doi: 10.1186/s12889-015-1483-1 [published Online First: 2015/04/18]
643 30. Campbell OM, Benova L, Macleod D, et al. Who, What, Where: an analysis of private
644 sector family planning provision in 57 low- and middle-income countries. *Trop Med*
645 *Int Health* 2015;20(12):1639-56. doi: 10.1111/tmi.12597 [published Online First:
646 2015/09/29]
647 31. Ali MM, Cleland J. Long term trends in behaviour to protect against adverse
648 reproductive and sexual health outcomes among young single African women.

649 *Reprod Health* 2018;15(1):136. doi: 10.1186/s12978-018-0576-6 [published Online
650 First: 2018/08/16]
651 32. Asiimwe JB, Ndugga P, Mushomi J. Socio-demographic factors associated with
652 contraceptive use among young women in comparison with older women in Uganda.
653 *DHS Working Papers No 95* 2013
654 33. Asiimwe JB, Ndugga P, Mushomi J, et al. Factors associated with modern contraceptive
655 use among young and older women in Uganda; a comparative analysis. *BMC Public*
656 *Health* 2014;14:926. doi: 10.1186/1471-2458-14-926 [published Online First:
657 2014/09/10]
658 34. Ngome E, Odimegwu C. The social context of adolescent women's use of modern
659 contraceptives in Zimbabwe: a multilevel analysis. *Reprod Health* 2014;11:64. doi:
660 10.1186/1742-4755-11-64 [published Online First: 2014/08/12]
661 35. Shahabuddin AS, Nöstlinger C, Delvaux T, et al. What Influences Adolescent Girls'
662 Decision-Making Regarding Contraceptive Methods Use and Childbearing? A
663 Qualitative Exploratory Study in Rangpur District, Bangladesh. *PLoS One*
664 2016;11(6):e0157664. doi: 10.1371/journal.pone.0157664 [published Online First:
665 2016/06/24]
666 36. Mardi A, Ebadi A, Shahbazi S, et al. Factors influencing the use of contraceptives
667 through the lens of teenage women: a qualitative study in Iran. *BMC Public Health*
668 2018;18(1):202. doi: 10.1186/s12889-018-5116-3 [published Online First:
669 2018/02/01]
670 37. Tamang L, Raynes-Greenow C, McGeechan K, et al. Factors associated with
671 contraceptive use among sexually active Nepalese youths in the Kathmandu Valley.

672 *Contracept Reprod Med* 2017;2:13. doi: 10.1186/s40834-017-0040-y [published
673 Online First: 2017/12/05]
674 38. Hounton S, Barros AJ, Amouzou A, et al. Patterns and trends of contraceptive use
675 among sexually active adolescents in Burkina Faso, Ethiopia, and Nigeria: evidence
676 from cross-sectional studies. *Glob Health Action* 2015;8:29737. doi:
677 10.3402/gha.v8.29737 [published Online First: 2015/11/13]
678 39. Marrone G, Abdul-Rahman L, De Coninck Z, et al. Predictors of contraceptive use among
679 female adolescents in Ghana. *Afr J Reprod Health* 2014;18(1):102-9. [published
680 Online First: 2014/05/07]
681 40. de Vargas Nunes Coll C, Ewerling F, Hellwig F, et al. Contraception in adolescence: the
682 influence of parity and marital status on contraceptive use in 73 low-and
683 middle-income countries. *Reprod Health* 2019;16(1):21. doi:
684 10.1186/s12978-019-0686-9 [published Online First: 2019/02/23]
685 41. Ahinkorah BO. Predictors of unmet need for contraception among adolescent girls and
686 young women in selected high fertility countries in sub-Saharan Africa: A multilevel
687 mixed effects analysis. *PLoS One* 2020;15(8):e0236352. doi:
688 10.1371/journal.pone.0236352 [published Online First: 2020/08/08]
689 42. Boko I, Baxerres C, Ouattara F, et al. Interroger au Bénin les usages populaires d'un
690 médicament abortif, le misoprostol. *Rev med perinat* 2017;9(1):20-24. doi:
691 10.1007/s12611-016-0388-2
692 43. Bradley SE, Casterline JB. Understanding unmet need: history, theory, and measurement.
693 *Stud Fam Plann* 2014;45(2):123-50. doi: 10.1111/j.1728-4465.2014.00381.x
694 [published Online First: 2014/06/17]

44. International Planned Parenthood Federation (IPPF). Provide: Strengthening youth friendly services. London: IPPF, 2008.

45. UNESCO. UN urges Comprehensive Approach to Sexuality Education. <https://en.unesco.org/news/urges-comprehensive-approach-sexuality-education> [Accessed November 2020].


SOCIO-CULTURAL NORMS
To be a pregnant before marriage
Perceptions of fertility
Contraception depends on her partner
Sexual intercourse is prohibited until marriage.
Contraception is only for married women.
Decision of contraception use is made by the husband.
It is the husband who decides the number of children in a family.
High number of children is appreciated.
Abstinence is better in order to avoid pregnancy.
To be a pregnant before marriage is a shame.
Contraception is considered as abortion, which is a sin.
Aspirin is one of effective contraceptive methods.
Misuse of medicine
Natural methods (cycle method) is simple and useful.
Traditional methods (herbs, decoctions, and items) are useful.
Perceived effectiveness of natural methods
Modern methods have side effects, which could be fatal.
Perceived effectiveness of modern methods
Contraception is for women with psychological disorders.
Perception of modern method users
PERCEPTIONS OF CONTRACEPTION