Title

Relationship between the kinematics of wrist/finger joint movements and

muscle loading during typing in people with different skill levels

Author names

Takanori Ito M.Sc.[a,b], Akira Kobayashi M.Sc.[a], Yuka Matsumoto M.Sc.[a], Hayase

Funakoshi[a], Mio Yoshida[a], Naohiko Kanemura Ph.D.[a,c], Takanori Kokubun

Ph.D.[a,c]

Author affiliations

[a]Graduate School of Health and Social Services, Saitama Prefectural University,

Saitama 343-8540 Japan.

[b]Department of Rehabilitation, Kawagoe Rehabilitation Hospital, Saitama 350-

1138 Japan.

[c]Department of Physical Therapy, School of Health and Social Services, Saitama

Prefectural University, Saitama 343-8540, Japan

Corresponding author

Takanori Kokubun Ph.D., Department of Health and Social Services, Health and

Social Services, Graduate School of Saitama Prefectural University, Saitama

343-8540 Japan. E-mail; kokubun-takanori@spu.ac.jp

Author's E-mail

Takanori Ito; 2391002b@spu.ac.jp, Akira Kobayashi; 1991003n@spu.ac.jp, Yuka

Matsumoto; 2191005s@spu.ac.jp, Hayase Funakoshi; 2381302r@spu.ac.jp, Mio

Yoshida;   2381309c@spu.ac.jp,   Naohiko   Kanemura;   kanamura-

naohiko@spu.ac.jp, Kokubun Takanori; kokubun-takanori@spu.ac.jp

Word counts

Abstract: 248

Main text: 3,577 (excluding references and figure captions)

Abstract

Background

Musculoskeletal disorders caused by computer use are associated with socio-

economic problems. Reducing the biomechanical loading caused by hand

movements could reduce the occurrence of such disorders. However, most

participants recruited for past experimental studies had a certain level of typing

skill, and differences in the biomechanical loading of those with different typing

skills have not been examined. This study aimed to quantify the relationship

between the movements of the wrist and finger joints and the loading on the

forearm muscles and to clarify the differences between typists of different skill

levels.

Methods

A 3D motion capture system measured wrist and index finger joint movements,

and surface electromyography measured muscle activities for the right hand

during a typing task. We compared wrist and finger joint movements and forearm

muscle loading during typing, keystroke time, and key release time between

skilled and unskilled typists.

Findings

Skilled typists move their wrists and fingers faster with less muscle activity

during typing, the active tension of their finger flexor muscles during keystrokes

are high, and they have higher mechanical stresses on the finger flexor tendons

during keystrokes. Unskilled typists develop strategies to prevent excessive

stiffness in their wrist joints when making keystrokes. They are expected to have

a higher cumulative loading on the extensor muscles of the wrist during key

release.

Interpretation

Biomechanical loading in typing skill is different, which may make it possible to

predict disability. This could provide information on the changes in physical

parameters and environments to prevent disability.

Keywords

Typing; Upper extremity musculoskeletal disorders (UE-MSDs); Motion capture

system; Electromyography; wrist/finger joint movements; Muscle loading

1. Introduction

Computers are essential tools in modern society and are frequently used for long periods of time. The intensive use of computer keyboards is associated with increased discomfort in the wrist and fingers (So et al., 2017), reducing worker productivity. Additionally, computer use for more than 4 hours per day or 20 hours per week increases the risk of upper extremity musculoskeletal disorders (UE-MSDs) of the wrist and fingers, such as tendonitis and carpal tunnel syndrome (Bergqvist et al. 1995; Gerr et al. 2002; Jenssen et al. 2003). People are expected to spend more time working at computers as digital transformation continues, possibly increasing the incidence of UE-MSDs.

Most studies that have investigated the kinematics of keyboard typing have focused on wrist movements. Few studies (Sommerich et al. 1996; Baker et al. 2007) have focused on finger movements during typing, but there is no research on the relationship between finger movements and the muscle activities that facilitate these movements.

The motion of the sagittal plane of the finger/wrist joints is interrelated as described by the tenodesis effect (Zhao et al. 2002; Su et al. 2005). Evaluating only the finger/wrist joints and muscle activities may lead to misinterpretation of

mechanical loading on the body. Some UE-MSDs are thought to be related to

high muscle loading on a single keystroke, which is repeated over a long period

of time.

However, the relationship between these characteristic movements and

individual typists remains unclear: each typist has a unique pattern and style of

typing. Since typing is a complex task controlled by many joints and muscles,

the movement style may be affected by skill level; different styles of movement

may then carry different risks of disability.

This study aimed to quantify the relationship between the movements of the

wrist and finger joints and the loading of the forearm muscles and to clarify the

differences according to typing skill. This study hypothesized those skilled

typists exhibit increased immobilization of the wrist joints and high-speed wrist

and finger joint movements. This would increase the muscle loading on the

forearm muscles, resulting in different levels of risk for developing

musculoskeletal disorders of the hand. By clarifying these relationships, we

believe that we can deepen our understanding of the influence of subject

characteristics on biomechanical loading during typing and provide basic

knowledge for physical and environmental improvement approaches aimed at

maintaining healthy work environments and preventing the occurrence of UE-

MSDs.

2. Methods

2.1. Subjects

Twelve healthy volunteers (nine men and three women) participated in this experiment. All participants provided written informed consent, following a detailed explanation of the study's purpose and risks according to the Declaration of Helsinki. This study was approved by the ethics review committee at Saitama Prefectural University, Saitama, Japan (approval number: 20,508).

2.2. Equipment

We used a 3D motion capture system (Vicon Motion Systems Limited, Vicon, Oxford Metrics plc, London, UK; sampling frequency: 100 Hz) and wireless multipoint surface electromyograph (EMG) (DELSYS, Delsys Trigno Wireless System, Massachusetts, USA; sampling frequency: 1,000 Hz). The height of the table was 70 cm, and a monitor (Apple, MacBook Air) and QWERTY keyboard (Microsoft, Wired Keyboard 600 ANB-00040, Washington, USA) were placed on it. The chair was adjusted for each subject to their preferred configuration prior to beginning the tests.

2.3. Protocol

First, the height and weight of the subjects and the physical information necessary for modeling were measured. Forty four infrared-reflecting markers (diameters 4 and 14 mm) were attached to points on the participants (Fig.1)—according to the Vicon plug-in-gait full-body (PIG) and wrist/finger (Baker et al. 2007; Cook et al. 2007) models—to calculate the sagittal plane of the wrist and the metacarpophalangeal (MCP) joint of the index finger angle and velocity.

Surface EMGs were attached to the right side of the flexor carpi radialis (FCR), flexor carpi ulnaris (FCU), extensor carpi radialis (ECR), and extensor carpi ulnaris (ECU) muscles. Then, the maximum voluntary isometric contraction (MVC) of each muscle was measured.

Each participant practiced typing for 5 min to get accustomed to the test environment. During this practice, the height of the chair and the angle of the monitor were adjusted according to each participant's preference. As a test (pretest) for typing skills, we used an electronic keyboarding program (e-typing, e-typing Co., Ltd., Aichi, Japan; http://www.e-typing.ne.jp) which has a text display that advances as the user types. All subjects used the same sentence.

For the typing test, all subjects were asked to type the exact same sentences,

which consisted of approximately 5,000 Japanese characters. No instructions

were given regarding typing speed, rhythm, or time limit; all subjects typed

according to their usual natural style. After confirming that the subject's hands

and wrists were placed on the keyboard, we instructed them to begin the task,

whereupon they started typing the sentences displayed on the monitor. Each

participant performed the task three times, resting as necessary.

2.4. Data processing

We analyzed all data using Nexus 2.9 (Vicon Motion Systems Limited, Vicon,

Oxford Metrics plc, London, UK) and Python 3.7 (Python Software Foundation,

Beaverton, Orlando, USA). The wrist joint angle was calculated from the markers

attached to the index finger metacarpal head, radial styloid process, ulnar styloid

process, and forearm in the PIG model. For the wrist joint angle in this study, the

positive value is expressed as the dorsal flexion, and the negative value as the

palmar flexion angle. The MCP joint was used to define the hand as one rigid

body, and the angle of the proximal phalanx segment with respect to the hand

on the sagittal plane was calculated and used as the flexion/extension angle (Fig.

1). For the MCP angle in this study, the positive value is expressed as the flexion

angle and the negative value as the extension angle.

The kinematics data were filtered with a fourth-order Butterworth low-pass filter

with a cut-off frequency of 6 Hz. After filtering was completed, the mean joint

angle, joint angle change as the root-mean-square (RMS), and joint velocity were

calculated in each trial of the wrist and MCP joint.

The EMG data were passed through a fourth-order Butterworth bandpass filter

with a cutoff frequency of 20–480 Hz, full-wave rectified, and then smoothed at

10 Hz. The EMG data during the typing sessions were normalized to the

percentage of MVC.

2.5. Data analyses

2.5.1. Grouping by the pretest results

The accuracy (%) and typing speed (words per minute, WPM) in Japanese for all

subjects in the pretest were obtained. The accuracy is the percentage of keys

that can be input without mistakes. The reason for calculating this is to make

sure that we are not doing the task unsystematically in an attempt to accomplish

it quickly. The typing speed was calculated for all participants, and those whose

speed was higher than the average value were classified as skilled (in the skilled

typist [ST] group), and those whose speed was lower than the average value

were classified as unskilled (the unskilled typist [UT] group). In a study that

investigated the keystroke characteristics of typing novices and experts

(Takaoka et al., 2014), both were classified as having a correct response rate of

95% or higher, or close to it, as a condition for employment.

2.5.2. Analysis interval

For all data, the start of the analysis interval was defined as the time when the

first Enter key was pressed down after the examiner gave the subject the signal

to start. The end of the analysis interval was defined as the time when the Enter

key presented at the end of the task text was pressed down.

The keystroke for the Enter key on the keyboard was determined using the 3D

motion capture system: the change in position information of the Enter key when

the position of the 4 mm marker attached to it was lowered was measured. Using

this input, the timing of the input signal of the key and the start of measurement

by the 3D motion capture system and the EMG were matched, and all kinematic

data and EMG data were synchronized in time.

2.5.3. Analysis of wrist joint and metacarpophalangeal joint motion

The initial posture of the wrist joint at the beginning of typing was investigated

by calculating the mean and standard deviation (SD) values of the initial

dorsiflexion angle of the wrist joint at the start time. The mean and SD values of

the palmar flexion/dorsiflexion angles of the wrist and MCP joint angles of the

index finger at the analysis interval calculated to examine the average posture of

the wrist and the MCP joint during typing. Similarly, the mean and SD values of

the RMS of the palmar flexion/dorsiflexion angles of the wrist and index finger in

the analysis section were calculated. The angle velocity of the wrist and index

finger MCP joints was investigated by calculating the mean and SD values of the

angular change (angular velocity) per unit time.

2.5.4 Analysis of muscle loadings in the forearm

To investigate the muscle loading in each muscle of the forearm, we used the

amplitude probability distribution function analysis for the time series data

of %MVC. Amplitude probability distribution function analysis is a method that

expresses the percentage of the total time that the output was below a certain

level as the probability (P) of occurrence for that output during the entire time

that the EMG was measured. After plotting the function of power output and

appearance probability of each muscle in each trial for each subject, we

calculated the mean and SD values of the muscle loadings at $P = 0.1$, 0.5, and

0.9.

2.5.5. Identification of press and release times

Based on the subject's key input signals, we obtained the time when the input

of each key started and ended. The time at which the "U" key was pressed and

input started was defined as the press time, and the time when the fingertip left

the "U" key and no input was made, as the release time. The mean and SD

values of each data point were calculated and compared between the two

groups.

2.6. Statistics

After confirming the normality of all kinematics and muscle activity data using

the Shapiro-Wilk test, we investigated the homoscedasticity of the data using

the F-test. Since there was no difference in kinematics data and EMG data in

each trial (confirmed by one-way ANOVA; $P < 0.05$), the average values of the

three trials were used as representative values. The data of the two groups were

compared using the unpaired t-test ($P < 0.05$) or Wilcoxon signed-rank test.

3. Results

3.1. Subjects and environments

The subjects of this study were 12 healthy adults, nine men and three women.

The mean (SD) value of correct answers in the typing test showed in Table.1. The

mean value of WPM of all subjects was used as the benchmark for skill: subjects

who achieved a score higher than the mean were classified into the ST group,

and those who achieved a lower score than the mean into the UT group.

The mean (SD) age, height, weight, hand width, finger length and the male-to-

female ratio showed in Table.1. All subjects were right handed.

The mean (SD) chair height and screen angle showed in Table.1. The ratio of PC

used on a daily basis (desktop PC to notebook PC) was 1:11. The mean (SD)

values for frequency and duration of computer use were 5.2 (1.2) times per week

and 3.3 (1.8) hours per day, respectively. There were no significant differences

between the two groups in any of the data.

3.2. Wrist and metacarpophalangeal joint motion

There was no significant difference in the mean dorsiflexion angle of the wrist

joint at the beginning of the typing motion between the two groups.

The mean value (SD) of the wrist joint dorsiflexion angle, RMS, and angular

velocity showed in Fig.2. The angular velocity of flexion/extension was

predominantly higher in the ST group (P = 0.003).

The mean (SD) values of the index finger MCP joint of flexion angle, RMS, and

angular velocity showed in Fig.2. The mean angular velocity of the MCP in the

ST group was significantly higher than that in the UT group (P = 0.002).

3.3. Muscle loading

The mean values (SD) of %MVC of FCR, FCU, ECR, and ECU at the static activity

level (P = 0.1), the mean activity level (P = 0.5), the maximum activity level (P =

0.9) showed in Fig.3. The mean of %MVC of FCR at P = 0.9 was significantly

higher than that in the ST group (P = 0.023). The mean of %MVC of ECR at P =

0.1, 0.5, 0.9 were significantly higher than that in the UT group (P = 0.040, 0.020,

0.020).

3.4. Press and release time data

The mean (SD) of the release time was 0.12 (0.02) and 0.20 (0.14) s for the ST

group, 0.13 (0.01) and 0.11 (0.01) s for the UT group, and 0.11 (0.02) and 0.26

(0.15) s for the ST group. The comparison showed no significant difference

between the two groups.

The mean values (SD) of the wrist joint dorsiflexion angle, angular velocity, and

the index finger MCP joint flexion angle, angular velocity during the press time

and release time showed in Fig.4, Fig.5. The mean value of the wrist joint

dorsiflexion angular velocity during the release time in the ST group was

significantly higher than that in the UT group ($P = 0.005$).

The mean value of the index finger MCP joint flexion/extension angular velocity

during the press time and release time in the ST group was significantly higher

than that in the UT group ($P = 0.002, 0.006$).

The mean values (SD) of %MVC of FCR, FCU, ECR, and ECU during press time

and release time showed in Fig.4, Fig.5. The mean values (SD) of %MVC of ECR

during press time of the ST group was significantly lower than that of the UT

group ($P = 0.020$).

4. Discussion

In this study, we examined the typing skills, joint movements and muscle loading

of 12 volunteers to evaluate the differences between skilled and unskilled typists.

4.1. Subjects and environment

Although the ST and UT groups showed similar typing accuracy, the typing

speed between the two groups was significantly different ($P < 0.05$). In both

groups, the accuracy was around 95%, indicating that the subjects were able to

adapt to the test environment and reproduce their natural typing motion in it.

In typing experiments with English sentences (Rempel et al. 2007; Szeto et al.

2000), the average typing speed of the subjects ranged from 20 to 50 WPM. On

average, one word in English consists of five keystrokes: four keys and one

space key (Crump et al. 2011; Nagasawa 2017). The WPM for Japanese in this

study was equivalent to approximately 34 to 44 WPM in English, indicating that

the typing speeds of all participants in this study were within the range of

subjects in previous studies. Therefore, we believe that the validity of the design

for assessing typing skills is ensured.

4.2. Kinematics of finger and wrist joint

There were no significant differences in the dorsiflexion angle and RMS of the

wrist between the ST and UT groups, but the angular velocity of

flexion/extension was predominantly higher in the ST group. The kinematic

parameters of the MCP joint of the index finger showed no significant differences

in the joint angle and RMS, but the angular velocity of flexion/extension was

predominantly higher in the ST group. The higher typing speed of the ST group

was due to fast movements of the wrist joint and fingers, as well as coordinated

fast joint action in the joints.

Previous studies have shown that the dorsiflexion angle of the wrist joint

changes depending on the gable angle, split angle, and tilt angle of the keyboard

(Simoneau 2003; Rempel 2007). Since all participants in the present study

performed the task with the same keyboard at the same tilt angle, it seems that

typists adjust their posture and range of wrist motion to the characteristics of

the keyboard regardless of the level of skill.

4.3. Muscle loading and movement

The two groups showed similar muscle loading of the wrist flexors (FCR and

FCU) during the task and muscle activity during press and release times. It has

been reported that the flexion movement of the MCP joint during the typing

motion is synergistic with the flexion movement of the wrist joint (Dennerlein

2007). In the present study, the ST group performed faster finger and wrist joint

movements: the passive tension of the wrist flexors due to the dorsiflexion

motion of the wrist acted on the flexion motion of the MCP and wrist joints,

enabling efficient and energetically economical movement.

However, the ST group may have used excessive muscle activity of the wrist

flexors to accelerate the joint movement of the index finger during keystrokes.

Additionally, because the fingertips were in contact with the keys at a high speed

during key pressing, the finger flexors may have acted excessively to maintain

the posture of the hand against the reaction force from the keys.

The muscle activity of the wrist extensor muscle decreases as the dorsiflexion

angle of the wrist decreases during typing (Simoneau 2003), possibly due to

changes in muscle length and moment arm. In this study, there was no difference

in the angle of the wrist joint and the degree of change of angle between the two

groups, but the muscle loading of the static activity level of the ECR—an

extensor muscle of the wrist joint—was significantly higher in the UT group than

in the ST group. This may be due to excessive activity of the ECR in the UT group

to stiffen the wrist joint, even when the moment arm of the muscle was long.

This excessive contraction may reduce the sliding ability of the ECR tendon in

its sheath. Additionally, the ECU muscle acts during lifting of the hand after a

key press (Dennerlein 2007). In the UT group, when fingertips were released from

the keystroke, excessive lifting of the hand was observed during the movement

of the wrist joint, possibly increasing the muscle loading of the ECU muscle

during key release.

4.4. Limitations

The measurement conditions in this study were not restricted, as the posture of

the participants was not constrained during the test. Participants could place

their forearms on the desk or keep them floating. Since placing the forearm on

the desk may decrease the muscle loading of the extensor muscles of the wrist

joint, the distance between the upper body and the keyboard and the presence

or absence of space on the desk in front of the keyboard must be checked at

the beginning of the measurement, and the effects of differences in the contact

time and contact area between the desk and the forearm on the body during the

task should be determined.

The majority of the participants recruited for this study were male. Physical

parameters, such as hand size, range of motion of joints, and absolute muscular

strength, vary depending on sex, and this may have affected the physicality of

the typing test.

In this study, we examined the differences in physical parameters with regard to

skill level among healthy adult students. We used a younger age grouping,

assuming people of this age would form the core labor force. However, the

subjects in this study were not experienced at desk work, and we were not able

to verify the results for older typists with greater experience. Therefore, the

findings of this study should be generalized with caution because the

characteristics of the study population may differ from those who are at risk of

developing musculoskeletal disorders.

In this study, the press time of the "U" keystroke and the time from the release

of the "U" key to the start of the next keystroke were used as the analysis

intervals, and the movements of the index finger were analyzed in detail.

However, we were not able to confirm for all participants which finger was

actually used to input the "U" key. Thus, we may not have been able to evaluate

the movement of only the index finger. Additionally, because we did not measure

the fingertip force during keystrokes, we could not show a direct relationship

between the angular velocity of the MCP joint and the actual force generated

during keystrokes. For future investigations, we intend evaluating the

movements of fingers other than the index finger and examining the relationships

between changes in posture, muscle loading, and fingertip force during specific

keystrokes.

5. Conclusions

The posture of the wrist and MCP joints of the index finger during the typing

motion was similar, regardless of skill level. Skilled typists type faster with less

muscle activity because they can utilize the passive tension of the muscle

tendons for keystrokes. However, to achieve this high keystroke speed, they

exhibit high active tension of the finger flexors. The active tension of the finger

flexors may also be higher in response to the reaction force received from the

keys as the fingertips press the keys, possibly increasing the mechanical stress

on the finger flexor tendons and becoming a risk factor for the occurrence of

disorders.

Unskilled typists require more time to switch between flexion and extension of

the hand because of the high load on the extensor muscles caused by the

excessive immobilization of the wrist joint and their strategy to prevent

keystrokes. This style of movement has a high risk for developing extensor

muscle-tendon disorders of the wrist.

Acknowledgements

The authors would like to acknowledge the members of the SPU Fundamental

Research group for their assistance.

Sources of funding

There is no applicable funding.

References

1. Billy C L So, Andy S K Cheng, Grace P Y Szeto. Cumulative IT Use Is

Associated with Psychosocial Stress Factors and Musculoskeletal

Symptoms. *Int. J. Environ. Res. Public Health* **2017**; 14: 1541; doi:

10.3390/ijerph14z121541.

2. U Bergqvist, E Wolgast, B Nilsson, M Voss. Musculoskeletal disorders

among visual display terminal workers: individual, ergonomic, and work

organizational factors. *Ergonomics* **1995**; 38(4): 763–776; doi:

10.1080/00140139508925148.

3. F Gerr, M Marcus, C Ensor, D Kleinbaum, S Cohen, A Edwards, E Gentry, D

J Ortizl, C Monteilh. A prospective study of computer users: I. Study design

and incidence of musculoskeletal symptoms and disorders. *Am J Ind Med*.

2002; 41(4): 221–235; doi:10.1002/ajim.10066.

4. C Jensen. Development of neck and hand-wrist symptoms in relation to

duration of computer use at work. *Scand J Work Environ Health*. **2003**; 29(3):

197–205; doi: 10.5271/sjweh.722.

5. C M Sommerich, W S Marras, M Parnianpour. A quantitative description of

typing biomechanics. *J Occup Rehabil*. **1996**; 6(1): 33–56; doi:

10.1007/BF02110393.

6. N A Baker, R Cham, E H Cidboye, J Cook, M S Redfern. Kinematics of the

fingers and hands during computer keyboard use. *Clin Biomech*. **2007**; 22(1):

34–43; doi: 10.1016/j.clinbiomech.2006.08.008.

7. C Zhao, P C Amadio, M E Zobitz, T Momose, P Couvreur, K N An. Effect of

synergistic motion on flexor digitorum profundus tendon excursion. *Clinical*

Orthop Relat Res. **2002**; (396): 223–230; doi: 10.1097/00003086-200203000-00033.

8. F-C Su, Y L Chou, C S Yang, G T Lin, K N An. Movement of finger joints induced by synergistic wrist motion. *Clin Biomech*. **2005**; 20(5): 491–497; doi: 10.1016/j.clinbiomech.2005.01.002.

9. J R Cook, N A Baker, R Cham, E Hale, M S Redfern. Measurements of wrist and finger postures: A comparison of goniometric and motion capture techniques. J Appl Biomech. **2007**; 23(1): 70–78; doi: 10.1123/jab.23.1.70.

10. E Takaoka, M Sugiura, H Komiya. A study on typist characterization. *IPSJ SIG Technical Report*. **2014**; 125(9): 1–14.

11. D Rempel, A Barr, D Brafman, Ed Young. The effect of six keyboard designs on wrist and forearm postures. *Appl Ergon*. **2007**; 38(3): 293–298; doi: 10.1016/j.apergo.2006.05.001.

12. G P Y Szeto, J Kim F N. A comparison of wrist posture and forearm muscle activities while using an alternative keyboard and a standard keyboard. *J Occup Rehabili*. **2000**; 10(3): 189–197; doi: 10.1023/A:1026662302131.

13. Gordon D, Logan, M J C Crump, Hierarchical Control of Cognitive Processes: The Case for Skilled Typewriting. *Psychology of Learning and*

Motivation. **2011**; (54): 1–27; doi: 10.1016/B978-0-12-385527-5.00001-2.

14. N Nagasawa. How Japanese university students type on smartphone and PC. *Computer and Education*. **2017**; (43): 67–72; doi: 10.14949/konpyutariyoukyouiku.43.67.

15. G G Simoneau, R W Marklin, J E Berman. Effect of Computer Keyboard Slope on Wrist Position and Forearm Electromyography of Typists Without Musculoskeletal Disorders. *Phys Ther*. **2003**; 83(9): 816–830; doi: 10.1093/ptj/83.9.816.

16. J T Dennerlein, I Kingma, B Visser, J. H van Dieën. The contribution of the wrist, elbow and shoulder joints to single-finger tapping. *J Biomech*. **2007**; 40(13): 3013–3022; doi: 10.1016/j.jbiomech.2007.01.025.

Fig.1 Explanation of experimental equipment.
(A) The positions of the markers and EMGs
(B) Calculation of MCP joint angle

Fig.2 Wrist and metacarpophalangeal joint motion
The error bar represents the standard deviation between subjects.
* P < 0.05

Fig.3 Muscle loading
From left to right, the static activity level (P = 0.1), the mean activity
level (P = 0.5), the maximum activity level (P = 0.9) are listed.
The error bar represents the standard deviation between subjects.
* P < 0.05

Fig.4 Press time data
From left to right, the wrist joint dorsiflexion angle, angular velocity,
the index finger MCP joint flexion angle, angular velocity, %MVC of
FCR, FCU, ECR, and ECU are listed.
The error bar represents the standard deviation between subjects.
* P < 0.05

Fig.5 Release time data
From left to right, the wrist joint dorsiflexion angle, angular velocity, the index finger MCP joint flexion angle, angular velocity, %MVC of FCR, FCU, ECR, and ECU are listed.
The error bar represents the standard deviation between subjects.
* P < 0.05

Table.1 Subjects and Environments
The mean value of WPM of all subjects was used as the benchmark for skill: subjects who achieved a score higher than the mean were classified into the ST group, and those who achieved a lower score than the mean into the UT group.

	Skilled Typist (n = 5)	Unskilled Typist (n = 7)	p
correct answers(%)	94.0 (2.3)	95.1 (2.6)	0.513
WPM	219.5 (21.3)	170.6 (36.2)	*0.033
age	21.6 (0.8)	20.7 (0.5)	0.050
height (cm)	167.3 (5.0)	166.1 (4.2)	0.700
weight (kg)	63.2 (8.9)	56.6 (6.4)	0.202
hand width (cm)	7.0 (0.4)	6.8 (0.1)	0.466
finger length (cm)	10.2 (0.4)	9.9 (0.3)	0.305
chair height (cm)	42.4 (3.9)	45.7 (2.8)	0.127
screen angle (°)	129.0(12.4)	122.9(17.5)	0.552
male-to-female ratio	3:2	6:1	—
right-left handed ratio	5:0	7:0	—
desktop-notebook PC ratio	1:4	0:7	—
frequency; PC use times per week	5.6 (1.0)	4.86 (1.3)	0.340
frequency; PC use times hours per days	3.8 (1.5)	2.93 (1.9)	0.445