Fitness barriers to spread of colistin resistance overcome by first establishing niche in patients with enhanced colistin exposure

Zena Lapp, PhD;[1,2] Jennifer H Han, MD;[3] Divya Choudhary, BTech;[4] Stuart Castaneda, MS;[1]

Ali Pirani, MS;[5] Kevin Alby, PhD;[6] Pam C Tolomeo, MPH, CCRP;[7] Ellie JC Goldstein, MD;[8]

Ebbing Lautenbach, MD;[9] Evan S Snitkin, PhD[5,10]

[1]Department of Computational Medicine and Bioinformatics, University of Michigan,

Ann Arbor, MI 48109

[2]Duke Global Health Institute, Duke University, Durham, NC 27710 (Z.L. current institution)

[3]GlaxoSmithKline, Rockville, MD 20850

[4]Department of Biochemistry, University of Oxford, Oxford, United Kingdom OX2 6HD

[5]Department of Microbiology and Immunology, University of Michigan, Ann Arbor, Michigan

48109

[6]University of North Carolina, Chapel Hill, NC 27599

[7]Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania, Perelman School

of Medicine, Philadelphia, PA 19104

[8]R M Alden Research Laboratory; David Geffen School of Medicine, University of California, Los

Angeles, Santa Monica, CA 90404

[9]Department of Medicine (Infectious Diseases), Department of Biostatistics, Epidemiology, and

Informatics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

[10]Department of Internal Medicine/Division of Infectious Diseases, University of Michigan, Ann

Arbor, MI 48109

Corresponding author

Evan Snitkin

27   1520D MSRB I, 1150 W. Medical Center Dr., Ann Arbor, MI, 48109-5680

28   (734) 647-6472

29   esnitkin@umich.edu

Author Contributions

32   E.S.S., J.H.H., E.L., and Z.L. conceptualized the study and acquired funding to support the

33   project. K.A. performed MIC testing. Z.L, A.P., S.C., and D.C. performed data curation and

34   formal analysis. Z.L. performed investigation and visualization. E.S.S. and Z.L. developed

35   methodology. E.S.S. supervised the project. Z.L. and E.S.S. wrote the original draft. All authors

36   reviewed and edited the manuscript.

Competing Interests Statement

39   J.H.H. is currently an employee of, and holds shares in, the GSK group of companies.

Classification

42   Major classification: Biological sciences

43   Minor classification: Microbiology

Keywords

Antibiotic resistance, resistance evolution, colistin, multidrug resistant organism, *Klebsiella*

pneumoniae, healthcare-associated pathogen, pathogen transmission

Abstract

There is an urgent need to improve our understanding of how new antibiotic resistant organisms

emerge and spread. A high-priority resistance threat is the ST258 lineage of carbapenem-

resistant *Klebsiella pneumoniae*. Here, we studied resistance to the last-line drug colistin among

ST258 by tracking its evolution across 21 U.S. hospitals over the course of a year. Phylogenetic

analysis supported a significant fitness cost being associated with resistance, as resistance

emergence was common but resistance variants were rarely transmitted. Furthermore, several

resistance variants that were transmitted had acquired secondary variants that reverted the

strain to susceptible. The exceptions to the general pattern of instability associated with

resistance were two large clusters of resistant strains in one sublineage (clade IIB) present

across Southern California hospitals. Quantification of transmission fitness in the healthcare

environment indicated that, while resistant isolates from other clades were less fit than their

susceptible counterparts, clade IIB resistant isolates were more fit, despite having similar

resistance variants. Additional analyses supported the increased fitness of colistin-resistant

clade IIB isolates being driven by a lineage-defining variant that increased clade IIB's

association with patient subpopulations with enhanced colistin exposure. These results show

that a favorable genetic background and sustained selective pressure led to the emergence and

spread of a colistin-resistant ST258 sublineage across a regional healthcare network. These

findings highlight the utility of integrating pathogen genomic and corresponding clinical data

from regional healthcare networks to detect emerging antibiotic resistance threats and

understand the clinical practices and patient populations that drive their spread.

Significance Statement

Selective pressure in hospitals leads to frequent antibiotic resistance evolution. However,

emergent resistance alleles are often not transmitted to other individuals because of fitness

costs associated with resistance. Due to the difficulty of studying pathogen fitness in humans,

our understanding of how resistant organisms circumvent these costs is limited. We integrate

genomic and clinical data to understand the evolutionary trajectories leading to transmissible

resistance for the last-line antibiotic colistin. While colistin resistance is generally associated

with a fitness cost that hinders transmission, this cost was mitigated in a sublineage that had

previously acquired mutations increasing its association with patient populations more likely to

receive colistin, suggesting a key role for historical contingency in the emergence and spread of

stable resistance.

Introduction

Multidrug resistant organisms (MDROs) pose a significant threat to public health due to uncontrolled transmission and dwindling treatment options (1). Of greatest concern are epidemic MDRO lineages that have become adapted to healthcare settings and continually gain resistance due to selective antibiotic pressure (2, 3). As resistance evolution continues to outpace our ability to develop new therapies (3), there is a critical need to improve our understanding of the forces driving the emergence and spread of resistance so that we can maintain the efficacy of last-line antibiotics. While numerous studies have documented evolutionary trajectories leading to stable antibiotic resistance in laboratory settings (4–6), these findings may not translate to the real world due to a failure to capture relevant selective pressures associated with patient-to-patient transmission in healthcare settings (6). However, in part due to the difficulty of quantifying fitness effects outside of the lab, similar studies *in situ* are scarce.

When *de novo* evolution of antibiotic resistance occurs in an individual, there is often little to no subsequent transmission to others due to the fitness cost of maintaining resistance in the absence of antibiotic exposure (7, 8). However, in some cases this barrier is overcome, as evidenced by the proliferation of epidemic resistance clones (8, 9). This may happen when the resistance variant or gene has an inherently low fitness cost and is thus maintained in the population even in the absence of antibiotic exposure (10). Alternatively, the fitness cost of a given resistance element may depend on the genetic background in which it emerges, leading to dissemination of these elements only in certain lineages. Indeed, experimental evidence has shown that the genetic background of a strain interacts with resistance determinants to influence the fitness of the resistant strain (11, 12), highlighting the importance of historical evolutionary events in determining the potential ability of a resistant strain to spread. While the

preferential spread of resistance elements in specific strains lends support to the importance of

genetic background, in most instances it is unclear what the underlying epistatic interactions are

that lead to these different fitness effects, particularly *in situ* (8). Understanding how the strain

genetic background, in combination with selective pressures in healthcare settings, drives the

emergence and spread of antibiotic resistance is critical for both early detection of high-risk

clones and devising strategies to contain them.

Carbapenem-resistant *Klebsiella pneumoniae* (CRKP) is a global threat as it is resistant to

nearly all available antibiotics and is associated with high mortality rates (1). One of the few

remaining treatment options for CRKP is the antibiotic colistin (13). Concerningly, the viability of

colistin as a treatment option moving forward is threatened by the frequent emergence of

colistin resistance during treatment with this drug (14, 15). However, despite the accessibility of

these resistance variants to CRKP, their spread to other patients is rare, with most cases even

within a single healthcare setting stemming from parallel evolution (15). Thus, even though no

fitness effects for colistin resistance have been detected *in vitro* (16–18), there appears to be a

significant fitness cost *in situ*. Here, we document the emergence and spread of multiple colistin

resistance variants in CRKP sequence type (ST) 258 across a regional healthcare network.

Through integration of genomic and clinical data, we show how resistance spread was

enhanced in genetic backgrounds that mitigated the fitness cost of resistance, thereby enabling

regional dissemination of colistin-resistant lineages.

Results

Most resistant isolates contain variants in known resistance genes.

Antimicrobial susceptibility testing revealed that 118/337 (35%) CRKP ST258 isolates were

resistant to colistin, and that resistance was found consistently across long-term acute care

hospitals (LTACHs) over the yearlong study (**Fig S1**). To identify likely resistance variants, we

first searched all isolates for *mcr*-containing mobile genetic elements and variants in canonical

chromosomal genes known to confer resistance (*pmrA*, *pmrB*, *phoP*, *phoQ*, *crrA*, *crrB*, and

mgrB). We did not find *mcr* genes in any of the isolates, as expected since *mcr* genes have not

been found in CRKP isolates from the U.S. during the timeframe of the study (July 2014 to

August 2015) (19). However, the majority of colistin resistance we observed (103/118, 87%)

could be explained by known and putative resistance variants in canonical resistance genes

(**Fig S2**; **Table S1**; see methods for details on identification of resistance variants).

As not all resistance in our dataset could be explained by variants in known resistance genes,

we next performed a genome-wide association study (GWAS)(20) on the isolates with unknown

resistance determinants to identify putative non-canonical resistance-conferring variants. We

identified two additional putative resistance genes in this way: *qseC* and phosphotransferase

system sugar transporter subunit IIB (**Fig S2**; **Table S1**). These putative resistance variants

account for 6/15 (40%) of the resistant isolates lacking variants in known resistance genes in

our dataset. Notably, *qseC*, which explained resistance in 4 isolates, has been shown to confer

colistin resistance in an experimental evolution study using clinical isolates (21). We were

unable to identify putative colistin resistance-conferring variants for the other nine isolates. For

all subsequent analyses, we defined resistance genes as the set of canonical and GWAS-

identified resistance genes.

<u>Epistatic interactions appear to influence resistance in isolates with more than one variant in</u>

<u>resistance genes.</u>

To further explore the relationship between genetic variation at putative resistance loci and

colistin resistance we next examined the number of variants in resistance genes among

resistant and susceptible isolates. Supporting a causal role for genetic variation in putative

resistance loci, we observed that while only 9/154 (5.8%) isolates without a variant in a putative

resistance locus were phenotypically resistant, 93/122 (76.2%) isolates with a single variant in a

putative resistance locus were colistin resistant. However, when expanding to look at isolates

with two or more variants in putative resistance loci, the association with resistance inverted,

with 45/61 (73.8%) isolates being phenotypically susceptible and 16/61 (26.2%) resistant (**Fig**

1A). This unexpected relationship between accumulation of multiple resistance variants and

susceptibility was also observed at the level of colistin MIC, with susceptible isolates with two or

more variants in resistance genes having lower MICs than those with one or fewer (**Fig 1B**;

median MIC of 0.5 vs. 1, Wilcox $p = 0.0002$).

We hypothesized that the enrichment for susceptibility among isolates with multiple variants in

resistance loci could be a consequence of the fitness cost of resistance leading to selection for

secondary mutations that abrogated resistance, and thereby mitigated the fitness cost.

Supporting this hypothesis, we found that in 36/45 (80%) cases where susceptible isolates

harbored multiple variants in resistance genes, at least one of the variants or variant sites had

been previously shown to confer resistance *in vitro* (**Table S1**), suggesting that the secondary

variants present in resistance genes may be suppressor mutations that revert historically

resistant strains back to susceptible. This hypothesis was further corroborated by visualizing

these variants on the phylogeny (**Fig 2**). For instance, we observed several instances where

putative suppressor mutations have arisen in a lineage containing a resistance-conferring *phoQ*

mutation. These and other putative suppressor mutations usually occurred within the same

gene or in a downstream gene of the molecular pathway to resistance (**Fig S3**), and were

sometimes followed by re-acquisition of resistance through a different mechanism (**Fig 2**; **Fig**

S4). Taken together, these findings suggest that, while variants in resistance genes are often

associated with an increase in MIC (**Fig 1B**), there also exist epistatic interactions among

variants in these genes that may lead to colistin susceptibility, even in strains containing known

resistance-conferring mutations (22, 23).

<u>Colistin resistance exhibits patterns of *de novo* evolution, onward dissemination, and reversion</u>

<u>to susceptibility.</u>

To better understand the origin and fate of colistin resistance variants, we further investigated

the phylogenetic relationship among colistin-resistant and susceptible isolates (**Fig 2**; **Fig S5**).

Based on the observed resistance distribution on the phylogeny we classified each isolate into

one of four groups: resistant cluster (sub-clade harboring multiple resistant isolates), resistant

singleton (single resistant isolate with susceptible phylogenetic neighbors), susceptible revertant

(susceptible but contains a putative resistance-conferring variant), and susceptible non-

revertant (other susceptible isolates). Grouping resistance in this way revealed that, while 70%

(28/40) of resistance emergence events did not spread between patients (i.e. were resistant

singletons), the 12 that did spread accounted for 76% (90/118) of resistance across study

facilities. Visualization of these resistance clusters on the phylogeny revealed a striking

dichotomy between a clade II sublineage (referred to as clade IIB) and the other ST258

sublineages (clades I and IIA). Clade IIB contained two large clusters of colistin-resistant

isolates, while the other two sublineages mainly exhibited sporadic parallel evolution of

resistance. In total, these two clonal expansions in clade IIB, which occurred across 11/12

LTACHs in Southern California (10 in the Los Angeles area and 1 in San Diego; **Fig S6**),

accounted for over half (69/118, 58%) of resistance in the isolate collection. Each cluster

harbored a distinct resistance mutation, with one being associated with a nonsense mutation in

mgrB (Gln30*) and the other a missense mutation in *phoQ* (Thr244Asn), respectively. Notably,

these same types of variants occurred in the sporadic colistin-resistant isolates (**Fig 2**; **Fig S4**),

suggesting that these specific resistance variants were likely not inherently more fit. Within

these clonal expansions, many of the isolates appeared to regain susceptibility to colistin via the

accumulation of additional variants in resistance genes as described above, suggesting that

despite their proliferation, there is still a fitness cost associated with resistance.

<u>Resistant strains in clade IIB are more fit than their susceptible non-revertant counterparts.</u>

Next, we were interested in understanding whether the widespread dissemination of colistin

resistance variants in clade IIB could be explained by an increased ability to spread relative to

resistant strains in the other two clades. In particular, we hypothesized that the success of

resistance variants in clade IIB could be a consequence of a decreased fitness cost associated

with resistance, as compared to the cost of resistance in other clades. While previous studies

have not found a fitness cost associated with colistin resistance in *K. pneumoniae* (16–18), this

is inconsistent with observed lack of onward transmission of most resistant mutants (15). We

hypothesized that this was because the measurement of fitness in laboratory conditions fails to

capture the range of stresses encountered during colonization and transmission among patients

in healthcare settings. To overcome this issue and estimate *in situ* transmission fitness of

resistant and susceptible isolates from different lineages in the context of the healthcare

environment, we used time-scaled haplotypic density (THD), an analytic approach that uses the

genetic relatedness between isolates to quantify the extent of relative spread of each strain (24).

A higher THD value is consistent with higher transmission fitness (i.e. ability to spread).

In clades I and IIA we observed evidence of a significant transmission fitness cost for resistance

variants, as resistance clusters and susceptible revertants were less able to spread than

susceptible non-revertants from those clades (**Fig 3**; R cluster: Wilcox p = 0.04; S revertant:

Wilcox p = 0.02). . In contrast to clades I and IIA, resistance variants in clade IIB were

associated with a significant transmission fitness benefit, with resistant clusters and susceptible

revertants from clade IIB being more able to spread than susceptible non-revertants from that

clade (R cluster: Wilcox p = 0.03; S revertant: Wilcox p = 0.009). Taken together, these findings

suggest that resistance variants in isolates from clade IIB conferred a transmission fitness

advantage, which may account for the independent emergence and spread of two different

resistance alleles in this clade. In contrast, the accumulation of variants in resistance genes in

the other clades appears to be associated with a transmission fitness cost, which is consistent

with their limited clonal spread.

<u>Clade IIB isolates have more colistin exposure compared to isolates from other clades.</u>

Lastly, we investigated whether colistin use in study facilities influenced the emergence and

spread of stable resistance in clade IIB. Supporting the role of colistin use in the spread of

colistin resistance, we found a positive association between having a clade IIB isolate and both

prior exposure to colistin (15/120, 12.5% in clade IIB vs. 8/217, 3.7% in other clades; Fisher's

exact p = 0.003), as well as treatment with colistin (41/120, 34.2% in clade IIB vs. 39/217,

18.0% in other clades; Fisher's exact p = 0.001), regardless of colistin mutation status. It is

notable that most of the colistin exposure in patients harboring clade IIB isolates occurred after

patients acquired their isolates, which suggests that the association isn't simply a reflection of

antibiotic exposure increasing the risk for acquiring isolates resistant to the cognate antibiotic.

Rather, we hypothesized that this association between colistin use and clade IIB is mediated by

our previously reported observation that clade IIB is enriched in respiratory isolates (25) and the

frequent use of colistin to treat respiratory tract infections during the study period (**Fig 4A**). This

hypothesis is supported by a slight enrichment in colistin treatment for clade IIB respiratory

isolates compared to non-respiratory isolates (31/83, 37.3% for respiratory vs. 10/37, 27.0% for

non-respiratory; Fisher's exact p = 0.3), and a significant enrichment across all clades (53/183,

29% for respiratory vs. 27/154, 17.5% for non-respiratory; Fisher's exact p = 0.01). Furthermore,

in addition to potentially influencing the spread of colistin resistance, the extensive use of

colistin to treat infections in a setting of clonally disseminating colistin resistance led to 36/118

(30.5%) patients with colistin-resistant isolates being treated with this antibiotic, even though the

treatment was likely ineffective (**Fig 4B**).

In putting our past and present observations together, we noted that a defining feature of clade

IIB was disruption of the putative O-antigen glycosyltransferase *kfoC* (25). Therefore, we

hypothesize that disruption of *kfoC* increased the affinity of clade IIB for the respiratory tract,

which in turn resulted in the clade IIB lineage having increased rates of colistin exposure, and

thereby decreased the transmission fitness cost of colistin resistance, allowing colistin-resistant

clade IIB strains to spread across a regional healthcare network (**Fig 5**).

Discussion

Improving our understanding of how antibiotic resistance evolves and disseminates in the

healthcare setting is critical for optimizing surveillance, antibiotic stewardship, and infection

prevention strategies. Here, we used genomic data from a comprehensive regional sample of

clinical CRKP ST258 isolates to track the origin and fate of colistin resistance variants, and then

overlaid clinical data to elucidate the drivers of resistance spread. Our analysis identified a

circulating sublineage whose genetic background appears to decrease the fitness cost of

resistance *in situ* by increasing association with patient populations with increased colistin

exposure, thus allowing colistin-resistant strains to spread among regional healthcare facilities.

More broadly, these findings highlight how regional genomic surveillance can be applied to hone

in on emerging resistance threats with epidemic potential, as well as uncover patient

populations and antibiotic usage patterns accelerating their spread.

Examination of colistin resistance variants within the comprehensive and longitudinal context of

circulating CRKP strains allowed for inferences into their functional impact and clinical

12

significance. We discovered that resistant isolates with multiple variants in resistance genes

have increased resistance, but susceptible isolates with multiple variants in resistance genes

have decreased resistance. Additionally, we identified several instances where a known

resistance variant was present in susceptible isolates, likely due to a suppressor variant in either

the same resistance gene or a downstream resistance gene (22, 23). This finding has two

important implications. First, known resistance variants occurring in both resistant and

susceptible isolates complicates the computational identification of novel resistance

determinants using GWAS. Possible solutions include increasing sample size or removing

isolates with known resistance variants. Second, the presence of a colistin resistance variant

cannot necessarily be equated with *in vitro* phenotypic resistance. Clinically, this suggests that

testing for resistance variants may not be a substitute for testing for phenotypic resistance, as

colistin treatment may still be effective for some strains harboring resistance variants. In addition

to being phenotypically susceptible, revertants that harbor resistance variants may in general

have more difficulty becoming resistant once more, which may lend credence to treating these

patients with colistin (23).

Of particular genomic and clinical interest are two independent instances of emergence and

spread of colistin resistance variants in clade IIB, in stark contrast to the sporadic emergence of

resistance variants in the other clades. Notably, the types of resistance variants present in clade

IIB also occurred in other clades, implying that the nature of the variant itself is not what allowed

the clade IIB variants to spread. Instead, our analysis of transmission fitness using THD

suggests that these colistin resistance variants were able to spread in this clade because the

genetic background of the clade reduced the fitness cost of the variants. We previously showed

that a defining feature of clade IIB is that the majority of isolates contain a disruption in the

lipopolysaccharide (LPS) O locus gene *kfoC* (25), a putative glycosyltransferase. Given the key

role that LPS plays in both colistin-mediated killing and colistin resistance (14), it is possible that

inactivation of *kfoC* facilitates the spread of colistin resistance by altering the outer membrane in

a way that reduces the fitness cost of resistance-conferring LPS modifications. Furthermore, our

data supports that *kfoC* disruption may indirectly mitigate the fitness cost of colistin resistance

by increasing the association between clade IIB and patient populations in which colistin use is

more frequent. In particular, clade IIB isolates are significantly enriched in both association with

the respiratory tract and colistin exposure. Thus, increased colistin use in clade IIB may have

both set the stage for the emergence of multiple instances of stable resistance in this clade and

provided a continued selective pressure to maintain resistance. These findings highlight the

importance of continued real-time monitoring of resistance to improve antibiotic stewardship and

decrease the likely ineffective use of colistin to treat colistin-resistant CRKP strains.

One strength of our study is that we have comprehensive sampling of clinical CRKP isolates

from LTACHs over the course of a year. This not only allowed us to capture putative

transmission events of colistin resistance, but also allowed us to estimate the transmission

fitness of different ST258 sublineages *in situ*. Additionally, whole-genome sequencing of study

isolates permitted us to interrogate not only variants in known resistance genes, but also

investigate putative resistance-conferring variants in other genes using GWAS. Furthermore,

our access to clinical information about colistin use allowed us to identify a relationship between

this clinical variable and different ST258 sublineages, providing us with insight into the selective

pressures of colistin resistance evolution and spread in patients.

Our study also has several limitations. First, we do not have rectal surveillance cultures, which

limits our ability to fully capture the population of colistin-resistant and susceptible CRKP in the

LTACHs studied, and thus to fully define spread. While this could bias our clinical collection if

certain strains are more likely to present clinically and be cultured at extraintestinal sites (e.g.

clade IIB vs. other clades), the strains we have included are the most clinically relevant since

they were identified in clinically derived samples. Also, we cannot be certain that our

classification of isolates into different categories of colistin susceptibility and resistance is

entirely accurate. For instance, putative *de novo* evolution of resistance in resistant singletons

may not have occurred in the patient we sampled. However, our comprehensive sampling of

clinical isolates from the facilities in the study provides some confidence that resistance in

resistant singleton isolates did in fact evolve in that patient. Another limitation of our study is that

we do not have information about colistin exposure prior to the patient entering the healthcare

facility, and we do not have information about the dose or duration of colistin use. Even with this

limitation, we were able to identify significant associations between colistin use and different

ST258 sublineages.

In conclusion, we identified an emerging ST258 sublineage that is more fit and more amenable

to maintaining colistin resistance than other ST258 strains *in situ*, likely due to an altered

genetic background. This is of particular concern due to the already limited treatment options for

CRKP, and therefore merits further surveillance to determine the extent of spread. Furthermore,

our findings highlight how the inclusion of relevant clinical data amplifies the value of regional

genomic surveillance by enabling identification of novel resistance and virulence variants of

interest, as well as revealing the clinical practices and patient populations driving their spread.

Materials and Methods

Study isolates and metadata

We used whole-genome sequences of clinical CRKP isolates from a prospective longitudinal

study in 21 U.S. LTACHs over the course of a year (BioProject accession no. PRJNA415194)

(26). Isolates and metadata were collected, and isolates were sequenced, as described in Han

et al. (26). Prior patient use of colistin in the facility from which the isolate was taken (up to 30

days before the isolate collection date), as well as colistin use for empiric and definitive

treatment, was extracted from the electronic health record. We only used information on colistin,

not polymyxin B, for all analyses.

Antibiotic susceptibility testing

Custom Sensititre broth microdilution panels (ThermoFisher Scientific, Waltham, MA) were used

to determine colistin MICs. We used Clinical & Laboratory Standards Institute (CLSI) guidelines

for interpretations with an MIC cutoff of ≥4 characterized as resistant (27). CLSI guidelines

classify isolates with an MIC ≤2 as being of intermediate resistance and therefore are

technically considered non-resistant using this interpretation; however, the European Committee

on Antimicrobial Susceptibility Testing defines isolates with an MIC ≤2 as susceptible (28). For

clarity, we chose to use the term susceptible for isolates with an MIC ≤2.

Isolate selection

Multi-locus sequence types were called using ARIBA (29). Over 90% of CRKP isolates collected

over the course of the study belonged to ST258; therefore, we focus all of our analyses on this

sequence type. We ordered the isolates by resistance status followed by collection date. For all

analyses except GWAS, only the first patient ST258 isolate from this ordered list was used, thus

prioritizing resistant isolates over susceptible isolates. Sample sizes were too small to glean

insights into within-host colistin resistance evolution.

Single nucleotide variant calling, indel calling, and phylogenetic tree reconstruction

Variant calling was performed with a customized variant calling pipeline

(https://github.com/Snitkin-Lab-Umich/variant_calling_pipeline/) as follows. The quality of

sequencing reads was assessed with FastQC v0.11.9 (30), and Trimmomatic v0.39 (31) was

used for trimming adapter sequences and low-quality bases. Single nucleotide variants (SNVs)

were identified by (i) mapping filtered reads to the ST258 KPNIH1 reference genome

(BioProject accession no. PRJNA73191) using the Burrows-Wheeler short-read aligner (bwa

v0.7.17) (32), (ii) discarding polymerase chain reaction duplicates with Picard v2.24.1 (33), and

(iii) calling variants with SAMtools and bcftools v1.9 (34). Variants were filtered from raw results

using VariantFiltration from GATK v4.1.9.0 (35) (QUAL >100; MQ >50; ≥10 reads supporting

variant; and FQ < 0.025). Indels were called using the GATK HaplotypeCaller (36) with the

following filters: root mean square quality (MQ) > 50.0, GATK QualbyDepth (QD) > 2.0, read

depth (DP) > 9.0, and allele frequency (AF) > 0.9. In addition, a custom Python script was used

to filter out (mask) variants in the whole-genome alignment that were: (i) SNVs <5 base pairs

(bp) in proximity to indels, (ii) in a recombinant region identified by Gubbins v2.3.4 (37), in a

phage region identified by the Phaster web tool (38) or (iii) they resided in tandem repeats of

length greater than 20bp as determined using the exact-tandem program in MUMmer v3.23

(39). This whole-genome masked variant alignment was used to reconstruct a maximum

likelihood phylogeny with IQ-TREE v1.6.12 (40) using the general time reversible model GTR+G

and ultrafast bootstrap with 1000 replicates (-bb 1000) (41).

Insertion calling

Large insertions relative to the reference genome were called using panISa v0.1.4 (42).

Variant preprocessing

We preprocessed variants to include multiallelic sites and used the major allele method for

variant binarization, as described in Saund *et al.* (43). We used SnpEff to predict the functional

impact of single nucleotide variants and indels (high, moderate, low, modifier) (44). Additionally,

we considered all insertions in or upstream of genes as high impact, and those downstream of

17

genes as moderate impact. Only high and moderate impact variants were included in

downstream analyses.

<u>Identification of putative resistance variants</u>

Known resistance genes

We consider the following genes known (canonical) resistance genes: *mgrB*, *phoP*, *phoQ*, *crrA*,

crrB, *pmrA*, and *pmrB* (45). We group variants in known resistance genes into the following

categories, in order of decreasing confidence:

1. <u>Known:</u> experimentally confirmed resistance variants, including loss-of-function variants

in *mgrB* (45–48).

2. <u>Known site:</u> the variant occurs at a nucleotide site where there is an experimentally

confirmed resistance variant, but that specific amino acid change has not been

experimentally confirmed (22, 49–51).

3. <u>Putative:</u> nonsynonymous or disruptive variants in known resistance genes where >60%

of the variants are present in resistant isolates.

Genome-wide association study

We performed a burden test using treeWAS v1.0 (20), a convergence-based GWAS method, as

our sample size is relatively small and resistance is a very convergent phenotype. A burden test

increases power to detect resistance genes, and convergence-based methods control for the

structure of the phylogeny more than mixed model methods. For GWAS, we included only

isolates with no mutations in known resistance genes as we were most interested in identifying

novel resistance genes, and because the entire dataset contained a number of susceptible

isolates with known resistance variants that would have confounded the analysis. We used

pyseer v1.3.6 (52) to calculate the number of unique patterns and determine a p-value cutoff (p

< 9.47e-5). Putative resistance genes were considered ones that were identified as significant

by treeWAS, had more than one convergence event on the phylogeny, and >60% of all the

variants in the gene were found in resistant isolates. Variants within these genes were

considered putative resistance variants using the same definition as for putative resistance

variants in known resistance genes. Using these requirements, we included four putative

resistance variants from two putative resistance genes based on the GWAS results.

We define resistance genes as the set of all known and GWAS-identified resistance genes, and

resistance variants as the set of all known and putative resistance variants in those genes.

Identification of putative suppressor variants

We define putative suppressor variants as those in resistance genes where >60% of isolates

with the variant are susceptible and contain a resistance variant.

Comparison of isolate MICs

The significance of MIC differences between different groups of isolates was determined using

two-sided Wilcox tests.

Determination of isolate resistance group

We assigned each isolate to one of four categories (**Fig S3**):

1. Resistant singletons: resistant isolates that do not cluster on the phylogeny, or that

466   cluster but contain distinct resistance variants.

467   2. Resistant clusters: resistant isolates that cluster on the phylogeny and contain the same

468   resistance variant. Additionally, if a cluster on the phylogeny has unknown resistance

469   variants, we also defined it as a resistant cluster.

3. Susceptible revertants: susceptible isolates that contain a resistance variant.

4. Susceptible non-revertants: susceptible isolates that do not contain a resistance variant.

For the purposes of identifying resistant clusters that include susceptible revertants, we

classified an isolate as "quasi-resistant" if it contained a known resistance variant and/or was

resistant. Clusters of "quasi-resistant" isolates were identified using the get_clusters function in

regentrans v0.1 (https://github.com/Snitkin-Lab-Umich/regentrans) (53) with a pureness of 1.

Each isolate in these clusters was then denoted as a resistant cluster or susceptible revertant

isolate depending on their corresponding phenotype.

Calculation of transmission fitness using time-scaled haplotypic density

Even though there is a clear transmission fitness cost associated with colistin resistance in the

healthcare setting as evidenced by our finding that the vast majority of resistant strains do not

spread, previous experimental approaches to studying the fitness of colistin-resistant CRKP

have found no fitness cost to resistance (17, 54). Therefore, we define transmission fitness in

the healthcare setting as the epidemic success of the strain based on time-scaled haplotypic

density (THD) (24). Given a genetic distance matrix, a mutation rate, and a timescale, THD uses

kernel density estimation to estimate the frequency of transmission events during that time

period. We calculated the THD for each isolate with the R package thd v1.0.1 (24) using the

KPNIH1 reference genome length (5,394,056 base pairs), a mutation rate of 1.03e-6 (55), the

time scale parameter, and a timescale of one year. Pairwise single nucleotide variant distances

were calculated with the dist.dna function in ape v5.4.1(56) (model = 'N', pairwise.deletion =

TRUE). Prior to performing comparisons, we used the clusters identified with regentrans to

remove isolates that were the only representative of their resistance group in a given cluster (i.e.

all resistant singletons, single susceptible revertant isolates that were embedded in a resistant

cluster, and single resistant cluster isolates that were embedded in a susceptible revertant

cluster. We did this because comparisons of THD only make sense in the context of other

isolates. In other words, the use of genetic context for inferring transmission fitness prevents

capturing the fitness cost or benefit in these singleton isolates. Significant differences between

THD of different isolates were determined using two-sided Wilcox tests.

Calculation of previous colistin use and colistin treatment significance

We used two-sided Fisher's exact tests to calculate the significance of the association between

various groups of isolates and previous colistin use, as well as between various groups of

isolates and colistin treatment.

Data visualization

We performed all data visualization in R v4.0.2 (57) using the following packages: tidyverse

v1.3.0 (58), ggtree v2.2.4 (59, 60), pheatmap v1.0.12 (61), ggplotify v0.0.5 (62), and cowplot

v1.1.0 (63). Code and data corresponding to the manuscript figures can be found on GitHub

(https://github.com/Snitkin-Lab-Umich/ltach-crkp-colistin-ms).

Acknowledgements

We gratefully acknowledge Kindred Healthcare for their assistance in collecting data and

isolates used in this study. We also thank Sean Muldoon for his support and guidance

throughout the study. Finally, we thank the patients and staff of the long-term acute-care

hospitals (LTACHs) for their gracious participation in this study, and Mary Hayden and Robert

Weinstein for critical review of the manuscript.

Funding

This research was supported by a CDC Cooperative Agreement FOA no. CK16-004-Epicenters

for the Prevention of Healthcare Associated Infections and the National Institutes of Health R01

AI139240-01 and 1R01 AI148259-01. Z.L. received support from the National Science

Foundation Graduate Research Fellowship Program under grant no. DGE 1256260. Any

opinions, findings, and conclusions or recommendations expressed in this material are those of

the authors and do not necessarily reflect the views of the National Science Foundation. The

funding bodies had no role in the design of the study or collection, analysis, and interpretation of

data, or in writing the manuscript.

References

1.   E. Tacconelli, *et al.*, Discovery, research, and development of new antibiotics: the WHO

priority list of antibiotic-resistant bacteria and tuberculosis. *The Lancet Infectious*

Diseases (2017) https:/doi.org/10.1016/S1473-3099(17)30753-3 (February 11, 2018).

2.   R. C. MacLean, A. S. Millan, The evolution of antibiotic resistance. *Science* **365**, 1082–

1083 (2019).

3.   World Health Organization, Ed., *Antimicrobial resistance: global report on surveillance*

(World Health Organization, 2014).

4.   G. G. Perron, M. Zasloff, G. Bell, Experimental evolution of resistance to an antimicrobial

peptide. *Proceedings of the Royal Society B: Biological Sciences* **273**, 251–256 (2006).

5.   A. C. Palmer, R. Kishony, Understanding, predicting and manipulating the genotypic

evolution of antibiotic resistance. *Nat Rev Genet* **14**, 243–248 (2013).

6.   T. J. Kawecki, *et al.*, Experimental evolution. *Trends in Ecology & Evolution* **27**, 547–560

(2012).

7. ,   The cost of antibiotic resistance from a bacterial perspective. *Drug Resistance Updates*

3, 237–245 (2000).

8.   S. Baker, N. Thomson, F.-X. Weill, K. E. Holt, Genomic insights into the emergence and

spread of antimicrobial-resistant bacterial pathogens. *Science* **360**, 733–738 (2018).

9.   L. K. Logan, R. A. Weinstein, The Epidemiology of Carbapenem-Resistant

Enterobacteriaceae: The Impact and Evolution of a Global Menace. *J Infect Dis* **215**,

S28–S36 (2017).

10.   A. H. Melnyk, A. Wong, R. Kassen, The fitness costs of antibiotic resistance mutations.

Evol Appl **8**, 273–283 (2015).

11.   T. Vogwill, M. Kojadinovic, R. C. MacLean, Epistasis between antibiotic resistance

mutations and genetic background shape the fitness effect of resistance across species

of Pseudomonas. *Proc Biol Sci* **283** (2016).

12.   T. Vogwill, M. Kojadinovic, V. Furió, R. C. MacLean, Testing the Role of Genetic

Background in Parallel Evolution Using the Comparative Experimental Evolution of

Antibiotic Resistance. *Molecular Biology and Evolution* **31**, 3314–3323 (2014).

13.   Y. Doi, Treatment Options for Carbapenem-resistant Gram-negative Bacterial Infections.

Clinical Infectious Diseases **69**, S565–S575 (2019).

14.   L. Poirel, A. Jayol, P. Nordmann, Polymyxins: Antibacterial Activity, Susceptibility Testing,

and Resistance Mechanisms Encoded by Plasmids or Chromosomes. *Clin. Microbiol.*

Rev. **30**, 557–596 (2017).

15.   N. Macesic, *et al.*, Emergence of Polymyxin Resistance in Clinical Klebsiella pneumoniae

Through Diverse Genetic Adaptations: A Genomic, Retrospective Cohort Study. *Clinical*

Infectious Diseases **70**, 2084–2091 (2020).

16.   A. Cannatelli, *et al.*, In Vivo Evolution to Colistin Resistance by PmrB Sensor Kinase

Mutation in KPC-Producing Klebsiella pneumoniae Is Associated with Low-Dosage

Colistin Treatment. *Antimicrobial Agents and Chemotherapy* **58**, 4399–4403 (2014).

17.   A. Cannatelli, A. Santos-Lopez, T. Giani, B. Gonzalez-Zorn, G. M. Rossolini, Polymyxin

resistance caused by mgrB inactivation is not associated with significant biological cost in

Klebsiella pneumoniae. *Antimicrob Agents Chemother* **59**, 2898–2900 (2015).

18.   A. B. Janssen, *et al.*, Evolution of Colistin Resistance in the Klebsiella pneumoniae

Complex Follows Multiple Evolutionary Trajectories with Variable Effects on Fitness and

Virulence Characteristics. *Antimicrobial Agents and Chemotherapy* **65**, e01958-20.

19.   R. L. Skov, D. L. Monnet, Plasmid-mediated colistin resistance (mcr-1 gene): three

576   months later, the story unfolds. *Eurosurveillance* **21**, 30155 (2016).

577   20.   C. Collins, X. Didelot, A phylogenetic method to perform genome-wide association

578   studies in microbes that accounts for population structure and recombination. *PLOS*

579   *Computational Biology* **14**, e1005958 (2018).

580   21.   M. E. Pitt, *et al.*, Octapeptin C4 and polymyxin resistance occur via distinct pathways in

581   an epidemic XDR Klebsiella pneumoniae ST258 isolate. *J. Antimicrob. Chemother.*

582   (2018) https:/doi.org/10.1093/jac/dky458.

583   22.   M. Pitt, *et al.*, Multifactorial Chromosomal Variants Regulate Polymyxin Resistance In

584   Extensively Drug-Resistant Klebsiella pneumoniae (2017).

585   23.   E. S. Snitkin, *et al.*, Genomic insights into the fate of colistin resistance and Acinetobacter

586   baumannii during patient treatment. *Genome Res.* **23**, 1155–1162 (2013).

587   24.   J.-P. Rasigade, *et al.*, Strain-specific estimation of epidemic success provides insights

588   into the transmission dynamics of tuberculosis. *Scientific Reports* **7**, 45326 (2017).

589   25.   Z. Lapp, *et al.*, Patient and Microbial Genomic Factors Associated with Carbapenem-

590   Resistant Klebsiella pneumoniae Extraintestinal Colonization and Infection. *mSystems* **6**

591   (2021).

592   26.   J. H. Han, *et al.*, Whole-Genome Sequencing To Identify Drivers of Carbapenem-

593   Resistant Klebsiella pneumoniae Transmission within and between Regional Long-Term

594   Acute-Care Hospitals. *Antimicrobial Agents and Chemotherapy* **63** (2019).

595   27.   M. P. Weinstein, M100-performance standards for antimicrobial susceptibility testing, 30th

596   edition. (2020) (March 26, 2021).

597   28. ,   EUCAST: Clinical breakpoints and dosing of antibiotics (June 7, 2021).

598   29.   M. Hunt, *et al.*, ARIBA: rapid antimicrobial resistance genotyping directly from sequencing

599   reads. *Microb Genom* **3** (2017).

600   30.   S. Andrews, *s-andrews/FastQC* (2021) (March 10, 2021).

24

31.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* **30**, 2114–2120 (2014).

32.   H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

33. ,   *broadinstitute/picard* (Broad Institute, 2021) (February 20, 2021).

34.   H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

2078–2079 (2009).

35.   A. McKenna, *et al.*, The Genome Analysis Toolkit: a MapReduce framework for analyzing

next-generation DNA sequencing data. *Genome Res* **20**, 1297–1303 (2010).

36.   R. Poplin, *et al.*, Scaling accurate genetic variant discovery to tens of thousands of

samples. *bioRxiv*, 201178 (2018).

37.   N. J. Croucher, *et al.*, Rapid phylogenetic analysis of large samples of recombinant

bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* **43**, e15–e15

(2015).

38.   D. Arndt, *et al.*, PHASTER: a better, faster version of the PHAST phage search tool.

Nucleic Acids Res **44**, W16–W21 (2016).

39.   S. Kurtz, *et al.*, Versatile and open software for comparing large genomes. *Genome*

Biology **5**, R12 (2004).

40.   L.-T. Nguyen, H. A. Schmidt, A. von Haeseler, B. Q. Minh, IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol* **32**,

268–274 (2015).

41.   B. Q. Minh, M. A. T. Nguyen, A. von Haeseler, Ultrafast Approximation for Phylogenetic

Bootstrap. *Mol Biol Evol* **30**, 1188–1195 (2013).

42.   P. Treepong, *et al.*, panISa: ab initio detection of insertion sequences in bacterial

genomes from short read sequence data. *Bioinformatics* **34**, 3795–3800 (2018).

43.   K. Saund, Z. Lapp, S. N. Thiede, A. Pirani, E. S. Snitkin, prewas: data pre-processing for

more informative bacterial GWAS. *Microbial Genomics,* **6**, e000368 (2020).

44.   P. Cingolani, *et al.*, A program for annotating and predicting the effects of single

nucleotide polymorphisms, SnpEff. *Fly (Austin)* **6**, 80–92 (2012).

45.   B. Berglund, Acquired Resistance to Colistin via Chromosomal And Plasmid-Mediated

Mechanisms in Klebsiella pneumoniae. *Infectious Microbes & Diseases* **1**, 10–19 (2019).

46.   A. O. Olaitan, S. Morand, J.-M. Rolain, Mechanisms of polymyxin resistance: acquired

and intrinsic resistance in bacteria. *Front. Microbiol.* **5** (2014).

47.   A. Jayol, *et al.*, High-Level Resistance to Colistin Mediated by Various Mutations in the

crrB Gene among Carbapenemase-Producing Klebsiella pneumoniae. *Antimicrob Agents*

Chemother **61** (2017).

48.   Y.-H. Cheng, T.-L. Lin, Y.-T. Lin, J.-T. Wang, Amino Acid Substitutions of CrrB

Responsible for Resistance to Colistin through CrrC in Klebsiella pneumoniae.

Antimicrobial Agents and Chemotherapy **60**, 3709–3716 (2016).

49.   S. Matamouros, K. R. Hager, S. I. Miller, HAMP Domain Rotation and Tilting Movements

Associated with Signal Transduction in the PhoQ Sensor Kinase. *mBio* **6** (2015).

50.   S. S. Yadavalli, *et al.*, Functional Determinants of a Small Protein Controlling a Broadly

Conserved Bacterial Sensor Kinase. *Journal of Bacteriology* **202** (2020).

51.   A. O. Olaitan, *et al.*, Worldwide emergence of colistin resistance in Klebsiella pneumoniae

from healthy humans and patients in Lao PDR, Thailand, Israel, Nigeria and France

owing to inactivation of the PhoP/PhoQ regulator mgrB: an epidemiological and molecular

study. *International Journal of Antimicrobial Agents* **44**, 500–507 (2014).

52.   J. A. Lees, M. Galardini, S. D. Bentley, J. N. Weiser, J. Corander, pyseer: a

comprehensive tool for microbial pangenome-wide association studies. *Bioinformatics* **34**,

4310–4312 (2018).

53.   S. Hoffman, Z. Lapp, J. Wang, E. S. Snitkin, "regentrans: a framework and R package for

using genomics to study regional pathogen transmission" (2021).

54.   T. J. Kidd, *et al.*, A Klebsiella pneumoniae antibiotic resistance mechanism that subdues

654   host defences and promotes virulence. *EMBO Mol Med* **9**, 430–447 (2017).

655   55.   J. R. Bowers, *et al.*, Genomic Analysis of the Emergence and Rapid Global Dissemination

656   of the Clonal Group 258 Klebsiella pneumoniae Pandemic. *PLoS ONE* **10**, e0133727

657   (2015).

658   56.   E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and

659   evolutionary analyses in R. *Bioinformatics* **35**, 526–528 (2019).

660   57. ,   R: The R Project for Statistical Computing (April 15, 2020).

661   58.   H. Wickham, *et al.*, Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686

662   (2019).

663   59.   G. Yu, D. K. Smith, H. Zhu, Y. Guan, T. T.-Y. Lam, ggtree: an r package for visualization

664   and annotation of phylogenetic trees with their covariates and other associated data.

665   *Methods in Ecology and Evolution* **8**, 28–36 (2017).

666   60.   G. Yu, T. T.-Y. Lam, H. Zhu, Y. Guan, Two Methods for Mapping and Visualizing

667   Associated Data on Phylogeny Using Ggtree. *Mol Biol Evol* **35**, 3041–3043 (2018).

668   61.   R. Kolde, *pheatmap: Pretty Heatmaps* (2019) (April 15, 2020).

669   62.   G. Yu, *ggplotify: Convert Plot to "grob" or "ggplot" Object* (2020) (February 20, 2021).

670   63.   C. O. Wilke, *cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2"* (2019)

671   (April 15, 2020).

Figure 1: Epistatic interactions appear to influence the extent of colistin resistance in clinical CRKP isolates. (A) Most susceptible isolates have no variants in known resistance genes, while most resistant isolates have at least one variant in a known resistance gene. Many susceptible isolates also have two variants in known resistance genes, suggesting potential reversion events. (B) Colistin MIC split by total number of variants in resistance genes. Shapes indicate whether any of the variants are putative resistance variants (see methods for definition

of resistance variants). On average, resistant isolates with two or more variants in resistance

genes have a higher MIC than resistance isolates with one or fewer variants in resistance

genes, and susceptible isolates with two or more variants in resistance genes have a lower MIC

than susceptible isolates with one or fewer variants in resistance genes. Significance defined as

Wilcox p < 0.05. MIC=minimum inhibitory concentration.

Figure 2: Distinct resistance evolution patterns exist in different CRKP ST258

sublineages. Clade IIB exhibits extensive dissemination of resistance variants, as well as

several putative reversion events, while the other clades contain sporadic and less transmitted

resistance variants. Each grey column of the heatmap is a gene, where all variants of a given

type (resistance in dark grey, suppressor in light grey), are grouped for clarity. See Fig S5 for

more details on individual variants. The asterisks indicate the 2 clade IIB resistance clusters

mentioned in the text. Scale bar is in substitutions per site per year. R=resistant; S=susceptible.

Figure 3: Colistin resistance does not impart a fitness cost in clade IIB strains. In clade

IIB, resistant isolates are more fit than corresponding susceptible non-revertants. On the other

hand, resistant isolates in other clades are less fit than susceptible non-revertants in those

clades. Significance defined as Wilcox p < 0.05. R=resistant; S=susceptible; THD=time-scaled

haplotypic density.

Figure 4: Patients with colistin-resistant isolates were treated with colistin. (A) The

majority of clade IIB isolates are respiratory isolates, and many of the patients with those strains

were treated with colistin (empiric and/or definitive). Differences in transparency are aesthetic to

draw attention to clade IIB isolates treated with colistin. (B) Many colistin-resistant isolates were

treated with colistin.

Figure 5: Disruption of the putative O-antigen glycosyltransferase *kfoC* may lead to

decreased transmission fitness cost of colistin resistance. Created with BioRender.com

Supplementary Figures and Table

Figure S1: Colistin resistance occurs across time and geography. R=resistant,

S=susceptible.

Figure S2: The majority of colistin resistance can be explained by variants in known resistance genes. R = resistant; GWAS = genome-wide association study. Known means the mutation or variant was previously shown to confer resistance, known site means the site was shown to confer resistance under a different amino acid change, putative and GWAS means that >60% of the variant was present in resistant isolates in canonical and GWAS genes, respectively.

734 **Figure S3: Schematic of the molecular pathway of canonical colistin resistance genes.**

Figure S4: Putative resistance and suppressor variants in canonical and non-canonical resistance genes. Columns are variants and rows are isolates. All variants except the last panel are from canonical resistance genes. R=resistant, S=susceptible.

742 **Figure S5: Toy example of how resistance groups are defined.** Note that susceptible

743 revertants are defined by the presence of a resistance variant, rather than a putative suppressor

744 variant. See methods for more details. Heatmap columns are each variants found in a

745 resistance gene. R=resistant, S=susceptible.

Figure S6: The two clonally expanded resistance variants occur across multiple LTACHs

in California. 10/11 LTACHs in the Los Angeles area have the variant, as well as one LTACH

in the San Diego area.

761   **Table S1: Resistance variants**

Gene	Variant type	Impact	Site type	Variant	Isolate count
mgrB	SNV	high	known	Trp47*	1
mgrB	SNV	high	known	Lys43*	1
mgrB	SNV	high	known	Gln30*	41
mgrB	SNV	high	known	Gln22*	1
mgrB	SNV	high	known	Lys3*	6
mgrB	Indel	high	known	Ile12fs	6
mgrB	Indel	high	known	Leu8fs	2
mgrB	Insertion	high	known	*NA*	1
mgrB	Insertion	high	known	*NA*	1

mgrB	Insertion	high	known	*NA*	1
mgrB	Insertion	high	known	*NA*	2
mgrB	Insertion	high	known	*NA*	1
pmrB	SNV	moderate	known	Leu82Arg	1
crrB	SNV	moderate	known	Gln10Leu	1
crrB	SNV	moderate	known	Leu94Met	16
crrB	SNV	moderate	known	Pro151Ser	2
crrB	SNV	moderate	known	Pro151Ser	1
crrB	SNV	moderate	known	Gly183Val	1
mgrB	SNV	moderate	known_site	Asp31Asn	1
phoQ	SNV	moderate	known_site	Thr244Asn	85
pmrB	SNV	moderate	known_site	Pro158Gln	1

pmrA	SNV	moderate	known_site	Gly53Ser	1
mgrB	SNV	moderate	putative_r	Ile41Asn	1
mgrB	SNV	moderate	putative_r	Asn25Thr	1
mgrB	SNV	modifier	putative_r		3
mgrB	SNV	modifier	putative_r		3
mgrB	SNV	modifier	putative_r		1
mgrB	SNV	modifier	putative_r		9
phoP	SNV	moderate	putative_r	Gly53Val	1
phoQ	SNV	moderate	putative_r	Leu308Gln	1
phoQ	SNV	moderate	putative_r	Leu257Gln	4
phoQ	SNV	moderate	putative_r	Thr248Pro	1
phoQ	SNV	moderate	putative_r	Leu105Gln	2

phoQ	SNV	moderate	putative_r	Leu62Pro	2
pmrB	SNV	moderate	putative_r	Leu262Pro	1
pmrB	SNV	moderate	putative_r	Ser203Pro	1
crrB	SNV	moderate	putative_r	Ala200Pro	1
crrB	Insertion	high	putative_r	*NA*	1
mgrB	SNV	moderate	putative_re vert	Thr21Pro	1
phoP	SNV	moderate	putative_re vert	Gln202Pro	3
phoP	SNV	moderate	putative_re vert	Thr189Pro	1
phoP	SNV	moderate	putative_re vert	His188Gln	1
phoP	SNV	moderate	putative_re vert	His188Arg	1

phoP	SNV	moderate	putative_revert	Arg185Leu	3
phoP	SNV	moderate	putative_revert	Arg111His	11
phoP	SNV	moderate	putative_revert	Val18Ala	4
phoP	Indel	high	putative_revert	Thr189fs	1
phoP	Indel	high	putative_revert	His188fs	1
phoQ	SNV	moderate	putative_revert	Gln481Leu	1
phoQ	SNV	moderate	putative_revert	Ala451Thr	1
phoQ	SNV	moderate	putative_revert	Gly443Ser	1

phoQ	SNV	moderate	putative_re vert	Arg433Leu	13
phoQ	SNV	moderate	putative_re vert	Asp417Val	2
phoQ	SNV	moderate	putative_re vert	Leu322Met	1
phoQ	SNV	moderate	putative_re vert	Glu123Lys	2
phoQ	SNV	moderate	putative_re vert	Ala36Thr	1
phoQ	Indel	high	putative_re vert	Leu344fs	1
phoQ	Indel	moderate	putative_re vert	Thr248_Leu 254del	1
crrB	Indel	high	putative_re vert	Ser338fs	16

mgrB	SNV	moderate	other	Phe44Cys	1
mgrB	SNV	moderate	other	Leu24Pro	3
mgrB	SNV	moderate	other	Leu19Gln	1
phoP	SNV	moderate	other	Ala80Gly	3
phoQ	SNV	moderate	other	Ile462Ser	1
phoQ	SNV	moderate	other	Leu365His	8
phoQ	SNV	moderate	other	Pro343Leu	3
phoQ	SNV	moderate	other	Thr332Pro	9
phoQ	SNV	moderate	other	Arg50Leu	7
phoQ	SNV	moderate	other	Gly33Ser	3
phoQ	Insertion	high	other	*NA*	1
pmrB	SNV	moderate	other	Pro95Gln	3

pmrB	Insertion	high	other	*NA*	1
PTS sugar transporter	SNV	moderate	gwas_r	Ser72Ile	1
PTS sugar transporter	Insertion	high	gwas_r	*NA*	1
qseC	SNV	moderate	gwas_r	Val180Ala	9
qseC	Indel	moderate	gwas_r	Arg12_Leu13 del	1