Communication

VAERS data reveals no increased risk of neuroautoimmune adverse events from COVID-19 vaccines

Chris von Csefalvay[1]

1   Starschema Inc., Arlington, VA; csefalvayk@starschema.net

Abstract: Neuroautoimmune disorders, such as multiple sclerosis and Guillain-Barre syndrome, have been documented in relation to various vaccines in the past. This paper uses passive reporting information from the CDC/FDA's VAERS system to analyse whether neuroautoimmune presentations are reported at a relatively higher or lower rate, vis-a-vis other adverse effects, for COVID-19 vaccines than for other vaccines. Through computing the reporting odds ratios for a range of symptoms and comparator vaccines, a clear indication in favour of the safety of COVID-19 vaccines emerges, with reports of neuroautoimmune adverse events in relation to other adverse events being over 70% less likely for COVID-19 than for comparator vaccines (ROR : 0.292, $p < 0.0001$). In comparison with other vaccines given as part of routine care in adulthood, COVID-19 vaccines have the lowest reporting odds ratio of neuroautoimmune adverse effects (median ROR: 0.246).

Keywords: Neuroautoimmune disorders, demyelinating disorders, vaccine safety, pharmacovigilance, COVID-19 vaccines

1. Introduction

The global drive to curb the spread of COVID-19 through a concerted vaccination effort is undoubtedly one of the most ambitious public health projects ever undertaken. At the time of writing, only a little over six months after the FDA granted EUAs to the first two of eventually three SARS-CoV-2 vaccines in early December, over 43% of the U.S. population has been fully vaccinated (i.e. two doses of a two-dose vaccine or a single dose of a one-dose vaccine). For comparison, during the severe 2017-18 flu season, only 37.1% of adults were vaccinated.[1]

To safeguard this unprecedented accomplishment, it is crucial to exercise close surveillance of potential adverse events (AEFIs). At least partially owing to the fact that the Moderna and Pfizer/BioNTech vaccines are first-in-man applications of mRNA vaccine technology at scale, a significant number of eligible individuals have refused to be vaccinated.[2–4] The fear of adverse effects, both short and long term, is undoubtedly a driving factor that fuels these apprehensions and eventually results in an overall reluctance to be vaccinated. Among mid- to long-term AEFIs, neurological AEFIs hold a particular place, as is evident from studies on reasons behind vaccine refusal.[5] The bitter past experience with Guillain-Barre syndrome (GBS) as a side effect of certain influenza vaccines, esp. the much publicised association between 1976-77 A/New Jersey/8/76 influenza vaccines and GBS, is still within living memory.[6] Similarly well-known is the association between the recombinant hepatitis B vaccine and multiple sclerosis.[7] In addition, the emergence of various theories about a purported (and disproven) causal mechanism between autism spectrum disorders and vaccinations that hypothesises an autoimmune pathogenetic process has created a persistent point of rhetoric that erroneously portrays vaccines as inevitably likely to cause neuroautoimmunity.[8] Given the often devastating economic, social, psychological and QoL impact of neuroautoimmune diseases,[9–11] some degree of apprehension appears natural.

⁴⁰ It is beyond the scope of this paper to outline the range of potential pathophysio-
⁴¹ logical mechanisms that may underlie autoimmunity, including neuroautoimmunity,
⁴² following a vaccination (for such a review, see Wraith et al. (2003)[12]). Rather, we seek
⁴³ to identify evidence regarding the safety of COVID-19 vaccines approved in the United
⁴⁴ States (Pfizer/BioNTech, Moderna and Johnson & Johnson) with regard to neuroautoim-
⁴⁵ mune AEFIs through an analysis of VAERS.

⁴⁶ As all studies leveraging VAERS, its limitations must be read in conjunction with
⁴⁷ results derived from VAERS data. Like all passive pharmacovigilance systems, it re-
⁴⁸ lies on reporting. Under the terms of the EUAs granted to each of the three vaccine
⁴⁹ manufacturers, COVID-19 vaccination providers must report any

⁵⁰ • vaccine administration errors (regardless of consequence);
⁵¹ • serious AEFI (regardless of causal attribution), incl. any AEFI resulting in inpatient
⁵² admission, long-term disability or death; and
⁵³ • instances of multi-system inflammatory syndrome,

⁵⁴ along with serious immunisation failures (cases of COVID-19 in vaccinated individuals
⁵⁵ that result in inpatient admission or death). This goes beyond to the typical reporting
⁵⁶ regime, wherein reporting is encouraged but not mandatory unless the patient was a
⁵⁷ minor at the time. At the same time, VAERS remains open to reports from individuals,
⁵⁸ allied healthcare workers and even unrelated third parties.[13–15] In other words, while
⁵⁹ there is a risk of unreported AEFIs, there may also be a degree of over-reporting (the same
⁶⁰ AEFI, for instance, may be reported by the physician, the nurse and the patient as well,
⁶¹ without the knowledge of any of the other parties). Moreover, VAERS does not verify the
⁶² accuracy or veracity of reports, nor does it require a causal attribution. This is notably
⁶³ illustrated by the number of accidents, drownings and congenital diagnoses reported to
⁶⁴ VAERS, none of which could conceivably be a causal consequence of vaccination. Thus,
⁶⁵ VAERS data must be appropriately analysed to fulfill its function, which is to generate
⁶⁶ early potential safety signals rather than to substantiate a causal relationship.

⁶⁷ In this paper, we are using a reporting odds ratio (ROR) analysis[16] to compare
⁶⁸ the likelihood that an AEFI reported to VAERS is one of a number of neuroautoim-
⁶⁹ mune AEFIs between COVID-19 and non-COVID-19 vaccines, concluding that no such
⁷⁰ association appears to emerge from the evidence as it presently stands, and that the
⁷¹ odds of reporting a neuroautoimmune AEFI are significantly lower than for comparable
⁷² vaccines.

2. Materials and Methods

⁷⁴ Data was obtained for 2015 to 2021, inclusive, from the VAERS website (https:
⁷⁵ //vaers.hhs.gov) on 12 June 2021. The data comprises all reports that were received
⁷⁶ between 01 January 2015 and 28 May 2021, comprising altogether 2,528,763 individual
⁷⁷ reports. Of these, 1,323,178 (52.33%) pertained to a COVID-19 vaccine.

2.1. Categorisation of cases

⁷⁹ Following ingestion using Python v.3.7.5 and `pandas` v.1.2.4,[17] the resulting data
⁸⁰ frames were joined and reshaped to yield individual entries per reported symptom.
⁸¹ Using the symptom description, reports of the following VAERS symptoms (i.e. symp-
⁸² toms using the coding phraseology as present in VAERS) were coded as involving a
⁸³ neuroautoimmune AEFI:

⁸⁴ • Demyelinating polyneuropathy
⁸⁵ • Immune-mediated neuropathy
⁸⁶ • Axonal neuropathy
⁸⁷ • Axonal and demyelinating neuropathy
⁸⁸ • Chronic inflammatory demyelinating polyradiculoneuropathy
⁸⁹ • Subacute inflammatory demyelinating polyradiculoneuropathy
⁹⁰ • Autoimmune neuropathy

• Autonomic neuropathy
• Guillain-Barre syndrome
• Acute disseminated encephalomyelitis
• Demyelinating polyneuropathy
• Neuromyelitis optica spectrum disorder
• Neuromyelitis optica
• Myelitis transverse
• Multiple sclerosis relapse
• Multiple sclerosis
• Relapsing multiple sclerosis
• Progressive multiple sclerosis
• Relapsing-remitting multiple sclerosis
• Optic neuritis
• Immune-mediated encephalitis
• Autoimmune encephalopathy
• Encephalitis autoimmune
• Autoimmune demyelinating disease

Based on this encoding, the individual data was segmented and age distributions
were calculated (Figure 1). Subjects receiving the COVID-19 vaccine were on average
older ($\mu =50.2$, $\sigma = 17.8$) than those who received other vaccines ($\mu =36.9$, $\sigma = 28.3$), a
phenomenon attributable to the relatively large number of childhood vaccines in relation
to vaccines received in later life. Except for male recipients of the COVID-19 vaccine, age
distribution of neuroautoimmune disorders followed a bimodal pattern, peaking in the
third and eight decades of life. Male recipients of the COVID-19 vaccine who reported a
neuroautoimmune AEFI tended towards a single mode in the early seventh decade of
life.

Figure 1. Distribution of reports by age, partitioned by vaccine type (COVID-19 vs. non-COVID-19 vaccine) and neuroautoimmune disorder status (neuroautoimmune vs. non-neuroautoimmune symptoms).

117 Data was cross-tabulated to yield a 2x2 contingency table for COVID-19 vs. non-
118 COVID-19 vaccines (Table 1), and the odds ratio was calculated using Fisher's exact test.

119 In addition, a Pearson's χ^2 test was performed and the Pearson residuals were calculated
120 for each permutation, using `statsmodels` v.0.12.2.[18]

Table 1. 2x2 contingency table by vaccine type (COVID-19 vs. non-COVID-19 vaccine) and neuroautoimmune disorder status.

	Neuroautoimmune AEFI	Non-neuroautoimmune AEFI
COVID-19 vaccine	612	1908
Non-COVID-19 vaccine	1,322,566	1,203,677

Next, symptoms were broken down individually, and a χ^2 distribution fitted. Along
with the actual values, the expected values under the null hypothesis of independence
(H_0) were ascertained, as well as the Pearson residuals.
Finally, a large contingency table was constructed, calculating the reporting odds
ratio of each AEFI for each vaccine category (VAERS variable `VAX_TYPE`), which acts as
a superset comprising vaccines that have typically the same target and same valency
(e.g. the Johnson&Johnson COVID-19 vaccine and the Moderna vaccine are both in
the `COVID19` vaccine category, whereas a trivalent and a quadrivalent influenza vaccine
would be different vaccine categories due to their difference in valency). For the purposes
of constructing this table, a regular expression was used to filter out a number of
entities that are typically recorded in VAERS but which do not indicate an appropriate
denominator for the calculation of the odds ratio. These were used to exclude

• normal results (e.g. `.*(negative|normal)$`),
• mere tests and assays that do not disclose a result (e.g. `.*assay$`),
• procedures (e.g. `.*(plasty|insertion|tomy|ery...)`),
• management of an ongoing medical status or device,
• administrative flags (e.g. `Blood (group|don(or|ation))`), and
• COVID-19 related public health interventions (e.g. `COVID-19`
`(prophylaxis|immunisation|screening)`).

RORs were cross-tabulated by vaccine type and filtered only for symptoms that
were categorised as neuroautoimmmune symptomatic presentations.

3. Results

3.1. Association with neuroautoimmune AEFIs

Fisher's two-sided exact test was used to calculate the odds ratio to determine the
odds ratio for an association between reporting a neuroautoimmune AEFI and reporting
after/in relationship with a COVID-19 vaccine. This yielded an OR of 0.292 at $p < 0.0001$
- a statistically highly significant result that indicates COVID-19 vaccination predisposes
to a significantly lower risk – by over 70% – of reporting a neuroautoimmune side effect
rather than any other side effect. It is important to note that this does not mean that
neuroautoimmune side effects are 70% less frequent, but rather that the likelihood of
reporting a neuroautoimmune side effect over any other side effect is 70% lower.

Table 2. Pearson residuals by vaccine type (COVID-19 vs. non-COVID-19 vaccine) and neuroautoimmune disorder status.

	Neuroautoimmune AEFI	Non-neuroautoimmune AEFI
COVID-19 vaccine	-19.459	20.386
Non-COVID-19 vaccine	0.615	-0.644

152 This is confirmed by the Pearson χ^2 residuals that were calculated (see Table 2). The
153 residual of neuroautoimmune AEFIs following a COVID-19 vaccine (-19.459) indicates
154 that the actual value is significantly higher than the value expected value under the

assumption of the null hypothesis of independence. It is therefore quite evident that the odds of reporting a neuroautoimmune AEFI over any other AEFI are correlated with whether the preceding vaccine was a COVID-19 vaccine or not, with recipients of COVID-19 vaccines being significantly (70%) less likely to report a neuroautoimmune AEFI.

3.2. Association of individual neuroautoimmune AEFIs

Figure 2. Expected vs. actual number of reports for individual neuroautoimmune symptoms based on Pearson's χ^2 test along with Pearson residuals for COVID-19 vaccines.

Following on from these findings, it is not surprising to see that in most cases, the number of actual cases reported is significantly lower than would be expected under the null hypothesis of independence (see Figure 2). The notable exceptions are

- progressive MS (Pearson residual: 0.243),
- immune-mediated encephalitis (Pearson residual: 0.243),
- autoimmune demyelinating disease (Pearson residual: 0.864),
- autoimmune encephalopathy (Pearson residual: 0.627), and
- MS relapse (Pearson residual: 0.664).

While the residuals indicate that the deviation is quite small for all of the above, it appears that in contrast with other vaccines, COVID-19 vaccines appear to result in a relatively higher number than expected of these AEFIs. This must be considered in the context of actual figures, of course – altogether, these differences account for fewer than 0.25 additional cases of progressive MS and immune-mediated encephalitis, fewer than one case of autoimmune encephalopathy, fewer than 4 cases of MS relapse and fewer than 1.5 cases of autoimmune demyelinating disease. Of over 1.3m reports, these constitute an excess of six cases, or approx. one in 220,530 reports. Consequently, it remains to be seen (especially given the low absolute numbers) whether this association remains in evidence as more data accrues, or, as is more likely, it is part of the noisy undercurrent that is characteristic of passive reporting.

180 *3.3. Reporting odds ratios between vaccines*

Figure 3. Reporting Odds Ratios of neuroautoimmune conditions in a range of vaccines approved for adult populations.

As Figure 3 shows, the ROR of neuroautoimmune presentations is significantly
lower for COVID-19 vaccines than it is for any of the comparator vaccines. The group of
comparators was chosen to primarily include vaccines that have the same age range of
approved users to reduce the bias arising from the fact that some vaccines are typically
given in a paediatric context (e.g. MMR), while COVID-19 vaccination was consciously
designed to address older populations first due to their higher overall risk from COVID-
19 as well as the relatively higher rate of respiratory, cardiac and endocrine comorbidities
that governed vaccine allocation in the beginning.

4. Discussion

Based on early data from VAERS for reports about COVID-19 vaccinations received
on or before 28 May 2021, there is convincing evidence that COVID-19 vaccines compare
remarkably well with other vaccines in respect of neuroautoimmune AEFIs. Despite
the limitations of this study, which drew on passive reporting data with all its inherent
problems, such as reporting bias and the lack of a reliable denominator, we have found
evidence that supports the safety of COVID-19 vaccines with regard to neuroimmuno-
logical presentations. Statistical analysis indicated that in comparison to recipients of
other vaccines, those who reported in respect of a COVID-19 vaccines were over 70%
less likely to report a neuroimmunological presentation than a non-neuroimmunological
presentation.

Similarly, the number of reports fell significantly under what was expected in
all but two somemwhat cohesive symptom groups: relapses of MS on one hand and
autoimmmune encephalitis on the other. The fact that some forms of autoimmune
encephalitis are distributed bimodally by age with a peak in the third and the seventh
decade of life may support a hypothesis that the increase in cases of autoimmune
encephalitis is the consequence of statistical noise and/or sampling bias due to the
higher average age of COVID-19 vaccine recipients in the sample.[19] Similarly, multiple
sclerosis exhibits a similar bimodality of age distribution, to the point that respective
terms for early-onset (teens to early 30s) and late onset (50s and later) MS have become
established in the literature.[20] Nevertheless, clinicians may, as a matter of abundance
of caution, keep a closer eye on patients with known MS or risk factors for MS (such
as previous diagnosis of optic neuritis or CIS/RIS) for signs of progression or relapse.

This may be accompanied byy proactively explaining the relative risk of a relapse in the
context of potentially worse clinical outcomes with COVID-19, especially for patients
treated with systemic immunosupressants.
Since the start of the global vaccination campaign, evidence has been accumulating
for the overall tolerability and safety of the COVID-19 vaccines. This study seeks to
add to our current understanding by quantifying the highly favourable risk profile of
COVID-19 vaccines with regard to neuroautoimmune disorders. While some questions
remain open and conclusions made over early data may have to be reconfirmed through
ongoing analysis, the current data supports the assertion that the COVID-19 vaccine has
perhaps the lowest reporting odds ratio for neuroautoimmune disorders of any vaccine
routinely given in adulthood. This is a reassuring indication of safety at a time when
concerns about adverse events still keep most countries from attaining the threshold of
collective immunity.

Funding: This research was funded by Starschema Inc. under its intramural research funding
programme.

Data Availability Statement: VAERS reporting data is available from the CDC's website at
https://vaers.hhs.gov. All code and scripts supporting this manuscript are deposited at https:
//github.com/chrisvoncsefalvay/covid19-neuroautoimmune-aefis and are made available with
the DOI <to follow>.

Conflicts of Interest: CvC is a consultant to a company that may be affected by the research
reported in this paper. The funders had no role in the design of the study; in the collection,
analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish
the results.

Abbreviations

236 The following abbreviations are used in this manuscript:
237

AEFI	Adverse event following immmunization
CDC	Centers for Disease Control and Prevention
CIS	clinically isolated syndrome
EUA	Emergency Use Authorization
FDA	Food and Drug Administration
GBS	Guillain-Barre syndrome
mRNA	messenger RNA
MS	multiple sclerosis
RIS	radiologically isolated syndrome
ROR	Reporting Odds Ratio
VAERS	Vaccine Adverse Effect Reporting System

References

1.   CDC. Estimates of influenza vaccination coverage among adults—United States, 2017–18 flu
season. *Retrieved October* **2018**, *25*, 2018.
2.   Dror, A.A.; Eisenbach, N.; Taiber, S.; Morozov, N.G.; Mizrachi, M.; Zigron, A.; Srouji, S.; Sela,
E. Vaccine hesitancy: the next challenge in the fight against COVID-19. *European Journal of
Epidemiology* **2020**, *35*, 775–779.
3.   Robertson, E.; Reeve, K.S.; Niedzwiedz, C.L.; Moore, J.; Blake, M.; Green, M.; Katikireddi, S.V.;
Benzeval, M.J. Predictors of COVID-19 vaccine hesitancy in the UK household longitudinal
study. *Brain, Behavior, and Immunity* **2021**, *94*, 41–50.
4.   Troiano, G.; Nardi, A. Vaccine hesitancy in the era of COVID-19. *Public Health* **2021**.
5.   Berry, S.D.; Johnson, K.S.; Myles, L.; Herndon, L.; Montoya, A.; Fashaw, S.; Gifford, D.
Lessons learned from frontline skilled nursing facility staff regarding COVID-19 vaccine
hesitancy. *Journal of the American Geriatrics Society* **2021**.
6.   Haber, P.; DeStefano, F.; Angulo, F.J.; Iskander, J.; Shadomy, S.V.; Weintraub, E.; Chen, R.T.
Guillain-Barré syndrome following influenza vaccination. *JAMA* **2004**, *292*, 2478–2481.

7. Hernán, M.A.; Jick, S.S.; Olek, M.J.; Jick, H. Recombinant hepatitis B vaccine and the risk of multiple sclerosis: a prospective study. *Neurology* **2004**, *63*, 838–842.

8. Poland, G.A.; Jacobson, R.M. Understanding those who do not understand: a brief review of the anti-vaccine movement. *Vaccine* **2001**, *19*, 2440–2445.

9. Marck, C.H.; Aitken, Z.; Simpson, S.; Weiland, T.J.; Kavanagh, A.; Jelinek, G.A. Predictors of change in employment status and associations with quality of life: a prospective international study of people with multiple sclerosis. *Journal of Occupational Rehabilitation* **2020**, *30*, 105–114.

10. Miller, A.; Dishon, S. Health-related quality of life in multiple sclerosis: the impact of disability, gender and employment status. *Quality of Life Research* **2006**, *15*, 259–271.

11. Patwardhan, M.; Matchar, D.; Samsa, G.; McCrory, D.; Williams, R.; Li, T. Cost of multiple sclerosis by level of disability: a review of literature. *Multiple Sclerosis Journal* **2005**, *11*, 232–239.

12. Wraith, D.C.; Goldman, M.; Lambert, P.H. Vaccination and autoimmune disease: what is the evidence? *The Lancet* **2003**, *362*, 1659–1666.

13. Chen, R.T.; Rastogi, S.C.; Mullen, J.R.; Hayes, S.W.; Cochi, S.L.; Donlon, J.A.; Wassilak, S.G. The Vaccine Adverse Event Reporting System (VAERS). *Vaccine* **1994**, *12*, 542–550.

14. Shimabukuro, T.T.; Nguyen, M.; Martin, D.; DeStefano, F. Safety monitoring in the vaccine adverse event reporting system (VAERS). *Vaccine* **2015**, *33*, 4398–4405.

15. Singleton, J.A.; Lloyd, J.C.; Mootrey, G.T.; Salive, M.E.; Chen, R.T.; Ellenberg, S.; Rastogi, S.; Krueger, C.; Braun, M.; Wise, R.; others. An overview of the vaccine adverse event reporting system (VAERS) as a surveillance system. *Vaccine* **1999**, *17*, 2908–2917.

16. Rothman, K.J.; Lanes, S.; Sacks, S.T. The reporting odds ratio and its advantages over the proportional reporting ratio. *Pharmacoepidemiology and Drug Safety* **2004**, *13*, 519–523.

17. McKinney, W.; others. pandas: a foundational Python library for data analysis and statistics. *Python for High Performance and Scientific Computing* **2011**, *14*, 1–9.

18. Seabold, S.; Perktold, J. Statsmodels: Econometric and statistical modeling with python. Proceedings of the 9th Python in Science Conference. Austin, TX, 2010, Vol. 57, p. 61.

19. Shan, W.; Yang, H.; Wang, Q. Neuronal Surface Antibody-Mediated Autoimmune Encephalitis (Limbic Encephalitis) in China: A Multiple-Center, Retrospective Study. *Frontiers in Immunology* **2021**, *12*, 344.

20. Kis, B.; Rumberg, B.; Berlit, P. Clinical characteristics of patients with late-onset multiple sclerosis. *Journal of Neurology* **2008**, *255*, 697–702.