Factors affecting the transmission of SARS-CoV-2 in school settings

Connor Reynolds, MPH[1*]; Sydney Ng, MPH[2*]; Wan Yang, PhD[1#]

[1]Department of Epidemiology, [2]Department of Biostatistics, Mailman School of Public Health, Columbia University;

[*]Authors contributed equally;

[#]Correspondence to: wy2202@columbia.edu

Abstract

Background

Several studies have reported SARS-CoV-2 outbreaks in schools, with a wide range of secondary attack rate (SAR; range: 0-100%). We aimed to examine the key risk factors to better understand transmission in school settings.

Methods

We collected records of SARS-CoV-2 school outbreaks globally published from January 2020 to January 2021 and compiled information on hypothesized risk factors. We utilized the directed acyclic graph (DAG) to conceptualize the risk mechanisms, used logistic regression to examine each risk-factor group, and further built multiple variable models based on the marginal analysis. Adjusted odds ratios (aOR) and 95% confidence intervals (CI) were calculated.

Results

From 17 relevant articles, 26 school clusters were included for analysis. The best-fit model showed that the intensity of community transmission (aOR: 1.26; 95% CI: 1.22 - 1.30, for each increase of 10 cases per 100,000 persons per week), social distancing (aOR: 0.26; 95% CI: 0.18 - 0.37), mask-wearing (aOR: 0.52; 95% CI: 0.35 - 0.78) were associated the risk of SARS-CoV-2 infection in schools. Compared to students in pre-schools, the aOR was 0.12 (95% CI: 0.07 - 0.19) for students in primary schools and 1.31 (95% CI: 0.93 - 1.87) for students in high schools.

Conclusions

Preventive measures in both schools (e.g. social distancing and mask-wearing) and communities (additionally, vaccination) should be taken to collectively reduce transmission and protect children in schools. Flexible reopening policies may be considered for different levels of schools given their risk differences.

Keywords

COVID-19, transmission, children, school, secondary attack rate

Key messages

- We collected published records of COVID-19 school outbreaks globally to investigate the considerable heterogeneity in secondary attack rates (SAR) reported from school outbreaks and compiled information regarding potential risk factors.
- Higher community death and case rates are associated with higher SARs in children in school settings.
- Mask-wearing and social distancing are associated with lower SARs.
- When compared to pre-schools and early childhood education centers, primary schools have lower rates of transmission of SARS-CoV-2 however high schools have higher rates.

Introduction

Since the early stages of the COVID-19 pandemic, concerns have been raised about the impact of schools on community transmission and the well-being of students and staff, as well as the impact on the schedules of healthcare workers concerning childcare [1]. Out of an abundance of caution and fear that the SARS-CoV-2 virus would spread rapidly in schools much like influenza pandemics [2], countries globally decided to suspend in-person classes and begin online instruction. By April 2020, over 1.5 million students worldwide were affected by school closures in response to the COVID-19 pandemic [3]. In contrast to influenza pandemics where children are the key drivers of transmission, studies have indicated that children are likely less susceptible to SARS-CoV-2 infection, tend to experience less severe disease when infected, and likely have lower transmissibility [4, 5]. Given this new evidence, schools in many places have gradually reopened since the summer of 2020, while implementing varying level of preventive measures (e.g., mask wearing, distancing, limiting the number of students, rotating schedules, and viral testing) to reduce risk of transmission. Given these circumstances, the risk of SARS-CoV-2 outbreaks in school settings may differ substantially across space and time. Indeed, several studies have examined school outbreaks of COVID-19 and reported secondary attack rates (SAR) – i.e., the proportion of infected contacts of an index case out of all contacts of that index case [6] – ranging from 0 (i.e., no secondary infections) to 100% (i.e., infections among all contacts). However, this discrepancy is still not fully understood and a better understanding can inform better preventive measures for future outbreaks not limited to COVID-19 or school settings.

To identify the main factors that determine the transmission of SARS-CoV-2 in schools and inform strategies to prevent future school outbreaks, here we examined the associations between SARS-CoV-2 SAR in children and various potential risk factors. We compiled data from relevant studies in the literature reporting SARS-CoV-2 SAR in schools and for posited risk factors (e.g. incidence in the community) and further used regression models to examine key

risk factors of having high SAR in schools. We found that appropriate distancing and mask

wearing, school levels, and the magnitude of SARS-CoV-2 transmission in the community are

key determinants of SAR in school settings.

Methods

Data sources

Studies were searched for on the "Living Evidence for COVID-19" database [24], which retrieves

articles from EMBASE via Ovid, PubMed, BioRxiv, and MedRxiv. Any article within this database

was considered, from December 2019 up to February 17, 2021. The search terms used include

"transmission AND (school OR schools)" or "transmission AND children." When titles and

abstracts were identified as being potentially relevant, the articles were read to determine if

the outbreaks took place in a school setting and the number of infections and contacts were

reported. In addition, we extracted 11 observations included in a systematic review of evidence

regarding the ability of children to transmit SARS-CoV-2 in schools [7]. In total, 17 studies were

included in this analysis (see Appendix 1).

Relevant data, as deemed by an initial conceptual analysis using the directed acyclic graph

(DAG; see details below), were taken from the articles identified above. These included the

time period of the study, study design, location, season, age of children, whether the study

happened during a lockdown, type of school according to the International Standard

Classification of Education [8], reported SARs, number of contacts of the index case, testing

method (PCR vs. serology), level of surveillance (all contacts, some contacts, only symptomatic),

and whether masks and social distancing were required. In addition, we compiled additional

data for potential risk or confounding factors of SARS- CoV-2 transmission in schools for each

identified study as detailed in the next section.

Conceptual analysis and variable coding

The unit of analysis was individual study and in cases where several school types were covered

in one study the data were stratified by those school types. We first conducted a conceptual

analysis using the DAG and identified eight key components that may affect SARS-CoV-2

transmission in schools (Fig 1). Below we describe each of the eight components, rationale for

inclusion, and related variables examined.

1) School types, based on studies indicating differential transmission risk among different age

groups [20, 22]. Here, we examined this factor as a categorical variable including 4 levels, i.e.

pre-school or early childhood education center (ECEC), primary school, high school, and mixed-

level school. The first three levels were per reports in the included school studies. For studies

that examined several types of school but did not report school type specific SARs, we assigned

them to a "mixed-level school" category. For example, if a study gave the overall SAR combining
a pre-school and a primary school it was given the value "mixed-level school."
SARS-CoV-2 SAR among children in school settings is the number of infected contacts divided by
the total number of contacts at each school.
118
119 2) Physical school settings such as student density in the classroom and ventilation systems that
120 may affect the intensity of school contact and clearance of air. As it is difficult to obtain
121 information related to ventilation settings, here we included class size in our analysis based on
122 the average number of students per classroom in each country, as reported by the Organisation
123 for Economic Co-operation and Development (OECD) [12].
124
3) Cultural climates, which "represent independent preferences for one state of affairs over
another that distinguish countries (rather than individuals) from each other" [9] and may reflect
the collective risk tendency of a population. The Hofstede's cultural dimensions theory [9]
included 6 related measures including individualism, masculinity, uncertainty avoidance, long
term orientation, and indulgence. In particular, individualism is defined as the degree of
interdependence of society maintains among its members. We reasoned that the individualism
measure would be most relevant to the level of compliancy to public health interventions and
in turn the risk of SARS-CoV-2 transmission. Thus, here we included individualism in our analysis
and dichotomized the reported values for each country into high (≥ 50) or low.
134
135 4) Preventive measures, which may reduce outbreak risk. Here we included both mask-wearing
136 and social distancing. For these variables, we dichotomized based on as the use of face masks
137 (or social distancing) by either students, staff, or both. For instance, for mask-wearing, if either
138 group (students or staff) were required to wear face masks onsite, the school would receive a
139 value of 1 and 0 otherwise.
140
5) Surveillance and/or testing policies implemented in schools. On the one hand, testing policies
could affect the reported values of SAR; for instance, testing of all contacts regardless of
symptoms may lead to identification of more infections including those asymptomatic and
increase the numerator of SAR. On the other, frequent testing of all if combined with school
closure may serve as a containment measure to reduce the risk of onward transmission and in
turn reduce SAR. Here we thus included the reported testing practices for contacts in the
school outbreak clusters as a categorical ordinal variable. Three types of testing were reported
in the school studies, including testing only the symptomatic, both symptomatic and some
asymptomatic, and all contacts. However, due to the small sample size in "only symptomatic"
(n = 3), we dichotomized surveillance to testing "only symptomatic or some asymptomatic" and
"all contacts" of an index case in each school cluster.

6) Community transmission intensity, which may increase the introduction of infections into

school settings. To examine its impact, we included two measures, i.e., the weekly case rate per

100,000 for the study area for the study period and weekly death rate per 100,000 from COVID-

19; both measures were computed using data from the John Hopkins Coronavirus Resource

Center [10] and standardized by overall country population. See "Sensitivity analysis" below for

justification of these choices.

7) Seasonal changes such as humidity and temperature, which may affect the survival and

transmission of SARS-CoV-2. Here, we used specific humidity (a measure of absolute humidity)

to examine the potential impact from disease seasonality, as specific humidity and temperature

are highly correlated. Specifically, ground surface temperature and relative humidity for each

study location were extracted from the National Oceanic and Atmospheric Administration using

the "stationaRy" package [11]. Daily mean specific humidity in g H_2O/kg air was then computed

based on the temperature and relative humidity data and was averaged over the study period.

8) Indicators of socioeconomic status such as national income that reflect a country's ability to

mobilize resources to fight COVID-19. As such, we included measured national income for each

study in our analysis; specifically, national income is measured as the gross domestic product

(GDP) subtracting capital depreciation and adding net foreign income, using data from the

World Inequality Database [13].

Statistical Analyses

Marginal analysis.

Due to the low number of observations (n = 26 schools), we conducted an initial analysis to test

combinations of the DAG covariates described above. The goal was to examine the relationship

between SAR and only one group of variables at a time and then include the most relevant

predictors into the final model based on this analysis. For each test, we used a logistic

regression model of the following form:

$$logit(SAR) \sim X$$

where logit is the log-odds (i.e., $log\left(\frac{p}{1-p}\right)$, with p as the probability of event) and SAR

represents the SAR as reported from each of the 26 schools. X is one of the combinations of

variables we examined as follows:

1) Preventive measures (mask-wearing and social distancing) adjusting for seasonal changes

2) Community transmission (weekly death rate per 100,000 and weekly case rate per 100,000)

3) Seasonal changes

4) Classroom size, adjusting for national income

5) School type

190
Multi-risk factor analysis
All five variable groups were found to be associated with SAR in the marginal analysis (see
Results). We thus tested models including different combinations of these variables. For all
models, we included surveillance type to account for potential biases in reporting including
missing asymptomatic infections, which would underestimate SAR. We also assessed for
confounding between our variables of interest and SARS-CoV-2 SAR (see adjustments specified
above). This procedure began with including all of the significant variables from the marginal
analysis and removing one at a time to see the effect of its removal on model fit. We then
evaluated and selected the most parsimonious model with the best fit based on the Akaike
information criterion (AIC; Table S1). The best performing model took the following form:
201
202 $logit(SAR) \sim$ mask-wearing + social distancing + weekly case rate + school type + surveillance
203
204 All statistical analyses were performed in RStudio, a user interface for R (R Foundation for
205 Statistical Computing, Vienna, Austria). All models were fitted using the "glm" function from the
206 built-in "stats" library in R.
207
Sensitivity analysis
We tested different measures of community transmission, to examine the robustness of our
models to potential biases due to variations in case-ascertainment, mortality risk, and delay in
event occurrence (e.g. from infection to death) and reporting. Specifically, we additionally
examined two mortality-related measures in representing the intensity of community
transmission: 1) weekly death rate for the study period and 2) weekly death rate for the study
period plus one month. The additional month added accounts for the time lag that occurs as
death from COVID-19 may take several weeks [14]. Risk factors identified using these two
measures were similar to those in our main analysis (Table S2). As the risk of death due to
COVID-19 reduced substantially over time [15], likely due to improved medical treatment and
management, such a change may bias the death rates recorded over time for the included
studies spanning the initial phase in March 2020 to a later phase in Dec 2020. For this reason,
we used weekly case rate per 100,000 to represent the intensity of community transmission in
our main analysis.
222
Results
Summary statistics
We identified 26 reported SARS-CoV-2 outbreaks in schools, totaling 630 secondary cases in
children among 8322 contacts. These outbreaks occurred in 12 countries, spanning 4 WHO
regions including the Americas, Western Pacific, European, and Eastern Mediterranean Region.

Table 1 shows the frequencies and summary statistics for SARS-CoV-2 SAR and other variables
included. SARS-CoV-2 SAR and weekly cases per 100,000 population were both right skewed.
While the reported SAR ranged from 0 to 100%, the majority of schools reported very low SAR
(median: 0.7%, interquartile range: 0 – 17%). The majority of school settings included in these
studies required mask-wearing (19/26, or 73%) and distancing (14/26, or 54%). Around half of
the studies tested all contacts of the index case (12/26, or 46%). The studies covered roughly
even proportion for different school types: 5 (19%) were pre-schools, 9 (35%) were primary
schools, 7 (27%) were high schools, and 5 (19%) were mixed schools.

Marginal analysis
The marginal analysis (Figure 2), adjusting for surveillance, indicated that, for the intensity of
community transmission, each 10-unit increase in weekly case rate was associated with an
increased risk of contracting SARS-CoV-2 in schools [odds ratio (OR): 1.11; 95% confidence
interval (CI): 1.08 - 1.14]; and each 10-unit increase in weekly death rate was associated with an
increased risk of contracting SARS-CoV-2 in schools (OR: 1.47; 95% CI: 1.40 - 1.54). For
preventive measures, social distancing (OR: 0.21, 95% CI: 0.16 - 0.29) was associated with a
lower SAR in schools. For school type, compared to pre-schools, being in primary schools (OR:
0.69, 95% CI: 0.49 - 0.98) or mixed schools (OR: 0.69; 95% CI: 0.49 - 0.98) was associated with a
lower risk of SARS-CoV-2 infection whereas being in high schools (OR: 1.96; 95% CI: 1.44 - 2.69)
was associated with a higher risk of SARS-CoV-2 infection. For the school physical setting
measure, each 1-person increase in the national average class size was associated with an
increased risk of contracting SARS-CoV-2 in schools (OR: 1.26; 95% CI: 1.21 - 1.30). For disease
seasonality, each 1 g H_2O/kg increase in specific humidity was associated with a decreased risk
of contracting SARS-CoV-2 in schools (OR: 0.88, 95% CI: 0.85, 0.92). Estimates were similar
when the model did not adjust for surveillance (Figure 2).

Multi-risk factor analysis
Among all models tested (Table S1), the best-performing model with the lowest AIC included
three key groups of risk factors, namely, intensity of community transmission, preventive
measures, and school type, adjusting for surveillance. Overall, these risk factors in combination
were able to explain 57.7% of the variance in the reported SARS-CoV-2 SAR (McFadden's
pseudo R^2 = 0.577; Fig 3) [21, 23]. Fig 4 and Table 2 show the estimated adjusted ORs (aORs)
for each risk factor.

The multi-risk factor model showed that every 10-unit increase in COVID-19 cases rate in the
community was associated with an increased risk of SARS-CoV-2 infection in schools (aOR: 1.26;
95% CI: 1.22 - 1.30). Conversely, both social distancing (aOR: 0.26; 95% CI: 0.18 - 0.37) and

mask-wearing (aOR: 0.52; 95% CI: 0.35 - 0.78) were associated a decreased risk of SARS-CoV-2

infection in schools. Compared to students in pre-schools, the aOR was 0.12 (95% CI: 0.07, 0.19)

for students in primary schools, 1.31 (95% CI: 0.93, 1.87) for students in high schools, and 0.26

(95% CI: 0.18, 0.39) for students in mixed schools.

Discussion

Leveraging available data on multiple reported SARS-CoV-2 outbreaks in schools and potential

risk factors, we have examined main factors associated with the risk of SARS-CoV-2

transmission in schools. Our analyses suggest that the most significant predictors of SARS-CoV-

2 SAR in schools are the level of community transmission (measured as weekly COVID-19 case

rate) and preventative measures, once adjusted for surveillance and school type.

The model identified community transmission as the most significant variable associated with

the risk of SARS-CoV-2 transmission in schools. In particular, the multi-risk factor model

estimated that, in a community of 100,000 persons, every 10 additional cases per week in the

community increases the odds of infection in schools by 26%. This finding highlights the

potential community-to-school importation of cases and subsequent risk of outbreak in schools.

As such, care must be applied when reopening or operating schools in areas with high levels of

community transmission. In addition, reversing some original fears about school-to-community

SARS-CoV-2 transmission, it is likely that the community transmission drives outbreaks in

school, not the reverse. Further, with the availability of COVID-19 vaccines, predominantly to

adults at present, it is paramount that all eligible adults get vaccinated promptly to lower the

risk of transmission in the community and in turn to provide indirect protection to children via

the increased population immunity.

Our model suggests that distancing policies have the greatest impact on reducing transmission

in schools (aOR: 0.26, 95% CI: 0.18 - 0.37). This finding is likely a combined outcome of reduced

number of contacts and reduced short-range transmission when social distancing policies were

followed. Maintaining distance has necessitated fewer people in a room at the same time,

leading to fewer contacts. In addition, the increased personal space in classroom enables

students to avoid the likely higher viral concentration within short-range of the emitter (either

via aerosols, droplets, or in combination) when far apart.

Nevertheless, it is important to note that social distancing measures may be more difficult to

achieve fully in disadvantaged communities with underfunded and overcrowded schools,

further exacerbating the COVID-19 crises in these communities, usually of color [16, 17].

Furthermore, racially motivated structural factors prevent these disadvantaged communities

from practicing social distancing policies outside of the school. For example, these communities

tend to make up most of essential workers and thus have higher rates of transmission in their
community [19], increasing potential introduction of infections into schools.
305
Another preventive measure, mask-wearing, was also found to be significantly protective
against SARS-CoV-2 infection in schools. While the association was not as substantial as for
social distancing, a reduction by half (aOR: 0.52; 95% CI: 0.35 - 0.78) is still considerable. In
addition to mitigating the risk of infection, this protective association may also come from
masks reducing the emission of viral particles of SARS-CoV-2 infected wearers [18]. This latter
means of protection is particularly important in settings like schools, as infections in children
are more likely to be asymptomatic or mild and thus more difficult to identify [23].
Nevertheless, concerns have been raised regarding the potential developmental and health
harms of masks on very young children. Given this trade-off, more effective use of other
preventive measures such as vaccination of all those eligible especially care takers, improved
ventilation, and physical distancing may serve to reduce the need for long-term use of masks
among very young children for their protection.
318
In comparison to pre-schools, students in both primary and mixed-level schools had a lower risk
of SARS-CoV-2 infection whereas those in high schools had a higher risk. This finding is
consistent with previous studies indicating the likely lower susceptibility to SARS-CoV-2
infection and transmissibility among young children [23]. In addition, it is also likely in part due
to the greater ability of older children to follow directions regarding preventive measures but
with less compliance among high school students. The similarity of the estimates between
primary school and mixed-level schools is likely due to the number of primary schools that were
folded into the mixed-level school category – three of the five schools that made up this group
included primary schools. Overall, these findings support flexible reopening policies for
different levels of schools given the risk differences.
329
330 When schools test all contacts of the index case, this effectively functions as a control measure
331 in that more cases will be detected and is more likely to result in a school closure. This may
332 explain why the OR estimate for all contacts surveillance is protective, as the outbreak was
333 identified and able to be contained before the asymptomatic individuals were able to infect
334 others.
335
This study has several limitations. First, the analyses included a limited number of studies (n =
26), due to the inclusion criteria and the dearth of published reports on this topic to date.
Future research with additional relevant studies may enhance the statistical power of inference
for the risk factors examined here. Second, due to a lack of detailed information for each
specific school setting, we used proxy measures in the analyses (e.g., class size at the national

level was used rather than for each reporting school), which may have limited the ability of the

models to identify the association of these factors with SARS-CoV-2 transmission risk. Similarly,

due to the lack of data, we were not able to examine other key factors such as ventilation in

classrooms, social economic status of individual students and their households, and potential

differences in susceptibility and transmissibility by age group. Future work with comprehensive

study designs and data collection is warranted to provide further insights into how infections,

not limited to SARS-CoV-2, spread in schools and the broad, bi-directional impact of school and

community transmission. This would be invariable to inform better strategies to combat future

infectious disease outbreaks.

Funding

WY was supported by the National Institute of Allergy and Infectious Diseases (AI145883).

Author contributions

CR and SN contributed to study design, conducted the literature review and data extraction,

and performed the statistical analysis; WY conceived and supervised the study; CR, SN, and WY

wrote the manuscript.

Conflict of interest

None.

References

1.   Bayham J, Fenichel EP. The impact of school closure for COVID-19 on the US healthcare

workforce and the net mortality effects [Internet]. medRxiv; 2020. p.

2020.03.09.20033415. Available from: https://doi.org/10.1101/2020.03.09.20033415

2.   Worby CJ, Chaves SS, Wallinga J, Lipsitch M, Finelli L, Goldstein E, et al. On the relative

role of different age groups in influenza epidemics. Epidemics [Internet]. 2015

Dec;13:10–6. Available from: /pmc/articles/PMC4469206/

/pmc/articles/PMC4469206/?report=abstract

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4469206/

3.   UNESCO I of S. Education: From disruption to recovery [Internet]. 2021. Available from:

https://en.unesco.org/covid19/educationresponse

4.   Viner RM, Mytton OT, Bonell C, Melendez-Torres GJ, Ward J, Hudson L, et al.

Susceptibility to SARS-CoV-2 Infection among Children and Adolescents Compared with

Adults: A Systematic Review and Meta-Analysis [Internet]. Vol. 175. American Medical

Association; 2021. p. 143–56. Available from: https://jamanetwork.com/

5. Otte im Kampe E, Lehfeld A-S, Buda S, Buchholz U, Haas W. Surveillance of COVID-19
school outbreaks, Germany, March to August 2020. Eurosurveillance [Internet]. 2020
Sep;25(38):2001645. Available from:
https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2020.25.38.2001645
6. CDC D of SE&PD. Principles of epidemiology in public health practice, lesson 3: Measures
of risk [Internet]. 2012. Available from:
https://www.cdc.gov/csels/dsepd/ss1978/lesson3/section2.html
7. Xu W, Li X, Dozier M, He Y, Kirolos A, Lang Z, et al. What is the evidence for transmission
of COVID-19 by children in schools? A living systematic review [Internet]. Cold Spring
Harbor Laboratory Press; 2020 Oct p. 2020.10.11.20210658. Available from:
https://www.medrxiv.org/content/10.1101/2020.10.11.20210658v1
https://www.medrxiv.org/content/10.1101/2020.10.11.20210658v1.abstract
8. UNESCO I of S. International standard classification of education (ISCED) [Internet].
2011. Available from: http://uis.unesco.org/en/topic/international-standard-
classification-education-isced
9. Hofstede I. Hofstede insights: Country comparison [Internet]. 2020. Available from:
https://www.hofstede-insights.com/country-comparison/the-usa/
10. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real
time. Lancet Infectious Dis. 2020 Jan;
11. Iannone R. stationaRy: Detailed meteorological data from stations all over the world
[Internet]. 2020. Available from: https://CRAN.R-project.org/package=stationaRy
12. Organisation for Economic Co-operation and Development. Education indicators in
focus [Internet]. 2012. Available from: http://www.oecd.org/education/skills-beyond-
school/EDIF%202012--N9%20FINAL.pdf
13. Alvaredo F, Chancel L, Piketty T, Saez E, Zucman G. World inequality database [Internet].
2018. Available from: wid.world
14. Chen Y, Li T, Ye Y, Chen Y, Pan J. Impact of fundamental diseases on patients with
COVID-19. Disaster Medicine and Public Health Preparedness. 2020;
15. James N, Menzies M, Radchenko P. COVID-19 second wave mortality in europe and the
united states. Chaos: An Interdisciplinary Journal of Nonlinear Science. 2021;
16. Baker BD, Srikanth A, Cotto Jr R, Green III PC. School funding disparities and the plight of
latinx children. education policy analysis archives. 2020;28(135):n135.
17. Anyane-Yeboa A, Sato T, Sakuraba A. Racial disparities in COVID-19 deaths reveal harsh
truths about structural inequality in america. Journal of internal medicine.
2020;288(4):479–80.
18. Asadi S, Cappa CD, Barreda S, Wexler AS, Bouvier NM, Ristenpart WD. Efficacy of masks
and face coverings in controlling outward aerosol particle emission from expiratory
activities. Scientific reports. 2020;10(1):1–3.

19.   Dorn A van, Cooney RE, Sabin ML. COVID-19 exacerbating inequalities in the US. The
Lancet. 2020 Apr;395(10232):1243–4.
20.   Davies NG, Klepac P, Liu Y, Prem K, Jit M, Eggo RM. Age-dependent effects in the
transmission and control of COVID-19 epidemics. Nature medicine. 2020
Aug;26(8):1205-11.
21.   Mittlböck M, Schemper M. Explained variation for logistic regression. Statistics in
medicine. 1996 Oct 15;15(19):1987-97.
22.   Galanti M, Birger R, Ud-Dean M, Filip I, Morita H, Comito D, Anthony S, Freyer GA,
Ibrahim S, Lane B, Matienzo N, et al.. Rates of asymptomatic respiratory virus infection
across age groups. Epidemiology & Infection. 2019;147.
23.   McFadden, D., *Conditional logit analysis of qualitative choice behavior*, in *Frontiers in
econometrics*, P. Zarembka, Editor. 1973, Academic Press: New York. p. 105 - 142.
24.   COVID-19 Open Access Project. (2021). *Living Evidence on COVID-19*. Retrieved
from https://ispmbern.github.io/covid-19/living-review/
429

Table and Figure Captions

Table 1. Characteristics of collected data from literature review.

Table 2. Results of the best-fit multi-risk factor model for the identification of factors associated with SARS-CoV-2 SAR in schools. Adjusted odds ratio (aOR) estimates and 95% confidence intervals are given from the logistic regression model including surveillance to control for differences in testing school clusters, school type due to inconsistent reporting of age groups in the literature, mask-wearing and social distancing for preventative measures, and weekly case rates per 100,000.

Figure 1. Directed acyclic graph (DAG) describing the relationship among variables. This DAG represents the meaningful relationships between the variables relevant to SARS-CoV-2 SAR among children in school settings and informs all further analyses. The outcome measurement, SAR, is presented in red while risk factors are in black and surveillance in green.

Figure 2. Odds ratio estimates from the marginal models. Left panel shows results from models without adjusting for surveillance and right panel shows results from corresponding models additionally adjusting for surveillance. Black points show the mean odds ratio estimates and horizontal black bars show the 95% confidence intervals. The vertical black bar in each plot indicate the null value of 1.0. Each set of models is delineated by alternating the shaded regions.

Figure 3. Model fit of the best-performing multi-risk model. Black dots show the fitted SAR for each study (y-axis), compared to the observed SAR (x-axis); the red bars around each point show the 95% confidence intervals of model-estimates. The McFadden's pseudo-R^2 is computed using Eqn 30 in [23].

Figure 4. Odds ratio estimates from the best-performing multi-risk model. Black points show the mean odds ratio estimates and horizontal black bars show the 95% confidence intervals. The vertical black bar indicates the null value of 1.0. Each variable type is delineated by the shaded regions.

Supplemental Tables

Table S1. Performance of different models. In the marginal analysis, 5 models for 5 groups of risk factors are tested. In the multi-risk factor analysis, 11 models with all possible combinations of 4 groups of risk factors are tested (i.e., excluding models with only one group tested in the marginal analysis). All models adjusted for surveillance. Note that although class size was identified as a significant risk factor in the marginal analysis, further including this variable in

the multi-risk models gave impossible values for parameter estimates; we therefore excluded

class size in the multi-risk model analysis. The best-performing model with the lowest AIC is

bolded.

Table S2. Sensitivity analysis for both cases and deaths, with and without a 1-month extension

to the study time period.

Tables

Table 1. Characteristics of collected data from literature review

Variable	Summary
Surveillance	n (%)
All contacts	12 (46.2%)
Some asymptomatics	11 (42.3%)
Only symptomatics	3 (11.5%)
School type	n (%)
Mixed	5 (19.2%)
Secondary	7 (26.9%)
Primary	9 (34.6%)
Pre-school/ECEC	5 (19.2%)
Mask-wearing	n (%)
Yes	7 (26.9%)
No	19 (73.1%)
Social distancing	n (%)
Yes	12 (46.1%)
No	14 (53.9%)
High individualism (≥ 50)[a]	n (%)
	22 (88%)
Secondary attack rate	median (IQR)[b]
	0.007 (0 - 0.17)
Weekly case rate (per 100,000)	median (IQR)
	7.42 (0.29, 26.4)
Daily mean specific humidity (g H2O/kg air)	median (IQR)
	8.00 (4.95, 9.83)
Average class size	median (IQR)
	21.55 (20.00, 23.80)
National income (in thousands)	median (IQR)
	38,942 (37,248, 50,752)

[a] Individualism was accounted for in 25 studies, had no significant effect and was not included for final study.

[b] IQR, interquartile range

Table 2. Results of the best-fit multi-risk factor model for the identification of factors associated
with SARS-CoV-2 SAR in schools. Adjusted odds ratio (aOR) estimates and 95% confidence
intervals are given from the logistic regression model including surveillance to control for
differences in testing school clusters, school type due to inconsistent reporting of age groups in
the literature, mask-wearing and social distancing for preventative measures, and weekly case
rates per 100,000.

Variable	aOR (95% CI)
Surveillance	
All contacts	0.542 (0.454, 0.646)
Only symptomatics or some asymptomatics	(reference)
School Type	
Mixed	0.261 (0.176, 0.388)
Secondary	1.310 (0.925, 1.860)
Primary	0.115 (0.068, 0.196)
Pre-school/ECEC	(reference)
Mask wearing	
Yes	0.523 (0.351, 0.779)
No	(reference)
Social distancing	
Yes	0.260 (0.184, 0.368)
No	(reference)
Weekly case rate (per 100,000)	
	1.260 (1.220, 1.300)

485
486

487   Figures

488   **Figure 1.** Directed acyclic graph (DAG) describing the relationship among variables. This DAG
489   represents the meaningful relationships between the variables relevant to SARS-CoV-2 SAR
490   among children in school settings and informs all further analyses. The outcome measurement,
491   SAR, is presented in red while risk factors are in black and surveillance in green.

492
493

Figure 2. Odds ratio estimates from the marginal models. Left panel shows results from models
without adjusting for surveillance and right panel shows results from corresponding models
additionally adjusting for surveillance. Black points show the mean odds ratio estimates and
horizontal black bars show the 95% confidence intervals. The vertical black bar in each plot
indicate the null value of 1.0. Each set of models is delineated by alternating the shaded
regions.

500
501

Figure 3. Model fit of the best-performing multi-risk model. Black dots show the fitted SAR for

each study (y-axis), compared to the observed SAR (x-axis); the red bars around each point

show the 95% confidence intervals of model-estimates. The McFadden's pseudo-R^2 is

computed using Eqn 30 in [23].

Figure 4. Odds ratio estimates from the best-performing multi-risk model. Black points show
the mean odds ratio estimates and horizontal black bars show the 95% confidence intervals.
The vertical black bar indicates the null value of 1.0. Each variable type is delineated by the
shaded regions.

514   **for**

515   **Factors affecting the transmission of SARS-CoV-2 in school settings**

516   Connor Reynolds, MPH; Sydney Ng, MPH; Wan Yang, PhD

518 This document includes: 1) Appendix 1 listing all SARS-CoV-2 school outbreak studies included

519 in our analysis; and 2) all supplemental tables.

521 **Appendix 1: Studies Included**

522   1.   J. P. Armann, M. Unrath, C. Kirsten, et al. "Anti-SARS-CoV-2 IgG antibodies in adolescent

523   students and their teachers in Saxony, Germany (SchoolCoviDD19): very low

524   seropraevalence and transmission rates." 2020.

525   2.   A. Fontanet, R. Grant, L. Tondeur, et al. "SARS-CoV-2 infection in primary schools in

526   northern France: A retrospective cohort study in an area of high transmission." In:

527   *medRxiv* (Jun. 2020), p. 2020.06.25.20140178. DOI: 10.1101/2020.06.25.20140178.

528   <URL: https://doi.org/10.1101/2020.06.25.20140178>.

529   3.   C. Atherstone, M. Siegel, E. Schmitt-Matzen, et al. "SARS-CoV-2 Transmission Associated

530   with High School Wrestling Tournaments Florida, December 2020-January 2021." In:

531   *Morbidity and Mortality Weekly Report* 70.4 (2021), p. 141.

532   4.   L. T. Brandal, T. S. Ofitserova, H. Meijerink, et al. "Minimal transmission of SARS-CoV-2

533   from paediatric COVID-19 cases in primary schools, Norway, August to November 2020."

534   In: *Eurosurveillance* 26.1 (2021), p. 2002011.

535   5.   D. Buonsenso and B. Graglia. "High rates of SARS-CoV-2 transmission in a high-school

536   class." In: *Journal of Paediatrics and Child Health* (2021).

537   6.   K. Danis, O. Epaulard, T. Benet, et al. "Cluster of coronavirus disease 2019 (COVID-19) in

538   the French Alps, February 2020." In: *Clinical Infectious Diseases* 71.15 (2020), pp. 825-

539   832.

540   7.   S. Desmet, E. Ekinci, I. Wouters, et al. "No SARS-CoV-2 carriage observed in children

541   attending daycare centers during the first weeks of the epidemic in Belgium." In:

542   *medRxiv* (May. 2020), p. 2020.05.13.20095190. DOI: 10.1101/2020.05.13.20095190.

543   <URL: https://doi.org/10.1101/2020.05.13.20095190>.

544   8.   T. Dub, E. Erra, L. Hagberg, et al. "Transmission of SARS-CoV-2 following exposure in

545   school settings: experience from two Helsinki area exposure incidents." In: *medRxiv* (Jul.

546   2020), p. 2020.07.20.20156018. DOI: 10.1101/2020.07.20.20156018. <URL:

547   https://doi.org/10.1101/2020.07.20.20156018>.

548   9.   A. Fontanet, L. Tondeur, Y. Madec, et al. "Cluster of COVID-19 in Northern France: A

549   Retrospective Closed Cohort Study." In: *SSRN Electronic Journal* (Apr. 2020), p.

550   2020.04.18.20071134. ISSN: 1556-5068. DOI: 10.1101/2020.04.18.20071134. <URL:
551   https://doi.org/10.1101/2020.04.18.20071134>.

10. L. Heavey, G. Casey, C. Kelly, et al. "No evidence of secondary transmission of COVID-19
552   from children attending school in Ireland, 2020." In: *Eurosurveillance* 25.21 (May. 2020),
553   p. 1. ISSN: 15607917. DOI: 2807/1560-7917.ES.2020.25.21.2000903. <URL:
554   /pmc/articles/PMC7268273/?report=abstract
555   https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7268273/>.
556

11. National Centre for Immunisation Research and Surveillance. *COVID-19 in schools the
557   experience in NSW*. 2020. <URL:https://www.ncirs.org.au/sites/default/files/2020-
558   04/NCIRS%20NSW%20Schools%20COVID_Summary_FINAL%20public_26%20April%202
559   020.pdf}.>
560

12. E. Larosa, O. Djuric, M. Cassinadri, et al. "Secondary transmission of COVID-19 in
561   preschool and school settings after their reopening in northern Italy: a population-based
562   study." In: *medRxiv* (2020).
563

13. A. S. Lopez, M. Hill, J. Antezano, et al. "Transmission Dynamics of COVID-19 Outbreaks
564   Associated with Child Care Facilities" Salt Lake City, Utah, April"July 2020." In: *MMWR.
565   Morbidity and Mortality Weekly Report* 69.37 (Sep. 2020), pp. 1319-1323. ISSN: 0149-
566   2195. DOI: 10.15585/mmwr.mm6937e3. <URL:
567   http://www.cdc.gov/mmwr/volumes/69/wr/mm6937e3.htm?s{_}cid=mm6937e3{_}w
568   >.
569

14. K. Macartney, H. E. Quinn, A. J. Pillsbury, et al. "Transmission of SARS-CoV-2 in
570   Australian educational settings: a prospective cohort study." In: *The Lancet Child and
571   Adolescent Health* 4.11 (Nov. 2020), pp. 807-816. ISSN: 23524642. DOI: 10.1016/S2352-
572   4642(20)30251-0.
573

15. C. Stein-Zamir, N. Abramson, H. Shoob, et al. "A large COVID-19 outbreak in a high
574   school 10 days after schools' reopening, Israel, May 2020." In: *Eurosurveillance* 25.29
575   (Jul. 2020), p. 2001352. ISSN: 1560-7917. DOI: 10.2807/1560-
576   7917.ES.2020.25.29.2001352. <URL:
577   https://www.eurosurveillance.org/content/10.2807/1560-
578   7917.ES.2020.25.29.2001352>.
579

16. J. P. Torres, C. Piñera, V. De La Maza, et al. "SARS-CoV-2 antibody prevalence in blood in
580   a large school community subject to a Covid-19 outbreak: a cross-sectional study." In:
581   *Clinical Infectious Diseases* (Jul. 2020). ISSN: 1058-4838. DOI: 10.1093/cid/ciaa955.
582   <URL:https://academic.oup.com/cid/advance-
583   article/doi/10.1093/cid/ciaa955/5869860>.
584

17. C. F. Yung, K. Kam, K. D. Nadua, et al. "Novel coronavirus 2019 transmission risk in
585   educational settings." In: *Clinical Infectious Diseases* 72.6 (2020), p. 1055-1058. <URL:
586   https://academic.oup.com/cid/article/72/6/1055/5862649>
587

Supplemental Tables

Table S1. Performance of different models. In the marginal analysis, 5 models for 5 groups of
risk factors are tested. In the multi-risk factor analysis, 11 models with all possible combinations
of 4 groups of risk factors are tested (i.e., excluding models with only one group tested in the
marginal analysis). All models adjusted for surveillance. Note that although class size was
identified as a significant risk factor in the marginal analysis, further including this variable in
the multi-risk models gave impossible values for parameter estimates; we therefore excluded
class size in the multi-risk model analysis. The best-performing model with the lowest AIC is
bolded.

Analysis	Model	AIC
Marginal analysis	Mask wearing, distancing, seasonal changes, and surveillance	1106.60
	Weekly cases, weekly deaths, surveillance	890.34
	Specific humidity and surveillance	1278.58
	Class size, national income, and surveillance	1028.87
	School type and surveillance	1177.23
Multi-risk factor analysis (all adjusted for surveillance)	Mask wearing, distancing, and specific humidity	1106.59
	School type, and specific humidity	1035.03
	Mask wearing, distancing, and weekly cases	950.51
	Mask wearing, distancing, and school type	795.92
	Specific humidity and weekly cases	1106.99
	School type and weekly cases	893.05
	Mask wearing, distancing, specific humidity, and school type	794.60
	Mask wearing, distancing, specific humidity, and weekly cases	951.39
	Specific humidity, school type, and weekly cases	794.85
	Mask wearing, distancing, weekly cases, school type, and specific humidity	598.01
	Mask wearing, distancing, weekly cases, and school type	**596.33**

598 **Table S2.** Sensitivity analysis for both cases and deaths, with and without a 1-month extension
599 to the study time period.

		With extension		Without extension
Cases	*Variable*	*aOR (95% CI)*	*Variable*	*aOR (95% CI)*
	mask wearing	0.559 (0.365-0.858)	mask wearing	0.523 (0.351-0.779)
	distancing	0.126 (0.0869-0.183)	distancing	0.260 (0.184-0.368)
	weekly cases	2.12 (1.88-2.40)	weekly cases	9.83 (7.21-13.4)
	surveillance	0.499 (0.417-0.597)	surveillance	0.542 (0.454-0.646)
	primary		primary	0.115 (0.0678-
	school	0.264 (0.171-0.406)	school	0.196)
	high school	1.46 (1.03-2.09)	high school	1.31 (0.925-1.86)
	mixed school	0.193 (0.130-0.288)	mixed school	0.261 (0.176-0.388)
	AIC	642.04	AIC	596.33
Deaths	*Variable*	*aOR (95% CI)*	*Variable*	*aOR (95% CI)*
	mask wearing	0.74 (0.50-1.11)	mask wearing	0.96 (0.64-1.44)
	distancing	0.28 (0.2-0.404)	distancing	0.23 (0.16-0.319)
			weekly	
	weekly deaths	16.7 (10.2-27.3)	deaths	21.5 (11.1-41.7)
	surveillance	1.02 (0.82-1.28)	surveillance	1.01 (0.79-1.30)
	primary		primary	
	school	0.27 (0.18-0.42)	school	0.29 (0.19-0.439)
	high school	1.15 (0.82-1.62)	high school	0.73 (0.5-1.06)
	mixed school	0.28 (0.19-0.409)	mixed school	0.26 (0.18-0.384)
	AIC	663.71	AIC	714.65

600
601
602