Estimating Vaccination Effects and Variant Strains on COVID-19 outbreak course in Japan, as of June, 2021

Junko Kurita[1], Tamie Sugawara[2], Yasushi Ohkusa[2]

1) Department of Nursing, Tokiwa University, Ibaraki, Japan

2) National Institute of Infectious Diseases, Tokyo, Japan

Corresponding author: Junko Kurita, kuritaj@tokiwa.ac.jp

ICMJE Statement

Contributors JK was responsible for the coordination of the study and responsible for the data setting. YO developed the model and TS illustrated the results. All authors contributed to the writing of the final manuscript.

Keywords: COVID-19, effective reproduction number, variant strain, vaccine coverage

Abstract

Background: Japan started vaccinations for COVID-19 in late February, 2021, mainly

using BNT162b2 mRNA COVID-19 vaccine. At that time, N501Y strain was emerging.

Later, L452R strain appeared in Japan.

Object: We evaluated effects of COVID-19 vaccination and variant strains on

infectiousness in Japan.

Method: The effective reproduction number $R(t)$ was regressed on vaccine coverage and

the shares of variant strains along with data of temperature, humidity, mobility, and

countermeasures. We presumed two definitions for vaccine coverage: at least a single

dose with a 12-day delay and a second dose. The study period was February, 2020

through June 10, as of July 1, 2021.

Results: Estimation results indicate that vaccine coverage for both first and second

doses was not significant. Moreover, infectiousness of the mutated strain N501Y

decreased significantly, but that of L452R was not significant.

Discussion and Conclusion: Vaccination effects and effects of mutated strain with

higher infectiousness were not confirmed. Close and continuous monitoring is

necessary.

Introduction

Wide coverage of COVID-19 vaccination has changed the outbreak situation in European countries and the US. Unfortunately, vaccination started only in February, 2021 in Japan, among the latest of starting dates in economically developed countries. As of June 10, 2021, the rate of completion for second dose was only about 3% in Japan (Figure 1) [1,2].

Fortunately, prevalence of COVID-19 has been much lower in Japan than in European countries and the US. For instance, prevalence in the US was approximately 10% as of June 6, 2021, but only 0.6% in Japan, which has used no lock down. Although reasons for such differences remain unknown, vaccine efficacy in Japan might be different from efficacy in US and European countries.

Vaccination provides personal protection, and eventually herd immunity to unvaccinated persons. The former was investigated in clinical trials [3], but the latter might depend on the surrounding society. Therefore, the present study examined vaccine effectiveness, including herd immunity in Japan. To measure the vaccine effectiveness, we estimated the effects of vaccine coverage on SARS-CoV-2 infectiousness in Japan.

At the time of starting vaccination in Japan, the mutant strain N501Y (alha strain) had been emerging and expanding to dominate infection. Subsequently, the new mutant

strain L452R, delta strain, also appeared in May. Its infectiousness and pathogenicity were estimated as 35–90% higher than those of the original strain, mainly in the UK, which was circulating before the emerging variant strain [4–7]. Therefore, we consider the prevalence of these mutated strains together when evaluating vaccine effects.

As countermeasures against the COVID-19 outbreak in Japan, school closure and voluntary event cancellation were adopted from February 27, 2020 through the end of March. Large commercial events were cancelled. Subsequently, a state of emergency was declared for April 7 through 25 May, stipulating voluntary restrictions against leaving home. Consumer businesses such as retail shops and restaurants were shut down. During this period, the first peak of infection was reached on April 3. Infections subsequently decreased until July 29. The so-called "Go To Travel Campaign" (GTTC) started on July 22 as a 50% subsidized travel program aimed at supporting sightseeing businesses with government-issued coupons for use in shopping at tourist destinations. It was expected that the campaign might expand the outbreak. Thereafter, GTTC continued to the end of December, by which time a third wave of infection had emerged. The third wave in December, which was larger than either of the preceding two waves, reached its highest peak at the end of December. Therefore, GTTC was inferred as the main reason underlying the third wave [8].

To force the third wave lower, a second state of emergency was declared: from

January 8, 2021 through March 15, 2021. However, the fourth wave emerged at the end

of February, probably because of the spread of variant strains. Moreover, to support

hosting of the Olympics and Paralympics games in Tokyo in July, a third emergency

state was declared on April 25, 2021.

Although results have been mixed, some results of earlier studies suggest that

COVID-19 might be associated with climate conditions [9–12]. If that were true for

Japan, then GTTC might not be the main reason for the third wave. Moreover, mobility

was inferred as the main cause of the outbreak dynamics in the first wave in Japan [13]

and throughout the world [14–17]. This study was conducted to estimate vaccine

coverage effects on SARS-CoV-2 infectiousness for the outbreak in Japan.

Methods

This study examined numbers of symptomatic patients reported by the Ministry of

Health, Labour and Welfare (MHLW) for January 14 – June 10, published [18] as of

July 1, 2021. Some patients were excluded from data for Japan: those presumed to be

persons infected abroad or infected as Diamond Princess passengers. Those patients

were presumed not to represent community-acquired infection in Japan. For some

symptomatic patients with unknown onset dates, we estimated onset dates from an empirical distribution with duration extending from onset to the report date among patients for whom the onset date had been reported.

Onset dates among patients who did not report this information and a reporting delay were adjusted using the same procedures as those used for our earlier research [19,20]. As described hereinafter, we estimated the onset dates of patients for whom onset dates had not been reported. Letting $f(k)$ represent this empirical distribution of incubation period and letting N_t denote the number of patients for whom onset dates were not published and available at date t, then the number of patients for whom the onset date was known is t-1. The number of patients with onset date t-1 for whom onset dates were not available was estimated as $f(1)N_t$. Similarly, patients with onset date t-2 and for whom onset dates were not available were estimated as $f(2)N_t$. Therefore, the total number of patients for whom the onset date was not available, given an onset date of s, was estimated as $\Sigma_{k=1}f(k)N_s+k$ for the long duration extending from s.

Moreover, the reporting delay for published data from MHLW might be considerable. In other words, if $s+k$ is larger than that in the current period t, then $s+k$ represents the future for period t. For that reason, N_{s+k} is not observable. Such a reporting delay engenders underestimation of the number of patients. For that reason, it

must be adjusted as $\Sigma_{k=1}^{t-s}f(k)N_s+k\ /\Sigma_{k=1}^{t-s}f(k)$. Similarly, patients for whom the onset

dates were available are expected to be affected by the reporting delay. Therefore, we

have $M_s|_t\ /\Sigma_{k=1}^{t-s}f(k)$, where $M_s|_t$ represents the reported number of patients for whom

onset dates were period s as of the current period t.

We defined $R(t)$ as the number of infected patients on day t divided by the number

of patients who were presumed to be infectious. The number of infected patients was

calculated from the epidemic curve by the onset date using an empirical distribution of

the incubation period, which is $\Sigma_{k=1}f(k)E_{t+k}$, where E_t denotes the number of patients for

whom the onset date was period t. The distribution of infectiousness in symptomatic

and asymptomatic cases $g(k)$ was assumed to be 30% on the onset day, 20% on the

following day, and 10% for the subsequent five days [21]. Then the number of

infectiousness patients was $\Sigma_{k=1}g(k)E_{t-k}$. Therefore, R(t) was defined as

$\Sigma_{k=1}f(k)E_{t+k}/\Sigma_{k=1}g(k)E_{t-k}$.

Data indicating the shares of mutated variants among all cases were published by

the Tokyo Metropolitan Government. Unfortunately, detailed information about mutated

strains has not been published for the entirety of Japan. We used two measures for the

mutant strains as shares, respectively, of N52Y and L452R [22].

We use average temperature and relative humidity data for Tokyo during the day as

climate data because national average data are not available. We obtained data from the

Japan Meteorological Agency (https://www.data.jma.go.jp/gmd/risk/obsdl/index.php).

Additionally, we identified several remarkable countermeasures in Japan: two

emergency status declarations, GTTC, and school closure and voluntary event

cancellation (SCVEC). The latter, SCVEC, extended from February 27 through March:

this countermeasure required school closure and cancellation of voluntary events, and

even cancellation of private meetings. Then the first state of emergency was declared

April 7. It ceased at the end of May. It required voluntary restriction against going out,

school closures, and shutdown of businesses. To subsidize travel and shopping at tourist

destinations, GTTC started on July 22. It was halted temporarily at the end of December.

The second state of emergency was declared on January 7, 2021 for the 11 most

affected prefectures. This countermeasure required restaurant closure at 8:00 p.m., with

voluntary restrictions against going out, but it did not require school closure. It

continued until March 21, 2021. The third state of emergency was declared on April 25,

2021 for four prefectures: Tokyo, Osaka, Hyogo, and Kyoto. Later, the application areas

were extended gradually. They never covered the entirety of Japan.

To clarify associations among R(t) and vaccine coverage in addition to the mutant

strains, climate, mobility, and countermeasures, we regressed the daily R(t) on dummy

variables for daily vaccine coverage, weekly shares of N501Y and L452R strains, daily

climate, mobility, and countermeasures using ordinary least squares regression.

Temperatures were measured in degrees Celsius, with humidity, and mobility as

percentages in regression, not as standardized. When some variables were found to be

not significant, we excluded them as explanatory variables and estimated the equation

again.

We examine two definitions for vaccine coverage: rate of receiving at least a single

dose with a 12 day delay and the completion rate of the second dose. Delay in the

former was the response period for increasing antibody [23]. If a vaccine perfectly

protects the recipient from infection, the estimated coefficient of vaccine coverage

would be 0.01 if one assumes an average of $R(t)$ with no vaccination in the study period.

That would mean that increasing vaccine coverage by one percentage point can be

expected to reduce $R(t)$ by one percentage point. If the estimated coefficient of vaccine

coverage were smaller than 0.01, then it might reflect imperfect personal prevention.

Conversely, if the estimated coefficients of vaccine coverage were smaller than 0.01, it

would mean that herd immunity contributed to prevention of infection among

non-recipients. We adopted 5% as the level at which to infer significance.

Results

Figure 1 depicts vaccine coverage for the first dose and second dose as scatter diagrams. It also shows the shares of mutant strains N501Y and L452R as bars. These are increasing almost monotonically during the period. Figure 2 depicts $R(t)$ during the study period.

Figure 3 presents an empirical distribution of the duration of onset to reporting in Japan. The maximum delay was 31 days. Figure 4 presents an empirical distribution of incubation periods among 91 cases for which the exposed date and onset date were published by MHLW in Japan. The mode was six days. The average was 6.6 days. The calculated $R(t)$ is shown in Figure 1.

Table 1 presents estimation results of two definitions of vaccine coverage to $R(t)$. In both definitions of vaccine coverage, climate conditions, the third emergency status, vaccine coverage and the share of L452R were not significant. Therefore, we deleted these variables from the explanatory variables. All remaining variables in the specification were found to be significant. However, the sign of N501Y was negative. One can readily infer that N501Y was less infectious than the original strain.

Mobility was found to be positive and significant. The first two emergency status periods and GTTC were found to be negative and significant. However, SCVEC was

found to be significant and positive. Vaccine coverage was significantly negative, as expected. However, its estimated coefficients were approximately 0.5 for specification (2) and one for specification (4). These were 50–100 times larger than expectations from personal prevention.

Discussion

Estimation results showed that vaccine coverage was not significant in either definition. Moreover, the share of the mutant strain did not increase infectiousness significantly. It was probably attributable to multicollinearity among vaccine coverage and the share of the mutated strain. However, even in the final specification, which is shown in the right-most column in Table 1, H501N was found to be significantly negative, which was inconsistent with results reported from earlier studies. It might be attributable to the time during which N501Y had been spreading.

The third emergency status was not significant. That finding is probably attributable to an overlap of the period of the third emergency and initiation of the vaccine campaign or when the mutated strain had been becoming dominant among cases of infection. Consequently, multicollinearity might occur among these variables. In fact, if vaccine coverage and the share of the mutant strains is dropped from the full

specification, then the estimated coefficient of the third emergency status was -0.97147, with a *p* value was 0.000. That value is comparable to those of the first two emergency status periods.

The present study has some limitations. First, it was conducted only a short time after vaccination was started and the mutant strains were emerging. Accumulation of the data can be expected to affect the estimation results heavily.

Secondly, readers must be reminded when interpreting the obtained results that the obtained results do not indicate causality. This study demonstrated that negative association exists between N501Y and infectiousness. That finding does not necessarily mean that N501Y reduces infectiousness.

Conclusion

The estimation results yielded no evidence that vaccine coverage has reduced the mutation strain infectiousness. That finding is probably attributable to the data being obtained when vaccination was started, the rapid spread of the mutant strain, and the monotonically decreasing phase of $R(t)$ during the same period. Accumulation of more data and longer periods of data accumulation might decrease that misleading effect.

The present study is based on the authors' opinions: it does not reflect any stance or

policy of their professionally affiliated bodies.

Acknowledgments

We acknowledge the great efforts of all staff at public health centers, medical

institutions, and other facilities who are fighting the spread and destruction associated

with COVID-19.

Ethical considerations

All information used for this study was published data on the web were used. There

is therefore no ethical issue related to this study.

Competing Interest

No author has any conflict of interest, financial or otherwise, to declare in relation

to this study.

Reference

1. Prime Minister and his Cabinet. Novel Coronavirus Vaccines.

   https://www.kantei.go.jp/jp/headline/kansensho/vaccine.html (in Japanese)

   [accessed on June 15, 2021]

2. Japan Ministry of Health, Labour and Welfare. Situation of vaccine coverage for

   COVID-19.

   https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/vaccine_sesshujisseki.html   (in

   Japanese) [accessed on June 15, 2021]

3. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, Perez JL,

   Perez Marc G, Moreira ED, Zerbini C, Bailey R, Swanson KA, Roychoudhury S,

   Koury K, Li P, Kalina WV, Cooper D, Frenck RW Jr, Hammitt LL, Tureci O, Nell

   H, Schaefer A, Unal S, Tresnan DB, Mather S, Dormitzer PR, ?ahin U, Jansen KU,

   Gruber WC; C4591001 Clinical Trial Group. Safety and Efficacy of the BNT162b2

   mRNA Covid-19 Vaccine. N Engl J Med. 2020;383:2603-15.

4. Leung K, Shum MHH, Leung GM, Lam TTY, Wu JT. Early transmissibility

   assessment of the N501Y mutant strains of SARS-CoV-2 in the United Kingdom,

   October to November 2020. Euro Surveill 2021;26:2002106.   doi:

   10.2807/1560-7917.ES.2020.26.1.2002106.

5. Graham MS, Sudre CH, May A, Antonelli M, Murray B, Varsavsky T, Klaser K, Canas LS, Molteni E, Modat M, Drew DA, Nguyen LH, Polidori L, Selvachandran S, Hu C, Capdevila J; COVID-19 Genomics UK (COG-UK) Consortium, Hammers A, Chan AT, Wolf J, Spector TD, Steves CJ, Ourselin S. Changes in symptomatology, reinfection, and transmissibility associated with the SARS-CoV-2 variant B.1.1.7: an ecological study. Lancet Public Health 2021;6:e335-e345.

6. Davies NG, Abbott S, Barnard RC, Jarvis CI, Kucharski AJ, Munday JD, Pearson CAB, Russell TW, Tully DC, Washburne AD, Wenseleers T, Gimma A, Waites W, Wong KLM, van Zandvoort K, Silverman JD; CMMID COVID-19 Working Group; COVID-19 Genomics UK (COG-UK) Consortium, Diaz-Ordaz K, Keogh R, Eggo RM, Funk S, Jit M, Atkins KE, Edmunds WJ. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. Science 2021;372(6538):eabg3055. doi: 10.1126/science.abg3055.

7. Zhao S, Lou J, Cao L, Zheng H, Chong MKC, Chen Z, Chan RWY, Zee BCY, Chan PKS, Wang MH. Quantifying the transmission advantage associated with N501Y substitution of SARS-CoV-2 in the UK: an early data-driven analysis. J Travel Med 2021;28:taab011.

8. Japan Times. Japan to suspend Go To Travel program nationwide from Dec. 28 to

Jan. 11

https://www.japantimes.co.jp/news/2020/12/14/national/suga-go-to-travel-coronavirus/ [accessed on June 5,2021]

9. Shi P, Dong Y, Yan H, Zhao C, Li X, Liu W, He M, Tang S, Xi S. Impact of temperature on the dynamics of the COVID-19 outbreak in China. Sci Total Environ. 2020 ;728:138890.

10. Tobias A, Molina T. Is temperature reducing the transmission of COVID-19 ? Environ Res. 2020 ;186:109553.

11. Yao Y, Pan J, Liu Z, Meng X, Wang W, Kan H, Wang W. No association of COVID-19 transmission with temperature or UV radiation in Chinese cities. Eur Respir J. 2020 ;55:2000517.

12. Walrand S. Autumn COVID-19 surge dates in Europe correlated to latitudes, not to temperature-humidity, pointing to vitamin D as contributing factor. Scientific Reports volume 11, Article number: 1981 (2021) Cite this article

https://www.nature.com/articles/s41598-021-81419-w

13. Kurita J, Sugishita Y, Sugawara T, Ohkusa Y. Mobility data can reveal the entire COVID1-19 outbreak course in Japan. JMIR Public Health & Surveillance 2021,7.

https://publichealth.jmir.org/2021/2/e20335

14. Bergman N, Fishman R. Mobility Reduction and Covid-19 Transmission Rates.**doi:**

    https://doi.org/10.1101/2020.05.06.20093039

15. Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, Whittaker C,

    Zhu H, Berah T, Eaton JW, Monod M, Imperial College COVID-19 Response

    Team; Ghani AC, Donnelly CA, Riley S, Vollmer MAC, Ferguson NM, Okell LC,

    Bhatt S. Estimating the effects of non-pharmaceutical interventions on COVID-19

    in Europe. Nature 2020 ;584(7820):257-61. doi: 10.1038/s41586-020-2405-7.

16. Li Y, Campbell H, Kulkarni D, Harpur A, Nundy M, Wang X, Nair H, for theUsher

    Network for COVID-19 Evidence Reviews (UNCOVER) group. The temporal

    association of introducing and lifting non-pharmaceutical interventions with the

    time-varying reproduction number (R) of SARS-CoV-2: a modelling study across

    131 countries. *Lancet Infect Dis* **2021; 21: 193–202**

    DOI:https://doi.org/10.1016/S1473-3099(20)30785-4?

    https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(20)30785-4/fullte

    xt

17. Larrosa JMC. SARS-CoV-2 in Argentina: Lockdown, mobility, and contagion. J

    Med Virol 2020. doi: 10.1002/jmv.26659.

    https://onlinelibrary.wiley.com/doi/10.1002/jmv.26659

18. Japan Ministry of Health, Labour and Welfare. Press Releases.

    https://www.mhlw.go.jp/stf/newpage_10723.html (in Japanese) [accessed on June 6,

    2021].

19. Kurita J, Sugawara T, Ohkusa Y. Estimated effectiveness of school closure and

    voluntary event cancellation as COVID-19 countermeasures in Japan. J Infect

    Chemother 2021, 27:62-4.   doi: 10.1016/j.jiac.2020.08.012.

20. Sugishita Y, Kurita J, Sugawara T, Ohkusa Y. Effects of voluntary event

    cancellation and school closure as countermeasures against COVID-19 outbreak in

    Japan. Plos one 2020. https://doi.org/10.1371/journal.pone.0239455

21. Kimball A, Hatfield KM, Arons M, James A, Taylor J, Spicer K, Bardossy AC,

    Oakley LP, Tanwar S, Chisty Z, Bell JM, Methner M, Harney J, Jacobs JR, Carlson

    CM, McLaughlin HP, Stone N, Clark S, Brostrom-Smith C, Page LC, Kay M,

    Lewis J, Russell D, Hiatt B, Gant J, Duchin JS, Clark TA, Honein MA, Reddy SC,

    Jernigan JA; Public Health ? Seattle & King County; CDC COVID-19

    Investigation Team. Asymptomatic and Presymptomatic SARS-CoV-2 Infections in

    Residents of a Long-Term Care Skilled Nursing Facility - King County,

    Washington, March 2020. Morb Mortal Wkly Rep. 2020 Apr 3;69(13):377-381.

    doi: 10.15585/mmwr.mm6913e1.

22. Tokyo metropolitan Government. Data of COVID-19 monitoring meeting in Tokyo metropolitan.

    https://www.bousai.metro.tokyo.lg.jp/taisaku/saigai/1013388/index.html (in Japanese) [accessed on July 2,2021]

23. Hunter PR, Brainard J. Estimating the effectiveness of the Pfizer COVID-19 BNT162b2 vaccine after a single dose. A reanalysis of a study of 'real-world' vaccination outcomes from Israel.

    https://www.medrxiv.org/content/10.1101/2021.02.01.21250957v1 doi: https://doi.org/10.1101/2021.02.01.21250957

Figure 1: Vaccine coverage and shares of mutant strains N501Y and L452R in 2021

share of the mutant strain (%)   vaccine coverage (%)

Note: Blue bars represent shares of N501Y. Light blue bars represent shares of L452R, in Tokyo measured at the left-hand side. Red scattered points denote vaccine coverage as defined by the first dose with a 12-day delay. Black scattered points denote vaccine coverage defined by the second dose. The two vaccine coverage data were measured at the right-hand side. Because daily vaccine coverage was not reported on weekends and national holidays, data of vaccine coverage are missing for these days.

Figure 2: Effective reproduction number from February, 2020 through June 10, 2021.

R(*t*)

(date)

Figure 3: Empirical distribution of duration from onset to report by MHLW, Japan.

(%)

Note: Bars represent the probability of duration from onset to report based on 657

patients for whom the onset date was available in Japan. Data were obtained from

MHLW, Japan.

Figure 4: Empirical distribution of the incubation period published by MHLW, Japan.

(%)

(days)

Notes: Bars show the distribution of incubation periods for 91 cases for which the exposure date and onset date were published by MHLW, Japan. Patients for whom incubation was longer than 14 days are included in the bar shown for day 14.

Table 1: Estimation results of R(t) on vaccine coverage with the climate condition, mobility, and countermeasures

Specification	First dose		Second dose		Final form	
Explanatory variable	Estimated coefficient	p-value	Estimated coefficient	p-value	Estimated coefficient	p-value
Temperature	-0.00552	0.206	-0.00578	0.190	N.A.	N.A.
Humidity	-0.00187	0.279	-0.00172	0.322	N.A.	N.A.
Mobility	0.00954	0.000	0.00961	0.000	0.00893	0.000
SCVEC	0.72630	0.000	0.72397	0.000	0.82686	0.000
1st Emergency status	-0.84730	0.000	-0. 84562	0.000	-0. 82230	0.000
GTTC	-0.92010	0.000	-0.92214	0.000	-0.88165	0.000
2nd Emergency status	-1.06823	0.000	-1.06986	0.000	-0.91042	0.000
3rd Emergency status	0.33644	0.180	0.31917	0.31917	N.A.	N.A.
Vaccine	0.03087	0.429	0.09913	0.09913	N.A.	N.A.

coverage (%)

N501Y(%)	-0.02050	0.000	-0.02080	0.000	-0.01497	0.000
L452R(%)	-0.00181	0.967	-0.00113	0.978	N.A.	N.A.
Constant	1.12603	0.000	1.11259	0.000	0.91848	0.000
Adjusted R^2	0.5766		0.5657		0.5562	
Number of observations	467		460		485	

Note: The dependent variable was R(t): GTTC stands for "Go To Travel Campaign";

SCVEC denotes school closure and voluntary event cancellation. The sample period

was from February 1, 2021 through June 10, 2021 as of July 3, 2021.