

1 **Detection of B.1.351 and B.1.1.7 SARS-CoV-2 variants in Monterrey metropolitan**
2 **area in Northeast Mexico**

3 Kame A. Galán-Huerta, MD/PhD¹, Eduardo Garza-de-la-Peña, MD/PhD², Gabriela V.
4 Elizondo-Valdez, Bs², María F. Herrera-Saldivar, Bs¹, Ana M. Rivas-Estilla, PhD¹, Sonia
5 A. Lozano-Sepúlveda, PhD¹, Natalia Martínez-Acuña, PhD¹, Daniel Arellanos-Soto, PhD¹,
6 Javier Ramos-Jiménez MD FIDSA^{1,3}

7

8 ¹Centro de Investigación e Innovación en Virología Médica, Departamento de Bioquímica
9 y Medicina Molecular, Facultad de Medicina, Universidad Autónoma de Nuevo León;

10 ²Laboratorio de Patología Clínica y Genética Molecular;

11 ³Servicio de Infectología, Hospital Universitario “Dr. José Eleuterio González”,
12 Universidad Autónoma de Nuevo León;

13

14 Corresponding author:

15 Javier Ramos-Jiménez

16 Avenida Francisco I. Madero S/N, Colonia Mitras Centro, Monterrey, NL. 64460. Mexico.

17 +52(81)8348-5013

18 javier.ramosjm@uanl.edu.mx

19 **Abstract word count: 45**

20 **Text word count: 768**

21 **Article summary line:** Introduction of B.1.351 and B.1.1.7 SARS-CoV-2 variants in
22 Monterrey, Mexico.

23 **Running title:** SARS-CoV-2 variants of concern, Monterrey, Mexico

24 **Keywords:** SARS-CoV-2, Variant of concern, Phylogeny, Mexico

25

26 **Detection of B.1.351 and B.1.1.7 SARS-CoV-2 variants in Monterrey metropolitan**
27 **area in Northeast Mexico**

28 **Authors:**

29 Kame A. Galán-Huerta, Eduardo Garza-de-la-Peña, Gabriela V. Elizondo-Valdez, María F.
30 Herrera-Saldivar, Ana M. Rivas-Estilla, Sonia A. Lozano-Sepúlveda, Natalia Martínez-
31 Acuña, Daniel Arellanos-Soto, and Javier Ramos-Jiménez

32

33 **Author affiliations:**

34 Universidad Autónoma de Nuevo León, Monterrey, Mexico (K.A. Galán-Huerta, M.F.
35 Herrera-Saldivar, A.M. Rivas-Estilla, S.A. Lozano-Sepúlveda, N. Martínez-Acuña, D.
36 Arellanos-Soto, J. Ramos-Jiménez)

37 Laboratorio de Patología Clínica y Genética Molecular, Monterrey, Mexico (E. Garza-de-
38 la-Peña, G.V. Elizondo-Valdez)

39

40

41

42 **Abstract**

43 SARS-CoV-2 variants of concern (VOC) have spread throughout the world. In
44 Mexico, VOC B.1.351 has been detected on one occasion and B.1.1.7 several times. We
45 detected lineages B.1.351 and B.1.1.7 in patients who traveled to USA and B.1.1.7 in a
46 patient with no travel history.

47

48 Since late 2020, several variants of SARS-CoV-2 have emerged. These variants,
49 named variants of concern (VOC), are monitored because they are associated with an
50 increase in transmissibility, increase in severity, and vaccine effectiveness reduction (1-3).
51 VOC Alpha (Lineage B.1.1.7) was first detected in Mexico in December 2020, and Nuevo
52 Leon in January 2021. VOC Beta (Lineage B.1.351) was detected in Mexico in April 2021
53 and not previously detected in Nuevo Leon.

54 We randomly selected, via randomizer.org, residual RNA samples derived from
55 nasopharyngeal swabs from patients living in Nuevo Leon, Mexico. Samples were positive
56 for SARS-CoV-2 and had Ct values <30 in TaqPath COVID-19 CE-IVD RT-PCR Kit
57 (Thermo Fisher Scientific, Waltham, MA). We used samples after a two-week retention
58 period. To maintain anonymity, samples were deidentified and only age and gender were
59 obtained. We sequenced the whole genome with ARTIC Network's amplicon sequencing
60 protocol (4) in a MinION (Oxford Nanopore Technologies, Oxford, UK).

61 Global reference sequences were selected from Nextstrain's global dataset (n=3644)
62 and all the B.1.1.7 and B.1.351 genomes from Mexico until April 29th, 2021 (n=77) and

63 retrieved them from the GISAID database. We did a BLAST search for each of the recently
64 obtained sequences and retrieved the five most similar sequences.

65 Afterwards, we used Nextstrain (5) to analyze the newly obtained sequences and to
66 get the global perspective of our data. Posteriorly, we made two alignments. One with all
67 the sequences, from lineage B.1.1.7, that were included in the clades where the sequenced
68 samples were grouped. The other alignment included all the sequences belonging to the
69 lineage B.1.351. We constructed a maximum-likelihood tree in IQ-TREE v2.0.3 (6) for
70 each alignment, searched for temporal data in TempEst v1.5.3 and eliminated outlier
71 sequences shown in the root-to-tip plot.

72 Next, we inferred maximum clade credibility trees for each alignment with BEAST
73 v1.10.4 (7). We used a strict molecular clock, HKY+G substitution model, and a coalescent
74 exponential growth tree prior. We used a chain length of 100 million and sampled every
75 10,000 steps. Chain convergence was evaluated using Tracer v.1.7.1 and trees were
76 summarized using TreeAnnotator discarding 10% as burn-in. Sequences generated in this
77 report were deposited in GISAID with the following numbers: EPI_ISL_1789696 -
78 EPI_ISL_1789699.

79 The Ethics Committee from Facultad de Medicina y Hospital Universitario
80 Universidad Autónoma de Nuevo León reviewed and approved this work with the
81 following registration number: BI20-0004.

82 We detected lineage B.1.351 in one patient and lineage B.1.1.7 in three patients.
83 The four patients were males and three of them had traveled to USA in the previous two
84 weeks. Lineage B.1.351 was detected in a middle-aged man with recent travel to

85 Pennsylvania, USA. He had no contact with similar cases and had onset of symptoms the
86 second week of April. Lineage B.1.1.7 was detected in two teenagers and one middle-aged
87 man. The two teenagers had recent travel history to Colorado, USA, contact with similar
88 cases and had onset of symptoms the first week of April. The middle-aged man had no
89 recent travel history, no contact with similar cases and had onset of symptoms the second
90 week of April. These four patients were treated ambulatory.

91 Initial phylogenetic analysis places the sample NLE-UANL-074 in lineage B.1.351
92 and samples NLE-UANL-067, 068 and 075 in lineage B.1.1.7 (Figure panel A). Until May
93 2021, there were only three B.1.351 viruses detected in Mexico: two from Campeche, and
94 one from Baja California Norte. Sample NLE-UANL-074 groups in a well-supported clade
95 with viruses isolated in Pennsylvania state (Figure panel B and D). This clade does not
96 group with the previously reported samples from Mexico, confirming an introduction.

97 B.1.1.7 lineage was previously introduced to Mexico and explains the samples
98 dispersed though the phylogenetic tree (Figure panel C). Further analysis revealed that
99 samples NLE-UANL-067 and 068 grouped with samples isolated in Florida in a well-
100 supported clade (Figure panel E). These viruses differ from previously reported in Mexico,
101 suggesting an introduction from USA. Sample NLE-UANL-075, isolated from the patient
102 without travel history, groups with viruses isolated in Texas in a well-supported clade
103 (Figure panel F). This finding could be explained if the patient had contact with people who
104 traveled to Texas and were pre-symptomatic.

105 Even though samples NLE-UANL-067, 068 and 075 belong to lineage B.1.1.7 and
106 were isolated in Nuevo Leon, they are not related because they group in different well-

107 supported clades. Even though lineage B.1.1.7 was previously detected in Nuevo Leon,
108 there was no widespread transmission throughout the state.

109 This is the first report of B.1.351 and the third detection of B.1.1.7 in Nuevo Leon
110 state. We need posterior surveillance to evaluate community transmission and association
111 of these lineages with a new wave of COVID-19 cases.

112 **Acknowledgments**

113 This work was funded by Consejo Nacional de Ciencia y Tecnología (CONACyT)
114 with the following grant number 312328, awarded to Kame A. Galán-Huerta. The authors
115 gratefully acknowledge the authors and submitting laboratories that generated genome data
116 and shared it via GISAID. A full list acknowledging the authors submitting data used in
117 this report can be found in Appendix Table.

118 **Author bio**

119 Dr. Galán-Huerta is an assistant professor and founder member of the Center for
120 Investigation and Innovation in Medical Virology, in the Faculty of Medicine of the
121 Autonomous University of Nuevo Leon. His primary research interests are emerging and
122 reemerging viral diseases and epidemiology.

123

124 **References**

125 1. World Health Organization. Tracking SARS-CoV-2 variants. [cited 2021 Jun 03].
126 <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>

- 127 2. Challen R, Brooks-Pollock E, Read JM, Dyson L, Tsaneva-Atanasova K, Danon L.
128 Risk of mortality in patients infected with SARS-CoV-2 variant of concern
129 202012/1: matched cohort study. *BMJ*. 2021 Mar 9;372:n579. doi:
130 10.1136/bmj.n579. PubMed PMID: 33687922; PubMed Central PMCID:
131 PMC7941603.
- 132 3. Madhi SA, Baillie V, Cutland CL, Voysey M, Koen AL, Fairlie L, et al. Efficacy of
133 the ChAdOx1 nCoV-19 Covid-19 Vaccine against the B.1.351 Variant. *N Engl J*
134 *Med*. 2021 May 20;384(20):1885-1898. doi: 10.1056/NEJMoa2102214. Epub ahead
135 of print. PubMed PMID: 33725432.
- 136 4. Josh Quick 2020. nCoV-2019 sequencing protocol v3 (LoCost). protocols.io. [cited
137 2021 Jun 03]. [https://protocols.io/view/ncov-2019-sequencing-protocol-v3-locost-](https://protocols.io/view/ncov-2019-sequencing-protocol-v3-locost-bh42j8ye)
138 [bh42j8ye](https://protocols.io/view/ncov-2019-sequencing-protocol-v3-locost-bh42j8ye)
- 139 5. Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, et al.
140 Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics*. 2018 Dec
141 1;34(23):4121-4123. doi: 10.1093/bioinformatics/bty407. PubMed PMID:
142 29790939; PubMed Central PMCID: PMC6247931.
- 143 6. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler
144 A, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
145 in the Genomic Era. *Mol Biol Evol*. 2020 May 1;37(5):1530-1534. doi:
146 10.1093/molbev/msaa015. PubMed PMID: 32011700; PubMed Central PMCID:
147 PMC7182206.
- 148 7. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian
149 phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evol*.

150 2018 Jun 8;4(1):vey016. doi: 10.1093/ve/vey016. PubMed PMID: 29942656;

151 PubMed Central PMCID: PMC6007674.

152 8. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. ggtree: an R package for visualization
153 and annotation of phylogenetic trees with their covariates and other associated data.
154 Methods Ecol Evol. 2017 Jan 8(1):28-36. doi: 10.1111/2041210X.12628






155

156 Address for correspondence: Javier Ramos-Jiménez, Servicio de Infectología, Hospital
157 Universitario “Dr. José Eleuterio González”, Facultad de Medicina, Universidad Autónoma
158 de Nuevo León, Avenida Francisco I. Madero S/N, Colonia Mitras Centro, Monterrey, NL,
159 Mexico. CP: 64460; +52(81)8348-5013; e-mail: javier.ramosjm@uanl.edu.mx

160

161 Figure. Phylogenetic relationship of SARS-CoV-2, 2019-2021. A) Global context tree
162 highlighting lineages B.1.351 (blue) and B.1.1.7 (green). Black boxes show sequences that
163 were analyzed. B) Phylogenetic tree of B.1.351 lineage showing the relationship of samples
164 from this report with samples from Mexico and USA. The black box shows posterior
165 magnification. C) Phylogenetic tree of B.1.1.7 lineage showing the relationship of samples
166 from this report with samples from Mexico and USA. Black boxes show posterior
167 magnification. D) Magnification of B.1.351 tree showing the clade in which the sequenced
168 virus groups. E) Magnification of the discontinuous box in the B.1.1.7 tree showing the
169 relationship of the sequenced viruses. F) Magnification of the continuous box in the B.1.1.7
170 tree showing the relationship of the sequenced virus. Posterior probabilities are shown in
171 the nodes of the magnifications. Viruses reported in this research letter are in red.

172 Phylogenetic trees were visualized in ggtree (8). Sequences used in this analysis were
173 downloaded from GISAID. The authors of sequences used are listed in the Appendix Table.

A**B****C****D****E****F**