COVID-19 pandemic dynamics in India, the SARS-CoV-2 Delta variant, and implications for vaccination

Wan Yang[1*] and Jeffrey Shaman[2]

[1]Department of Epidemiology, [2]Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, New York, NY, USA

*Correspondence to: wy2202@cumc.columbia.edu

Abstract

Background: The COVID-19 Delta pandemic wave in India surged and declined within 3 months; cases then remained low despite the continued spread of Delta elsewhere. Here we aim to estimate key epidemiological characteristics of the Delta variant based on data from India and examine the underpinnings of its dynamics.

Methods: We utilize multiple datasets and model-inference methods to reconstruct COVID-19 pandemic dynamics in India during March 2020 – June 2021. We further use model estimates to retrospectively predict cases and deaths during July – mid-Oct 2021, under various vaccination and vaccine effectiveness (VE) settings to estimate the impact of vaccination and VE for non-Delta-infection recoverees.

Findings: We estimate that Delta escaped immunity in 34.6% (95% CI: 0 – 64.2%) of individuals with prior wildtype infection and was 57.0% (95% CI: 37.9 – 75.6%) more infectious than wildtype SARS-CoV-2. Models assuming higher VE among those with prior non-Delta infection, particularly after the 1st dose, generated more accurate predictions than those assuming no such increases (best-performing VE setting: 90/95% vs. 30/67% baseline for the 1st/2nd dose). Counterfactual modeling indicates that high vaccination coverage for 1st vaccine-dose in India (~50% by mid-Oct 2021) combined with the boosting of VE among recoverees averted around 60% of infections during July – mid-Oct 2021.

Interpretation: Non-pharmaceutical interventions, infection seasonality, and high coverage of

1-dose vaccination likely all contributed to pandemic dynamics in India during 2021. Given the

shortage of COVID-19 vaccines globally and boosting of VE, for populations with high prior

infection rates, prioritizing the first vaccine-dose may protect more people.

Key words: COVID-19; Delta SARS-CoV-2 variant; India; vaccination effectiveness; boosting;

prior infection

Research in context

Evidence before this study

We searched PubMed for studies published through Nov 3, 2021 on the Delta (B.1.617.2) SARS-

CoV-2 variant that focused on three areas: 1) transmissibility [search terms: ("Delta variant" OR

"B.1.617") AND ("transmission rate" OR "growth rate" OR "secondary attack rate" OR

"transmissibility")]; 2) immune response ([search terms: ("Delta variant" OR "B.1.617") AND

("immune evas" OR "immune escape")]; and 3) vaccine effectiveness ([search terms: ("Delta

variant" OR "B.1.617") AND ("vaccine effectiveness" OR "vaccine efficacy" OR "vaccination")].

Our search returned 256 papers, from which we read the abstracts and identified 54 relevant

studies.

Forty-two studies addressed immune evasion and/or vaccine effectiveness. Around half (n=19)

of these studies measured the neutralizing ability of convalescent sera and/or vaccine sera

against Delta and most reported some reduction (around 2- to 8-fold) compared to ancestral

variants. The remainder (n=23) used field observations (often with a test-negative or cohort-

design) and reported lower VE against infection but similar VE against hospitalization or death.

Together, these laboratory and field observations consistently indicate that Delta can evade

preexisting immunity. In addition, five studies reported higher B-cell and/or T-cell vaccine-

induced immune response among recovered vaccinees than naïve vaccinees, suggesting

potential boosting of pre-existing immunity; however, all studies were based on small samples

(n = 10 to 198 individuals).

Sixteen studies examined transmissibility, including 1) laboratory experiments (n=6) showing

that Delta has higher affinity to the cell receptor, fuses membranes more efficiently, and/or

replicates faster than other SARS-CoV-2 variants, providing biological mechanisms for its higher

transmissibility; 2) field studies (n=5) showing higher rates of breakthrough infections by Delta

and/or higher viral load among Delta infections than other variants; and 3) modeling/mixed

studies (n=5) using genomic or case data to estimate the growth rate or reproduction number,

reporting a 60-120% increase. Only one study jointly estimated the increase in transmissibility

(1.3-1.7-fold, 50% CI) and immune evasion (10-50%, 50% CI); this study also reported a 27.5%

(25/91) reinfection rate by Delta.

Added value of this study

We utilize observed pandemic dynamics and the differential vaccination coverage for two

vaccine doses in India, where the Delta variant was first identified, to estimate the

epidemiological properties of Delta and examine the impact of prior non-Delta infection on

immune boosting at the population level. We estimate that Delta variant can escape immunity

from prior wildtype infection roughly one-third of the time and is around 60% more infectious

than wildtype SARS-CoV-2. In addition, our analysis suggests the large increase in population

receiving their first vaccine dose (~50% by end of Oct 2021) combined with the boosting effect

of vaccination for non-Delta infection recoverees likely mitigated epidemic intensity in India

during July – Oct 2021.

Implications of all the available evidence

Our analysis reconstructs the interplay and effects of non-pharmaceutical interventions,

infection seasonality, Delta variant emergence, and vaccination on COVID-19 pandemic

dynamics in India. Modeling findings support prioritizing the first vaccine dose in populations

with high prior infection rates, given vaccine shortages.

INTRODUCTION

The Delta (PANGO lineage: B.1.617.2) SARS-CoV-2 variant of concern (VOC)[1-4] has spread quickly to over 170 countries (GISAID,[5] as of 11/3/2021). Several lines of evidence have indicated that Delta is able to evade immunity from prior infection by preexisting variants; these include reduced neutralizing ability of convalescent sera and vaccinee sera against Delta,[6-9] reduced vaccine effectiveness (VE) against infection,[10-13] and reduced VE against symptomatic disease after 1-dose of vaccine (but only slight reduction for full vaccination).[14-16] In addition, studies have found a higher secondary attack rate, growth rate, or reproduction number for Delta than prior variants including Alpha (range of the mean estimates: 60-120%).[2,17-21] In particular, a recent study, fitting a model to mortality data in Delhi, India, estimated a 1.3-1.7-fold (50% CI) increase in transmissibility and 10-50% (50% CI) immune evasion for Delta; however, the authors noted large uncertainty in their estimates.[22] Further, factors such as host behavioral changes and seasonal modulation of risk due to changes in environmental conditions are difficult to account for and could confound these estimates. As a result, estimates of prior immunity evasion and relative transmissibility for Delta and the contributions of these properties to the rapid spread of this variant remain uncertain.

India, where Delta was first identified, experienced an intensive pandemic wave in late March 2021. However, unlike many places seeing a prolonged Delta pandemic wave, the Delta wave in India only lasted 3 months and declined rapidly after peaking mid-May. Since June 2021 cases have remained low. A high infection rate after the Delta wave has been cited as a reason for this dramatic epidemic decline, as vaccination coverage was low at the time (4.2% fully vaccinated at the end of June 2021). However, given an estimated basic reproduction number (R_0) of 6-7,[20] roughly 83-86% ($1 - 1/R_0$) of the population would need immunity for the Delta epidemic to subside. Assuming 10-50% immunity escape[22] and a 25-35% infection rate prior to the Delta wave,[23] this implies that 53-73% of India's 1.4 billion people would have been infected by Delta within the span of 3 months, despite a national lockdown at the time.

To better understand COVID-19 pandemic dynamics in India and the epidemiological

characteristics of Delta, here we utilize a model-inference method recently developed for SARS-

CoV-2 VOCs. The model-inference method incorporates epidemiological, population mobility,

and weather data to model SARS-CoV-2 transmission dynamics, while accounting for case

under-ascertainment, impacts of non-pharmaceutical interventions (NPIs) and vaccination,

infection seasonality, and new variants.[24] Applying this method, we have jointly estimated the

immune escape potential and change in transmissibility for Alpha, Beta, and Gamma,

separately, using data from countries where these three VOCs were first reported.[24] In

addition, several laboratory studies have reported stronger vaccine-induced immune responses

among recovered vaccinees than naïve vaccinees, suggesting potential boosting of pre-existing

immunity. In India, while only 23% of the population have received two vaccine doses, 53%

have received their first vaccine dose, as of the end of Oct 2021. This large discrepancy in one-

and two dose coverage, combined with a likely high population infection rate, offers an

opportunity to examine the boosting effect of prior non-Delta infection on vaccine-induced

immunity at the population level. Therefore, in this study, we first reconstruct the pandemic

dynamics in India during March 2020 – June 2021 and estimate key epidemiological

characteristics of Delta. We then further use our model estimates to retrospectively predict

cases and deaths during July – Oct 2021, under various vaccination and VE scenarios, and

compare these simulations to observations in order to estimate the impact of vaccination and

VE for those with prior non-Delta infection.

METHODS

Data sources and processing

We used reported COVID-19 case and mortality data to capture transmission dynamics,

weather data to estimate infection seasonality, mobility data to represent concurrent NPIs, and

vaccination data to account for changes in population susceptibility due to vaccination in the

model-inference system. COVID-19 case and mortality data from the week of March 8, 2020

(the first week COVID-19 deaths were reported in India) to the week of Oct 17, 2021 came from

the COVID-19 Data Repository of the Center for Systems Science and Engineering (CSSE) at

Johns Hopkins University.[25,26] Surface station temperature and humidity data were accessed

using the "rnoaa" R package.[27] We then aggregated these data for all weather stations in India

(n = 390 stations) with measurements from Jan 2020 to Oct 2021 and calculated the average for

each week of the year. Mobility data were derived from Google Community Mobility Reports;[28]

we aggregated all business-related categories (i.e., retail and recreational, transit stations, and

workplaces) in all locations in India to weekly intervals. Vaccination data (1st and 2nd dose) were

obtained from Our World in Data.[29,30]

Model-inference system

The model-inference system was developed and described in detail in our previous study.[31]

Below we describe each component in brief.

Epidemic model

The epidemic model follows an SEIRSV (susceptible-exposed-infectious-recovered-susceptible-

vaccination) construct per Eqn 1:

$$\begin{cases} \dfrac{dS}{dt} = \dfrac{R}{L_t} - \dfrac{b_t e_t m_t \beta_t IS}{N} - \varepsilon - v_{1,t} - v_{2,t} \\[2mm] \dfrac{dE}{dt} = \dfrac{b_t e_t m_t \beta_t IS}{N} - \dfrac{E}{Z_t} + \varepsilon \\[2mm] \dfrac{dI}{dt} = \dfrac{E}{Z_t} - \dfrac{I}{D_t} \\[2mm] \dfrac{dR}{dt} = \dfrac{I}{D_t} - \dfrac{R}{L_t} + v_{1,t} + v_{2,t} \end{cases}$$

where S, E, I, R are the number of susceptible, exposed (but not yet infectious), infectious, and

recovered/immune/deceased individuals; N is the population size; and ε is the number of

travel-imported infections. In addition, the model includes the following key components:

1) Virus-specific properties, including the time-varying variant-specific transmission rate

β_t, latency period Z_t, infectious period D_t, and immunity period L_t. Note the subscript, t, denotes

time in week, as all parameters are estimated for each week as described below.

2) The impact of NPIs. Specifically, we use relative population mobility (see data above)

to adjust the transmission rate via the term m_t. To further account for potential changes in

effectiveness, the model additionally includes a parameter, e_t, to scale NPI effectiveness.

3) The impact of vaccination, via the terms $v_{1,t}$ and $v_{2,t}$. Specifically, $v_{1,t}$ is the number of

individuals successfully immunized after the first dose of vaccine and is computed using

vaccination data and vaccine efficacy for 1st dose; and $v_{2,t}$ is the additional number of

individuals successfully immunized after the second vaccine dose (i.e., excluding those

successfully immunized after the first dose).

4) Infection seasonality, computed using temperature and specific humidity data as

described previously (see supplemental material of Yang and Shaman[24]). The estimated relative

seasonal trend, b_t, is used to adjust the relative transmission rate at time t.

Observation model to account for under-detection and delay

Using the model-simulated number of infections occurring each day, we further computed the

number of cases and deaths each week to match with the observations, as done in Yang et al.[32]

For example, for case data, we include 1) a time-lag from infectiousness to detection (i.e., an

infection being diagnosed as a case) to account for delays in detection; and 2) an infection-

detection rate (r_t), i.e. the fraction of infections (including subclinical or asymptomatic

infections) reported as cases, to account for under-detection. To compute the model-simulated

number of new cases per week, we multiply the model-simulated number of new infections per

day by the infection-detection rate, and further distribute these simulated cases in time per the

distribution of time-from-infectiousness-to-detection. We then aggregate the daily lagged,

simulated cases to weekly totals for model inference (see below).

Model inference and parameter estimation

The inference system uses the EAKF,[33] a Bayesian statistical method, to estimate model state

variables (i.e., S, E, I, R from Eqn 1) and parameters (i.e., β_t, Z_t, D_t, L_t, e_t, from Eqn 1 as well as r_t

and other parameters from the observation model). Briefly, the EAKF uses an ensemble of

model realizations (n=500 here), each with initial parameters and variables randomly drawn

from a *prior* range (see Table S1). After model initialization, the system integrates the model

ensemble forward in time for a week (per Eqn 1) to compute the prior distribution for each

model state variable and parameter, as well as the model-simulated number of cases and

deaths for that week. The system then combines the prior estimates with the observed case

and death data for the same week to compute the posterior per Bayes' theorem.[33] During this

filtering process, the system updates the posterior distribution of all model variables and

parameters for each week.

Estimating the immune escape potential and changes in transmissibility for Delta

To identify the most plausible combination of changes in transmissibility and level of immune

evasion, per methods developed in ref[24], we ran the model-inference, repeatedly and in turn,

to test 14 major combinations of these two quantities and select the best performing run.

Based on the best-performing model estimates, we then computed the variant-specific

transmissibility as the product of the variant-specific transmission rate (β_t) and infectious

period (D_t). To reduce uncertainty, we averaged transmissibility estimates over the first

pandemic wave and the period when Delta is dominant, separately. We then computed the

average change in transmissibility due to Delta as the ratio of the two averaged estimates (i.e.,

after: before the rise of Delta). To quantify immune evasion, we recorded the changes in

susceptibility over time and then computed the level of immune evasion as the ratio of the

total increase in susceptibility due to immune evasion during the second wave to the model-

estimated population immunity at the end of the first wave.

To account for model stochasticity, we repeated the model-inference process 300 times, each

with 500 model realizations and summarized the results from all 150,000 model estimates. As a

sensitivity test and part of the effort to examine the impact of prior non-Delta infection on VE,

we performed the analysis using 12 different VE settings (see details below).

Model validation using independent data

To compare model estimates with independent observations not assimilated into the model-

inference system, we identified three measurements of cumulative infection rates from three

nationwide serology surveys in India: i) the first national serosurvey conducted during May 11 –

June 4, 2020 (n = 28,000 adults 18 years or older);[34] ii) the second national serosurvey

conducted during August 18 – September 20, 2020 (n = 29,082 individuals 10 years or older);[35]

and iii) the third national serosurvey conducted during December 18, 2020 – January 6, 2021 (n

= 28,598 individuals 10 years or older).[36] To account for the delay in antibody generation, we

shifted the timing of each serosurvey 14 days when comparing survey results to model-

inference system estimates of cumulative infection rates in Fig 1B.

Estimating the impact of vaccination and prior non-Delta infection on boosting vaccine-

induced immunity

We generated retrospective projections of cases and deaths from the week starting 7/4/2021

to the week starting 10/17/2021 (i.e., 16 weeks following the model-inference period), under

various vaccination and VE settings. We considered four levels of VE for those recovered from

non-Delta infection: 1) No boosting effect, i.e., using the same VE values as those without prior

infection. Here, we set VE at fourteen days after the 1st dose (VE1) to 30% and at seven days

after the 2nd dose (VE2) to 67%, based on data for the AstraZeneca vaccine against Delta;[15] 2)

Higher VE for the 1st dose but no future boosting for the 2nd dose (here, VE1 is set to 40%, 50%,

or 60%, and VE2 fixed at 67%); 3) Higher VE for the 2nd dose but not 1st dose (here, VE1 is fixed

at 30% and VE2 set to 75%, 85%, or 95%); and 4) Higher VE for both doses (here, VE1/VE2 are

set to 50%/75%, 60%/80%, 70%/85%, 80%/90%, or 90%/95%). To test the impact of

vaccination, in addition to projections using reported vaccination rates, we also generated

counterfactual projections assuming no further vaccination during the 16-week period.

For all projections, the model was initiated using model-inference estimates made at the week

of 6/27/2021, except for the infection-fatality risk (IFR). For IFR, estimates were decreasing

during June 2020 (Fig S1B) and model-inference extended to the end of July 2021 showed

continued decreases, likely due to improved healthcare and increased protection from prior

infection or vaccination. We thus assumed that IFR would decrease linearly for the first 6 weeks

of the projection period and then flatten and remain at that low IFR until the week of

10/17/2021. To account for NPIs, we used mobility data during the week of 7/4/2021 – the

week of 10/17/2021. As for the model-inference runs, we repeated the projections for each

scenario 300 times (each with 500 model realizations) and summarized the projections from all

runs. To evaluate the projection accuracy, we computed the relative root-mean-square-error

(RRMSE) and correlation between the projected and observed values for cases and deaths,

respectively.

RESULTS

The first COVID-19 pandemic wave in India, March 2020 – January 2021

From March 2020 to January 2021 India recorded over 10 million COVID-19 cases (0.77% of its

population); however, a nationwide serology survey suggested that ~24% of its population had

been infected by December 2020.[23] Accounting for under-detection of infection (Fig S1),

implemented non-pharmaceutical interventions (NPIs), seasonality, and vaccination, we used

the model-inference system to reconstruct pandemic dynamics in India since March 2020 (Fig

1A). Model-estimated infection rates closely match with measurements from three nationwide

serologic surveys conducted during the early, mid, and late phases of the first pandemic wave

(Fig 1B). Our analysis indicates that the 2-month long national lockdown (March 24 – May 31,

2020) and the less favorable weather conditions during pre-monsoon season (i.e., March –

May) likely contributed to initial low infection rates. By mid-May 2020, the model-inference

system estimates that only 0.43% (95% CrI: 0.19 – 1.7%) of the population had been infected

[vs. 0.73% (95% CI: 0.34%, 1.13%) among adults estimated by serosurvey[34]]. As the country

lifted its lockdown in June 2020 and entered the monsoon season (June – September), when

conditions are likely more favorable for transmission (Fig 1C), the first pandemic wave began.

Nevertheless, continued regional restrictions during June – November 2020 and less favorable

weather conditions during the autumn (October – November; see mobility and seasonal trends

in Fig 1C) likely mitigated pandemic intensity. The estimated mean of the reproduction number

R_t (i.e., average number of secondary infections per primary infection) was above 1 but less

than 1.35 from June to mid-September; in addition, R_t dropped below 1 during October –

November (Fig 1D). By the end of January 2021 when case rates reached a minimum following

the first wave, the model-inference system estimates that 26.1% (95% CrI: 19.9 – 33.0%) of the

population had been infected (Fig 1B).

The second pandemic wave in India and estimated epidemiological characteristics of Delta

Infections resurged dramatically in late March 2021, largely due to the rise of the Delta variant.

Despite a weeks-long second national lockdown implemented beginning April 20, 2021, India

reported another 19 million cases during late March – June 2021, about twice the number

reported during the previous 12 months. Accounting for under-detection (Fig S1), we estimate

that 32.3% (95% CrI: 22.4 – 46.5%) of the population were infected during this 3-month period,

including reinfections. This intense transmission was likely facilitated by the higher

transmissibility and immune evasive capabilities of the Delta variant. Estimated transmissibility

increased substantially during the second pandemic wave (Fig 1E). In addition, estimated

population susceptibility increased at the start of the second pandemic wave (Fig 1F),

suggesting loss of population immunity against Delta. Due to this immune escape, an estimated

50.5% (95% CrI: 21.8 – 79.0%) of the population remained susceptible at the end of June 2021,

despite two large pandemic waves and rollout of mass vaccination (of note, 19% of the

population had received at least 1 dose of vaccine by the end of June 2021). These findings

along with the seasonal trends described above suggest that the decline of the second wave

was largely due to the NPIs implemented and less favorable weather conditions during March –

May, rather than high population immunity.

Combining the model-inference estimates during the first and second pandemic waves in India,

we estimated that Delta was able to escape immunity among 34.6% (95% CI: 0 – 64.2%) of

individuals with prior wildtype infection and was 57.0% (95% CI: 37.9 – 75.6%) more

transmissible than wildtype SARS-CoV-2. Estimates are similar under different VE settings (Fig

S2).

Impact of vaccination and prior non-Delta infection on boosting vaccine-induced immunity

Despite the likely conducive conditions during the monsoon season (June – September), easing

of NPIs, and relatively high susceptibility estimated at the end of June 2021, cases and deaths in

India remained at relatively low levels during July – Oct 2021. Counterfactual modeling suggests

that the faster rollout of vaccination during this period substantially mitigated the epidemic risk

(Fig 2). Projected cases and deaths assuming no further vaccination uptake are much higher

than observed; In contrast, models including the reported vaccination rates more closely match

reported cases and deaths (Fig 2). Further, models assuming higher VE for non-Delta infection

recoverees generated more accurate projections than those assuming no boosting effect (Fig

3). The boosting effect appears to be more pronounced for the 1st vaccine dose (see, e.g., Fig 3A

where larger dots, representing higher VE after the 1st dose, had smaller errors). Overall, the

model assuming 90%/95% VE for the 1st/2nd dose of vaccine for non-Delta infection recoverees

generated the most accurate projections. These projections estimate that vaccination rollout

combined with the boosting effect averted 57% of infections during July – mid-Oct 2021.

Discussion

Combining epidemiological, behavioral, and weather observational data with a comprehensive

model-inference system, we estimate the Delta SARS-CoV-2 variant escaped immunity in

roughly one-third of individuals with wildtype infection during the previous year and was

around 60% more infectious than wildtype SARS-CoV-2. In addition, our analysis suggests the

large increase in population receiving their first vaccine dose (~50% by end of Oct 2021)

combined with the boosted VE for non-Delta infection recoverees likely mitigated the epidemic

intensity in India in recent months.

Previously, we have estimated the changes in transmissibility and immune escape potential for

three other major SARS-CoV-2 VOCs: namely, a 46.6% (95% CI: 32.3 – 54.6%) increase in

transmissibility but nominal immune escape for Alpha (i.e., B.1.1.7), a 32.4% (95% CI: 14.6 –

48.0%) increase in transmissibility and 61.3% (95% CI: 42.6 – 85.8%) immune escape for Beta

(i.e., B.1.351), and a 43.3% (95% CI: 30.3 – 65.3%) increase in transmissibility and 52.5% (95%

CI: 0 – 75.8%) immune escape for Gamma (i.e., P.1). Compared with Alpha, data from the UK

have shown that the secondary attack rate for contacts of cases with Delta was around 1.5

times higher than Alpha (12.4% vs. 8.2%), during March 29 – May 11, 2021.[2] In a partially

immunized population, the secondary attack rate reflects the combined outcome of the

transmissibility of the etiologic agent and population susceptibility to that agent. Consistent

with the UK data, our estimates of the relative transmissibility and immune escape potential

combine to a 44.1% (95% CI: 4.2 – 86.6%) higher secondary attack rate by Delta than Alpha [i.e.,

(1+57%)/(1+46.6%) × (1+34.6%) – 1 = 44.1% increase]. This higher competitiveness of Delta

over Alpha explains the rapid variant displacement observed in regions previously dominated

by Alpha (e.g. the UK and the US).

In addition, we estimate that 34.6% (95% CI: 0 – 64.2%) of individuals with acquired immunity

from wildtype infection would be susceptible to Delta due to immune escape. This estimate is

also in line with a recent study[22] reporting a 27.5% reinfection rate during the Delta pandemic

wave in Delhi, India, based on a small subset of people with repeated serology measures. In

addition to immune escape from wildtype infection, studies have also reported reduced ability

of sera from Beta- and Gamma-infection recoverees to neutralize Delta,[8,37] suggesting Delta can

also escape immunity conferred by those two VOCs. Such immune escape ability would also

allow Delta to rapidly replace Beta and Gamma in regions previously hard-hit by those two

VOCs, as has been observed in many countries in Africa and South America.[5] More

fundamentally, these findings highlight the complex, non-linear immune landscape of SARS-

CoV-2 and the importance to monitor the immune escape potential of new variants against

both previous and concurrent circulating variants.

Despite the successful development of multiple vaccines, shortage of supplies – particularly in

resource-limited countries – remains an impediment to global mass vaccination.[38] In response,

researchers have proposed dose sparing strategies such as fractionation[39] and 1-dose

vaccination for recoverees.[40] The latter 1-dose strategy draws on laboratory studies showing

higher vaccine-induced immune response among recovered vaccinees than naïve vaccinees

(i.e., boosting of pre-existing immunity).[41-44] Here, we utilized model-inference estimates and

vaccination data in India to test the impact of boosting at the population level. The findings

further support the effectiveness of 1-dose vaccination for recoverees. In light of continued

vaccine shortages, prioritizing first-dose vaccination thus may be an effective strategy for

mitigating COVID-19 burden in countries with high underlying SARS-CoV-2 infection rates.

Due to a lack of detailed epidemiological data (e.g., age-specific and subnational) and thus

model simplification, our estimates have uncertainties as indicated by the large credible

intervals. Nevertheless, these estimates are in line with independent data from three

nationwide serology surveys conducted at three time points during the first pandemic wave in

India (Fig 1B), as well as Delta-related epidemiological data from the UK[2] and Delhi, India,[22] as

discussed above; these consistencies support the accuracy of our estimates. Unlike estimates

from the contact tracing data, however, here we are able to separately quantify the changes in

transmissibility and immune escape potential of the Delta variant. In addition, our analysis also

suggests high 1-dose vaccine effectiveness among those with prior infection. These findings

and the methods used to generate them could support better understanding of future SARS-

CoV-2 variant dynamics given local prior infection rates, variant prevalence, and vaccination

coverage.

Data Availability: All data used in this study are publicly available as described in the "Data

sources and processing" section.

Code availability: All source code and data necessary for the replication of our results and

figures are made publicly available at https://github.com/wan-yang/covid_voc_delta.

Acknowledgements: This study was supported by the National Institute of Allergy and

Infectious Diseases (AI145883 and AI163023), the National Science Foundation Rapid Response

Research Program (RAPID; DMS-2027369) and a gift from the Morris-Singer Foundation.

Author contributions: WY designed the study, performed the analysis, and wrote the first draft;

JS provided input to the analysis and contributed to the final draft.

Competing interests: JS and Columbia University disclose partial ownership of SK Analytics. JS

discloses consulting for BNI.

References:

1. World Health Organization. Tracking SARS-CoV-2 variants. 2021.

https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/.

2. Public Health England. SARS-CoV-2 variants of concern and variants under investigation in

England. Technical briefing 14. 2021.

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachmen

t_data/file/991343/Variants_of_Concern_VOC_Technical_Briefing_14.pdf (accessed

6/16/2021 2021).

3. National Collaborating Centre for Infectious Diseases. Updates on COVID-19 Variants of

Concern. 4/26/2021 2021. https://nccid.ca/covid-19-variants/ (accessed 5/3/2021 2021).

4. Centers for Disease Control and Prevention. SARS-CoV-2 Variant Classifications and

Definitions. 2021. https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html

(accessed 6/17/2021 2021).

5. Global Initiative on Sharing All Influenza Data (GISAID). Tracking of Variants. 6/16/2021

2021. https://www.gisaid.org/hcov19-variants/ (accessed 6/16/2021 2021).

6. Mlcochova P, Kemp SA, Dhar MS, et al. SARS-CoV-2 B.1.617.2 Delta variant replication and

immune evasion. *Nature* 2021.

7. Wall EC, Wu M, Harvey R, et al. Neutralising antibody activity against SARS-CoV-2 VOCs

B.1.617.2 and B.1.351 by BNT162b2 vaccination. *Lancet* 2021.

8. Liu C, Ginn HM, Dejnirattisai W, et al. Reduced neutralization of SARS-CoV-2 B.1.617 by

vaccine and convalescent serum. *Cell* 2021; **184**(16): 4220-+.

9.   Arora P, Sidarovich A, Krüger N, et al. B.1.617.2 enters and fuses lung cells with increased
efficiency and evades antibodies induced by infection and vaccination. *Cell Reports* 2021;
37(2): 109825.

10.   Chemaitelly H, Tang P, Hasan MR, et al. Waning of BNT162b2 Vaccine Protection against
SARS-CoV-2 Infection in Qatar. *N Engl J Med* 2021.

11.   Goldberg Y, Mandel M, Bar-On YM, et al. Waning Immunity after the BNT162b2 Vaccine in
Israel. *N Engl J Med* 2021.

12.   Puranik A, Lenehan PJ, Silvert E, et al. Comparison of two highly-effective mRNA vaccines
for COVID-19 during periods of Alpha and Delta variant prevalence. *medRxiv* 2021:
2021.08.06.21261707.

13.   Elliott P, Haw D, Wang H, et al. Exponential growth, high prevalence of SARS-CoV-2, and
vaccine effectiveness associated with the Delta variant. *Science* 2021: eabl9551.

14.   Bernal JL, Andrews N, Gower C, et al. Effectiveness of COVID-19 vaccines against the
B.1.617.2 variant. *medRxiv* 2021: 2021.05.22.21257658.

15.   Bernal JL, Andrews N, Gower C, et al. Effectiveness of Covid-19 Vaccines against the
B.1.617.2 (Delta) Variant. *New England Journal of Medicine* 2021; **385**(7): 585-94.

16.   Seppala E, Veneti L, Starrfelt J, et al. Vaccine effectiveness against infection with the Delta
(B.1.617.2) variant, Norway, April to August 2021. *Eurosurveillance* 2021; **26**(35).

17.   Allen H, Vusirikala A, Flannagan J, et al. Household transmission of COVID-19 cases
associated with SARS-CoV-2 delta variant (B.1.617.2): national case-control study. *The*
Lancet regional health Europe 2021: 100252.

18.   Challen R, Dyson L, Overton CE, et al. Early epidemiological signatures of novel SARS-CoV-
2 variants: establishment of B.1.617.2 in England. *medRxiv* 2021: 2021.06.05.21258365.

19.   Earnest R, Uddin R, Matluk N, et al. Comparative transmissibility of SARS-CoV-2 variants
Delta and Alpha in New England, USA. *medRxiv* 2021.

20.   Vohringer HS, Sanderson T, Sinnott M, et al. Genomic reconstruction of the SARS-CoV-2
epidemic in England. *Nature* 2021.

21.   Campbell F, Archer B, Laurenson-Schafer H, et al. Increased transmissibility and global
spread of SARS-CoV-2 variants of concern as at June 2021. *Eurosurveillance* 2021; **26**(24).

22.   Dhar MS, Marwal R, Vs R, et al. Genomic characterization and epidemiology of an

453   emerging SARS-CoV-2 variant in Delhi, India. *Science* 2021: eabj9932.

454   23.   Murhekar MV, Bhatnagar T, Thangaraj JWV, et al. SARS-CoV-2 seroprevalence among the

455   general population and healthcare workers in India, December 2020–January 2021. *Int J*

456   *Infect Dis* 2021; **108**: 145-55.

457   24.   Yang W, Shaman J. Development of a model-inference system for estimating

458   epidemiological characteristics of SARS-CoV-2 variants of concern. *Nature*

459   *Communications* 2021; **12**: 5573.

460   25.   COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at

461   Johns Hopkins University. 2021. https://github.com/CSSEGISandData/COVID-19.

462   26.   Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real

463   time. *Lancet Infect Dis* 2020; **20**(5): 533-4.

464   27.   Chamberlain S, Anderson B, Salmon M, et al. Package 'rnoaa'. 2021.

465   28.   Google Inc. Community Mobility Reports. 2020.

466   https://www.google.com/covid19/mobility/.

467   29.   Data on COVID-19 (coronavirus) vaccinations by Our World in Data. 2020.

468   https://github.com/owid/covid-19-data/tree/master/public/data/vaccinations.

469   30.   Mathieu E, Ritchie H, Ortiz-Ospina E, et al. A global database of COVID-19 vaccinations.

470   *Nat Hum Behav* 2021; **5**(7): 947-53.

471   31.   Yang W, Shaman J. Epidemiological characteristics of three SARS-CoV-2 variants of

472   concern and implications for future COVID-19 pandemic outcomes. *medRxiv* 2021:

473   2021.05.19.21257476.

474   32.   Yang W, Kandula S, Huynh M, et al. Estimating the infection-fatality risk of SARS-CoV-2 in

475   New York City during the spring 2020 pandemic wave: a model-based analysis. *The Lancet*

476   *Infectious diseases* 2021; **21**(2): 203-12.

477   33.   Anderson JL. An ensemble adjustment Kalman filter for data assimilation. *Mon Weather*

478   *Rev* 2001; **129**(12): 2884-903.

34.   Murhekar MV, Bhatnagar T, Selvaraju S, et al. Prevalence of SARS-CoV-2 infection in India:

Findings from the national serosurvey, May-June 2020. *Indian J Med Res* 2020; **152**(1): 48-

57.

35.   Murhekar MV, Bhatnagar T, Selvaraju S, et al. SARS-CoV-2 antibody seroprevalence in

India, August-September, 2020: findings from the second nationwide household

serosurvey. *The Lancet Global Health* 2021; **9**(3): e257-e66.

36.   Murhekar MV, Bhatnagar T, Thangaraj JWV, et al. SARS-CoV-2 seroprevalence among the

general population and healthcare workers in India, December 2020 - January 2021. *Int J*

Infect Dis 2021; **108**: 145-55.

37.   de Oliveira T, Lessells R. Update on Delta and other variants in South Africa and other

world. 2021. https://www.krisp.org.za/manuscripts/DeltaGammaSummary_NGS-

SA_6JulV2.pdf.

38.   Padma TV. COVID vaccines to reach poorest countries in 2023 — despite recent pledges.

7/5/2021 2021. https://www.nature.com/articles/d41586-021-01762-w (accessed

11/11/2021 2021).

39.   Cowling BJ, Lim WW, Cobey S. Fractionation of COVID-19 vaccine doses could extend

limited supplies and reduce mortality. *Nature Medicine* 2021; **27**(8): 1321-3.

40.   Sasikala M, Shashidhar J, Deepika G, et al. Immunological memory and neutralizing activity

to a single dose of COVID-19 vaccine in previously infected individuals. *International*

Journal of Infectious Diseases 2021; **108**: 183-6.

41.   Andreano E, Paciello I, Piccini G, et al. Hybrid immunity improves B cells and antibodies

against SARS-CoV-2 variants. *Nature* 2021.

42.   Lucas C, Vogels CBF, Yildirim I, et al. Impact of circulating SARS-CoV-2 variants on mRNA

vaccine-induced immunity. *Nature* 2021.

43.   Sokal A, Barba-Spaeth G, Fernández I, et al. mRNA vaccination of naive and COVID-19-

recovered individuals elicits potent memory B cells that recognize SARS-CoV-2 variants.

Immunity 2021.

44. Sapkal GN, Yadav PD, Sahay RR, et al. Neutralization of Delta variant with sera of

Covishield™ vaccinees and COVID-19-recovered vaccinated individuals. *J Travel Med* 2021;

28(7).

Fig 1. Model-inference estimates and validation. (A) Model fit. (B) Model validation. (C) Observed relative mobility and estimated disease seasonal trend, compared to case and death rates over time. Key model-inference estimates are shown for the real-time reproduction number R_t (D), transmissibility (E), and population susceptibility (F). Blue lines and surrounding areas show the estimated mean, 50% (dark) and 95% (light) CrIs. Boxes and whiskers show the estimated mean, 50% and 95% CrIs for weekly cases and deaths in (A) and infection rates in (D) – (F). Grey shaded areas indicate the timing of national lockdowns (darker) or local restrictions (lighter); horizontal arrows indicate the timing of variant identification and vaccination rollout. In (C), for mobility (blue line; y-axis), values below 1 (dashed horizontal line) indicate reductions due to public health interventions. For the disease seasonal trend (orange line; y-axis), values above 1 indicate weather conditions more conducive for transmission than the yearly average and *vice versa*. *Note that the transmissibility estimates have removed the effects of changing population susceptibility, NPIs, and disease seasonality; thus, the trends are more stable than the reproduction number (R_t; left column) and reflect changes in variant-specific properties.*

Fig 2. Impact of vaccination. Model projections of weekly number of reported cases (A) and reported deaths (B) for India during July – mid-Oct 2021, compared to reported data. Crosses ('x') show reported data (left y-axis). Red dashed lines show median counterfactual model projections assuming no further vaccination uptake during the 16-week period. Blue dashed lines show median model projections using reported vaccination rates and assuming 90%/95% vaccine effectiveness for individuals with prior non-Delta infection after the 1st/2nd vaccine dose. Shaded areas with the same color show projected interquartile ranges. For comparison, estimated seasonality (orange lines), reported mobility (dark-blue lines), and cumulative vaccination uptake (full bar for 1st dose and filled section for 2nd dose) are overlaid (see right y-axis scale). All numbers are scaled per one million people.

Fig 3. Impact of prior non-Delta infection on immune boosting. Model projections under different vaccine effectiveness (VE) settings are used to examine the most plausible VE for individuals with prior non-Delta infection, based on projection accuracy: (A) the relative root-mean-square-error (RRMSE) and (B) correlation between the projected and observed values for cases and deaths, respectively. The size of the dots represents VE for recoverees after the 1st vaccine dose and the color represents VE after the 2nd vaccine dose. The dots with the black circles represent the baseline VE setting (i.e. 30%/67% for the 1st/2nd dose). For comparison, projected weekly numbers of reported cases (C) and reported deaths (D) under different VE settings are plotted along with the weekly actuals. For clarity, only median projections are shown here; see example projections including interquartile ranges in Fig 2.

Supplemental Table and Figures

Table S1. Prior ranges for the parameters used in the model-inference system.

Parameter/ variable	Symbol	Prior range	Source/rationale
Initial exposed	E(t=0)	500 – 1000 times of the reported number of cases on the first week of simulation (i.e. the week of 3/8/2020)	Low infection-detection rate in first weeks.
Initial infectious	I(t=0)	Same as for E(t=0)	
Initial susceptible	S(t=0)	99 – 100% of the population	Almost everyone is susceptible initially
Population size	N	N/A	Based on data
Variant-specific transmission rate	β	[0.4, 0.7]	Based on R_0 estimates of around 1.5-4 for SARS-CoV-2.[1-3] Slightly lower ranges are used here, as initial testing showed that use of higher values tended to overestimate the observations.
Scaling of effectiveness of NPI	e	[0.5, 1.5]	Around 1, with a large bound to be flexible.

Latency period	Z	[2, 5] days	Incubation period: 5.2 days (95% CI: 4.1, 7)[1]; latency period is likely shorter than the incubation period
Infectious period	D	[2, 5] days	Time from symptom onset to hospitalization: 3.8 days (95% CI: 0, 12.0) in China,[4] plus 1-2 days viral shedding before symptom onset. We did not distinguish symptomatic/asymptomatic infections.
Immunity period	L	[730, 1095] days	Assuming immunity lasts for 2-3 years
Mean of time from viral shedding to diagnosis	T_m	[5, 8] days	From a few days to a week from symptom onset to diagnosis/reporting,[4] plus 1-2 days of viral shedding (being infectious) before symptom onset. There may be a slightly longer delay for South Africa and Brazil.
Standard deviation (SD) of time from viral shedding to diagnosis	T_{sd}	[1, 3] days	To allow variation in time to diagnosis/reporting

Infection-detection rate	r	Starting from U[0.0001, 0.02] at time 0 and allowed to increase over time using space re-probing[5] with values drawn from U[0.015, 0.08] starting at week 8 (the week of 4/26/2020), U[0.02, 0.08] starting at week 15 (the week of 6/14/2020), U[0.01, 0.08] starting 1/17/2021 (around the end of the first pandemic wave), U[0.01, 0.05] starting 5/23/2021 (around the end of the Delta wave).	Large uncertainties; therefore, in general we use large prior bounds and large bounds for space re-probing (SR). Note that SR is only applied to 3-10% of the ensemble members and r can migrate outside either the initial range or the SR ranges during EAKF update. In India, due to the younger population age structure, infection detection rates were likely low throughout.
Infection fatality risk (IFR)		Starting from U[0.0001, 0.003] at time 0 and allowed to change over time using space re-probing[5] with values drawn from U[0.0001, 0.0015] starting at week 8 (the week of 4/26/2020), values drawn from U[0.0001, 0.0008] during the week of 7/05/2020 to the week of 3/21/2021 when case fatality risk was lower as computed from the data, values drawn from U[0.0001, 0.0015] starting the week of 4/4/2021 with the rise of Delta, values drawn from U[0.0001, 0.003] starting the week of 5/2/2021 when the healthcare systems began to be overwhelmed, and values drawn from U[0.0001, 0.0005] starting the week of 6/6/2021 when many have been infected or vaccinated and reported cases and deaths declined to low levels.	Based on previous estimates[6] but extend to have wider ranges. Note that SR is only applied to 3-10% of the ensemble members and IFR can migrate outside either the initial range or the SR ranges during EAKF update. In India, due to the younger population age structure, infection fatality risk was likely low throughout.
Vaccine efficacy (VE)		VE = 30% fourteen days after the 1st dose, and 67% seven days after the 2nd dose.	During 1/15/2021 – 6/27/2021, India used the Oxford/AstraZeneca and Covaxin vaccines; during 6/28/2021 – 10/31/2021,

Sputnik V was also used. Here we set the baseline VE based on data for the Oxford/AstraZeneca vaccine.[7]

References:

1.   Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *New Engl J Med* 2020.

2.   Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet* 2020; **395**(10225): 689-97.

3.   Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* 2020; **368**(6490): 489-93.

4.   Zhang J, Litvinova M, Wang W, et al. Evolving epidemiology and transmission dynamics of coronavirus disease 2019 outside Hubei province, China: a descriptive and modelling study. *The Lancet Infectious diseases* 2020; **20**(7): 793-802.

5.   Yang W, Shaman J. A simple modification for improving inference of non-linear dynamical systems. *arXiv* 2014: 1403.6804.

6.   Verity R, Okell LC, Dorigatti I, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *The Lancet Infectious diseases* 2020; **20**(6): 669-77.

7.   Bernal JL, Andrews N, Gower C, et al. Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant. *New England Journal of Medicine* 2021; **385**(7): 585-94.

Fig S1. Estimated infection-detection rate (A) and infection-fatality risk (B) during each week of the study period. For comparison, estimated weekly infection rates are superimposed in each plot (right y-axis). Blue lines and surrounding areas show model estimated mean, 50% and 95% CrIs. Boxes and whiskers show model-estimated weekly infection rates (mean, 50% and 95% CrIs). Grey shaded boxes indicate the timing of lockdowns (darker) or local restrictions (lighter); horizontal arrows indicate the timing of variant identification and vaccination rollout. *Note that infection-fatality risk estimates were based on reported COVID-19 deaths and may not reflect the true values due to the likely under-reporting of COVID-19 deaths.*

Fig S2. Estimated change in transmissibility and immune escape potential for the Delta variant, using different vaccine effectiveness (VE) settings for those with prior non-Delta infection. The baseline VE (i.e. 30%/67% for the 1st/2nd dose) was used for those without any prior infection; those infected by Delta were assumed to acquire immunity against this variant after recovery (see Eqn 1 in the main text). Twelve VE settings were tested (see x-axis label for VE after 1st/2nd dose). Model-inference was done 300 times (each with 500 ensemble members) for each VE setting. The boxplots summarize the distribution of estimates from the 300 runs: middle bar = median, edges = interquartile range, whiskers = 95% CI, red 'x' = mean.