Title

Genomic evidence for prisons as amplifiers of community tuberculosis epidemics

Short title: Prison amplifiers of tuberculosis epidemics

Authors

Katharine S. Walter PhD[1]*, Paulo César Pereira dos Santos BS[2], Thais Oliveira Gonçalves

BS[3], Brunna Oliveira da Silva BS[2], Andrea da Silva Santos PhD[2], Alessandra de Cássia Leite

MSc[2], Alessandra Moura da Silva BA[4], Flora Martinez Figueira Moreira MSc[2], Roberto Dias de

Oliveira PhD[5], Everton Ferreira Lemos PhD[4], Eunice Cunha PhD[3], Yiran E. Liu BS[1,6], Albert Ko

MD[7], Caroline Colijn PhD[8], Ted Cohen MD[7], Barun Mathema PhD[9], Julio Croda MD[4,7,10]†, Jason

R. Andrews MD[1]†

†These authors contributed equally to this work.

*Corresponding author. Email: kwalter@stanford.edu

Affiliations

[1]Division of Infectious Diseases and Geographic Medicine, Stanford University School of

Medicine; Stanford, United States.

[2] Health Sciences Research Laboratory, Federal University of Grande Dourados; Dourados,

Brazil.

[3] Laboratory of Bacteriology, Central Laboratory of Mato Grosso do Sul; Campo Grande, Brazil.

[4] School of Medicine, Federal University of Mato Grosso do Sul, School of Medicine; Campo

Grande, Brazil.

[5] Nursing Course, State University of Mato Grosso do Sul; Dourados, Brazil.

[6]Cancer Biology Graduate Program, Stanford University School of Medicine; Stanford, United

States.

[7]Department of Epidemiology of Microbial Diseases, Yale School of Public Health; New Haven,

United States.

[8]Department of Mathematics, Simon Fraser University; Burnaby, Canada.

[9]Department of Epidemiology, Columbia University Mailman School of Public Health; New

York, United States.

[10] Mato Grosso do Sul Office, Oswaldo Cruz Foundation; Campo Grande, Brazil.

Abstract

Prisons are high-incidence settings for tuberculosis around the world, yet the contribution of spillover from prisons in driving community epidemics has not been quantified. *We* whole genome sequenced 1,152 *M. tuberculosis* isolates from participants diagnosed with tuberculosis within prisons and in the community in *Mato Grosso do Sul, Brazil from 2014 to 2019*. By integrating timed phylogenies and detailed location data, we reconstructed probabilistic transmission histories. *M. tuberculosis* sequences from incarcerated and non-incarcerated people were closely phylogenetically related. We found that 57% of recent community-wide tuberculosis cases were attributable to transmission from individuals with an incarceration history, 2.6% of the population. Further, we find genomic evidence that the prison system disseminates *M. tuberculosis* genotypes through frequent transfers across the state. This population-wide genomic transmission reconstruction framework can be applied to identify key environments amplifying infectious disease transmission to prioritize public health interventions.

Teaser

Prison spillover plays an outsized role in community tuberculosis transmission.

MAIN TEXT

Introduction

Tuberculosis remains one of the leading causes of death by an infectious disease and led to 1.4 million deaths in 2019.(1) Locating where *M. tuberculosis* transmission occurs is critical for public health interventions, yet in high-incidence settings, where most tuberculosis is transmitted remains unknown.*(1–6)* Prisons, hospitals, or mines are known to be high-risk environments*(7–9)* and may have epidemiological impacts that transcend their walls, serving as institutional amplifiers of infection.*(10, 11)* Identifying institutional amplifiers of tuberculosis is of particular concern in Central and South America, where the incarcerated population has increased by over

200% since 2000 and 11% percent of tuberculosis cases now occur among incarcerated people,

who comprise <1% of the total population.*(12)*

The full impact of increasing incarceration rates on the tuberculosis epidemic in the

Americas and elsewhere has been difficult to quantify. Many people are infected in prison, but

only diagnosed later.*(11, 13, 14)* Further, spillover may occur when incarcerated people, prison

staff, or visitors are infected within high-incidence prison environments and transmit *M.*

tuberculosis onwards in communities outside prison. Previous studies have found that the excess

risk of tuberculosis within prisons extends to surrounding communities*(15, 16)* and that *M.*

tuberculosis genotypes responsible for jail and prison outbreaks are also found in the surrounding

communities.*(13, 15, 17, 18)*

Pathogen genomes can be powerfully harnessed to infer high-resolution transmission

histories, including who-infected-whom.*(19)* However, studies have not yet combined *M.*

tuberculosis genomic and epidemiologic data to reconstruct transmission linkages between

potential institutional amplifiers and the community, nor have they estimated the fraction of

community transmission attributable to such spillover. To quantify the role of prisons in

population-wide tuberculosis transmission, we conducted prospective genomic surveillance for

tuberculosis in the Brazilian state of Mato Grosso do Sul and used full *M. tuberculosis* genomes,

combined with detailed meta-data on incarceration history, to reconstruct tuberculosis

transmission chains spanning prisons and the community.

Results

Tuberculosis and incarceration trends in Mato Grosso do Sul

Brazil's incarcerated population grew by 107% (from 361,402 to 748,009) from 2005 to

2019, an increase closely paralleled in the Central West state of Mato Grosso do Sul, where the

incarcerated population has grown by 115% (7,891 to 16,976) over the same period (Fig. 1a-c). In

2019, prisons in the state were at 203% occupancy. While the state's notification rate of new and

retreatment tuberculosis in the general population (38 per 100,000) is similar to Brazil's national

tuberculosis notification rate (40 per 100,000) in 2019,[20] the notification rate within prisons

was 43 times as high (1,666 per 100,000), again similar to Brazil's national notification rate

within prisons (1,596 per 100,000) in 2019 (Fig. 1d,e).

Population-based genomic surveillance across Mato Grosso do Sul state

From 2014 to 2019, 3,491 new and retreatment cases of tuberculosis were reported in the

Campo Grande and Dourados, the two largest cities in Mato Grosso do Sul, Brazil. Of these,

1,249 had positive cultures and we sequenced 787 *M. tuberculosis* isolates. We sequenced

additional isolates from three other cities in the state and from earlier years in Dourados for a total

of 1,090 isolates from unique tuberculosis notifications (Fig. 1b). Whole genome sequences

(WGS) for 1,043 isolates met our coverage and quality criteria (Methods). We excluded 10%

(108/1043) of isolates with mixed infection, resulting in 935 high-quality genomes from 935

unique tuberculosis episodes from 918 individuals.

Of the 935 isolates in our analyses, 50% (465/935) were incarcerated; 16% (150/935)

were from patients who were formerly incarcerated; and 34% (320/935) of the study population

did not have an incarceration history at the time of tuberculosis notification. Among those who

did not have an incarceration history, 32 people reported contact with incarcerated individuals.

Additional population characteristics are in Table S1. Isolates were largely from *M. tuberculosis*

lineage 4 and predominantly fell into three sublineages: lineage 4.1, 4.3, and 4.4 (Fig. 2). Overall,

we found a low prevalence of antibiotic resistance across isolates, with 93.7% of isolates

susceptible to all antibiotics (Fig. 2).

Phylogenetic structure of *M. tuberculosis* from prisons and the community

A maximum likelihood phylogeny (Fig. 2) demonstrates that *M. tuberculosis* isolates

sampled from incarcerated and non-incarcerated people do not form distinct clades and are

closely phylogenetically related. Terminal branch lengths, a proxy for the degree of recent

transmission,[21] are significantly longer for isolates from people with no incarceration history at

the time of tuberculosis notification (9.4×10^{-4} substitutions/site) than isolates from incarcerated

individuals (2.2×10^{-4}; two-sample t-test p-value = 1.14×10^{-7}) or formerly incarcerated

individuals (3.8×10^{-4}; p = 1.6×10^{-4}), suggesting that individuals with an incarceration history

were more likely to be recently infected (Fig. S1).

Genetic clustering of *M. tuberculosis* from prisons and the community

To identify potential clusters of recent transmission, we applied a commonly used 12-SNP

threshold,[22] including all isolates within the threshold distance of at least one other clustered

isolate. Eighty-three percent (777/935) of the isolates fell into 84 genomic clusters (each

including 2 to 170 isolates), providing evidence that notifications were largely attributable to

recent, local transmission rather than travel-associated importation or re-activation of genetically

distinct latent infections (Fig. 2). Isolates from incarcerated people were more frequently

clustered (93.3%, 434/465), than those from formerly incarcerated (86.0%, 129/150; p <0.0001),

or never incarcerated people (66.9%, 214/320; p <0.0001), again suggesting more recent

transmission within prisons.

We predicted that if prison and community-associated epidemics were distinct, isolates

from the community would be most closely related to and cluster with other isolates from the

community and vice versa. Of the 46 clusters with three or more isolates, including 703

participants, 40 included isolates from community members who had no reported incarceration

history. Of those potential transmission clusters, 85.0% (34/40) included isolates from people who

were currently or previously incarcerated.

We found a similar pattern of clustering of isolates from community members and

individuals with an incarceration history when using an alternative 5-SNP threshold for genomic

clustering[23]. With this threshold, of the 80 clusters with three or more isolates, 56 included

isolates from people with no incarceration history and 78.6% (44/56) also included isolates from people with an incarceration history.

A *M. tuberculosis* clone spans prisons and the community across the state

The largest potential transmission cluster, including 170 isolates sampled from September 2010 to April 2019, was distributed across the state, including cases found across state's two largest cities, Campo Grande and Dourados, as well as the smaller cities Corumbá (on the Bolivian border) and Ponta Porã (on the Paraguayan border) (Fig. 1b, Fig. 3a). The cluster had a most recent common ancestor in 1996 (95% HPD: 1989-2003; Fig. 3a), indicating that the drug-susceptible clone had circulated for approximately 23 years before the most recent samples were collected. 103 isolates were from people incarcerated at the time of diagnosis, 35 from people formerly incarcerated, and 32 from people with no incarceration history at the time of tuberculosis notification.

As observed in other clusters, isolates from people who are currently and formerly incarcerated were closely genetically related—and often, identical—to isolates from people who were not incarcerated at the time of notification, suggesting they are linked through recent transmission (Fig. 3b).

We hypothesized that the criminal justice system's frequent transfer of people between prisons and jails could disseminate TB across the state. Brazil's national criminal law defines different "regimes" or stages of incarceration(30), which may facilitate the potential for spillover and dissemination of infection. People are incarcerated under closed regimes, within prisons; semi-open regimes, in which people may work outside prisons and return at night; and open regimes, under which people serve sentences outside of prison, but are required to make periodic court appearances.(30) We analyzed movements in the state's incarceration database and found that the average duration of incarceration was 230 days in a prison and 25 days in a jail. The incarceration database documents 7,982 mean yearly transfers between prisons (including closed

and semi-open prisons) from 2015-2018 in a prison population of 17,221 in August 2018,

including frequent transfers between cities across the state (Fig. 4a).

To investigate the role of the criminal justice system in disseminating infection, we

tracked the spread of the largest sampled *M. tuberculosis* cluster (Fig. 3) across prisons in Mato

Grosso do Sul (Fig. 4b). From early 2011 to 2014, sampled isolates from the cluster were

concentrated in the maximum security prison in Dourados, after which it was identified within

two prisons in Campo Grande and then exported to prisons and jails in other cities. The cluster

was not contained within the state's prison network; notified tuberculosis cases among

community members outside of prison occurred over the duration of the clusters' state-wide

spread.

Transmission trees reveal who-infected-whom, where, and when transmission occurred

We further investigated genomic clusters by integrating timed phylogenies with

epidemiological information to infer transmission trees, allowing us to reconstruct not only

transmission linkages of who-infected-whom, but also locate when and where transmission

occurred. Reconstructed transmission linkages revealed frequent transmission within prisons and

spillover from prisons to the community. For example, we inferred likely transmission (89%

posterior probability) between two individuals (Fig. 5a,b) and occurred between July 2014 and

July 2015 with 94% posterior probability (Fig. 5c). Both the recipient and predicted infector were

incarcerated in the same prison in Campo Grande at the time of the recipient's tuberculosis

notification and that transmission occurred within this prison with a posterior probability of 97%.

Spillover from prisons plays a disproportionate role in community transmission

To estimate the total contribution of different populations to community-wide tuberculosis

transmission, we inferred transmission trees for all clustered isolates. We identified the location of

both the infector and recipient over the distribution of likely infection times, allowing us to

determine location-specific transmission probabilities. Finally, we summed transmission

probabilities over all potential transmission pairs by the incarceration status of both the infector

and recipient at the time of transmission, giving a population-wide WAIFW matrix (Fig. 5d).

Because we conducted active surveillance inside prisons and not in the general population, we

corrected pairwise transmission probabilities to account for higher genomic sampling proportion

of incarcerated populations (Fig. S2, Methods).

The population-wide WAIFW matrix reveals that a small proportion of the state's

population, 0.84% (22,706 incarcerated people of a total population of 2,713,147 in 2017), were

associated with 18% of observed transmissions to people with no incarceration history (Fig. 5d).

When additionally including people who were previously incarcerated, 2.6% of the population

(71,703 of 2,713,147) were associated with 39% of the transmission to people with no

incarceration history (Fig. 5d), demonstrating the outsized impact of prisons on tuberculosis

transmission in the state. In total, we found that 57% of all infections captured in our study

(including incarcerated, formerly incarcerated and never incarcerated individuals), including 46%

of infections in the community (among formerly incarcerated and never incarcerated individuals),

were attributable to transmission from incarcerated or formerly incarcerated individuals, who

comprise a small fraction of the state's population.

Transmission inferences are robust to epidemiological priors

We conducted a sensitivity analysis to determine the effect of epidemiological priors

(Table S2) on transmission inferences for the largest genomic cluster (170 isolates). Transmission

probabilities inferred for pairs of individuals were closely correlated across generation time priors

(Pearson's r: 0.94 – 0.98, Fig. S3) and sampling time priors (Pearson's r: 0.84 – 0.98, Fig. S4),

although the total number of observed transmission events was strongly influenced by both priors

(Figure S5). This suggests that, as expected, epidemiological priors impact the magnitude of

inferred transmission probabilities rather than the identity of predicted transmission pairs. Further,

alternative priors did not strongly influence the population-wide WAIWF matrix for the largest

genomic cluster (Fig. S6). The fraction of transmission to those with no incarceration history

attributable to spillover (from infectors who are currently or previously incarcerated) in the largest

cluster ranges from 63.5% (medium sampling and generation times) to 72.3% (short generation

time and long sampling time).

Discussion

Here, we find genomic evidence that spillover from a key, high-risk environment can

drive tuberculosis transmission risk in the broader population. Controlling tuberculosis thus

urgently requires reducing risk created by institutional amplifiers, which have outsized effects on

the general population.*(11)* Our population-level transmission reconstruction framework can be

applied to identify other high-risk environments or populations and quantify their contribution to

community tuberculosis transmission, informing targeted public health interventions.*(11)*

The disproportionate role that prisons play in driving community tuberculosis

transmission is not unique to Brazil. Incidence of tuberculosis in prisons is extremely high

globally.*(7, 8)* The rapid growth in incarceration rates across Central and South America, in

particular, puts an increasing population at heightened risk of tuberculosis—risk that we have

found can extend into neighboring communities. Although the prevalence of antibiotic resistance

is low in Central West, Brazil, prisons have also amplified drug-resistant tuberculosis in

Russia*(24)* and elsewhere. Further, the role of prisons and other detention centers as disease

reservoirs is not unique to tuberculosis. Previous studies have identified spillover of

meningococcal disease*(25)* and SARS-CoV-2*(26)* from prisons to the community.

Our results emphasize that reducing the extraordinarily high transmission risk created by

prison environments is an urgent priority. Public health programs need to work to reduce the

extreme overcrowding within prisons; improve the unsanitary, often inhumane conditions of

incarceration; and expand access to primary healthcare and nutrition. Tuberculosis control

programs should expand routine active screenings during*(11, 27)* and following incarceration so

that cases can be diagnosed early and linked to treatment. Individuals leaving prison are at

heightened risk of tuberculosis and may be a particularly vulnerable population. Further,

preliminary evidence suggests that isoniazid protective therapy is protective against tuberculosis

infection and should be considered as a protective intervention in high-risk prison

environments.*(28)*

Reducing this excess burden of disease will require work that extends beyond biomedical

interventions. The most direct way to mitigate the excess tuberculosis risk created by prisons is

for governments to identify alternatives to detention so that incarceration is used only as a last

resort. Prison releases have already begun as a means of reducing infectious disease transmission

risk:*(29)* in an attempt to reduce the risk of COVID-19 in prisons, Brazil released more than thirty

thousand people early by mid-2020.*(30)*

Our study has several limitations. First, while we were able to enroll a majority of *M.*

tuberculosis isolates from incarcerated people with tuberculosis in the city of Dourados, overall

sampling of the state's ongoing tuberculosis epidemic was incomplete. To minimize sampling

bias, we determined genomic sampling probabilities and adjusted transmission probabilities to

account for over-sampling of prisons. Our probabilistic approach allows us to account for

uncertainty in transmission linkages; however, we still cannot infer the identity or incarceration

history of unsampled people who contribute to transmission. Terminal branch lengths may

similarly be affected by sampling bias or pathogen population size and are thus an imperfect

proxy of recency of transmission.

Second, transmission trees reflect information from timed *M. tuberculosis* phylogenies

and epidemiological priors. It is difficult to distinguish genomic sampling proportion from

generation time and sampling time distributions. Because we conducted transmission inference in

a tuberculosis-endemic setting, rather than in a discrete outbreak, our rate of genomic sampling

was lower than in some previous tuberculosis transmission studies. To reduce uncertainty in

sampling proportion, we conducted transmission inference on recent subtrees (MRCA > 2012) for

which we had sampled a greater proportion of isolates. To investigate the influence of

epidemiological priors on transmission inferences, we conducted a sensitivity analysis (Table S2)

and found that our conclusions were largely unchanged (Supplementary Text, Figs. S3-S6).

Third, we conducted whole genome sequencing of *M. tuberculosis* isolates cultured from

sputum and generated single consensus genomes from each participant. Excluding within-host

diversity could lead to some incorrect transmission inferences. For example, if we sampled an *M.*

tuberculosis genome that was genetically distant from the genome that was transmitted, we could

incorrectly exclude transmission.

Finally, we employed an existing bioinformatic approach to identify *M. tuberculosis*

variants and excluded repetitive genomic regions as is common practice.*(31, 32)* Future studies

that leverage within-host *M. tuberculosis* diversity and/or include the diverse PE/PPE genes could

provide greater resolution and reduce the uncertainty of transmission inferences.*(33)*

In this study, we present genomic evidence that prisons act as tuberculosis reservoirs in

Central West, Brazil. The dramatic expansion of incarceration in recent decades has put an

increasing population at extremely high risk of tuberculosis; this risk extends to surrounding

communities. Reducing the excess tuberculosis transmission risk within prisons and other

detention centers is an urgent public health priority. Our findings additionally highlight the

potential for pathogen genomic surveillance to identify other key environments or core

populations with a disproportionate role in pathogen transmission and guide public health

priorities.

Materials and Methods

Study design

We conducted population-based tuberculosis surveillance in the state of Mato Grosso do Sul in Central West Brazil from January 2014 through May 2019. Surveillance included active screening in three of the largest prisons in the state as well as ongoing passive surveillance focused on the two largest cities in the state, Campo Grande and Dourados, and three cities at the state's border with Paraguay and Bolivia (Fig. 1a,b; **Supplementary Methods**). All participants provided written consent, and this study was conducted with the approval of the Research Ethics Committee from the Federal University of Grande Dourados, Federal University of Mato Grosso do Sul and National Research Ethics Committee (CONEP) (CAAE 37237814.4.0000.5160, 2676613.3.1001.5160, and 26620619.6.0000.0021) and Stanford University Institutional Review Board (IRB-40285).

Incarceration history

To investigate the incarceration history of tuberculosis patients more closely, we obtained permission from the Mato Grosso do Sul state prison administration agency, Agência Estadual de Administração do Sistema Penitenciário, to access a database of all prison entries, exits, and transfers within the criminal justice record system, Sistema Integrado de Gestão Operacional, from January 1, 2005 through December 31, 2018. Brazil's national criminal law defines different "regimes" or stages of incarceration*(34)*, which may facilitate the potential for spillover of infection. People are incarcerated under closed regimes, within prisons; semi-open regimes, in which people may work outside prisons and return at night; and open regimes, under which people serve sentences outside of prison, but are required to make periodic court appearances.*(34)*

Whole genome sequencing

We sequenced whole genomes from cultured isolates on an Illumina NextSeq (2 x 151-bp). Sequence data is available on the Sequence Read Archive (SRA), in BioProject PRJNA671770. We applied variant calling methods closely following those described in Menardo

et al.*(31)* to be consistent with the methods used for molecular clock estimation (**Supplementary**

Methods).

Phylogenetic and Bayesian evolutionary analysis

We fit maximum likelihood trees with RAxML-ng 1.0.1.*(35)* For each cluster including

three or more isolates, we fit Bayesian trees with BEAST 2.6.2*(36)* with a strict clock, constant

coalescent population size model using tuberculosis notification dates to calibrate tips

(**Supplementary Methods**).

Transmission inference

M. tuberculosis phylogenetic trees represent patterns of evolutionary relatedness between

the consensus bacterial genomes sampled from different individuals. Because most outbreaks are

incompletely sampled (i.e. an *M. tuberculosis* sequence is not available for every case) and

individuals may be infected with diverse populations of *M. tuberculosis*, phylogenies do not

represent the underlying history of transmission.*(19)* We used TransPhylo*(19)* to infer

transmission linkages—including unsampled hosts—that were consistent with the underlying

timed phylogenies and transmission process (i.e. generation time and sampling intervals). To

focus on densely sampled trees that would not be dominated by unsampled hosts, we sliced

phylogenies at 2012 in order to generate subtrees with a most recent common ancestor of 2012 or

later for transmission inference, including 122 subtrees each with 2 to 23 tips and comprising

56% of isolates (528/935).

By sampling from the posterior distribution of transmission trees, we estimated the

posterior probability that each individual within a transmission tree was infected by each other

individual in the tree. We summarized the posterior set of transmission trees in TransPhylo with

who acquired infection from whom (WAIFW) matrices, W, where $W_{i,j}$ is the posterior probability

of transmission from individual j to individual i. Many transmission events are unobserved

because of incomplete sampling; therefore, the sum of transmission probabilities to any individual

is often less than one. We combined transmission trees with incarceration histories to quantify the

proportion of observed transmission attributable to people with a history of incarceration. An

individual's location at the time of their tuberculosis notification may not reflect where

transmission occurred. To understand where infector-recipient pairs were at the time transmission

occurred, we integrated transmission trees with incarceration information (Fig. 4c). First, we used

the transmission trees to infer the posterior distribution of infection times for each recipient i. We

then partitioned the infector-recipient transmission probability $W_{i,j}$ across the recipient's infection

time distribution, resulting in W_{i_u,j_v}, the probability of transmission from individual j to

individual i while j has incarceration status v and i has incarceration status u. Incarceration status

was trichotomized as incarcerated, formerly incarcerated, and no incarceration history.

We then summed pairwise, location-specific transmission probabilities over all individual

subtrees with respect to the incarceration status of both infector j and recipient i. We defined the

population wide transmission matrix, $T_{u,v} = \sum_i \sum_j W_{i_u,j_v}$, where $T_{u,v}$ is the sum of the posterior

probabilities of observed transmission events to individuals in population u from individuals in

population v. We normalized $T_{u,v}$ by dividing by the sum of posterior transmission probabilities

to each recipient population u, $N_u = \sum_v T_{u,v}$. We then defined the uncorrected proportion of the

total posterior transmission probabilities to population u from population v, or the transmission

fraction, as $p_{u,v} = \frac{T_{u,v}}{N_u}$.

Genomic sampling varied across populations and specifically, we predicted that

incarcerated people might have been sampled at a higher rate due to our active case finding and

greater culture coverage in three prisons in the state. To correct for sampling bias, we determined

the genomic sampling proportion r_k, which we defined as the proportion of notified tuberculosis

cases for which a high-quality genomic sequence was available, for population k at the time of

tuberculosis notification. We calculated genomic sampling proportion as the number of sequenced

genomes for each group divided by the number of tuberculosis notifications for each group from

2014-2019 in Campo Grande and Dourados, the two major cities in our prospective study (Fig.

S2). We adjusted each pairwise transmission probability by the sampling rate of the infector,

based on the incarceration status of the infector at the time of tuberculosis notification, for $j \in$

$k: W'_{i,j} = W_{i,j} \frac{1}{r_k}$ and calculated the sampling-adjusted transmission fractions as described above.

Role of the funding source

The funding source had no role in study design, data collection, data analysis, data interpretation,

or writing of the report.

Statistical methods

We inferred transmission trees with TransPhylo*(19)* as described above. We compared

phylogenetic tree terminal branch lengths between populations with two-sample t-tests and

compared the rate of clustering between populations with two-sample proportion tests.

References

1. G. Churchyard, P. Kim, N. S. Shah, R. Rustomjee, N. Gandhi, B. Mathema, D. Dowdy, A.

Kasmar, V. Cardenas, What We Know about Tuberculosis Transmission: An Overview, *J. Infect.*

Dis. **216**, S629–S635 (2017).

2. S. C. Auld, N. Sarita Shah, B. Mathema, T. S. Brown, N. Ismail, S. V. Omar, J. C. M. Brust, K.

N. Nelson, S. Allana, A. Campbell, K. Mlisana, P. Moodley, N. R. Gandhi, Extensively drug-

resistant tuberculosis in South Africa: Genomic evidence supporting transmission in communities,

Eur. Respir. J. **52** (2018), doi:10.1183/13993003.00246-2018.

3. K. Middelkoop, B. Mathema, L. Myer, E. Shashkina, A. Whitelaw, G. Kaplan, B. Kreiswirth,

R. Wood, L. G. Bekker, Transmission of tuberculosis in a south African community with a high

prevalence of HIV infection, *J. Infect. Dis.* **211**, 53–61 (2015).

4. J. R. Andrews, C. Morrow, R. P. Walensky, R. Wood, Integrating social contact and

environmental data in evaluating tuberculosis transmission in a South African township., *J. Infect.*

Dis. **210**, 597–603 (2014).

5. T. A. Yates, P. Y. Khan, G. M. Knight, J. G. Taylor, T. D. McHugh, M. Lipman, R. G. White,

T. Cohen, F. G. Cobelens, R. Wood, D. A. J. Moore, I. Abubakar, The transmission of

Mycobacterium tuberculosis in high burden settings, *Lancet Infect. Dis.* **16**, 227–238 (2016).

6. J. R. Andrews, C. Morrow, R. Wood, Modeling the role of public transportation in sustaining

tuberculosis transmission in South Africa, *Am. J. Epidemiol.* **177**, 556–561 (2013).

7. K. Dolan, A. L. Wirtz, B. Moazen, M. Ndeffo-mbah, A. Galvani, S. A. Kinner, R. Courtney,

M. McKee, J. J. Amon, L. Maher, M. Hellard, C. Beyrer, F. L. Altice, Global burden of HIV,

viral hepatitis, and tuberculosis in prisoners and detainees, *Lancet* **388**, 1089–1102 (2016).

8. O. Cords, L. Martinez, J. Warren, J. O'Marr, K. S. Walter, T. Cohen, J. Zheng, A. I. Ko, J.

Croda, J. R. Andrews, Incidence and Prevalence of Tuberculosis in Incarcerated Populations: A

Systematic Review and Meta-analysis, *Lancet Public Heal.* .

9. J. Urrego, A. I. Ko, A. Da Silva Santos Carbone, D. S. G. Paião, R. V. E. Sgarbi, C. W. Yeckel,

J. R. Andrews, J. Croda, The impact of ventilation and early diagnosis on tuberculosis

transmission in Brazilian prisons, *Am. J. Trop. Med. Hyg.* **93**, 739–746 (2015).

10. S. Basu, D. Stuckler, M. McKee, Addressing institutional amplifiers in the dynamics and

control of tuberculosis epidemics, *Am. J. Trop. Med. Hyg.* **84**, 30–37 (2011).

11. T. S. Mabud, M. de Lourdes Delgado Alves, A. I. Ko, S. Basu, K. S. Walter, T. Cohen, B.

Mathema, C. Colijn, E. Lemos, J. Croda, J. R. Andrews, J. Z. Metcalfe, Ed. Evaluating strategies

for control of tuberculosis in prisons and prevention of spillover into communities: An

observational and modeling study from Brazil, *PLoS Med.* **16**, e1002737 (2019).

12. K. Walter, L. Martinez, Arakaki-Sanchez, V. Sequera, S. G. Estigarribia, T. Cohen, A. Ko, A.

García-Basteiro, Z. Rueda, O. R. Lopez, M. Espinal, J. Croda, J. Andrews, The escalating

tuberculosis crisis in Central and South American prisons. In press., *Lancet* (2020),

doi:10.1016/S0140-6736(20)32578-2.

13. F. P. C. Sacchi, R. M. Praça, M. B. Tatara, V. Simonsen, L. Ferrazoli, M. G. Croda, P. N.

Suffys, A. I. Ko, J. R. Andrews, J. Croda, Prisons as reservoir for community transmission of

tuberculosis, brazil, *Emerg. Infect. Dis.* **21**, 452–455 (2015).

14. R. Coker, M. McKee, R. Atun, B. Dimitrova, E. Dodonova, S. Kuznetsov, F. Drobniewski,

Risk factors for pulmonary tuberculosis in Russia: Case-control study, *Br. Med. J.* **332**, 85–87

(2006).

15. J. L. Warren, L. Grandjean, D. A. J. Moore, A. Lithgow, J. Coronel, P. Sheen, J. L. Zelner, J.

R. Andrews, T. Cohen, Investigating spillover of multidrug-resistant tuberculosis from a prison: a

spatial and molecular epidemiological analysis, *BMC Med.* **16**, 122 (2018).

16. H. E. Jenkins, V. Plesca, A. Ciobanu, V. Crudu, I. Galusca, V. Soltan, A. Serbulenco, M.

Zignol, A. Dadu, M. Dara, T. Cohen, Assessing spatial heterogeneity of multidrug-resistant

tuberculosis in a high-burden country, *Eur. Respir. J.* **42**, 1291–1301 (2013).

17. M. N. Séraphin, X. Didelot, D. J. Nolan, J. R. May, M. S. R. Khan, E. R. Murray, M. Salemi,

J. G. Morris, M. Lauzardo, Genomic investigation of a Mycobacterium tuberculosis outbreak

involving prison and community cases in Florida, United States, *Am. J. Trop. Med. Hyg.* **99**, 867–

874 (2018).

18. T. F. Jones, C. L. Woodley, F. F. Fountain, W. Schaffner, Increased incidence of the outbreak

strain of Mycobacterium tuberculosis in the surrounding community after an outbreak in a jail.

(Original Article), *South. Med. J.* **96**, 155–158 (2003).

19. X. Didelot, C. Fraser, J. Gardy, C. Colijn, H. Malik, Genomic infectious disease epidemiology

in partially sampled and ongoing outbreaks, *Mol. Biol. Evol.* **34**, 997–1007 (2017).

20. World Health Organization, *Global tuberculosis report 2020* (Geneva, 2020).

21. K. E. Holt, P. Mcadam, P. Vuong, K. Thai, N. Thuy, T. Thuong, D. Thi, M. Ha, N. N. Lan, N.

H. Lan, N. Thi, Q. Nhu, H. T. Hai, V. Thi, N. Ha, G. Thwaites, D. J. Edwards, A. P. Nath, K.

Pham, D. B. Ascher, J. Farrar, C. C. Khor, Y. Y. Teo, M. Inouye, Frequent transmission of the

Mycobacterium tuberculosis Beijing lineage and positive selection for the EsxW Beijing variant

in Vietnam, *Nat. Genet.* (2018), doi:10.1038/s41588-018-0117-9.

22. T. M. Walker, T. A. Kohl, S. V. Omar, J. Hedge, C. Del Ojo Elias, P. Bradley, Z. Iqbal, S.

Feuerriegel, K. E. Niehaus, D. J. Wilson, D. A. Clifton, G. Kapatai, C. L. C. Ip, R. Bowden, F. A.

Drobniewski, C. Allix-Béguec, C. Gaudin, J. Parkhill, R. Diel, P. Supply, D. W. Crook, E. G.

Smith, A. S. Walker, N. Ismail, S. Niemann, T. E. A. Peto, J. Davies, C. Crichton, M. Acharya, L.

Madrid-Marquez, D. Eyre, D. Wyllie, T. Golubchik, M. Munang, Whole-genome sequencing for

prediction of Mycobacterium tuberculosis drug susceptibility and resistance: A retrospective

cohort study, *Lancet Infect. Dis.* **15**, 1193–1202 (2015).

23. T. M. Walker, C. L. C. Ip, R. H. Harrell, J. T. Evans, G. Kapatai, M. J. Dedicoat, D. W. Eyre,

D. J. Wilson, P. M. Hawkey, D. W. Crook, J. Parkhill, D. Harris, A. S. Walker, R. Bowden, P.

Monk, E. G. Smith, T. E. A. Peto, Whole-genome sequencing to delineate Mycobacterium

tuberculosis outbreaks: A retrospective observational study, *Lancet Infect. Dis.* **13**, 137–146

(2013).

24. F. Drobniewski, Y. Balabanova, V. Nikolayevsky, M. Ruddy, S. Kuznetzov, S. Zakharova, A.

Melentyev, I. Fedorin, Drug-resistant tuberculosis, clinical virulence, and the dominance of the

Beijing strain family in Russia, *J. Am. Med. Assoc.* **293**, 2726–2731 (2005).

25. J. W. Tappero, R. Reporter, J. D. Wenger, B. A. Ward, M. W. Reeves, T. S. Missbach, B. D.

Plikaytis, L. Mascola, A. Schuchat, Meningococcal Disease in Los Angeles County, California,

and among Men in the County Jails, *N. Engl. J. Med.* **335**, 833–841 (1996).

26. E. Reinhart, D. L. Chen, Incarceration And Its Disseminations: COVID-19 Pandemic Lessons

From Chicago's Cook County Jail, *Health Aff.* **39**, 1412–1418 (2020).

27. A. da S. Santos, R. D. de Oliveira, E. F. Lemos, F. Lima, T. Cohen, O. Cords, L. Martinez, C.

Gonçalves, A. I. Ko, J. R. Andrews, J. Croda, Yield, Efficiency, and Costs of Mass Screening

Algorithms for Tuberculosis in Brazilian Prisons, *Clin. Infect. Dis.* (2020),

doi:10.1093/cid/ciaa135.

28. R. D. de Oliveira, A. da Silva Santos, C. B. Reis, A. de Cássia Leite, F. P. C. Sacchi, R. C. P.

de Araujo, P. C. P. dos Santos, V. C. Rolla, L. Martinez, J. Andrews, J. Croda, Primary

prophylaxis to prevent tuberculosis infection in prison inmates: A randomized, double-blind,

placebo-controlled trial, *Am. J. Trop. Med. Hyg.* **103**, 1466–1472 (2020).

29. E. A. Wang, B. Western, D. M. Berwick, COVID-19, Decarceration, and the Role of

Clinicians, Health Systems, and Payers: A Report From the National Academy of Sciences,

Engineering, and Medicine., *Jama* **06511** (2020), doi:10.1001/jama.2020.22109.

30. R. De Oliveira Andrade, Covid-19: Prisons exposed in Brazil's crisis*BMJ* **370** (2020),

doi:10.1136/bmj.m2884.

31. F. Menardo, S. Duchêne, D. Brites, S. Gagneux, The molecular clock of mycobacterium

tuberculosis, *PLoS Pathog.* **15** (2019), doi:10.1371/journal.ppat.1008067.

32. C. J. Meehan, G. A. Goig, T. A. Kohl, L. Verboven, A. Dippenaar, M. Ezewudo, M. R.

Farhat, J. L. Guthrie, K. Laukens, P. Miotto, B. Ofori-Anyinam, V. Dreyer, P. Supply, A. Suresh,

C. Utpatel, D. van Soolingen, Y. Zhou, P. M. Ashton, D. Brites, A. M. Cabibbe, B. C. de Jong, M.

de Vos, F. Menardo, S. Gagneux, Q. Gao, T. H. Heupink, Q. Liu, C. Loiseau, L. Rigouts, T. C.

Rodwell, E. Tagliani, T. M. Walker, R. M. Warren, Y. Zhao, M. Zignol, M. Schito, J. Gardy, D.

M. Cirillo, S. Niemann, I. Comas, A. Van Rie, Whole genome sequencing of Mycobacterium

tuberculosis: current standards and open issues, *Nat. Rev. Microbiol.* **17**, 533–545 (2019).

33. K. S. Walter, C. Colijn, T. Cohen, B. Mathema, Q. Liu, J. Bowers, D. M. Engelthaler, A.

Narechania, D. Lemmer, J. Croda, J. R. Andrews, Genomic variant-identification methods may

alter mycobacterium tuberculosis transmission inferences, *Microb. Genomics* **6**, 1–16 (2020).

34. J. Mariner, J. Cavallaro, *Behind Bars in Brazil* (Human Rights Watch, New York, 1998).

35. A. M. Kozlov, D. Darriba, T. Flouri, B. Morel, A. Stamatakis, J. Wren, Ed. RAxML-NG: A

fast, scalable, and user-friendly tool for maximum likelihood phylogenetic inference,

Bioinformatics (2019), doi:10.1093/bioinformatics/btz305.

36. R. Bouckaert, T. G. Vaughan, J. Barido-Sottani, S. Duchêne, M. Fourment, A. Gavryushkina,

J. Heled, G. Jones, D. Kühnert, N. De Maio, M. Matschiner, F. K. Mendes, N. F. Müller, H. A.

Ogilvie, L. Du Plessis, A. Popinga, A. Rambaut, D. Rasmussen, I. Siveroni, M. A. Suchard, C. H.

Wu, D. Xie, C. Zhang, T. Stadler, A. J. Drummond, BEAST 2.5: An advanced software platform

for Bayesian evolutionary analysis, *PLoS Comput. Biol.* **15**, e1006650 (2019).

37. A. da S. Santos, R. D. de Oliveira, E. F. Lemos, F. Lima, T. Cohen, O. Cords, L. Martinez, C.

Gonçalves, A. Ko, J. R. Andrews, J. Croda, Yield, Efficiency and Costs of Mass Screening

Algorithms for Tuberculosis in Brazilian Prisons., *Clin. Infect. Dis.* (2020),

doi:10.1093/cid/ciaa135.

38. D. Robinson, fuzzyjoin: Join Tables Together on Inexact Matching (2020) (available at

https://github.com/dgrtwo/fuzzyjoin).

39. F. Krueger, Trim Galore (2019) (available at

https://github.com/FelixKrueger/TrimGalore/blob/master/Docs/Trim_Galore_User_Guide.md).

40. M. Martin, Cutadapt Removes Adapter Sequences From High-Throughput Sequencing Reads,

EMBnet.journal **17** (2011), doi:https://doi.org/10.14806/ej.17.1.200.

41. D. E. Wood, S. L. Salzberg, Kraken: Ultrafast metagenomic sequence classification using

exact alignments, *Genome Biol.* **15**, R46 (2014).

42. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler transform,

Bioinformatics **25**, 1754–1760 (2009).

43. D. Brites, C. Loiseau, F. Menardo, S. Borrell, M. B. Boniotti, R. Warren, A. Dippenaar, S. D.

C. Parsons, C. Beisel, M. A. Behr, J. A. Fyfe, M. Coscolla, S. Gagneux, A new phylogenetic

framework for the animal-adapted mycobacterium tuberculosis complex, *Front. Microbiol.* **9**,

2820 (2018).

44. C. Anyansi, A. Keo, B. J. Walker, T. J. Straub, A. L. Manson, A. M. Earl, T. Abeel,

QuantTB- A method to classify mixed Mycobacterium tuberculosis infections within whole

genome sequencing data, *BMC Genomics* **21**, 80 (2020).

45. A. J. Page, S. R. Harris, T. Seemann, B. Taylor, A. J. Delaney, J. A. Keane, J. Soares, SNP-

sites: rapid efficient extraction of SNPs from multi-FASTA alignments, *Microb. Genomics* **2**, 1–5

(2016).

46. P. Bradley, N. C. Gordon, T. M. Walker, L. Dunn, S. Heys, B. Huang, S. Earle, L. J.

Pankhurst, L. Anson, M. De Cesare, P. Piazza, A. A. Votintseva, T. Golubchik, D. J. Wilson, D.

H. Wyllie, R. Diel, S. Niemann, S. Feuerriegel, T. A. Kohl, N. Ismail, S. V Omar, E. G. Smith, D.

Buck, G. McVean, A. S. Walker, T. E. A. Peto, D. W. Crook, Z. Iqbal, Rapid antibiotic-resistance

predictions from genome sequence data for Staphylococcus aureus and Mycobacterium

tuberculosis, *Nat. Commun.* **6** (2015), doi:10.1038/ncomms10063.

47. C. Allix-Béguec, I. Arandjelovic, L. Bi, P. Beckert, M. Bonnet, P. Bradley, A. M. Cabibbe, I.

Cancino-Muñoz, M. J. Caulfield, A. Chaiprasert, D. M. Cirillo, D. A. Clifton, I. Comas, D. W.

Crook, M. R. De Filippo, H. de Neeling, R. Diel, F. A. Drobniewski, K. Faksri, M. R. Farhat, J.

Fleming, P. Fowler, T. A. Fowler, Q. Gao, J. Gardy, D. Gascoyne-Binzi, A. L. Gibertoni-Cruz, A.

Gil-Brusola, T. Golubchik, X. Gonzalo, L. Grandjean, G. He, J. L. Guthrie, S. Hoosdally, M.

Hunt, Z. Iqbal, N. Ismail, J. Johnston, F. M. Khanzada, C. C. Khor, T. A. Kohl, C. Kong, S.

Lipworth, Q. Liu, G. Maphalala, E. Martinez, V. Mathys, M. Merker, P. Miotto, N. Mistry, D. A.

J. Moore, M. Murray, S. Niemann, R. T. H. Ong, T. E. A. Peto, J. E. Posey, T. Prammananan, A.

Pym, C. Rodrigues, M. Rodrigues, T. Rodwell, G. M. Rossolini, E. S. Padilla, M. Schito, X.

Shen, J. Shendure, V. Sintchenko, A. Sloutsky, E. G. Smith, M. Snyder, K. Soetaert, A. M.

Starks, P. Supply, P. Suriyapol, S. Tahseen, P. Tang, Y. Y. Teo, T. N. T. Thuong, G. Thwaites, E.

Tortoli, S. V. Omar, D. van Soolingen, A. S. Walker, T. M. Walker, M. Wilcox, D. J. Wilson, D.

Wyllie, Y. Yang, H. Zhang, Y. Zhao, B. Zhu, Prediction of susceptibility to first-line tuberculosis

drugs by DNA sequencing, *N. Engl. J. Med.* **379**, 1403–1415 (2018).

48. J. E. Phelan, D. M. O'Sullivan, D. Machado, J. Ramos, Y. E. A. Oppong, S. Campino, J.

O'Grady, R. McNerney, M. L. Hibberd, M. Viveiros, J. F. Huggett, T. G. Clark, Integrating

informatics tools and portable sequencing technology for rapid detection of resistance to anti-

tuberculous drugs, *Genome Med.* **11**, 41 (2019).

49. E. Paradis, K. Schliep, R. Schwartz, Ed. ape 5.0: an environment for modern phylogenetics

and evolutionary analyses in R, *Bioinformatics* **35**, 526–528 (2019).

50. A. D. Leaché, B. L. Banbury, J. Felsenstein, A. Nieto-Montes De Oca, A. Stamatakis, New

Acquisition Bias Corrections for Inferring SNP Phylogenies, *Syst. Biol* **64**, 1032–1047 (2015).

51. A. Rambaut, A. J. Drummond, D. Xie, G. Baele, M. A. Suchard, Posterior summarization in

Bayesian phylogenetics using Tracer 1.7, *Syst. Biol.* **67**, 901–904 (2018).

52. A. Rieux, C. E. Khatchikian, tipdatingbeast: an r package to assist the implementation of

phylogenetic tip-dating tests using beast, *Mol. Ecol. Resour.* **17**, 608–613 (2017).

53. Y. I. Xu, I. Cancino-Muñoz, M. I. Torres-Puente, L. M. Villamayor, R. I. Borrá, M. Borrá s-

Má ñez, M. Bosque, J. J. Camarena ID, E. Colomer-Roig, J. I. Colomina, I. Escribano, O. I.

Esparcia-Rodríguez, A. I. Gil-Brusola, R. Moreno-Muñoz, D. NavarroID, M. Navarro, N. Orta, E.

Pé rez, J. Prat, J. I. Carlos Rodríguez, M. Montserrat Ruiz-García, H. Vanaclocha, C. ColijnID, I.

I. Comas, I. Cancino-Munoz, M. I. Torres-Puente, L. M. Villamayor, R. Borrás, M. Borrás-

Máñez, M. Bosque, J. J. Camarena, E. Colomer-Roig, J. I. Colomina, I. Escribano, O. I. Esparcia-

Rodríguez, A. I. Gil-Brusola, C. Gimeno, A. Gimeno-Gascón, B. Gomila-Sard, D. González-

Granda, N. Gonzalo-Jiménez, M. R. Guna-Serrano, J. L. López-Hontangas, C. Martín-González,

R. Moreno-Munoz, D. Navarro, M. Navarro, N. Orta, E. Pérez, J. Prat, J. C. Rodríguez, M. M.

Ruiz-García, H. Vanaclocha, C. Colijn, I. I. Comas, High-resolution mapping of tuberculosis

transmission: Whole genome sequencing and phylogenetic modelling of a cohort from Valencia

Region, Spain, *PLoS Med.* **16** (2019), doi:10.1371/journal.pmed.1002961.

54. D. Ayabina, J. O. Ronning, K. Alfsnes, N. Debech, O. B. Brynildsrud, T. Arnesen, G.

Norheim, A.-T. T. Mengshoel, R. Rykkvin, U. R. Dahle, C. Colijn, V. Eldholm, Genome-based

transmission modelling separates imported tuberculosis from recent transmission within an

immigrant population, *Microb. Genomics* **4**, 1–13 (2018).

Acknowledgments

We would like to acknowledge the contributions of AGEPEN, LACEN, and the Mato

Grosso do Sul State Health Department. All authors had full access to all the data in the

study and had final responsibility for the decision to submit

Funding:

National Institutes of Health grant R01AI130058 (JRA)

Brazil's National Council for Scientific and Technological Development

grant 404237/2012-6 (JC)

Author contributions:

Conceptualization: JRA, JC, KSW, CC, TC, BM, AIK

Methodology: JRA, JC, KSW, CC, TC, BM, AIK

Investigation: PCPS, TOG, BOS, ASS, ACL, AMS, FMFM, RDO, EFL, EC, YL

Visualization: KSW, JRA

Funding acquisition: JRA, JC

Project administration: JRA, JC

Supervision: JRA, JC

Writing – original draft: KSW, JRA

Writing – review & editing: all authors

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: Raw Illumina sequence data are available on the

Sequence Read Archive under BioProject PRJNA671770.

Figures and Tables

Figure 1. Tuberculosis is increasingly concentrated within prisons in Mato Grosso do Sul,

Brazil. (a) Map of Brazil with states colored by the 2019 incarceration rate with Mato Grosso do

Sul outlined in black. (b) Map of Mato Grosso do Sul state with the cities with passive

surveillance only and where active surveillance was additionally conducted within prisons. (c)

The incarcerated population size in Mato Grosso do Sul grew by 142% from 2005 to 2020,

increasing from 7,891 to 19,065 people. (d) Mato Grosso do Sul state's annual new and

retreatment tuberculosis notifications, colored by incarceration status and (e) the notification rate

per 100,000 people for the incarcerated population and non-incarcerated populations from 2009 to

2019. The y-axis is log-scaled. The mean tuberculosis notification rate was more than 46 times

greater among the incarcerated population compared to the non-incarcerated population. The

green and yellow bars (d) and (e) indicate the period of active surveillance in prisons in Campo

Grande and Dourados respectively.

Figure 2. *M. tuberculosis* isolates from incarcerated and non-incarcerated people are closely

related in Mato Grosso do Sul, Brazil. A maximum likelihood phylogeny of 932 tuberculosis

isolates from Lineage 4 inferred from a multiple sequence alignment of 19,753 SNPs. Branch

lengths are in units of substitutions per site. From the inside, rings are colored by incarceration

status, antimicrobial resistance prediction, and genomic cluster (Methods). "Other" resistance

indicates resistance to at least one antibiotic other than isoniazid or rifampicin. Branches are

colored by sublineage. The clade containing the largest predicted genomic cluster (Cluster 5),

including 170 isolates, is highlighted in purple.

Figure 3. A single, recently emerged *M. tuberculosis* clone spans prisons and the community

across the major cities Mato Grosso do Sul state. (a) A Bayesian time-calibrated phylogeny of

the largest sampled genomic cluster of 170 isolates. The cluster emerged approximately 23 years

before the most recently sampled isolate, with a most recent common ancestor in 1996 (95%

HPD: 1989-2003). Tip point color indicates patient's incarceration status at the time of

tuberculosis notification. Annotation bar colors indicate city. Grey error bars indicate the 95%

Bayesian highest posterior density intervals for node date. Clade posterior support values are

shown on the middle of branches for clades with posterior support > 0.5. (b) A haplotype network

of the single largest genomic cluster, including 170 isolates. Nodes represent unique haplotypes

and are scaled to size. Points along branches indicate SNP distances between haplotypes. Node

color indicates incarceration status at the time of diagnosis.

Figure 4. Prisons disseminate tuberculosis through the frequent movement of incarcerated

people between prisons. (a) Incarceration locates people within highly connected contact

networks. Map of Mato Grosso do Sul state with orange points indicating the 21 prisons which

most frequently transfer people between prisons. Edges are colored by the mean number of yearly

transfers made between prisons from 2015-2018, including transfers in both directions. Edges

were drawn between prison pairs with more than 10 transfers in the study period. Transfers

between prisons within a city are not shown. (b) An *M. tuberculosis* clone (depicted in Figure 3)

spreads throughout multiple prisons, jails, and the community. Each line represents an individual

patient infected with an isolate in the largest genomic cluster, ordered by notification date. Line

color indicates patient location, determined by matching patient names with the state incarceration

database (Methods). Points indicate TB notification dates; point color indicates patient location at

the time of TB notification. Individuals with notification dates after December 31, 2018 are not

shown.

Figure 5. Frequent transmission within prisons and spillover of infection from prisons

perpetuates the tuberculosis epidemic in Mato Grosso do Sul, Brazil. We integrated *M.*

tuberculosis phylogenetic trees with epidemiologic data to infer probable transmission linkages

among sampled patients as demonstrated by an example cluster (a-c). (a) Transmission histories

can be represented by a phylogenetic trees with branches colored by host in which tips represent

sampled isolates and transmission events are indicated by asterisks. (b) We summarized sampled

transmission trees into who-acquired-infection-from-whom matrices in which rows represent

individual recipients, columns represent individual infectors, and each cell is colored by pairwise

transmission probability. (c) The posterior distribution of infection times (grey histogram) for

individual d with bars below representing location information for individual d and individual c,

the predicted infector. Bars are colored by location over time and circles indicate date of

tuberculosis notification. (d) The population-wide WAIFW matrix describing the proportions of

observed transmissions to each recipient population from each infector population, corrected for

genomic sampling (Methods). Rows represent recipient populations, columns represent infector

populations, and cells indicate the proportion of total infections to each recipient population from

each infector population defined by their incarceration status at the time of transmission and

corrected for sampling rate.

654
655

0.003

a   b   c