Full Title: The role of prisons in disseminating tuberculosis in Brazil: a genomic epidemiology study

Short Title: Prison dissemination of tuberculosis

Authors:

Katharine S. Walter PhD[1], Paulo César Pereira dos Santos BS[2], Thais Oliveira Gonçalves BS[3], Brunna

Oliveira da Silva BS[2], Andrea da Silva Santos PhD[2], Alessandra de Cássia Leite MSc[2], Alessandra Moura

da Silva BA[4], Flora Martinez Figueira Moreira MSc[2], Roberto Dias de Oliveira PhD[5], Everton Ferreira

Lemos PhD[4], Eunice Cunha PhD[3], Yiran E. Liu BS[1,6], Albert Ko MD[7], Caroline Colijn PhD[8], Ted Cohen

MD[7], Barun Mathema PhD[9], Julio Croda MD[4,7,10]†, Jason R. Andrews MD[1]†

†These authors contributed equally to this work.

[1]Division of Infectious Diseases and Geographic Medicine, Stanford University School of Medicine;

Stanford, United States.

[2] Health Sciences Research Laboratory, Federal University of Grande Dourados; Dourados, Brazil.

[3] Laboratory of Bacteriology, Central Laboratory of Mato Grosso do Sul; Campo Grande, Brazil.

[4] School of Medicine, Federal University of Mato Grosso do Sul, School of Medicine; Campo Grande,

Brazil.

[5] Nursing Course, State University of Mato Grosso do Sul; Dourados, Brazil.

[6]Cancer Biology Graduate Program, Stanford University School of Medicine; Stanford, United States.

[7]Department of Epidemiology of Microbial Diseases, Yale School of Public Health; New Haven, United

States.

[8]Department of Mathematics, Simon Fraser University; Burnaby, Canada.

[9]Department of Epidemiology, Columbia University Mailman School of Public Health; New York,

United States.

[10] Mato Grosso do Sul Office, Oswaldo Cruz Foundation; Campo Grande, Brazil.

Corresponding Author: kwalter@stanford.edu (KSW)

Word count: 4585

Abstract

Background: Globally, prisons are high-incidence settings for tuberculosis. Yet the role of prisons as

reservoirs of *M. tuberculosis*, propagating epidemics through spillover from prisons to surrounding

communities, has been difficult to measure directly.

Methods and Findings: To quantify the role of prisons in driving wider community *M. tuberculosis*

transmission, we conducted prospective genomic surveillance in Central West Brazil from 2014 to 2019.

We whole genome sequenced 1,152 *M. tuberculosis* isolates collected during active and passive

surveillance inside and outside prisons and linked genomes to detailed incarceration histories. We applied

multiple phylogenetic and genomic clustering approaches and inferred timed transmission trees. We

found that *M. tuberculosis* sequences from incarcerated and non-incarcerated people were closely related

in a maximum likelihood phylogeny. The majority (70.8%; 46/65) of genomic clusters including people

with no incarceration history also included individuals with a recent history of incarceration. Among

cases in individuals with no incarceration history, 50.6% (162/320) were in clusters that included

individuals with recent incarceration history, suggesting that transmission chains often span prisons and

communities. We identified a minimum of 18 highly probable spillover events, *M. tuberculosis*

transmission from people with a recent incarceration history to people with no prior history of

incarceration, occurring in the state's four largest cities and across sampling years. We additionally found

that frequent transfers of people between the state's prisons creates a highly connected prison network

that likely disseminates *M. tuberculosis* across the state.

Conclusions: We developed a framework for measuring spillover from high-incidence environments to

surrounding communities by integrating genomic and spatial information to reconstruct tuberculosis

transmission chains. Our findings indicate that, in this setting, prisons serve not only as disease reservoirs,

but also disseminate *M. tuberculosis* across highly connected prison networks, both amplifying and

propagating *M. tuberculosis* risk in surrounding communities.

Abstract word count: 294

Key words: tuberculosis, epidemiology, genomics, disease reservoir, spillover, prisons

Introduction

Tuberculosis remains one of the leading causes of death by an infectious disease and led to 1.4 million deaths in 2019.[1] Locating where *M. tuberculosis* transmission occurs is critical for developing public health interventions to block transmission and prevent future infections. Yet in high-incidence settings, where most tuberculosis is transmitted remains unknown.[2–7] Built environments often shape *M. tuberculosis* transmission risk, and high-transmission settings like prisons, hospitals, and mines may have epidemiological impacts that transcend their walls[8–10], serving as institutional amplifiers of infection.[11,12] In particular, the conditions of detention, including severe overcrowding, poor ventilation, poor nutrition, and limited access to healthcare, put incarcerated people at elevated risk of many infectious diseases, including tuberculosis.[8–10]

Globally, incidence of tuberculosis in incarcerated populations is more than ten times that in the general population.[9,13] The role of prisons in driving community tuberculosis epidemics is of particular concern in Central and South America, where the incarcerated population has increased by over 200% since 2000 and tuberculosis cases among the incarcerated population have increased by over 269%[14]. Alarmingly, increasing tuberculosis notifications within prisons more than offset declines in tuberculosis outside prisons across the region[14].

The full impact of increasing incarceration rates on the tuberculosis epidemic in the Americas and elsewhere is underestimated by total incident cases within prisons. Many people are infected in prison, but only diagnosed later.[12,15,16] Further, spillover may occur when incarcerated people, prison staff, or visitors are infected within high-incidence prison environments and transmit *M. tuberculosis* onwards in communities outside prison. Previous studies have found that the excess risk of tuberculosis within prisons extends to surrounding communities[17,18] and that *M. tuberculosis* genotypes responsible for jail and prison outbreaks are also found in the surrounding communities[15,17,19,20]. Because transmission is rarely observed, however, it is often difficult to link community cases to transmission from prisons and previous studies have largely been associative.

Pathogen genomes can be powerfully harnessed to infer high-resolution evolutionary and

transmission histories, including who-infected-whom.[21] One previous study in Georgia used *M.*

tuberculosis genomes to identify frequent transmission of multi-drug resistant tuberculosis from prisons

to surrounding communities.[22] However, studies have not yet combined *M. tuberculosis* genomic,

epidemiologic, and location data to reconstruct transmission linkages between potential institutional

amplifiers and the community. To quantify the contribution of prison spillover to tuberculosis

transmission in the community, we conducted prospective genomic surveillance for tuberculosis in the

Central West Brazilian state of Mato Grosso do Sul. We integrated *M. tuberculosis* genomes with detailed

incarceration histories and took multiple phylogenetic and genomic clustering approaches to reconstruct

spillover events, transmission of tuberculosis from prisons to the community.

Methods

Study design

We conducted population-based tuberculosis surveillance in the state of Mato Grosso do Sul in

Central West Brazil from January 2014 through May 2019. Surveillance included active screening in

three of the largest prisons in the state as well as ongoing passive surveillance focused on the two largest

cities in the state, and three cities at the state's border with Paraguay and Bolivia (Fig. 1a,b; **S1 Text**). All

participants provided written consent, and this study was conducted with the approval of the Research

Ethics Committee from the Federal University of Grande Dourados, Federal University of Mato Grosso

do Sul and National Research Ethics Committee (CONEP) (CAAE 37237814.4.0000.5160,

2676613.3.1001.5160, and 26620619.6.0000.0021) and Stanford University Institutional Review Board

(IRB-40285) (**S1 Text**).

Incarceration history

We obtained permission from the Mato Grosso do Sul state prison administration agency,

Agência Estadual de Administração do Sistema Penitenciário, to access a database of all prison entries,

exits, and transfers within the criminal justice record system from January 1, 2005 through December 31,

2018 (**S1 Text**).

Laboratory diagnosis and culture

All sputum specimens collected in the participating laboratories were examined by microscopy or

GeneXpert® MTB/RIF, decontaminated with sodium hydroxide (NaOH), and inoculated in Ogawa-

Kudoh culture medium. Cultures were incubated at 37˚C for up to eight weeks and checked weekly for

visible colonies at the participating laboratories. Microbial species determination was made with the

MPT64 protein detection-based immunochromatographic rapid test (SD Bioline Kit, Standard

Diagnostics, Inc., Korea). DNA was extracted from cultured isolates with the manual CTAB (Cetyl

trimethylammonium bromide) method and whole genome sequenced on an Illumina NextSeq (2 x 151-

bp).

Whole genome sequencing

We sequenced whole *M. tuberculosis* genomes from cultured isolates on an Illumina NextSeq (2

x 151-bp). Sequence data is available on the Sequence Read Archive, in BioProject PRJNA671770. We

applied variant calling methods closely following those described in Menardo et al.[23] to be consistent

with the methods used for molecular clock estimation (**S1 Text**).

Phylogenetic and Bayesian evolutionary analysis

We fit a maximum likelihood tree with RAxML-ng 1.0.1 and rooted the tree on two Lineage 1

isolates from this study.[24] We clustered genomes using a 12-SNP threshold. For each cluster including

three or more isolates, we fit Bayesian trees with BEAST 2.6.2[25] with a strict clock, constant coalescent

population size model using tuberculosis notification dates to calibrate tips (**S1 Text**).

Transmission inference

M. tuberculosis phylogenetic trees represent patterns of evolutionary relatedness between the

consensus bacterial genomes sampled from different individuals. Because most outbreaks are

incompletely sampled and individuals may be infected with diverse populations of *M. tuberculosis*,

phylogenies do not represent the underlying history of transmission.[21] We used TransPhylo[21] to infer

transmission linkages—including unsampled hosts—that were consistent with the underlying timed

phylogenies and epidemiology of tuberculosis (i.e. generation time and sampling intervals). TransPhylo

samples transmission trees that are consistent with timed phylogenetic trees and epidemiological priors

governing the distribution of likely *M. tuberculosis* generation times (time between subsequent infections)

and sampling time (time from an individual's infection to diagnosis) (**S1 Text**).

We then combined transmission trees with incarceration histories to identify spillover events from

prisons to surrounding communities. We defined spillover events as transmission that occurred from an

individual with a history of current or former incarceration to the community, including individuals with

no incarceration history and prior incarceration. We considered transmission probabilities >50% as

evidence of spillover and included spillover events no more than two degrees removed from prisons.

TransPhylo infers transmission linkages including transmission to and from unsampled hosts and thus

does not require that all transmission events be observed. Therefore, we are able to quantify the minimum

number of transmission events, but not estimate the total role of spillover on community transmission.

Results

Tuberculosis and incarceration trends in Mato Grosso do Sul

Brazil's incarcerated population grew by 107% (from 361,402 to 748,009) from 2005 to 2019, an

increase closely paralleled in the Central West state of Mato Grosso do Sul, where the incarcerated

population grew by 115% (7,891 to 16,976) over the same period (Fig. 1a-c). In 2019, prisons in the state

were at 203% occupancy. While the state's notification rate of new and retreatment tuberculosis in the

general population (38 per 100,000) is similar to Brazil's national tuberculosis notification rate (40 per

100,000) in 2019,[26] the notification rate within prisons was 43 times as high (1,666 per 100,000), again

similar to Brazil's national notification rate within prisons (1,596 per 100,000) in 2019 (Fig. 1d,e).

Fig 1. Tuberculosis is increasingly concentrated within prisons in Mato Grosso do Sul, Brazil. (a)

Map of Brazil with states colored by the 2019 incarceration rate with Mato Grosso do Sul outlined in

black. (b) Map of Mato Grosso do Sul state with the cities with passive surveillance only and where

active surveillance was additionally conducted within prisons. (c) The incarcerated population size in

Mato Grosso do Sul grew by 142% from 2005 to 2020, increasing from 7,891 to 19,065 people. (d) Mato

Grosso do Sul state's annual new and retreatment tuberculosis notifications, colored by incarceration

status and (e) the notification rate per 100,000 people for the incarcerated population and non-

incarcerated populations from 2009 to 2019. The y-axis is log-scaled. The mean tuberculosis notification

rate was more than 46 times greater among the incarcerated population compared to the non-incarcerated

population. The green and yellow bars (d) and (e) indicate the period of active surveillance in prisons in

Campo Grande and Dourados respectively.

Population-based genomic surveillance across Mato Grosso do Sul state

169   From 2014 to 2019, 3,491 new and retreatment cases of tuberculosis were reported in the Campo

Grande and Dourados, the two largest cities in Mato Grosso do Sul, Brazil. Of these, 1,249 had positive

cultures and we sequenced 787 *M. tuberculosis* isolates. We sequenced additional isolates from three

other cities in the state and from earlier years in Dourados for a total of 1,090 isolates from unique

tuberculosis notifications (Fig. 1b). Whole genome sequences (WGS) for a total of 1,043 isolates met our

coverage and quality criteria (Methods). We excluded 10% (108 of 1043) of isolates with mixed

infection, resulting in 935 high-quality genomes from 935 unique tuberculosis episodes from 918

individuals.

177   Of the 935 isolates in our analyses, 50% (465 of 935) were from patients who were incarcerated

at the time of tuberculosis notification; 16% (150 of 935) were formerly incarcerated; and 34% (320 of

935) of the study population did not have an incarceration history. Among those who did not have an

incarceration history, 32 people reported contact with incarcerated individuals. Additional population

characteristics are in Table E1. Isolates were largely from *M. tuberculosis* lineage 4 (932 of 935) and

predominantly fell into three sub-lineages: lineage 4.1, 4.3, and 4.4 (Fig. 2). We focus subsequent

analyses of transmission on lineage 4. Overall, we found a low prevalence of antibiotic resistance across

isolates, with 93.7% of isolates susceptible to all antibiotics (Fig. 2).

Fig 2. *M. tuberculosis* isolates from incarcerated and non-incarcerated people are closely related in

Mato Grosso do Sul, Brazil. A maximum likelihood phylogeny of 932 tuberculosis isolates from

Lineage 4 inferred from a multiple sequence alignment of 19,753 SNPs and rooted on two Lineage 1

isolates from this study. Branch lengths are in units of substitutions per site. Branches are colored by sub-

lineage. From the inside, rings are colored by incarceration status, antimicrobial resistance prediction, and

genomic cluster (Methods). "Other" resistance indicates resistance to at least one antibiotic other than

isoniazid or rifampicin. The clade containing the largest predicted genomic cluster (Cluster 5), including

170 isolates, is highlighted in purple.

Phylogenetic structure of *M. tuberculosis* from prisons and the community

In a maximum likelihood phylogeny (Fig. 2), *M. tuberculosis* isolates sampled from incarcerated

and non-incarcerated people are distributed across the tree and do not form distinct clades, indicating

recent shared evolutionary history of isolates sampled from prisons and the community. Terminal branch

lengths, a proxy for the degree of recent transmission,[27] were significantly longer for isolates from

people with no incarceration history at the time of tuberculosis notification (9.4×10^{-4} substitutions/site)

compared to isolates from incarcerated individuals (2.2×10^{-4}; $p < 0.001$) or formerly incarcerated

individuals (3.8×10^{-4}; $p=0.019$), suggesting that individuals with an incarceration history were more

likely to be recently infected, although this could alternatively reflect a higher rate of genomic sampling

(Fig. S1; permutation test in **S1 Text**).

We tested for phylogenetic signal in incarceration status of sampled tuberculosis patients—

whether incarceration status was distributed randomly across the tree—with Fritz's D[28], a test

"clumping" in observed traits. The *D* statistic is equal to 1 if the trait is distributed randomly across tree

tips with respect to the phylogeny and 0 if the trait is consistent with a pattern of clumping consistent with

Brownian motion of the trait along the phylogenetic tree.

We predicted that if prison and community epidemics were distinct, incarceration status would

exhibit an extremely clumped pattern on the *M. tuberculosis* phylogeny, with separate clades including

samples from incarcerated people and community members. Conversely, if transmission between prisons

and the community occurred just as frequently as transmission within prisons, there would be a random

pattern of incarceration status across the phylogeny. We found that incarceration status had a moderate

phylogenetic signal (Fritz's D=0.59), indicating a pattern of shared evolutionary history of isolates

collected from individuals with and without an incarceration history, with some degree of "clumping"

with respect to incarceration status.

Genomic clustering of *M. tuberculosis* from prisons and the community

Patterns of genomic relatedness are often used to infer potential recent transmission; *M.*

tuberculosis isolates that are more closely genetically related are hypothesized to be more likely linked

through recent transmission[29]. To identify potential transmission clusters, we applied a commonly used

12-SNP threshold [29], including all isolates within the threshold distance of at least one other clustered

isolate. Eighty-three percent (777 of 935) of the isolates fell into 84 genomic clusters (each including 2 to

170 isolates), providing evidence that notifications were largely attributable to recent, local transmission

rather than travel-associated importation or re-activation of genetically distinct latent infections (Fig. 2).

Isolates from incarcerated people were more frequently clustered (93.3%, 434/465), than those from

formerly incarcerated (86.0%, 129 of 150; p <0.0001), or never incarcerated people (66.9%, 214 of 320; p

<0.0001), again suggesting more recent transmission within prisons.

We predicted that if prison and community-associated epidemics were distinct, isolates from the

community would be most closely related to and cluster with other isolates from the community and vice

versa. The majority (70.8%; 46/65) of genomic clusters including people with no incarceration history

also included individuals with a recent history of incarceration. Among cases in individuals with no

incarceration history, 50.6% (162/320) were in clusters that included individuals with recent incarceration

history, suggesting that transmission chains often span prisons and communities.

We found a similar pattern of clustering of isolates from community members and individuals

with an incarceration history when using an alternative 5-SNP threshold for genomic clustering[30]. With

this threshold, of the 80 clusters, 56 included isolates from people with no incarceration history and

78.6% (44 of 56) also included isolates from people with an incarceration history. Among all tuberculosis

cases occurring among individuals with no incarceration history, 46.9% (150 of 320) were in 5-SNP

clusters that included individuals with incarceration history.

A *M. tuberculosis* clone spans prisons and the community across the state

The largest potential transmission cluster, including 170 isolates sampled from September 2010

to April 2019, was distributed across the state, including cases found across state's two largest cities,

Campo Grande and Dourados, as well as the smaller cities Corumbá (on the Bolivian border) and Ponta

Porã (on the Paraguayan border) (Fig. 1b, Fig. 3a). The cluster had a most recent common ancestor in

1996 (95% HPD: 1989-2003; Fig. 3a), indicating that it had circulated for approximately 23 years before

the most recent samples were collected. 103 isolates were from people incarcerated at the time of

diagnosis, 35 from people formerly incarcerated, and 32 from people with no incarceration history at the

time of tuberculosis notification.

As observed in other clusters, isolates from people who are currently and formerly incarcerated

were closely genetically related—and often, identical—to isolates from people who were not incarcerated

at the time of notification, suggesting they are linked through recent transmission (Fig. 3b). The three

largest haplotypes (identical sequences), including 14, 9, and 8 samples, were dominated by isolates from

people who were incarcerated. Each haplotype also included isolates from people who were formerly

incarcerated and from people with no incarceration history, indicating very recent transmission between

incarcerated individuals and individuals in the community.

Fig 3. A single, recently emerged *M. tuberculosis* clone spans prisons and the community across the

major cities Mato Grosso do Sul state. (a) A Bayesian time-calibrated phylogeny of the largest sampled

genomic cluster of 170 isolates. The cluster emerged approximately 23 years before the most recently

sampled isolate, with a most recent common ancestor in 1996 (95% HPD: 1989-2003). Tip point color

indicates patient's incarceration status at the time of tuberculosis notification. Annotation bar colors

indicate city. Grey error bars indicate the 95% Bayesian highest posterior density intervals for node date.

Clade posterior support values are shown on the middle of branches for clades with posterior support >

0.5. (b) A haplotype network of the single largest genomic cluster, including 170 isolates. Nodes

represent unique haplotypes and are scaled to size. Points along branches indicate SNP distances between

haplotypes. Node color indicates incarceration status at the time of diagnosis.

Prison networks disseminate *M. tuberculosis* across space

We hypothesized that the criminal justice system's frequent transfer of people between prisons

and jails could disseminate TB across the state. We analyzed movements in the state's incarceration

database and found that the average duration of incarceration was 230 days in a prison and 25 days in a

jail. The incarceration database documents 7,982 mean yearly transfers between prisons (including closed

and semi-open prisons) from 2015-2018 in a prison population of 17,221 in August 2018, including

frequent transfers between cities across the state (Fig. 4a).

To investigate the role of the criminal justice system in disseminating infection, we tracked the

spread of the largest sampled *M. tuberculosis* cluster (Fig. 3) across prisons in Mato Grosso do Sul (Fig.

4b). From early 2011 to 2014, sampled isolates from the cluster were concentrated in the maximum

security prison in Dourados, after which it was identified within two prisons in Campo Grande and then

exported to prisons and jails in other cities. The cluster was not contained within the state's prison

network; notified tuberculosis cases among community members outside of prison occurred over the

duration of the clusters' state-wide spread.

Fig 4. Prisons disseminate tuberculosis through the frequent movement of people between prisons.

(a) Incarceration locates people within highly connected contact networks. Map of Mato Grosso do Sul

state with orange points indicating the 21 prisons which most frequently transfer people between prisons.

Edges are colored by the mean number of yearly transfers made between prisons from 2015-2018,

including transfers in both directions. Edges were drawn between prison pairs with more than 10 transfers

in the study period. Transfers between prisons within a city are not shown. (b) An *M. tuberculosis* clone

(depicted in Figure 3) spreads throughout multiple prisons, jails, and the community. Each line represents

an individual patient infected with an isolate in the largest genomic cluster, ordered by notification date.

Line color indicates patient location, determined by matching patient names with the state incarceration

database (Methods). Points indicate TB notification dates; point color indicates patient location at the

time of TB notification. Individuals with notification dates after December 31, 2018 are not shown.

Transmission trees reveal frequent spillover of infection

We further investigated recent transmission in the state by inferring probabilistic transmission

trees representing who-infected-whom for the 84 identified *M. tuberculosis* genomic clusters with

TransPhylo[21]. Because our sampling of the TB epidemic in Mato Grosso do Sul was incomplete and we

did not observe the majority of transmission events resulting in sampled cases, we quantified the

minimum number of recent infections attributable to prison spillover rather than the total role of spillover

on the community epidemic. We identified a minimum of 16.8% (18 of 107) infections among never

incarcerated individuals were directly attributable to observed spillover from prisons. Observed spillover

of infection to never or formerly incarcerated individuals occurred across the state in the four largest cities

sampled (Campo Grande, Dourados, Ponta Pora, and Corumba) and every year, from 2014-2019.

Together, these findings indicate that spillover of *M. tuberculosis* infection from prisons to surrounding

communities is frequent. The result is that the 0.84% of the state's population incarcerated at a given time

(22,706 of 2,713,147 in 2017) or 2.6% currently or formerly incarcerated at a given time (71,703 of

2,713,147), have a disproportionate effect on TB transmission outside of prisons.

Transmission inferences are sensitive to epidemiological priors

We conducted a sensitivity analysis to determine the effect of epidemiological priors (Table S2)

on transmission inferences for the largest genomic cluster (170 isolates). Transmission probabilities

inferred for pairs of individuals were closely correlated across generation time priors (Pearson's r: 0.94 –

0.98, Fig. S3) and sampling time priors (Pearson's r: 0.84 – 0.98, Fig. S4), although the total number of

observed transmission events was strongly influenced by both priors (Figure S5). This suggests that, as

expected, epidemiological priors impact the magnitude of inferred transmission probabilities rather than

the identity of predicted transmission pairs. As expected, the minimum number of observed spillover

events in the largest cluster were sensitive to epidemiological priors, when transmission was defined as a

transmission probability greater than or equal to 50%, and ranged from 0 (short sampling and generation

times) to 28 (medium generation time and very long sampling time) (Fig. S6).

Discussion

Here, we linked intensive *M. tuberculosis* genomic surveillance to detailed individual-level

incarceration histories, allowing us to reconstruct high-resolution population-wide tuberculosis

transmission in Central West Brazil, a setting with an extreme disparity in tuberculosis incidence between

prisons and surrounding communities. We found that *M. tuberculosis* sequences from incarcerated and

non-incarcerated individuals were closely evolutionarily related and sometimes identical. Locally

circulating *M. tuberculosis* clones spanned prisons and the community, and the majority of genomic

clusters included individuals with and without a recent history of incarceration. In reconstructed

transmission trees, we found evidence of frequent spillover of infection in all the major cities in the state

and across sampling years, indicating that spillover is not the effect of a single prison but a widespread

phenomenon. We found that frequent transfers of people between prisons in Mato Grosso do Sul, Brazil

creates a closely connected prison contact network that likely facilitates the spread of *M. tuberculosis*

between prisons and may disseminate infection risk across the state.

While alarming disparities in tuberculosis incidence between prisons and surrounding

communities are well-documented,[9,13] it has been more challenging to directly link tuberculosis cases

in the community to spillover from prisons. Previous studies of the broader effects of prisons on general

TB epidemics have largely been associative. Here, we describe a probabilistic framework for identifying

spillover events that can be applied more generally to measure the impact of high-transmission

environments such as mines or hospitals on tuberculosis transmission in the general population. Our

finding of frequent and widespread spillover events demonstrates the outsized effect of prison

environments on community-wide *M. tuberculosis* transmission. This also suggests that the introduction

of *M. tuberculosis* lineages or resistance mutations into prisons is likely to be a source of community

epidemics. Our finding of the likely role of prison transfers in disseminating *M. tuberculosis* highlights an

additional role that carceral systems may play in propagating infectious diseases both within highly-

connected prison systems and across space.

The rapid rise in incarceration rates in many countries[31] and stark global disparities in

tuberculosis incidence inside compared to outside prisons[8,9,13,32] suggests that spillover of infection

may contribute to community TB transmission outside of Brazil. A study of multi-drug resistant (MDR)

TB transmission found that up to 31% of MDR TB in the country of Georgia was directly or indirectly

linked to prisons.[22] Our transmission inference framework is conservative in that it identifies a

minimum contribution of spillover to total community transmission, and therefore, is consistent with

these findings. The role of prisons and other detention centers as disease reservoirs is not unique to

tuberculosis. Previous studies have identified spillover of meningococcal disease[33] and SARS-CoV-

2[34] from prisons to the community. Prison transfers have also spread of SARS-CoV-2 across prisons in

California, resulting in widespread outbreaks.[35]

The frequent spillover of infection from prisons to surrounding communities indicates that for TB

control to be achieved in Central West Brazil, TB control efforts must focus on reducing the

extraordinarily high transmission risk within prisons. Tuberculosis control programs should expand

routine active screenings during[12,36] and following incarceration so that cases can be diagnosed early

and linked to treatment. Individuals leaving prison are at heightened risk of tuberculosis and are a

particularly vulnerable population. Preventive therapy is not provided in prisons in Brazil and many other

low- and middle-income countries and could represent an effective means for tuberculosis prevention,

particularly among individuals exiting prisons for community environments where the risk of reinfection

is lower. More broadly, improving the unsanitary, inhumane conditions of incarceration and expanding

access to primary healthcare and nutrition are critical.

Reducing this excess burden of disease will require work that extends beyond biomedical

interventions. The most direct way to mitigate the excess tuberculosis risk created by prisons is for

governments to rapidly reduce incarcerated populations, as advocated by the American Public Health

Association, to reduce the broader harms associated with incarceration.[37] Prison releases have already

begun as a means of reducing infectious disease transmission risk. [38] In an attempt to reduce the risk of

COVID-19 in prisons, for example, Brazil released more than thirty thousand people by mid-2020.[39]

Our study has several limitations. First, our study highlights the uncertainty inherent in

transmission inferences drawn from genomic surveillance data, which almost always constitutes an

incomplete sample of population-wide infections. We addressed the incomplete sampling of *M.*

tuberculosis genomes in two ways. First, we took a probabilistic transmission inference approach that

accounts for uncertainty in transmission linkages and also includes unsampled individuals contributing to

transmission.[21] Second, in order to collect the most comprehensive set of *M. tuberculosis* genomes

possible, we combined isolates collected during active mass screening within prisons and passive routine

surveillance in the community. The result was that isolates from prisons are overrepresented in our

genomic collection. Because of the over-sampling within prisons, we could not directly quantify the

contribution of transmission within prisons and spillover from prisons on overall recent infections in the

study area. However, by taking a probabilistic transmission inference approach that accounted for

uncertainty in transmission linkages as well as unsampled individuals contributing to transmission, we

were able to quantify the minimum number of reconstructed spillover events consistent with inferred

transmission trees. Because we cannot infer the identity or incarceration history of unsampled people who

contribute to transmission, our estimates of the contribution of spillover are conservative and likely

underestimate the full impact of spillover on infections in the community. However, our finding of

geographically widespread spillover occurring over several years suggests that spillover is common and

an important contributor to population-wide *M. tuberculosis* transmission.

Additionally, transmission trees reflect information from timed *M. tuberculosis* phylogenies and

epidemiological priors and it remains difficult to distinguish genomic sampling proportion from

generation time and sampling time distributions. Because we conducted transmission inference in a

tuberculosis-endemic setting, rather than in a discrete outbreak, our rate of genomic sampling was lower

than in some previous tuberculosis transmission studies. To reduce uncertainty in sampling proportion,

we conducted transmission inference on recent subtrees for which we had sampled a greater proportion of

isolates. To investigate the influence of epidemiological priors on transmission inferences, we conducted

a sensitivity analysis (Table E2) and found that our conclusions about transmission linkages were largely

unchanged (Supplementary Text, Figs. S3-S4), although, as expected, the number of transmission events

and thus spillover events was dependent on epidemiological priors (Figs. S5-S6).

Finally, we conducted whole genome sequencing of *M. tuberculosis* isolates cultured from

sputum and generated single consensus genomes from each participant, excluding within-host diversity.

We also employed an existing bioinformatic approach to identify *M. tuberculosis* variants and excluded

repetitive genomic regions as is common practice.[23,40] Future studies that leverage within-host *M.*

tuberculosis diversity and/or include the diverse PE/PPE genes could provide greater resolution and

reduce the uncertainty of transmission inferences.[41]

In this study, we present genomic evidence that spillover of tuberculosis infection risk from

prisons to surrounding communities is frequent in Central West Brazil. The dramatic expansion of

incarceration in recent decades has put an increasing population at extremely high risk of tuberculosis;

this risk extends to surrounding communities. Reducing the excess tuberculosis transmission risk within

prisons and other detention centers is therefore an urgent public health priority. Our findings additionally

highlight the potential for pathogen genomic surveillance to be used to identify other key environments

with a disproportionate role in pathogen transmission in order to prioritize public health interventions.

Acknowledgments

We would like to acknowledge the contributions of AGEPEN, the Mato Grosso do Sul state criminal

justice agency; LACEN, the state reference laboratory; and the Mato Grosso do Sul State Health

Department. All authors had full access to all the data in the study and had final responsibility for the

decision to submit.

References

1.	The World Health Organization, World Health Organization. Global Tuberculosis Report 2020. Geneva; 2020.

2.	Churchyard G, Kim P, Shah NS, Rustomjee R, Gandhi N, Mathema B, et al. What We Know about Tuberculosis Transmission: An Overview. J Infect Dis. 2017;216: S629–S635. doi:10.1093/infdis/jix362

3.	Auld SC, Sarita Shah N, Mathema B, Brown TS, Ismail N, Omar SV, et al. Extensively drug-resistant tuberculosis in South Africa: Genomic evidence supporting transmission in communities. Eur Respir J. 2018;52. doi:10.1183/13993003.00246-2018

4.	Middelkoop K, Mathema B, Myer L, Shashkina E, Whitelaw A, Kaplan G, et al. Transmission of tuberculosis in a south African community with a high prevalence of HIV infection. J Infect Dis. 2015;211: 53–61. doi:10.1093/infdis/jiu403

5.	Andrews JR, Morrow C, Walensky RP, Wood R. Integrating social contact and environmental data in evaluating tuberculosis transmission in a South African township. J Infect Dis. 2014;210: 597–603. doi:10.1093/infdis/jiu138

6.	Yates TA, Khan PY, Knight GM, Taylor JG, McHugh TD, Lipman M, et al. The transmission of Mycobacterium tuberculosis in high burden settings. Lancet Infect Dis. 2016;16: 227–238. doi:10.1016/S1473-3099(15)00499-5

7.	Andrews JR, Morrow C, Wood R. Modeling the role of public transportation in sustaining tuberculosis transmission in South Africa. Am J Epidemiol. 2013;177: 556–561. doi:10.1093/aje/kws331

8.	Dolan K, Wirtz AL, Moazen B, Ndeffo-mbah M, Galvani A, Kinner SA, et al. Global burden of HIV, viral hepatitis, and tuberculosis in prisoners and detainees. Lancet. 2016;388: 1089–1102. doi:10.1016/S0140-6736(16)30466-4

9.	Cords O, Martinez L, Warren JL, O'Marr JM, Walter KS, Cohen T, et al. Incidence and prevalence of tuberculosis in incarcerated populations: a systematic review and meta-analysis. Lancet Public Heal. 2021. doi:10.1016/S2468-2667(21)00025-6

10.	Urrego J, Ko AI, Da Silva Santos Carbone A, Paião DSG, Sgarbi RVE, Yeckel CW, et al. The impact of ventilation and early diagnosis on tuberculosis transmission in Brazilian prisons. Am J Trop Med Hyg. 2015;93: 739–746. doi:10.4269/ajtmh.15-0166

11.	Basu S, Stuckler D, McKee M. Addressing institutional amplifiers in the dynamics and control of tuberculosis epidemics. Am J Trop Med Hyg. 2011;84: 30–37. doi:10.4269/ajtmh.2011.10-0472

12.	Mabud TS, de Lourdes Delgado Alves M, Ko AI, Basu S, Walter KS, Cohen T, et al. Evaluating strategies for control of tuberculosis in prisons and prevention of spillover into communities: An observational and modeling study from Brazil. Metcalfe JZ, editor. PLoS Med. 2019;16: e1002737. doi:10.1371/journal.pmed.1002737

13.	Baussano I, Williams BG, Nunn P, Beggiato M, Fedeli U, Scano F. Tuberculosis incidence in prisons: A systematic review. PLoS Med. 2010;7: 1–10. doi:10.1371/journal.pmed.1000381

14.	Walter KS, Martinez L, Arakaki-Sanchez D, Sequera VG, Estigarribia Sanabria G, Cohen T, et al. The escalating tuberculosis crisis in central and South American prisons. Lancet. 2021 [cited 10 Apr 2021]. doi:10.1016/s0140-6736(20)32578-2

15.	Sacchi FPC, Praça RM, Tatara MB, Simonsen V, Ferrazoli L, Croda MG, et al. Prisons as reservoir for community transmission of tuberculosis, brazil. Emerg Infect Dis. 2015;21:

452–455. doi:10.3201/eid2103.140896

16. Coker R, McKee M, Atun R, Dimitrova B, Dodonova E, Kuznetsov S, et al. Risk factors
for pulmonary tuberculosis in Russia: Case-control study. Br Med J. 2006;332: 85–87.
doi:10.1136/bmj.38684.687940.80

17. Warren JL, Grandjean L, Moore DAJ, Lithgow A, Coronel J, Sheen P, et al. Investigating
spillover of multidrug-resistant tuberculosis from a prison: a spatial and molecular
epidemiological analysis. BMC Med. 2018;16: 122. doi:10.1186/s12916-018-1111-x

18. Jenkins HE, Plesca V, Ciobanu A, Crudu V, Galusca I, Soltan V, et al. Assessing spatial
heterogeneity of multidrug-resistant tuberculosis in a high-burden country. Eur Respir J.
2013;42: 1291–1301. doi:10.1183/09031936.00111812

19. Séraphin MN, Didelot X, Nolan DJ, May JR, Khan MSR, Murray ER, et al. Genomic
investigation of a Mycobacterium tuberculosis outbreak involving prison and community
cases in Florida, United States. Am J Trop Med Hyg. 2018;99: 867–874.
doi:10.4269/ajtmh.17-0700

20. Jones TF, Woodley CL, Fountain FF, Schaffner W. Increased incidence of the outbreak
strain of Mycobacterium tuberculosis in the surrounding community after an outbreak in a
jail. (Original Article). South Med J. 2003;96: 155–158. Available:
https://go.gale.com/ps/i.do?p=AONE&sw=w&issn=00384348&v=2.1&it=r&id=GALE%
7CA98828112&sid=googleScholar&linkaccess=fulltext

21. Didelot X, Fraser C, Gardy J, Colijn C, Malik H. Genomic infectious disease
epidemiology in partially sampled and ongoing outbreaks. Mol Biol Evol. 2017;34: 997–
1007. doi:10.1093/molbev/msw275

22. Gygli SM, Loiseau C, Jugheli L, Adamia N, Trauner A, Reinhard M, et al. Prisons as
ecological drivers of fitness-compensated multidrug-resistant Mycobacterium
tuberculosis. Nat Med 2021 277. 2021;27: 1171–1177. doi:10.1038/s41591-021-01358-x

23. Menardo F, Duchêne S, Brites D, Gagneux S. The molecular clock of mycobacterium
tuberculosis. PLoS Pathog. 2019;15. doi:10.1371/journal.ppat.1008067

24. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: A fast, scalable,
and user-friendly tool for maximum likelihood phylogenetic inference. Wren J, editor.
Bioinformatics. 2019 [cited 13 May 2019]. doi:10.1093/bioinformatics/btz305

25. Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A,
et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis.
PLoS Comput Biol. 2019;15: e1006650. doi:10.1371/journal.pcbi.1006650

26. World Health Organization. Global tuberculosis report 2020. Geneva; 2020.

27. Holt KE, Mcadam P, Vuong P, Thai K, Thuy N, Thuong T, et al. Frequent transmission of
the Mycobacterium tuberculosis Beijing lineage and positive selection for the EsxW
Beijing variant in Vietnam. Nat Genet. 2018. doi:10.1038/s41588-018-0117-9

28. Fritz SA, Purvis A. Selectivity in mammalian extinction risk and threat types: A new
measure of phylogenetic signal strength in binary traits. Conserv Biol. 2010;24: 1042–
1051. doi:10.1111/j.1523-1739.2010.01455.x

29. Walker TM, Kohl TA, Omar S V., Hedge J, Del Ojo Elias C, Bradley P, et al. Whole-
genome sequencing for prediction of Mycobacterium tuberculosis drug susceptibility and
resistance: A retrospective cohort study. Lancet Infect Dis. 2015;15: 1193–1202.
doi:10.1016/S1473-3099(15)00062-6

30. Walker TM, Ip CLC, Harrell RH, Evans JT, Kapatai G, Dedicoat MJ, et al. Whole-
genome sequencing to delineate Mycobacterium tuberculosis outbreaks: A retrospective

observational study. Lancet Infect Dis. 2013;13: 137–146. doi:10.1016/S1473-
3099(12)70277-3

31.   Walmsley R. World Prison Population List (twelfth edition). Inst Crim Policy Res. 2018.
Available: www.prisonstudies.org

32.   Walter K, Martinez L, Arakaki-Sanchez D, Sequera V, G ES, T C, et al. The escalating
tuberculosis crisis in Central and South American prisons. Lancet.

33.   Tappero JW, Reporter R, Wenger JD, Ward BA, Reeves MW, Missbach TS, et al.
Meningococcal Disease in Los Angeles County, California, and among Men in the County
Jails. N Engl J Med. 1996;335: 833–841. doi:10.1056/nejm199609193351201

34.   Reinhart E, Chen DL. Incarceration and its disseminations: COVID-19 pandemic lessons
from Chicago's cook county jail. Health Aff. 2020;39: 1412–1418.
doi:10.1377/hlthaff.2020.00652

35.   Sandra McCoy, Stefano M. Bertozze, David Sears, Catherine Duarte, Drew Cameron,
Brie Williams. Urgent Memo: COVID-19 Outbreak: San Quentin Prison. San Francisco,
CA; 2020. Available: https://www.documentcloud.org/documents/6956448-06-15-20-San-
Quentin-Urgent-Memo.html

36.   Santos A da S, Oliveira RD de, Lemos EF, Lima F, Cohen T, Cords O, et al. Yield,
Efficiency and Costs of Mass Screening Algorithms for Tuberculosis in Brazilian Prisons.
Clin Infect Dis. 2020 [cited 3 Apr 2020]. doi:10.1093/cid/ciaa135

37.   American Public Health Association. Advancing Public Health Interventions to Address
the Harms of the Carceral System. 2021. Available: https://www.apha.org/policies-and-
advocacy/public-health-policy-statements/policy-database/2021/01/14/advancing-public-
health-interventions-to-address-the-harms-of-the-carceral-system

38.   Wang EA, Western B, Berwick DM. COVID-19, Decarceration, and the Role of
Clinicians, Health Systems, and Payers: A Report From the National Academy of
Sciences, Engineering, and Medicine. Jama. 2020;06511. doi:10.1001/jama.2020.22109

39.   De Oliveira Andrade R. Covid-19: Prisons exposed in Brazil's crisis. The BMJ. 2020.
doi:10.1136/bmj.m2884

40.   Meehan CJ, Goig GA, Kohl TA, Verboven L, Dippenaar A, Ezewudo M, et al. Whole
genome sequencing of Mycobacterium tuberculosis: current standards and open issues.
Nat Rev Microbiol. 2019;17: 533–545. doi:10.1038/s41579-019-0214-5

41.   Walter KS, Colijn C, Cohen T, Mathema B, Liu Q, Bowers J, et al. Genomic variant-
identification methods may alter mycobacterium tuberculosis transmission inferences.
Microb Genomics. 2020;6: 1–16. doi:10.1099/mgen.0.000418

a) Incarceration status
- No incarceration history
- Formerly incarcerated
- Incarcerated

b) Antimicrobial resistance
- Isoniazid
- MDR
- Other
- Rifampicin
- Susceptible

c) Genomic cluster
- Other
- 1
- 2
- 5
- 4
- 3
- Unique

0.005

a   b   c

Lineage
- 4.1
- 4.2
- 4.3
- 4.4
- 4.7
- 4.8
- 4.9

a

b

Location at TB notification
Community
Jail
Prison: Campo Grande
Prison: Dourados
Prison: Other