1
2
3

Novelty-induced frontal-STN networks in Parkinson's disease
Running Title: Novelty-induced networks in Parkinson's disease

Rachel C Cole[1], Arturo I Espinoza[1], Arun Singh[2], Joel I Berger[3], James F Cavanagh[4],
Jeremy D Greenlee[3], Nandakumar S Narayanan[1]

[1]Department of Neurology, University of Iowa, Iowa City, IA, 52242, [2]Division of Basic
Biomedical Sciences, Sanford School of Medicine, University of South Dakota, Vermillion, SD,
57069, [3]Department of Neurosurgery, University of Iowa, Iowa City, IA, 52242, [4]Department of
Psychology, University of New Mexico, Albuquerque, New Mexico, 87131

*Corresponding Author
Nandakumar Narayanan
nandakumar-narayanan@uiowa.edu
169 Newton Road
Pappajohn Biomedical Discovery Building—5336
University of Iowa, Iowa City, 52242
319-353-5698

46 pages, 7 figures, 2 tables

Abstract: 135 words; Introduction, 617 words; Discussion: 1,207 words

The authors declare no competing financial interests.

Acknowledgements

We would like to acknowledge Haiming Chen, Youngcho Kim, Derrick Okine, Jan Wessel,
Darcy Diesberg, and Tobin Dykstra for technical contributions and helpful discussions. This
work was funded by NINDS R01 NS100849-01A1 to NN/JG/JFC and NIA F32 AG069445-01
to RC. This study was supported in part by the University of Iowa Institute for Clinical and
Translational Science, which is granted with Clinical and Translational Science Award funds
from the National Institutes of Health (UL1TR002537).

Abstract

Evaluating and responding to new information requires cognitive control. Here, we studied novelty-response mechanisms in Parkinson's disease (PD). In PD patient-volunteers, we recorded from cortical circuits with scalp-based electroencephalography (EEG) and from subcortical circuits using intraoperative neurophysiology during surgeries for implantation of deep-brain stimulation (DBS) electrodes. We report three major results. First, novel auditory stimuli triggered midfrontal ~4-Hz rhythms, which were attenuated in PD patients but were not linked with cognitive function or novelty-associated slowing. Second, 32% of subthalamic nucleus (STN) neurons were response-modulated; nearly all (94%) of these were also modulated by novel stimuli. Finally, response-modulated STN neurons were coherent with midfrontal low-frequency activity. These findings link scalp-based measurements of neural activity with neuronal activity in the STN. Our results provide insight into midfrontal cognitive control mechanisms and how hyperdirect circuits evaluate new information.

Significance statement

Parkinson's disease (PD) is a neurodegenerative disorder with motor and cognitive symptoms. Here we studied how the brain recruits cognitive control in response to new information. We found that PD patients have decreased cortical low-frequency oscillations linked to cognitive control. We also examined single neurons in PD patients in a deep brain structure called the subthalamic nucleus, and found that response-modulated neurons were uniquely modulated by novelty and they fired in-phase with these same low-frequency brain oscillations. Our work helps illustrate how neural circuits evaluate new information, which could be relevant for PD as well as other diseases that disrupt cognition.

INTRODUCTION

New information requires careful consideration. Indeed, when the brain evaluates novel information it recruits cognitive-control processes reflected by bursts of midfrontal low-frequency 1-8 Hz activity (Cavanagh & Frank, 2014; Solís-Vivanco et al., 2018). In the prefrontal cortex, our group and others have shown that these bursts of midfrontal activity engage single neurons encoding cognitive functions (Kim et al., 2017; Kim & Narayanan, 2019; Mikell et al., 2014; Narayanan et al., 2013). This interaction between cortical oscillations and single neurons may orchestrate the arc from extrinsic perceptions to intrinsic responses. Human diseases such as Parkinson's disease (PD) involve slowed or profoundly disrupted movements; however, it is unknown how stimulus-related factors such as novelty interact with motor responses. In this study, we studied midfrontal cognitive-control processes in PD via novelty-related processes.

One paradigm used to study the link between novel stimuli and responses is the oddball task. In this task, participants must respond to a commonly-occurring, standard stimulus on most trials. When novel stimuli are presented, the orienting response causes motor slowing (Sokolov, 1962) and elicits control processes (Fan, 2014). In electroencephalography (EEG) studies, novel stimuli and the subsequent orienting response evoke characteristic event-related potential (ERP) signatures in the midfrontal cortex (Debener et al., 2005; Polich, 2007; Ranganath & Rainer, 2003). Novelty-induced orienting can be considered a sub-feature of control. Cortical responses to novel stimuli involve midfrontal low-frequency oscillations between 1-8 Hz. Cavanagh and Frank (2014) proposed that these midfrontal oscillations reflect the recruitment of cognitive-control processes, however, the precise bands and topography of novelty-related slowing, particularly in PD patients, is unclear. Frontal activity may serve as a mechanism by which

neurons exert top-down control across the brain (Cavanagh & Frank, 2014), including

subcortical networks in the basal ganglia (Zink et al., 2006). These subcortical basal ganglia

networks can be modulated by novelty; indeed, a human intraoperative study by Mikell and

colleagues (2014) found that subcortical neurons in the substantia nigra increased firing more

following infrequent novel sounds than frequent standard sounds.

We and others have found consistent evidence that PD attenuates midfrontal 1-8 Hz

activity during attentional orienting, conflict, interval timing, and responses to startling stimuli

(Cavanagh et al., 2018; K.-H. Chen et al., 2016; Kelley et al., 2018; Lange et al., 2016; Parker et

al., 2015; Singh et al., 2018, 2021; Solís-Vivanco et al., 2018). Furthermore, we have found that

during interval timing midfrontal ~4-Hz rhythms are coherent with field potentials in the

subthalamic nucleus (STN) (Kelley et al., 2018). Notably, Singh and colleagues (2021) found

that attenuated ~4-Hz EEG oscillatory activity during interval timing was predictive of cognition

as measured by the Montreal Cognitive Assessment (MoCA), suggesting that the recruitment of

midfrontal ~4-Hz oscillations reflects general cognitive-control processes. These data lead to the

hypothesis that midfrontal ~4-Hz rhythms engage subcortical networks in response to novelty.

We tested this idea by first comparing midfrontal EEG during an oddball task between PD

patients and healthy older adult subjects (Andrés et al., 2006; Escera et al., 1998; Parmentier et

al., 2008, 2010). In addition, we recorded midfrontal EEG and intracranial STN neurons during

an intraoperative oddball task in PD patients undergoing DBS implantation surgery. We report

three main findings. First, we found that PD patients had decreased midfrontal ~4-Hz rhythms

compared to controls; however, these rhythms did not correlate with cognitive status or novelty-

related slowing. Second, we found that neurons in the STN were strongly response-modulated

rather than cue-modulated, and most response-modulated STN neurons were influenced by

novelty. Third, we found that midfrontal low-frequency oscillations were coherent with

response-modulated STN neurons. These data provide insight into how novel information

engages a midfrontal cognitive control system, and how this system in turn contributes to a

hyperdirect subcortical mechanism for response control.

MATERIALS AND METHODS

Participants

This investigation included both an EEG experiment and an intraoperative neurophysiology

experiment. For the EEG experiment, we recruited PD patient-volunteers and demographically-

similar control participants (see Table 1). PD patients were recruited through the Movement

Disorders clinic at University of Iowa Hospitals and Clinics (UIHC). Healthy older adults were

recruited to serve as control participants from either the Seniors Together in Aging Research

(STAR) registry or a list of people who previously participated in research in our lab and agreed

to be contacted for new research opportunities. Participants were considered healthy if they did

not have any neurological and psychological diseases or disorders. Participants were recruited by

email or phone and received compensation of $30/hour. All procedures were approved by the

University of Iowa (UI) Institutional Review Board (IRB) (#201707828).

In parallel with the EEG experiment, we recruited 18 PD patient-volunteers who had elected

to undergo bilateral STN deep brain stimulation (DBS) implantation surgery (see Table 2). These

participants were recruited over a time period of 1.5 years (July 2019–August 2020). Within the

study period there was a 5-month pause in enrollment due to research suspension during the

COVID-19 crisis. Patients were asked during preoperative clinical sessions (1–14 days prior to

surgery) whether they were interested in participating in research during their DBS implantation

surgery. If they agreed, a researcher separate from the surgical team discussed the research

protocols with them, obtained informed consent, and completed a brief practice session of the

task. All intraoperative research procedures were approved by the UI IRB (#201402730). All

patients were tested on levodopa as our past work has shown that levodopa does not reliably

modulate midfrontal ~4 Hz rhythms (Singh et al., 2018, 2021). Levodopa-equivalent doses

(LEDD) are presented in Table 2.

Oddball task for EEG experiment

We assessed the response to novelty using a version of the cross-modal oddball distractor task

(Andrés et al., 2006; Escera et al., 1998; Parmentier et al., 2008, 2010; Parmentier & Hebrero,

2013). This task was presented using the PsychToolbox-3 functions (Brainard, 1997; Kleiner et

al., 2007; Pelli, 1997, http://psychtoolbox.org/) in MATLAB 2018b on either a Linux or

Windows computer. Task-specific audio was played through Dell Rev A01 speakers positioned

on either side of the monitor. Responses were made with the left and right index fingers on a

standard QWERTY USB-keyboard. In addition to the cross-modal oddball distractor task,

participants completed additional cognitive and motor tasks, as well as detailed

neuropsychological and neuropsychiatric assessments as part of other protocols (Singh et al.,

2018, 2021).

Our version of the cross-modal oddball distractor task involved a choice reaction-time task in

which a white arrow appeared in the center of a black screen, and the participant was required to

press the key that corresponded with the direction of the arrow ("q" for left arrow, "p" for right

arrow) as quickly as possible. The appearance of the arrow was preceded by an audio-visual cue

by 500 ms (either the standard cue or the distractor cue). Participants were instructed that this

cue would appear 500 ms before the target stimulus (white arrow), and they were told that the

cue would be a green circle and a short tone (600-Hz sine wave tone lasting 200 ms). The audio-

visual cue was followed by the target arrow (**Figure 1A**). Participants had to respond within 1 s,

after which the fixation cross reappeared and the next trial started after a variable inter-trial

interval between 500–1000 ms.

Participants completed a brief practice of the task (10 trials), which had all the familiar

standard cues described above. Following the practice, participants completed four blocks of 60

trials each, for a total of 240 trials. Of the 240 trials, 80% contained the standard cues as

described above, 10% contained an unexpected auditory oddball cue (non-repeating, randomly-

created sine wave that sounded like a birdcall or robotic noise lasting 200 ms in duration) in

place of the expected tone, and 10% contained an unexpected visual oddball cue (unique

shape/color combination) in place of the green circle. Because visual stimuli were not presented

during intraoperative research, only auditory oddball trials were analyzed.

EEG recordings

EEG recordings were performed according to methods described in detail previously (Singh

et al., 2020, 2021). Briefly, we used a 64-channel EEG actiCAP (Brain Products GmbH) with a

0.1-Hz high-pass filter and a sampling frequency of 500 Hz. We used electrode Pz as reference

and electrode FPz as the ground. EEG activity was referenced according to the procedures

described in Singh et al., (2020, 2021). Briefly, an additional channel was recorded at the mid-

inion region (Iz), and we removed unreliable Fp1, Fp2, FT9, FT10, TP9, and TP10 channels,

resulting in 59 channels for pre- and post-processing. Data were epoched around the cues from -

1000 ms to 2500 ms peri-cue.

Bad channels and bad epochs were identified using the FASTER algorithm (Nolan et al.,

2010) and the pop_rejchan function from EEGLAB, and were then interpolated and rejected

respectively. On average, 1.6 ± 0.9 channels per subject were removed, and Cz was never

removed during preprocessing. Eye blink contaminants were removed following independent

component analysis (ICA).

Event-related potentials (ERPs) were low-pass-filtered at 20 Hz for analyses. Our primary

interest was in ~4-Hz midfrontal rhythms; consequently, we utilized time-frequency analyses.

We computed spectral measures by multiplying the fast Fourier transformed (FFT) power

spectrum of single-trial EEG data with the FFT power spectrum of a set of complex Morlet

wavelets (defined as a Gaussian-windowed complex sine wave: $e^{i2\pi tf}e^{-t^2/(2 \times \sigma^2)}$, where t=time and

f=frequency). Wavelets increased from 1–50 Hz in 50 logarithmically-spaced steps, which

defined the width of each frequency band, increasing from 3–10 cycles between 1–50 Hz and

taking the inverse FFT (Cohen, 2014). The end result of this process was identical to time-

domain signal convolution, and it resulted in estimates of instantaneous power (the magnitude of

the analytic signal) and phase angle (the arctangent of the analytic signal). We then cut the length

of each signal accordingly for each trial (-500 ms to 1000 ms). These short temporal epochs

reflect the wavelet-weighted influence of longer time and frequency periods. Power was

normalized by converting to a decibel (dB) scale ($10*log10(power_t/power_{baseline})$), allowing us to

directly compare the effects across frequency bands. The baseline for each frequency was

calculated by averaging power from -300 ms to -200 ms prior to cue onset (Singh et al., 2018,

2020, 2021). A 100 ms duration is often used as an effective baseline since pixel-wise time-

frequency data points have already been resolved over smoothed temporal and frequency

dimensions with the wavelets. For plotting purposes only, statistical-cluster-based thresholding

(minimum cluster-size of 500 pixels) was used to visualize areas where temporal-frequency

power differed between controls and PD patients (**Figure 2D,** black contours). Our previous

work has focused on diminished midfrontal cue-triggered ~4-Hz rhythms in PD, although we

have observed differences across 1-8 Hz depending on the specific demands elicited by the task

(e.g. 1-4 Hz novelty vs. 4-8 Hz control-related processes) (K.H. Chen et al., 2016; Kim et al.,

2017; Narayanan et al., 2013; Singh et al., 2018, 2020, 2021).

Oddball task for intraoperative neurophysiology

We used an auditory version of a 3-tone oddball task during intraoperative human

neurophysiology to capture the response to novelty in the operating room during DBS-electrode

implantation surgeries. This task is ideal for research for intraoperative patients whose heads are

in a stereotaxic frame during multielectrode recordings because, 1) it is in the auditory domain

and does not require a monitor for visual stimuli, 2) it is relatively simple in that it requires a

motor response to tones, and 3) several trials can be collected in a few minutes. Similar 3-tone

designs have been used previously (Debener et al., 2005), in which novelty is probed with novel

tones. All stimuli were auditory and presented at an appropriate volume through earbuds or

external speaker to be clearly heard over operating room noise. The task and stimuli were

presented using PsychToolbox-3 in MATLAB 2018b on a Windows laptop. Responses were

made with a Kinesis pedal that the patients held in their hands. Participants pressed buttons with

their thumbs to respond.

For the 3-tone oddball task, each trial consisted of one of the following sounds being played: a

standard tone that occurred frequently (500 Hz: 50% of trials), a target sound that occurred

infrequently (the word "Go": 30% of trials; this trial type was used to maintain attention), or a

novel tone (unique birdcall: 20% of trials; **Figure 3A**). Patients were instructed to press with

their left hand (typically thumb) if they heard the word "Go," and to press with their right

hand/thumb if they heard anything else. We focused our analyses on the standard and the novel

tones, as patients responded with their right hand for both the standard tones and the novel

oddball tones, facilitating a direct comparison of novelty induced motor initiation. Patients

practiced the task with 30 trials preoperatively and again on the morning of surgery. The

intraoperative experiment was conducted after the clinically-necessary microelectrode recording

electrodes were confirmed to be in the STN based on clinical recordings.

Intraoperative neurophysiology recordings

Bilateral DBS electrodes were implanted sequentially during a single stereotactic procedure.

Participants received short-acting pain relief and sedative medications such as remifentanil

and/or dexmedetomidine; these were stopped >1 hour prior to the task for participants to be

maximally awake for necessary clinical testing and participation in research. Patients underwent

standard bilateral DBS lead implantations using indirect framed stereotactic STN targeting,

refined by microelectrode recordings from 0.4 to 0.8 mΩ tungsten electrodes (Alpha-Omega,

Inc). Three simultaneous microelectrode recording tracks were used, consisting of anterior,

middle, and posterior trajectories, separated by 2 mm center-to-center from an entry point near

the coronal suture. STN margins were defined by the functional and electrical properties from

these microelectrode recordings, in line with standard clinical practice. Because microelectrode

recordings were clinically necessary, participants were not exposed to extra electrode

penetrations to participate in the study.

In addition to STN recordings, scalp EEG was also simultaneously recorded in the operating

room during the oddball task. Prior to the start of surgery, three midfrontal EEG electrodes

(Cadwell Industries, Inc) were placed over frontal locations on the forehead at the hairline with

right and left electrodes approximating F1 and F2 and midfrontal lead approximating Fz (**Figure**

4A). A reference electrode was placed on the left mastoid. We recorded EEG because this

location indexes midfrontal ~4-Hz rhythms and is in line with prior work (K.-H. Chen et al.,

2016; Kelley et al., 2018; Narayanan et al., 2013; Singh et al., 2021). EEG recordings were

recorded at 24 kHz and subsequently downsampled to 500 Hz. Signal was filtered below 50 Hz

with a low-pass filter, followed by a 1-Hz high-pass filter. Microelectrode recordings were

sampled at 24 kHz, amplified, and filtered for single neurons (1–8 kHz) and STN local field

potentials (<200 Hz). Neurophysiological and behavioral data were acquired simultaneously

using a Tucker-Davis Technologies multi-channel data acquisition system.

After the experiments, Offline Sorter (Plexon) was used to analyze STN activity and remove

artifacts. Spike activity was analyzed for all cells that fired at rates above 0.1 Hz. Principal

component analysis (PCA) and waveform shape were used for spike sorting. Single units were

defined as those 1) having a consistent waveform shape, 2) being a separable cluster in PCA

space, and 3) having a consistent refractory period of at least 2 ms in interspike interval

histograms. Isolated STN single units were included for analyses if 1) the recording location was

confirmed by the neurosurgeon to be in the STN based on clinically-necessary and observable

STN firing patterns during surgery, and 2) the unit was held throughout the duration of the task.

Experimental Design and Statistical Analyses

All data and statistical approaches were reviewed by the UI Biostatistics, Epidemiology, and

Research Design Core at the Institute for Clinical and Translational Science. All code and data

are available at narayanan.lab.uiowa.edu. For demographic summary statistics, medians and

interquartile ranges were calculated for continuous measures. Counts and percentages were

calculated for categorical measures. EEG participant demographics were stratified by group and

gender. Intraoperative patient-volunteer demographics were stratified by gender. We used the

nonparametric Wilcoxon rank sum test to test for differences in age, education, and MoCA

between strata. For behavioral data, we used a linear mixed-effects model with random intercept

for participants' response time to capture the effects of predictor variables (tone type: standard vs

novel tones and group: PD vs control) on response time. For behavioral data for the

intraoperative task, we used the nonparametric Wilcoxon rank sum test to test for differences

between different sound stimuli on accuracy and reaction time. For EEG data, we used a linear

mixed-effects model with random intercept for participants' time-frequency power to compare

average time-frequency-power between groups. The fixed-effect predictors included group, tone

type, MoCA score, and Unified Parkinson's Disease Rating Scale (UPDRS) Part 3 as control

variables and time-frequency power from the *a priori* region of interest (ROI) as the outcome

variable. We set our alpha for these analyses at 0.05. The MoCA score was included in the model

as a covariate given that it was different between groups. UPDRS Part 3 was included as a

covariate for PD subjects in order to account for degree of motor impairment. These analyses

were conducted in R.

Neuronal modulations were defined via a generalized linear model (GLM) for each neuron,

where the outcome variable was firing rate, and the predictor variable was cue or response,

consistent with past work (Emmons et al., 2019, 2020). For each neuron, we constructed a model

in which the outcome variable was the firing rate binned at 0.1 seconds, and the predictor

variable was either cue or response. Main effects comparisons were made stratified by neuron.

Across the ensemble, p values were corrected via Benjamini-Hochberg false-discovery-rate

(FDR; Benjamini & Yekutieli, 2001), with p values <0.05 considered significant modulation for

each neuron. We also analyzed neuronal patterns using PCA of peri-event histograms around

response binned at 0.1 second with kernel-density estimates (bandwidth 0.25). Finally, we

analyzed midfrontal EEG-STN coherence with the Neurospec 2.0 coherence toolbox using the

sp2a_m1 function with a segment power of 9 and a window size of 250 ms. To compare

coherence across neurons with different spike rates and trial numbers, spike-field coherence was

scaled such that the raw spike-field coherence of each neuron was divided by the 95%

confidence interval for each neuron. According to this metric, a "1" represents coherence at the

95% confidence interval. These analyses were performed in MATLAB.

RESULTS

Demographics

Table 1 shows demographics for the EEG study and Table 2 shows demographics for the

intraoperative study. For EEG participants, demographic results were stratified by group and

gender. Wilcoxon test results showed that the groups were not significantly different in terms of

age (p=0.38, effect size r=0.10) or years of education (p=0.09, effect size r=0.20). The MoCA

score was higher for the control subjects compared to PD subjects (p<0.01, effect size r=0.36),

thus we accounted for MoCA in our analyses.

PD patients had slowed reaction-times during novelty-related distraction

We studied how novelty engaged cortical cognitive control mechanisms by recording EEG

during the cross-modal oddball distractor task (**Figure 1**). This task asks participants to respond

to a target cue (arrow) following either a familiar stimulus (600-Hz tone) or a novel stimulus

consisting of complex and unique auditory features.

Accuracy was high overall (Median (1st Quartile–3rd Quartile); Control: standard 99.4%

(99%-100%), novel 100% (100%-100%); PD: standard 98.2% (93.3%-99.4%), novel 97.9% (92-

100%)), although control participants were more accurate than PD patients ($F_{(1,73)}$=11.1,

p=0.001). There was no main effect of tone type ($F_{(1,73)}$=0.39, p=0.54), and no interaction

between group and tone type ($F_{(1,73)}$=1.61, p=0.21). Regarding response speed, on trials with a

distracting novel tone, control subjects responded approximately the same as on trials with a

standard tone (Median (1st Quartile–3rd Quartile); Control: novel 454 ms (426-514), standard 454

ms (426-531 ms)), whereas PD patients responded more slowly compared to trials with a

standard tone (PD: novel 505 ms (439-571 ms), standard 462 ms (416-515 ms); **Figure 1B**).

There was a significant main effect of tone type ($F_{(1,73)}$=22.4 p<0.001), but this appears to be

driven by the effect within the PD group. There was no significant effect of Control vs. PD on

reaction time ($F_{(1,73)}$=1.0, p=0.32); however, there was a significant interaction between the

effects of group and tone type on reaction time ($F_{(1,73)}$=6.0, p=0.02). Post-hoc analyses revealed

that within the PD group, there was a significant difference between reaction times for the tone

types (t(73)=-6.2, p<0.0001). No other post hoc contrast was significant.

Finally, we found that novelty-related slowing was not related to cognitive function, as

measured by MoCA (Spearman ρ=-0.15, p=0.19; **Figure 1C**), or motor function, as measured by

UPDRS Part 3 in PD patients (Spearman ρ=0.10, p=0.50; **Figure 1D**). Taken together, these data

suggest that PD patients were more distracted by novel stimuli than controls.

PD patients had attenuated ~4-Hz midfrontal rhythms

Next, we examined cortical correlates of cognitive control in response to novelty. We found

that ERPs were distinct between controls and PD patients, with PD patients showing smaller

negative deflections around 100 ms and 300 ms for standard tone trials (**Figure 2A**, top panel)

and smaller negative deflections around 100 ms and 200 ms for novel tone trials (**Figure 2A**,

bottom panel). In all participants, novel tone trials showed a late (>300 ms) positive component

compared to the leveling off observed in the standard tone trials (e.g. the "P3a"). This positive

deflection appeared to be higher for PD subjects than control subjects; however, the possibility

of an underlying low-frequency contribution across the entire ERP in PD subjects suggests that

time-frequency analysis might be more effective in isolating unique temporal events that differ

between groups.

We examined power from an *a priori* tf-ROI (4-8 Hz for 300-400 ms post-tones; blue box in

Figure 2D). Our linear mixed-effects model revealed a significant effect of tone type on this tf-

ROI ($F(1,73)=51.1$, $p<0.001$), with novel trials showing higher 4-8 Hz band power. However,

there were no significant main effects of group, MoCA, or UPDRS Part 3, and there was not a

significant interaction between group and trial condition ($F(1,73)=3.7$, $p=0.06$). We found that

there was no significant relationship between midfrontal ~4-Hz power and novelty-related

slowing (Spearman $\rho=-0.16$, $p=0.17$), indicating that average ~4-Hz power across novelty trials

may not directly translate to performance on this task. Further, ~4-Hz power in our tf-ROI on

novel trials did not correlate significantly with MoCA (Spearman $\rho=0.13$, $p=0.28$) or UPDRS

Part 3 (for PD group only; Spearman $\rho=-0.24$, $p=0.10$), suggesting that midfrontal activity

during this task may not be associated with general cognitive functioning or affected by PD

motor symptoms. These data are convergent with our past work showing that that midfrontal ~4-

Hz rhythms signify cues and are distinct in PD, but are not specifically linked with response

control.

Human single STN neurons were modulated around responses to novelty

To comprehensively investigate the cortico-striatal pathways underling novelty-modulated

responses, we leveraged intraoperative neurophysiology to record from single STN neurons.

Participants performed the task with a median accuracy of 87% (Q1-Q3: 82%-92%); there was

no difference in median accuracy or reaction time between tone types (accuracy: standard 86%

(77%-93%), novel 90% (80%-95%), *p*=0.21, effect size r=0.71; reaction time: standard 877 ms

(721-1024 ms), novel 875 ms (761-936 ms), p=0.99, effect size r=0.71; **Figure 3B**).

With these 18 participants, we also performed intraoperative scalp EEG (**Figure 4A**).

Although these recording locations were anterior to the midfrontal areas which showed peak ~4-

Hz activity in the EEG experiment outside the operating room (i.e. **Figure 2E)**, we were able to

capture midfrontal cue-related ~4-Hz activity during trials with standard tones from the

intraoperative participants (**Figure 4B**). We did not identify strong novelty-related midfrontal

~4-Hz activity (**Figure 4C**), although a poorer signal-to-noise ratio due to intraoperative

limitations may have particularly affected responses to novel stimuli wherein trial numbers were

lower.

We identified 54 well-isolated STN neurons from 17 of the 18 participants (1 did not have

STN neuronal recordings; **Figure 4D-E**). We defined modulations via neuron-by-neuron GLMs

of firing rate vs. all cues and responses at a trial-by-trial level. We found that 5 of the 54 neurons

(9%) were significantly modulated around the cue. By contrast, we found significantly more

neurons (17 of 54, 32%) with response-related activity (χ^2=8.2; p=0.004; **Figure 4E & Figure**

5A–B). To further characterize the patterns of neuronal activity, we utilized PCA, a proven, data-

driven technique, to characterize neuronal activity (Chapin & Nicolelis, 1999; Narayanan &

Laubach, 2009). We found that the first component (principal component 1; PC1) explained

38.5% of variance and appeared to be modulated around response; PC2 explained 28.5% of

variance and was modulated prior to response (**Figure 5C-D**). These results provide convergent

evidence from two distinct analytic approaches (GLM and PCA) that STN neurons were strongly

response modulated.

Next, we were interested in how response-related activity was modulated by novel cues.

Strikingly, we found that nearly all (16 of 17; 94%; **Figure 6B**) response-related neurons had

differential activity with novel cues as quantified by an FDR-corrected neuron-by-neuron GLM

of event type on response-related activity. Thus, these data indicate that STN response-related

activity was powerfully modulated by novelty.

Response-related STN neurons had low-frequency coherence with midfrontal EEG

Finally, we were interested in the relationship between STN neuronal activity with scalp

EEG-related modulations. We examined spike-field coherence, through which we could link the

activity of midfrontal scalp electrodes to STN spiking (Kim & Narayanan, 2019; Narayanan et

al., 2013; Parker et al., 2014, 2015). For some STN neurons, we noticed that spikes fired in-

phase with low-frequency scalp EEG activity (**Figure 7A**). Among response-related STN

neurons, we noticed significant low-frequency spike-field scalp-STN coherence after the cue

around the time of responses (**Figure 7B**). Around responses, low-frequency coherence was

stronger for response-related neurons compared to non-response-related neurons (**Figure 7C vs 7B**; scaled spike-field coherence for response neurons: 1.1(0.5-1.8) vs. non-response neurons 0.3(0.1-0.7); Wilcoxon rank sum p=0.003; effect size r=0.40). These data provide evidence that STN response-related activity could reflect frontal top-down novelty-induced orienting and control. Together, these data support the idea that novelty can trigger midfrontal low-frequency rhythms, which in turn engage subcortical STN circuits involved in response control.

Discussion

In this study, we combined scalp-based EEG and intraoperative recordings to examine novelty-related brain activity in frontal-subcortical circuits. We report three main results: 1) Both PD and control participants engaged low-frequency power in response to novel stimuli, but, compared to control participants, PD patients had an attenuated midfrontal response ~4-Hz (although this activity was not linked with novelty-related slowing or cognitive status); 2) STN neurons were strongly modulated by novel responses, and 3) STN response-related neurons exhibited low frequency coherence with midfrontal EEG activity. Our findings provide a better understanding of midfrontal-STN cognitive control systems in PD and neuronal responses to novel information.

Midfrontal ~4-Hz rhythms are prominently triggered by events that recruit cognitive control such as errors, conflict, feedback, and novelty (Cavanagh & Frank, 2014; Cohen & Donner, 2013; Singh et al., 2018). Our past work found that this response is highly conserved in humans and rodents, and cortical dopamine is critical for coupling of cortical neurons with low-frequency oscillations (Narayanan et al., 2013; Parker et al., 2015). Taken together, our data support the

idea that midfrontal low-frequency rhythms recruit neuronal resources to respond to new

information.

Our work here is in line with past studies from our group and others showing that PD patients

have attenuated midfrontal rhythms between 1-8 Hz (Cavanagh et al., 2018; K.-H. Chen et al.,

2016; Giller et al., 2020; Singh et al., 2018, 2021; Solís-Vivanco et al., 2018). This line of work

includes data from the Simon reaction-time task (Singh et al., 2018), interval timing tasks (Singh

et al., 2021), and working memory manipulations (Itthipuripat et al., 2013). Our past work has

found that midfrontal 1-4 Hz frequencies are attenuated in PD (Kim et al., 2017; Parker et al.,

2014) and strongly correlated with cognitive function during an interval timing task (Singh et al.,

2021), although in the present study, these bands were novelty-specific but not reliably distinct

between control and PD patients (**Figure 2**, bottom panel and 2D). By contrast, we found that

novelty-triggered midfrontal 4-8 Hz oscillations were attenuated in PD but did not correlate with

novelty-related slowing. There are multiple generators of mid-frontal low-frequency activity

(Zuure et al., 2020), and it is possible that lower frequencies (1-4 Hz) are more commonly

evoked by an orienting response (Cavanagh et al., 2018; Parker et al., 2015; Wessel et al., 2016),

and higher frequencies (5-8 Hz) represent performance-related cognitive control (Cavanagh &

Frank, 2014; Cohen & Donner, 2013; Singh et al., 2018; Töllner et al., 2017; Zuure et al., 2020).

We found that response-modulated STN neuronal activity was coherent with these lower

frequencies 1-4 Hz. Our work is consistent with prior studies of STN neural activity; indeed,

Bockova and colleagues (2011) found that the STN was modulated by distractor stimuli with a

positive ERP peak around 200 ms, and Brittain and colleagues (2012) found that response

inhibition is associated with STN activity, which serves to suppress motor-related output from

the basal ganglia. The low-frequency coherence we found is consistent with recent studies that

revealed that oscillations in the STN during conflict are driven by midfrontal 2-7 Hz activity

(Cavanagh et al., 2011, 2017; Zavala et al., 2014; Zeng et al., 2021). It is possible that multiple

cortical 4-8 Hz theta features could couple with lower-frequency (~2 Hz) STN oscillations,

resulting in a novelty-induced slowing mechanism. This could suggest that the cortex has more

sophisticated information integration whereas the STN could more simply carry out reactive

responses to novelty (e.g. stop & orient). Multiple studies have indirectly demonstrated

connectivity between the frontal cortex and the STN in a pathway known as the hyperdirect

pathway (Brunenberg et al., 2012; Haynes & Haber, 2013). This pathway was described by

Nambu and colleagues (2002) as a cortico-subthalamo-pallidal pathway involved in motor

control and output. Intraoperative stimulation studies have demonstrated monosynaptic

hyperdirect connectivity (W. Chen et al., 2020; Kelley et al., 2018; Miocinovic et al., 2018;

Walker et al., 2012). It is possible that STN-midfrontal spike-field coherence is a result of either

hyperdirect connectivity or feed-forward activity via the basal ganglia indirect pathway, or both.

Experimentally teasing these contributions apart may require experiments in nonhuman systems

in which basal ganglia pathways can be reversibly inactivated while recording from STN

neurons.

Our data help us understand how PD impacts novelty-related responses. Previous studies have

shown that PD patients present with impaired habituation over repeated stimulus presentations,

and that attenuated midfrontal theta activity is related to the rate of startle habituation (Cavanagh

et al., 2018; K.H. Chen et al., 2016). Importantly, EEG habituation to novelty can effectively

classify PD patients (Cavanagh et al., 2018). In our study, PD patients were not as successful as

controls at quickly reorienting to the task at hand when presented with distracting novel stimuli.

We found that PD patients also demonstrated reduced novelty-related frontal theta oscillations

compared to controls (although theta power was not directly related to behavioral response

speed). These findings suggest that structural and functional changes related to PD affect the

circuitry that evaluates and responds to novel stimuli.

Our work has several limitations. First, intraoperative recordings present many challenges,

including a lack of control over the experimental environment. In particular, our intracranial

EEG used adhesive electrodes and the electrode placement was somewhat anterior to the peak of

midfrontal ~4-Hz activity our whole scalp EEG experiment captured (i.e. **Figure 2E**). These

factors may have affected our signal-to-noise ratio, particularly on novel trials with low trial

numbers. Because STN single-unit recordings occur in PD patients with attenuated novelty-

triggered midfrontal ~4-Hz rhythms, it is unknown how our STN findings generalize to non-PD

patients or patients with other brain diseases that disrupt frontostriatal circuits. Finally, our *a*

priori approach focused on a midfrontal ~4-Hz region-of-interest, which did not perfectly

overlap with data-driven novelty-related differences in controls vs. PD patients.

Overall, we found evidence for a ~4-Hz stimulus-response arc between the frontal cortex and

STN during novelty. Our findings align with recent work demonstrating clinical relevance of

midfrontal ~4-Hz activity in PD (Singh et al., 2021; Solís-Vivanco et al., 2018). Further, small

studies have revealed that STN low-frequency (~4–5 Hz) stimulation can beneficially impact

conflict (Scangos et al., 2017) and interval timing (Kelley et al., 2018). These findings in humans

are further supported by evidence from animal models of PD showing that highly-specific low-

frequency stimulation can improve interval timing (Kim et al., 2017; Kim & Narayanan, 2019).

It is possible, though it remains to be demonstrated in humans, that stimulating the STN at 4 Hz

resolves impaired midfrontal ~4-Hz oscillatory activity. Restoring behaviorally-relevant ~4-Hz

oscillations may contribute directly to improved cognitive performance. Future research should

systematically evaluate this mechanism in humans.

Overall, the current studies show that novelty-related distraction is more evident in PD

subjects compared to controls, that PD patients engage less novelty-related theta power, that

neurons in the STN are modulated around responses to novelty, and that neurons in the STN and

the midfrontal cortex have low-frequency coherence around responses to novelty. Our data

illuminate how novelty modulates ~4-Hz rhythms in hyperdirect circuits, which may provide

insight into neuronal responses to novelty and PD-related changes in cognitive control. This

work may be significant in the development of novel biomarkers or treatments for PD and other

brain diseases degrading basal ganglia circuits.

Contributions

RC, NN, JFC, and JG conceived and designed experiments; RC and AE recorded EEG data; RC
and JG performed intraoperative recordings; AS, RC, JFC, and NN wrote analysis code; RC,
JFC, NN, and JG wrote the paper; and JB independently checked the data.

References

Andrés, P., Parmentier, F. B. R., & Escera, C. (2006). The effect of age on involuntary capture of attention by irrelevant sounds: A test of the frontal hypothesis of aging. *Neuropsychologia*, *44*(12), 2564–2568. https://doi.org/10.1016/j.neuropsychologia.2006.05.005

Benjamini, Y., & Yekutieli, D. (2001). The Control of the False Discovery Rate in Multiple Testing under Dependency. *The Annals of Statistics*, *29*(4), 1165–1188. https://www.jstor.org/stable/2674075

Bočková, M., Chládek, J., Jurák, P., Halámek, J., Baláž, M., & Rektor, I. (2011). Involvement of the subthalamic nucleus and globus pallidus internus in attention. *Journal of Neural Transmission*, *118*(8), 1235–1245. https://doi.org/10.1007/s00702-010-0575-4

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Brittain, J. S., Watkins, K. E., Joundi, R. A., Ray, N. J., Holland, P., Green, A. L., Aziz, T. Z., & Jenkinson, N. (2012). A role for the subthalamic nucleus in response inhibition during conflict. *Journal of Neuroscience*, *32*(39), 13396–13401. https://doi.org/10.1523/JNEUROSCI.2259-12.2012

Brunenberg, E. J. L., Moeskops, P., Backes, W. H., Pollo, C., Cammoun, L., Vilanova, A., Janssen, M. L. F., Visser-Vandewalle, V. E. R. M., ter Haar Romeny, B. M., Thiran, J. P., & Platel, B. (2012). Structural and resting state functional connectivity of the subthalamic nucleus: Identification of motor stn parts and the hyperdirect pathway. *PLoS ONE*, *7*(6). https://doi.org/10.1371/journal.pone.0039061

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in Cognitive Sciences*, *18*(8), 414–421. https://doi.org/10.1016/j.tics.2014.04.012

Cavanagh, J. F., Kumar, P., Mueller, A. A., Richardson, S. P., & Mueen, A. (2018). Diminished

EEG habituation to novel events effectively classifies Parkinson's patients. *Clinical Neurophysiology*, *129*(2), 409–418. https://doi.org/10.1016/j.clinph.2017.11.023

Cavanagh, J. F., Sanguinetti, J. L., Allen, J. J. B., Sherman, S. J., & Michael, J. F. (2017). The Subthalamic Nucleus Contributes to Post-error Slowing. *Washington, DC: APA, Guideline Development Panel for the Treatment of Posttraumatic Stress Disorder in Adults.*, 139. https://doi.org/10.1162/jocn_a_00659

Cavanagh, J. F., Wiecki, T. V., Cohen, M. X., Figueroa, C. M., Samanta, J., Sherman, S. J., & Frank, M. J. (2011). Subthalamic nucleus stimulation reverses mediofrontal influence over decision threshold. *Nature Neuroscience*, *14*(11), 1462–1467. https://doi.org/10.1038/nn.2925

Chapin, J. K., & Nicolelis, M. A. L. (1999). Principal component analysis of neuronal ensemble activity reveals multidimensional somatosensory representations. *Journal of Neuroscience Methods*, *94*(1), 121–140. https://doi.org/10.1016/S0165-0270(99)00130-2

Chen, K.-H., Okerstrom, K. L., Kingyon, J. R., Anderson, S. W., Cavanagh, J. F., & Narayanan, N. S. (2016). Startle Habituation and Midfrontal Theta Activity in Parkinson Disease. *Journal of Cognitive Neuroscience*, *10*(4), 431–441. https://doi.org/10.1162/jocn_a_01012

Chen, W., de Hemptinne, C., Miller, A., Leibbrand, M., Little, S., Lim, D., Larson, P., & Starr, P. (2020). Prefrontal-Subthalamic Hyperdirect Pathway Modulates Movement Inhibition in Humans. *Neuron*, *106*(4), 579-588.e3. https://doi.org/10.1016/j.neuron.2020.02.012

Cohen, M. X. (2014). *Analyzing Neural Time Series Data: Theory and Practice*. The MIT Press.

Cohen, M. X., & Donner, T. H. (2013). Midfrontal conflict-related theta-band power reflects neural oscillations that predict behavior. *Journal of Neurophysiology*, *110*(12), 2752–2763. https://doi.org/10.1152/jn.00479.2013

Debener, S., Makeig, S., Delorme, A., & Engel, A. K. (2005). What is novel in the novelty

    oddball paradigm? Functional significance of the novelty P3 event-related potential as

    revealed by independent component analysis. *Cognitive Brain Research*, *22*(3), 309–321.

    https://doi.org/10.1016/j.cogbrainres.2004.09.006

Emmons, E., Kennedy, M., Kim, Y., & Narayanan, N. (2019). Corticostriatal stimulation

    compensates for medial frontal inactivation during interval timing. *Scientific Reports*, *9*(1),

    1–9. https://doi.org/10.1038/s41598-019-50975-7

Emmons, E., Tunes-Chiuffa, G., Choi, J., Bruce, R., Weber, M., Kim, Y., & Narayanan, N.

    (2020). Temporal Learning Among Prefrontal and Striatal Ensembles. *Cerebral Cortex*

    Communications, *1*(1), 1–12. https://doi.org/10.1093/texcom/tgaa058

Escera, C., Alho, K., Winkler, I., & Naatanen, R. (1998). *Neural Mechanisms of Involuntary*

    Attention. 590–604.

Fan, J. (2014). An information theory account of cognitive control. *Frontiers in Human*

    Neuroscience, *8*(SEP), 1–16. https://doi.org/10.3389/fnhum.2014.00680

Giller, F., Bensmann, W., Mückschel, M., Stock, A. K., & Beste, C. (2020). Evidence for a

    causal role of superior frontal cortex theta oscillations during the processing of joint

    subliminal and conscious conflicts. *Cortex*, *132*, 15–28.

    https://doi.org/10.1016/j.cortex.2020.08.003

Haynes, W. I. A., & Haber, S. N. (2013). The organization of prefrontal-subthalamic inputs in

    primates provides an anatomical substrate for both functional specificity and integration:

    Implications for basal ganglia models and deep brain stimulation. *Journal of Neuroscience*,

    33(11), 4804–4814. https://doi.org/10.1523/JNEUROSCI.4674-12.2013

Itthipuripat, S., Wessel, J. R., & Aron, A. R. (2013). Frontal theta is a signature of successful

working memory manipulation. *Experimental Brain Research*, *224*(2), 255–262.

https://doi.org/10.1007/s00221-012-3305-3

Kelley, R., Flouty, O., Emmons, E. B., Kim, Y., Kingyon, J., Wessel, J. R., Oya, H., Greenlee, J.

D., & Narayanan, N. S. (2018). A human prefrontal-subthalamic circuit for cognitive

control. *Brain*, *141*(1), 205–216. https://doi.org/10.1093/brain/awx300

Kim, Y., Han, S. W., Alberico, S. L., Ruggiero, R. N., De Corte, B., Chen, K.-H., & Narayanan,

N. S. (2017). Optogenetic Stimulation of Frontal D1 Neurons Compensates for Impaired

Temporal Control of Action in Dopamine-Depleted Mice. *Current Biology*, *27*(1), 39–47.

https://doi.org/10.1016/j.cub.2016.11.029

Kim, Y., & Narayanan, N. S. (2019). Prefrontal D1 Dopamine-Receptor Neurons and Delta

Resonance in Interval Timing. *Cerebral Cortex*, *29*(5), 2051–2060.

https://doi.org/10.1093/cercor/bhy083

Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3? *Perception 36*

ECVP Abstract Supplement.

Lange, F., Seer, C., Loens, S., Wegner, F., Schrader, C., Dressler, D., Dengler, R., & Kopp, B.

(2016). Neural mechanisms underlying cognitive inflexibility in Parkinson's disease.

Neuropsychologia, *93*(May), 142–150.

https://doi.org/10.1016/j.neuropsychologia.2016.09.021

Mikell, C. B., Sheehy, J. P., Youngerman, B. E., McGovern, R. A., Wojtasiewicz, T. J., Chan, A.

K., Pullman, S. L., Yu, Q., Goodman, R. R., Schevon, C. A., & McKhann, G. M. (2014).

Features and timing of the response of single neurons to novelty in the substantia nigra.

Brain Research, *1542*, 79–84. https://doi.org/10.1016/j.brainres.2013.10.033

Miocinovic, S., de Hemptinne, C., Chen, W., Isbaine, F., Willie, J., Ostrem, J., & Starr, P.

(2018). Cortical potentials evoked by subthalamic stimulation demonstrate a short latency

hyperdirect pathway in humans. *Journal of Neuroscience*, *38*(43), 9129–9141.

https://doi.org/10.1523/JNEUROSCI.1327-18.2018

Nambu, A., Tokuno, H., & Takada, M. (2002). Functional significance of the cortico-

subthalamo-pallidal "hyperdirect" pathway. *Neuroscience Research*, *43*(2), 111–117.

https://doi.org/10.1016/S0168-0102(02)00027-5

Narayanan, N. S., Cavanagh, J. F., Frank, M. J., & Laubach, M. (2013). Common medial frontal

mechanisms of adaptive control in humans and rodents. *Nature Publishing Group*, *16*(12),

1888–1895. https://doi.org/10.1038/nn.3549

Narayanan, N. S., & Laubach, M. (2009). Dynamic imaging of brain function. In F. Hyder (Ed.),

Dynamic Brain Imaging: Multi-Modal Methods and In Vivo Applications (pp. 135–165).

https://doi.org/10.1007/978-1-59745-543-5_1

Nolan, H., Whelan, R., & Reilly, R. B. (2010). FASTER: Fully Automated Statistical

Thresholding for EEG artifact Rejection. *Journal of Neuroscience Methods*, *192*(1), 152–

162. https://doi.org/10.1016/j.jneumeth.2010.07.015

Parker, K. L., Chen, K.-H., Kingyon, J. R., Cavanagh, J. F., & Narayanan, N. S. (2014). D1-

dependent 4 Hz oscillations and ramping activity in rodent medial frontal cortex during

interval timing. *Journal of Neuroscience*, *34*(50), 16774–166783.

https://doi.org/10.1523/JNEUROSCI.2772-14.2014

Parker, K. L., Chen, K.-H., Kingyon, J. R., Cavanagh, J. F., & Narayanan, N. S. (2015). Medial

frontal ∼4-Hz activity in humans and rodents is attenuated in PD patients and in rodents

with cortical dopamine depletion. *Journal of Neurophysiology*, *114*(2), 1310–1320.

https://doi.org/10.1152/jn.00412.2015

Parmentier, F. B. R., Elford, G., Escera, C., Andrés, P., & Miguel, I. S. (2008). The cognitive

locus of distraction by acoustic novelty in the cross-modal oddball task. *Cognition*, *106*(1),

408–432. https://doi.org/10.1016/j.cognition.2007.03.008

Parmentier, F. B. R., & Hebrero, M. (2013). Cognitive control of involuntary distraction by

deviant sounds. *Journal of Experimental Psychology: Learning Memory and Cognition*,

39(5), 1635–1641. https://doi.org/10.1037/a0032421

Parmentier, F. B. R., Maybery, M. T., & Elsley, J. (2010). The involuntary capture of attention

by novel feature pairings: A study of voice-location integration in auditory sensory

memory. *Attention, Perception, and Psychophysics*, *71*(7), 1439–1459.

https://doi.org/10.3758/APP

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. In *Spatial Vision* (Vol. 10, Issue 4, pp. 437–442).

https://doi.org/10.1163/156856897X00366

Polich, J. (2007). Updating P300: An Integrative Theory of P3a and P4b. *Clinical

Neurophysiology*, *118*(10), 2128–2148.

https://doi.org/10.1016/j.clinph.2007.04.019.Updating

Ranganath, C., & Rainer, G. (2003). Cognitive neuroscience: Neural mechanisms for detecting

and remembering novel events. *Nature Reviews Neuroscience*, *4*(3), 193–202.

https://doi.org/10.1038/nrn1052

Scangos, K. W., Carter, C. S., Gurkoff, G., Zhang, L., & Shahlaie, K. (2017). A pilot study of

subthalamic theta frequency deep brain stimulation for cognitive dysfunction in Parkinson's

disease. *Brain Stimulation*, *11*, 456–458. https://doi.org/10.1016/j.brs.2017.11.014

Singh, A., Cole, R. C., Espinoza, A. I., Evans, A., Cao, S., Cavanagh, J. F., & Narayanan, N. S.

(2021). Timing variability and midfrontal ~4 Hz rhythms correlate with cognition in

Parkinson's disease. *Npj Parkinson's Disease*, 2020.10.26.356154.

https://doi.org/10.1038/s41531-021-00158-x

Singh, A., Cole, R., Espinoza, A., Brown, D., Cavanagh, J., & Narayanan, N. (2020). Frontal

theta and beta oscillations during lower-limb movement in Parkinson's disease. *Clinical*

Neurophysiology, *131*(3), 694–702. https://doi.org/10.1016/j.clinph.2019.12.399

Singh, A., Richardson, S., Narayanan, N., & Cavanagh, J. (2018). Mid-frontal theta activity is

diminished during cognitive control in Parkinson's disease. *Neuropsychologia*, *117*(May),

113–122. https://doi.org/10.1016/j.neuropsychologia.2018.05.020

Sokolov, E. N. (1962). *Higher Nervous Functions: The Orienting Reflex*.

Solís-Vivanco, R., Rodríguez-Violante, M., Cervantes-Arriaga, A., Justo-Guillén, E., & Ricardo-

Garcell, J. (2018). Brain oscillations reveal impaired novelty detection from early stages of

Parkinson's disease. *NeuroImage: Clinical*, *18*(March), 923–931.

https://doi.org/10.1016/j.nicl.2018.03.024

Töllner, T., Wang, Y., Makeig, S., Müller, H. J., Jung, T.-P., & Gramann, K. (2017). Two

Independent Frontal Midline Theta Oscillations during Conflict Detection and Adaptation in

a Simon-Type Manual Reaching Task. *The Journal of Neuroscience*, *37*(9), 2504–2515.

https://doi.org/10.1523/jneurosci.1752-16.2017

Walker, H., Huang, H., Gonzalez, C., Bryant, J., Killen, J., Cutter, G., Knowlton, R.,

Montgomery, E., Guthrie, B., & Watts, R. (2012). Short latency activation of cortex during

clinically effective subthalamic deep brain stimulation for Parkinson's disease. *Movement*

Disorders, *27*(7), 864–873. https://doi.org/10.1002/mds.25025

Wessel, J. R., Jenkinson, N., Brittain, J. S., Voets, S. H. E. M., Aziz, T. Z., & Aron, A. R.

(2016). Surprise disrupts cognition via a fronto-basal ganglia suppressive mechanism.

Nature Communications, *7*, 1–10. https://doi.org/10.1038/ncomms11195

Zavala, B. A., Tan, H., Little, S., Ashkan, K., Hariz, M., Foltynie, T., Zrinzo, L., Zaghloul, K.

A., & Brown, P. (2014). Midline frontal cortex low-frequency activity drives subthalamic

nucleus oscillations during conflict. *Journal of Neuroscience*, *34*(21), 7322–7333.

https://doi.org/10.1523/JNEUROSCI.1169-14.2014

Zeng, K., Drummond, N. M. ., Ghahremani, A., Saha, U., Kalia, S. K., Hodaie, M., Lozano, A.

M., Aron, A. R., & Chen, R. (2021). Fronto-subthalamic phase synchronization and cross-

frequency coupling during conflict processing. *NeuroImage*, *238*(March), 118205.

https://doi.org/10.1016/j.neuroimage.2021.118205

Zink, C. F., Pagnoni, G., Chappelow, J., Martin-Skurski, M., & Berns, G. S. (2006). Human

striatal activation reflects degree of stimulus saliency. *NeuroImage*, *29*(3), 977–983.

https://doi.org/10.1016/j.neuroimage.2005.08.006

Zuure, M. B., Hinkley, L. B., Tiesinga, P. H. E., Nagarajan, S. S., & Cohen, M. X. (2020).

Multiple midfrontal thetas revealed by source separation of simultaneous MEG and EEG.

Journal of Neuroscience, *40*(40), 7702–7713. https://doi.org/10.1523/JNEUROSCI.0321-

20.2020

688
689

Figures and Tables

Figure 1. Novelty slows responses in control and PD. A) The cross-modal oddball distractor task consisted of trials with an arrow preceded by either a standard tone (80% of trials) or a novel tone (10%; of note, 10% of trials used a novel visual stimulus; these trials were not analyzed). The arrow required a left "p" or right "q" key press, based on the direction. B) Reaction time for Control and PD patients; each dot represents an individual's median RT and the thick horizontal bar represents group medians. Novel oddball stimuli disproportionately distracted PD subjects. Compared to Control participants, PD patients responded more slowly to the arrow on trials when novel distractors were presented relative to a standard tone. * indicates a significant main effect of tone type (standard vs. novel). # indicates a significant interaction between tone type

703   and group (control vs. PD) on reaction time via a linear mixed-effects model. C) Cognitive

704   function as measured by the Montreal Cognitive Assessment (MoCA) did not significantly

705   correlate with novelty-related slowing. D) UPDRS Part 3 did not significantly correlate with

706   novelty-related slowing in PD patients.

Figure 2. Novelty boosts midfrontal ~4-Hz power, which is attenuated in PD. A) Average

event-related potentials for trials with standard tones (top row) and novel tones (bottom row) for

controls (blue line) and PD patients (red line). B) Time-frequency power spectrograms for

standard tones (top panels), novel tones (middle panels), and novel – standard (bottom panels)

for Control and C) PD, and D) Subtraction of Control – PD. Black lines indicate areas of

significance as determined by cluster thresholding. The blue box represents the theta band

region-of-interest (ROI) (4-8 Hz) 300-400 ms after the cue. A linear mixed-effects model

revealed a significant main effect of tone type (novel vs. standard) on activity in this ROI, no

main effect of group (control vs. PD), and a trend towards interaction between tone type and

group (p<0.06). E) Topographical representation of the comparison of theta power for standard

721   tones (top panel) and novel tones (bottom panel) between controls and PD patients (Control –

722   PD). Data from 50 PD and 25 control participants.

Figure 3. Intraoperative recordings of novelty-related responses. A) Auditory 3-tone oddball

task performed by patients with Parkinson's disease in operating room during deep brain

stimulation surgery. Patients responded to standard (50% of trials) and novel tones (20% of

trials) with a right-hand response using a hand-held Kinesis pedal; an additional trial type with a

constant stimulus required a left response and was not analyzed. B) Reaction times for standard

and novel oddball tones; each dot represents individual median reaction times, and thicker

horizontal lines represent group median reaction times. No significant differences for reaction

times between groups were found by linear mixed-effect models. Data from 18 PD patient-

volunteers during intraoperative neurophysiology.

Figure 4. Recording simultaneously from frontal lobe and STN during surgery. A)

Schematic of surgical recording with three frontal scalp-based electrodes (center, right, and left

channels), and STN microelectrode recordings. B) Time-frequency plots of activity from the

center channel for standard and novel tones in the auditory 3-tone oddball task. Data from 18 PD

patient-volunteers undergoing DBS implantation surgery. D) Waveforms from a single STN

neuron indicating that neuronal waveforms of voltage vs. time were stable over ~6 minutes. E)

Response-locked peri-event raster of neuronal firing from an exemplar neuron; the top panel is a

peri-response raster, with each row representing one trial and each tick representing an action

potential. The bottom panel represents a peri-event histogram of firing rate around all responses.

Figure 5. STN neurons had prominent response modulations. A) Average z-scored activity of

all 54 STN neurons around responses. B) Trial-by-trial generalized linear models of firing rate

vs. cue and response revealed that STN neurons were more strongly modulated by response than

cue; *indicates significance via a $\chi 2$ test. C) Principal component analysis (PCA), a data-driven

approach to identify neuronal patterns across an ensemble, revealed that principal components 1

and 2 (PC1 and PC2) explained 64% of variance among STN ensembles. D) PC1 and PC2 were

modulated around response. Data from 54 STN neurons in 17 PD patient-volunteers.

Figure 6. Most response-related neurons in the subthalamic nucleus were modulated by

novelty. A) Example raster of neuronal spiking from one novelty-modulated neuron. B) We

recorded from 54 STN neurons. Of these, 17 STN neurons had response-modulations. 16 were

modulated by novelty and response (30% of the total), and 1 was modulated by response only

(2% of the total). Data recorded in 17 patient-volunteers.

Figure 7. Response-related STN neurons had low frequency coherence with frontal EEG.

A) An example time series from a single trial of center channel EEG and STN spiking activity;

cue is denoted via a green triangle, and response is denoted via a red triangle. This neuron

becomes coherent immediately with midfrontal low-frequency oscillations after the cue and prior

to response. B) Across all 17 response-modulated STN neurons, there was significantly more ~4-

Hz spike-field coherence with center-channel EEG electrodes than for C) 37 non-response

modulated STN neurons. Spike-field coherence is scaled with 1 representing the 95%

779   confidence-interval for coherence for comparison across neurons. Data from 54 STN neurons in

780   17 PD patient-volunteers.

782 **Table 1: EEG patient-volunteer demographics**

Group	Gender	N (%)	Age Median years (Q1-Q3)	Age range yrs	Right Handed (%)	Education median years (Q1-Q3)	MoCA median (Q1-Q3)	UPDRS Part 3 median (Q1-Q3)
Control	Total	25	70.0 (64.0-76.0)	54-85	22 (88%)	16.0 (16.0-18.0)	27.0 (26.0-28.0)	-
	F	12 (48%)	67.0 (60.3-72.3)	54-77	10 (83%)	16.0 (13.8-17.3	28.0 (27.0-28.3)	-
	M	13 (52%)	71.0 (64.0-78.0)	60-85	12 (92%)	17.0 (16.0-18.0)	26.0 (26.0-28.0)	-
PD	Total	50	68.0 (60.3 - 72)	52-86	45 (90%)	15.0 (13.0-18.0)	25.5 (22.3-27.0)	11.5 (7.0-17.0)
	F	19 (38%)	68.0 (66.0-71.5)	53-86	16 (84%)	16.0 (14.0-18.0)	27.0 (24.5-28.0)	12.0 (7.5-16.5)
	M	31 (62%)	66.0 (60.0-73.0)	52-84	29 (94%)	14.0 (12.0-17.8)	24.0 (22.0-26.0)	11.0 (6.0-17.0)
Control vs PD		p=0.41 (χ^2; M vs F;)	p=0.38 (Wilcoxon)		p=0.79 (χ^2)	p=0.09 (Wilcoxon)	p=0.002 (Wilcoxon)	

787   **Table 2. Intraoperative patient-volunteer demographics**

Gender	N	Age median yrs (Q1-Q3)	Age range yrs	Right Handed (%)	LEDD median (Q1-Q3)	Disease Duration median yrs (Q1-Q3)
Total	18	67.0 (62.3-68.5)	53-71	15 (83%)	975.0 (687.5-1175.0)	7.0 (3.5-10.8)
F	4 (22%)	62.5 (57.5-68.0)	53-71	3 (75%)	921.5 (819.8-987.5)	12.0 (10.8-13.5)
M	14 (78%)	67.0 (63.0-68.5)	53-71	12 (86%)	1018.5 (687.5-1218.8)	6.5 (3.0-7.0)

789