Structural disconnectome mapping of cognitive function in post-stroke patients

Knut K. Kolskår[1,2,3, *], Kristine M. Ulrichsen[1,2,3], Genevieve Richard[1], Erlend S. Dørum[1,2,3], Michel

Thiebaut de Schotten[4,5], Jaroslav Rokicki[1,2,14], Jennifer Monereo-Sánchez[1,6,7], Andreas Engvig[8,9],

Hege Ihle Hansen[10], Jan Egil Nordvik,[11,15] Lars T. Westlye[1,2,12], Dag Alnæs[1,13]

[1] NORMENT, Division of Mental Health and Addiction, Oslo University Hospital & Institute of

Clinical Medicine, University of Oslo, Norway

[2] Department of Psychology, University of Oslo, Norway

[3] Sunnaas Rehabilitation Hospital HT, Nesodden, Norway

[4] Brain Connectivity and Behaviour Laboratory, Sorbonne Universities, Paris, France

[5] Groupe d'Imagerie Neurofonctionnelle, Institut des Maladies Neurodégénératives- UMR 5293,

CNRS, CEA University of Bordeaux, Bordeaux, France

[6] Faculty of Health, Medicine and Life Sciences, Maastricht University, Netherlands

[7] Department of Radiology and Nuclear Medicine, Maastricht University Medical Center,

Netherlands

[8] Department of Nephrology, Oslo University Hospital, Ullevål, Norway

[9] Department of Medicine, Diakonhjemmet hospital, Norway

[10] Department of Neurology, Oslo University Hospital, Norway

[11] CatoSenteret Rehabilitation Center, Son, Norway

[12] KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, Norway

[13] Bjørknes College, Oslo, Norway

[14] Centre of Research and Education in Forensic Psychiatry, Oslo University Hospital, Oslo,

Norway

[15] Faculty of Health Sciences, Oslo Metropolitan University, Norway

* Corresponding author: Knut K. Kolskår (kolskaar@gmail.com), Sunnaas sykehus HF

Bjørnemyrveien 11, 1453 Bjørnemyr, Norway

Funding resources

The current study has been funded by European Research Council under the European Union's

Horizon 2020 research and innovation programme (ERC StG, grant agreement No. 818521/

802998); South-Eastern Norway Regional Health Authority, Grant/Award Numbers: 2014097,

2015044, 2015073, 2019107, 2020086; Norges Forskningsråd, Grant/Award Numbers: 249795,

262372; Norwegian ExtraFoundation for Health and Rehabilitation, Grant/Award Number:

2015/FO5146.

Conflict of Interest

We declare no competing financial interests. We obtained appropriate ethical approval from the

local ethics committee, and all procedures were in line with the declaration of Helsinki.

Open science and data availability statement

Relevant scripts and scrambled and anonymized data needed to reconstruct the reported analyses

and results are available on OSF (link below). While the sensitive nature of the data and our current

approvals do not allow for public sharing of real data, anonymized data may be available upon

request to the corresponding author following appropriate data transfer agreements.

OSF-repository: https://osf.io/juwhv/?view_only=1096394b4b2b4a51abceea5834d29969.

Keywords: DTI, stroke, prediction, cognitive sequelae

Abstract

Sequalae following stroke represents a significant challenge in current rehabilitation. The location

and size of focal lesions are only moderately predictive of the diverse cognitive outcome after

stroke. One explanation building on recent work on brain networks proposes that the cognitive

consequences of focal lesions are caused by damages to anatomically distributed brain networks

supporting cognition rather than specific lesion locations. To investigate the association between

post-stroke structural disconnectivity and cognitive performance, we estimated individual level

whole-brain disconnectivity probability maps based on lesion maps from 102 stroke patients using

normative data from healthy controls. Cognitive performance was assessed in the whole sample

using Montreal Cognitive Assessment, and a more comprehensive computerized test protocol was

performed on a subset (n=82). Multivariate analysis using Partial Least Squares on the

disconnectome maps revealed that higher disconnectivity in right insular and frontal operculum,

superior temporal gyrus and putamen was associated with poorer MoCA performance, indicating

that lesions in regions connected with these brain regions are more likely to cause cognitive

impairment. Furthermore, our results indicated that disconnectivity within these clusters was

associated with poorer performance across multiple cognitive domains. These findings demonstrate

that the extent and distribution of structural disconnectivity following stroke are sensitive to

cognitive deficits and may provide important clinical information predicting post stroke cognitive

sequalae.

Introduction

The magnitude and characteristics of cognitive impairments following stroke show substantial

individual differences across patients. While some patients show considerable deficits and

subsequent increased risk of neurodegeneration and dementia, others show no observable

impairments. This heterogeneity in cognitive deficits after stroke arise not only from differences in

the localization and extent of focal lesions, but also from potential secondary cascade effects in

terms of altered brain connectivity (Rehme & Grefkes, 2013), and alterations of the hierarchical

brain network structure (Stam, 2014). Currently, common predictors of cognitive deficits and

recovery include anatomical location, lesion severity, vascular risk factors, chronic brain pathology

and pre-stroke cognitive impairment (Bentley et al., 2014; Macciocchi, Diamond, Alves, & Mertz,

1998; Munsch et al., 2016; Pendlebury, 2009). However, recent studies indicate added predictive

value of connectivity-based measures, which capture perturbations of brain network connections or

dynamics beyond focal lesion site (Ktena Sofia et al., 2019; Lopes et al., 2021b).

  In the last decade, a large body of literature has characterized the brain as a complex

network consisting of nodes and their connections, collectively termed the brain connectome.

Building on functional imaging, a coarse parcellation of networks separate the brain in the cingulo-

opercular, frontoparietal, ventral attention and default mode networks, aiding cognitive control. The

cingulo-opercular network is associated with the ability to maintain focus, the ability to sustain top-

down cognitive control across cognitive tasks (Cai et al., 2016; Hilger, Ekman, Fiebach, & Basten,

2017; Uddin, Yeo, & Spreng, 2019; Wilk, Ezekiel, & Morton, 2012), whereas the frontoparietal

network is assumed to guide moment to moment-attentional control (Fassbender et al., 2006;

Majerus, Péters, Bouffier, Cowan, & Phillips, 2018). The ventral attention network serves

reorientation to relevant stimuli outside the scope of current attention (Maurizio Corbetta & Gordon

L Shulman, 2011; Vossel, Geng, & Fink, 2014), while default mode network is commonly

associated with self-referential internal activity, and is commonly downregulated during task

engagement (McKiernan, Kaufman, Kucera-Thompson, & Binder, 2003; Vatansever, Manktelow,

Sahakian, Menon, & Stamatakis, 2018). A central property of network nodes is how densely they

are connected. Highly connected nodes, or hubs, are critical for efficient information flow between

brain regions and are thought to play a crucial role in pivoting neural activity across distal brain

regions, allowing for the integration of information required to support cognitive operations (Cole

et al., 2013). Thus, focal damage to any part of a brain network might cause disruptions of distal but

intact brain regions. Indeed, even small lesions in a densely connected hub may cause connectome-

wide perturbations (Aben et al., 2019). Recent studies have therefore moved beyond traditional

lesion-symptom mapping, to include measures of network dysfunction to explain and predict stroke

sequelae (Lim & Kang, 2015; Ulrichsen et al., 2020). Recent findings using the lesion-network

mapping approach suggest that patients with overlapping symptoms have lesions in regions that are

functionally connected and that lesions to brain network hubs or their connecting white matter

pathways are associated with more symptoms (Fox, 2018). This network-based concept has also

been shown to apply to a wide range of brain disorders such as Alzheimer disease, schizophrenia

and multiple sclerosis (Crossley et al., 2014; Stam, 2014; van den Heuvel & Sporns, 2019).

Whereas association between functional connectivity following stoke has been investigated

in recent years (Klingbeil, Wawrzyniak, Stockert, & Saur, 2019; Lopes et al., 2021a; Ptak et al.,

2020), less is known regarding structural connectivity. The brain is connected by an intricate web of

white matter (WM) pathways, consisting of bundles of myelinated axons responsible for conveying

signals between brain regions, supporting functional networks. While measuring axonal

disconnections in the living human brain has remained a challenging task, a recent implementation

of diffusion tensor imaging (DTI) based tractography for assessing full-brain connection

probabilities has enabled opportunities for detailed estimation of the connections of one or several

lesions in individual patients (Foulon et al., 2018). Based on normative data from healthy controls,

voxel-wise disconnection probability maps for a particular patient can be derived based on a lesion

map. The extent and distribution of these disconnectivity maps have been shown to correlate with a

surrogate biomarker of neuronal damage in patients with MS (Rise et al., 2021) and shown promise

in predicting deficits following stroke (Salvalaggio, De Filippo De Grazia, Zorzi, Thiebaut de

Schotten, & Corbetta, 2020).

To test for associations between brain disconnection and cognitive performance, we used

Partial Least Squares analysis (PLS) to map common variance between voxel-wise structural

disconnection probability maps in 102 stroke survivors and their performance on the Montreal

Cognitive assessment (MoCA; Nasreddine et al., 2005). PLS (Krishnan, Williams, McIntosh, &

Abdi, 2011) is well-suited for investigating multivariate associations between neuroimaging

features and non-gaussian behavioral and clinical measures commonly obtained from stroke

patients (Blackburn, Bafadhel, Randall, & Harkness, 2013), and offers high sensitivity compared to

more traditional univariate approaches (Van Roon, Zakizadeh, & Chartier, 2014). Based on the

literature reviewed above, we hypothesized that patients with higher levels of stroke-induced brain

disconnectivity would show poorer cognitive performance, and further anticipated that this

association could not simply be explained by the size and location of the lesion itself. Since

previous studies and existing models of the distributed nature of the neuroanatomical basis of

cognitive functions are sparse, we remained agnostic about the anatomical distribution of the

associations with the disconnectome maps and performed an unbiased full brain analysis with

appropriate corrections for multiple comparisons.

Methods

The present cross-sectional study included participants previously described in detail (Dørum et al.,

2020; Kolskår et al., 2020; Richard et al., 2020; Ulrichsen et al., 2020). Briefly, the sample

comprised 102 stroke survivors from sub-acute (>24h post stroke and in clinical stable condition) to

chronic stage. Inclusion criteria were radiologically documented ischemic or hemorrhagic stroke,

and exclusion criteria were psychiatric conditions (bipolar disorder or schizophrenia), other

neurological conditions including known cognitive impairment pre-stroke, substance abuse, and

contraindications for MRI compatibility. All participants gave their written consent before

participating, and the study was approved by the Regional Committee for Medical and Health

Research Ethics South-East Norway (2014/694 and 2015/1282).

Table 1 displays sample demographics as well as time between stroke and MRI and

cognitive assessment.

	Mean	sd	min	max
Age	66.3	12.3	24	87
Sex	74.5% males			
Education (self-report, years)	14.6	3.3	7	30
Interval between stroke onset and MRI / MoCA assessment (days)	515	444	1	1399
Lesion size (2mm resolution)	7693	24723	5	30031
Number of lesions	3	2	1	Multiple*
Interval between stroke onset and CabPAD- assessment (days)	634	379	89	1428
NIHSS (score)	1.02	1.35	0	7
Hemispheric distribution Left:39 / Right:41 / Bilateral:19 / Subcortical: 2				
Time since stroke: Acute: 20 / Subacute: 25 / Chronic:57				

Table 1. Summary of clinical and demographic variables. NIHSS: National Institute of Stroke Scale. CapPad:

Cognitive Assessment at Bedside for iPad. Acute: Less than seven days since stroke. Subacute: between 7 and 180 days

*since stroke. Chronic: more than 180 days since stroke. * Based on radiological description. Number of lesions and*

lesion size is based on MNI- normalized lesion masks.

Cognitive assessment

Patients were assessed with MoCA (Nasreddine et al., 2005) at time of MRI scanning. At follow-up

(Table 1), a subsample (n=82) were also assessed using CabPad (Willer, Pedersen, Forchhammer,

& Christensen, 2016), a computerized test battery assessing a range of functions, including motor

speed (finger tapping) for dominant and non-dominant hand, verbal fluency (phonetic and semantic

word generation), attention span (symbol sequencing), working memory (reversed symbol

sequencing), a spatial Stroop / flanker task, spatial short-term memory and psychomotor speed

(symbol-digit coding task).

MRI acquisition

Patients were scanned at Oslo University Hospital on a 3T GE 750 Discovery MRI scanner

with a 32-channel head coil. T2-FLAIR images were acquired with the following parameters: TR:

8000 ms; TE: 127 ms, TI: 2240 ms; flip angle (FA): 90°; voxel size: 1x1x1 mm. T1-weighted scans

were collected using a 3D IR-prepared FSPGR (BRAVO) sequence (TR: 8.16 ms; TE: 3.18 ms; TI:

450 ms; FA: 12°; voxel size: 1×1×1 mm; FOV: 256 x 256, 188 sagittal slices) for co-registration.

Lesion delineation

Lesion delineation was performed on each participant's FLAIR image, using the semi-automated

Clusterize-Toolbox, implemented in MATLAB (de Haan, Clas, Juenger, Wilke, & Karnath, 2015),

guided by radiological descriptions. Normalization parameters were estimated by linear registration

of the structural T1-image to the MNI152-template and applied to the lesion masks using Flirt

(Jenkinson & Smith, 2001). Figure 1 displays overlap in lesion location across the sample.

Figure 1. Lesion overlap across 102 stroke patients. 12 transversal slices, with 5 mm thickness. Color scale indicates

number of participants overlapping. Z coordinates denotes transversal slices in the MNI152-coordinate system.

Group comparisons.

To asess goup differences in MoCA-score between the patients in acute, subacute and chronic

phase, a linear model was estimated with MoCA-score as dependent variable, and group as

independent variable. To assess impact of lesion load, a linear model was estimated with MoCA-

score as dependent variable and lesion size and number of lesions as independent variables.

Estimation of structural disconnectome maps

To estimate the extent of the structural disconnection for each patient, we employed an automated

tractography-based procedure (Foulon et al., 2018). Briefly, full-brain tractography data of 170

healthy controls from the 7T Human Connectome Project was used as a normative training set to

identify fibers passing through each lesion (Thiebaut de Schotten, Foulon, & Nachev, 2020). Using

affine and diffeomorphic deformations (Avants et al., 2011; Klein et al., 2009), individual patient's

lesion maps were registered to each control's native space and used as seeds for the probabilistic

tractography in Trackvis (Wang, Benner, Sorensen, & Wedeen, 2007). The resulting tractograms

were transformed to visitation maps, binarized, and registered to MNI152 space before a percentage

overlap map was produced by summarizing each point in the normalized healthy subject visitation

maps. The resulting disconnectome maps are whole-brain voxel-wise probability maps indicating

for each patient and for each voxel in the brain the probability that the voxels were disconnected.

Next, these individual-level disconnection maps were included in group-level analysis.

Statistical analysis

To assess associations between MoCA scores and disconnectome maps, we used non-rotated task-

based PLS using PLS Application (Krishnan, Williams Lj Fau - McIntosh, McIntosh Ar Fau - Abdi,

& Abdi, 2010) for MATLAB (MathWorks, 2018), entering female, age and MoCA score as

behavioral variables, with contrasts for mean effects for each (1 0 0; 0 1 0; 0 0 1). We performed

permutations (n=1000) to assess the significance of the estimated latent variables, while precision

was estimated using bootstrapping (n=1000) and used to calculate pseudo-z or bootstrap ratio brain

maps (McIntosh & Lobaugh, 2004).

To assess associations with cognitive performance, disconnectivity within each significant

PLS cluster associated with MoCA was extracted and correlated with CabPad performance.

Results

Figure 2 displays patient distribution of MoCA-scores. 35% of the patients fulfilled criteria for mild

cognitive impairment, based on a suggested cutoff at 26 (Nasreddine et al., 2005).

Figure 2. Histogram depicting the distribution of MoCA scores across the sample.

Investigation of association between group and MoCA performance revealed no significant

relationship (F=1.8, p=0.17, Figure 3). All participants were therefore pooled for further analysis.

Investigation of association between MoCA score and lesion load identified a significant

association between total lesion volume and MoCA score (t=-2.96, p=0.004), but not with number

of lesions (t=-1.502, p=0.13). However, after removal of one participant with an extreme lesion

size, the association between MoCA and lesion volume did not remain significant (t=-1.7, p=0.09).

Figure 3. Distribution of MoCA scores by patient group.

Figure 4 and Table 2 summarize the results from the PLS. Briefly, permutation testing revealed

three significant clusters (pseudo-z > 3) showing common variance between MoCA performance

and structural disconnectivity, including 1) right frontal operculum/insular cortex, 2) right superior

temporal gyrus, and 3) the right putamen. Of note, association between lesion size and individual

PLS brain scores remained highly similar after removal of an individual with an extreme lesion size

(included: t=-9.1, p= 7.6e-15, excluded: t=-7.4 p=5.13e-11), indicating that our main results within

the PLS was not strongly driven by the outlier.

Cluster-#	Location	Size mm^3	Max pseudo-Z	X, Y, Z (vox)
1	Right frontal operculum/insula	1063	4.09	[25, 75, 33]
2	Right superior temporal gyrus, posterior division	273	3.65	[15, 48, 41]
3	Right putamen	37	3.61	[34, 68 32]

249 **Table 2.** Disconnectome clusters significantly associated with MoCA performance, identified by PLS analysis.

Figure 4. Voxel-wise disconnectome bootstrap ratio maps for the MoCA association, thresholded at pseudo-z >3.

253   Figure 5 illustrates correlations between disconnectivity within each of the significant clusters

254   identified by MoCa-PLS and CabPad performance. Disconnectivity was negatively associated with

255   performance on all tests, with strongest correlations for semantic and phonetic fluency.

257
258   **Figure 5**. Correlation between disconnectivity within the significant clusters, and CabPad performance.

Discussion

In the current study, we investigated associations between individual structural disconnectome

maps and post-stroke cognitive performance in 102 stroke survivors. We also performed follow-up

investigation assessing cognitive associations with a broader cognitive battery on a subsample. Our

analysis revealed that structural disconnections implicating the right insula and frontal operculum,

along with right superior temporal gyrus and right putamen were associated with poorer general

cognitive performance as measured with MoCA. In a subsample comprising 82 of the patients, we

demonstrated that the association generalized to a range of cognitive domains, including word

generation, attention span, and speed. Together, our results support the relevance of altered

structural connectivity on cognitive impairments following stroke.

Stroke frequently cause cognitive impairment, and investigations of the underlying network

dynamics may add to our current understanding and prognostics. Structural disconnections caused

by stroke have been shown to influence functional connectivity both directly and indirectly (Griffis,

Metcalf, Corbetta, & Shulman, 2019, 2020). This suggests a mechanism of cognitive sequelae in

which altered functional connectivity due to structural disruptions cause aberrant function in distal

nodes, where lesions perturbating any part of a network supporting cognitive functions may cause

cognitive and behavioral impairment (Alstott, Breakspear, Hagmann, Cammoun, & Sporns, 2009)

(Chen et al., 2019; Lim & Kang, 2015).

A challenge in current healthcare is assessment of cognitive function with a sensitivity and

specificity able to detect and differentiate cognitive difficulties (Dong et al., 2010). MoCA

(Nasreddine et al., 2005) has been suggested as a feasible screening tool for detecting cognitive

impairment following stroke (Horstmann, Rizos T Fau - Rauch, Rauch G Fau - Arden, Arden C Fau

- Veltkamp, & Veltkamp, 2014; Julayanont & Nasreddine, 2017; Munthe-Kaas et al., 2021), where

key advantages are short administration time and multiple cognitive domains of assessment (Burton

& Tyson, 2015; Stolwyk Renerus, O'Neill Megan, McKay Adam, & Wong Dana, 2014).

Importantly, MoCA is sensitive to specific cognitive domains such as attention, executive and

visuospatial function when utilized in lesion-mapping studies (Shi et al., 2018; Zhao et al., 2017).

Of note, post stroke MoCA score has been demonstrated to differentiate cerebral blood flow in

anterior cingulate and prefrontal cortex (Nakaoku et al., 2018). Anterior cingulate is strongly

connected to insula and frontal operculum both structurally (Ghaziri et al., 2017) and functionally

(Horn et al., 2010), and is suggested to be among the core nodes of the cingulo-opercular network.

Importantly, these network nodes are suggested to mediate regulation and differentiation between

the frontoparietal, ventral attention and default mode networks (Goulden et al., 2014; Menon &

Uddin, 2010). The frontoparietal and default mode network are furthermore viewed as

anticorrelated networks reflecting cognitive state, where degree of this differentiation have been

shown to predict cognitive function in stroke survivors at the group (Geranmayeh, Leech, & Wise,

2016) and individual level (Lorenz et al., 2021). Indeed, lesions causing altered connectivity in

cingulo-opercular nodes have been linked to cognitive impairment across multiple domains (Siegel

et al., 2016; Warren et al., 2014), including general impairments as measured using MoCA

(Vicentini et al., 2021).

Our analysis revealed significant associations between MoCA and disconnectivity in the

right superior temporal gyrus (STG) and a small cluster in the putamen. STG is commonly

associated with language production and interpretation (Brugge, Volkov, Garell, Reale, & Howard,

2003), and has also been found to be activated in in conjunction with insula during evaluation of

responses in decision-making tasks (Megías, Cándido, Maldonado, & Catena, 2018; Paulus,

Feinstein, Leland, & Simmons, 2005), where information needs to be integrated and evaluated over

longer periods. STG has also been found to coactivate with visual attention , frontoparietal and

cingulo-opercular nodes during reorientation of visual attention (Vossel et al., 2014), suggesting a

more general role in attentional processing.

Putamen is involved in various aspects of motor functioning and learning, including

language functions and reward signaling. The putamen is connected to thalamic and motor cortices

(Jung et al., 2014; Leh, Ptito, Chakravarty, & Strafella, 2007), and is strongly connected to the

frontoparietal network where the integrity of the connecting pathway is correlated with executive

functioning in healthy adults (Bennett & Madden, 2014; Ystad et al., 2011). These results together

suggests that post-stroke cognitive dysfunction may partly arise from lack of differentiation

between large-scale brain networks. Our results align with this understanding, as structural

disconnectivity of the insula and operculum was associated with impaired MoCA performance,

potentially through disrupted network regulation.

Expanding our results, disconnectivity within the MoCA-associated clusters correlated with

performance across several cognitive domains as measured by CabPAD. While the apparent lack of

cognitive specificity within and across clusters may reflect that the disconnectome approach

captures a more over-arching impairment, strongest correlations were seen with phonetic and

semantic generation and visual attention span. Phonetic and semantic fluency reflect executive

functioning beyond pure language generation (Delis, Kramer, Kaplan, & Holdnack, 2004), and

greater functional connectivity within the cingulo-opercular and ventral attention network has been

associated with reading fluency.

Our results revealed significant associations within the right hemisphere only. A recent

investigation of structural connectivity in patients with ischemic leukoaraiosis found predominantly

right side altered graph metrics when compared to healthy controls, and reported associations with

cognitive performance (Lu et al., 2021). Comparably, right side functional abnormalities in cingulo-

opercular nodes during resting state have been associated with impaired MoCA performance

following a transient ischemic attack (Guo et al., 2014), and right side dominance in bilateral insula

activation is robustly found during task engagement in healthy controls (Rottschy et al., 2012).

Further, right hemisphere is to a larger degree linked to visual attention, reorientation, and bottom-

up processing, and damage to right side pathways associated with impaired spatial attention, target

detection and vigilance (Maurizio Corbetta & Gordon L. Shulman, 2011). Albeit speculative, our

results indicate a larger right-side vulnerability for abrupted structural connectivity augmenting post

stroke cognitive difficulties.

The current study has limitations. First, our cross-sectional design does not allow for

integrating premorbid cognitive function and chronic brain pathology, which is highly relevant as

an outcome predictor (Sagnier & Sibon, 2019). Further, the current sample represents a

heterogeneous group regarding stroke severity, we lack measures on vascular risk and the sample

size is moderate, factors that are relevant for the generalizability of the reported findings to the

stroke population in general (Marek et al., 2020). Further, jointly estimating all voxels in a single

PLS model increases sensitivity, however, comes with a trade-off in spatial specificity as we can

only make statements about significance for the spatial pattern as a whole. Still, the reliability of

each voxel's contribution to the observed pattern was assessed through bootstrapping

In conclusion, our study supports the relevance of investigating disrupted structural

connectivity following stroke. Although our study does not allow differentiation between proximal

and distal effects due to sample size and lesion homogeneity, our results highlight the relevance of

altered structural connectivity when investigating cognitive sequalae after stroke. In line with

previous studies, our results indicate lesions affecting the insula and the frontal operculum are

associated with cognitive impairment and support the inclusion of measures of structural

disconnection when evaluating cognitive and functional sequelae in stroke patients.

References

Aben, H. P., Biessels, G. J., Weaver, N. A., Spikman, J. M., Visser-Meily, J. M. A., de Kort, P. L. M., . .
. Jansen, B. P. W. (2019). Extent to Which Network Hubs Are Affected by Ischemic
Stroke Predicts Cognitive Recovery. *Stroke, 50*(10), 2768-2774.
doi:10.1161/STROKEAHA.119.025637
Alstott, J., Breakspear, M., Hagmann, P., Cammoun, L., & Sporns, O. (2009). Modeling the
Impact of Lesions in the Human Brain. *PLoS computational biology, 5*(6), e1000408.
doi:10.1371/journal.pcbi.1000408
Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A reproducible
evaluation of ANTs similarity metric performance in brain image registration.
NeuroImage, 54(3), 2033-2044.
doi:https://doi.org/10.1016/j.neuroimage.2010.09.025
Bennett, I. J., & Madden, D. J. (2014). Disconnected aging: Cerebral white matter integrity and
age-related differences in cognition. *Neuroscience, 276*, 187-205.
doi:https://doi.org/10.1016/j.neuroscience.2013.11.026
Bentley, P., Kumar, G., Rinne, P., Buddha, S., Kallingal, J., Hookway, C., . . . Beckmann, C. (2014).
Lesion locations influencing baseline severity and early recovery in ischaemic stroke.
European Journal of Neurology, 21(9), 1226-1232. doi:10.1111/ene.12464
Blackburn, D. J., Bafadhel, L., Randall, M., & Harkness, K. A. (2013). Cognitive screening in the
acute stroke setting. *Age and ageing, 42*(1), 113-116. doi:10.1093/ageing/afs116
Brugge, J. F., Volkov, I. O., Garell, P. C., Reale, R. A., & Howard, M. A. (2003). Functional
Connections Between Auditory Cortex on Heschl's Gyrus and on the Lateral Superior
Temporal Gyrus in Humans. *Journal of Neurophysiology, 90*(6), 3750-3763.
doi:10.1152/jn.00500.2003
Burton, L., & Tyson, S. F. (2015). Screening for cognitive impairment after stroke: A systematic
review of psychometric properties and clinical utility. *J Rehabil Med, 47*(3), 193-203.
doi:10.2340/16501977-1930
Cai, W., Chen, T., Ryali, S., Kochalka, J., Li, C.-S. R., & Menon, V. (2016). Causal Interactions
Within a Frontal-Cingulate-Parietal Network During Cognitive Control: Convergent
Evidence from a Multisite–Multitask Investigation. *Cerebral cortex, 26*(5), 2140-2153.
doi:10.1093/cercor/bhv046
Chen, H., Shi, M., Zhang, H., Zhang, Y.-D., Geng, W., Jiang, L., . . . Yin, X. (2019). Different Patterns
of Functional Connectivity Alterations Within the Default-Mode Network and
Sensorimotor Network in Basal Ganglia and Pontine Stroke. *Medical science monitor :
international medical journal of experimental and clinical research, 25*, 9585-9593.
doi:10.12659/MSM.918185
Cole, M. W., Reynolds, J. R., Power, J. D., Repovs, G., Anticevic, A., & Braver, T. S. (2013). Multi-
task connectivity reveals flexible hubs for adaptive task control. *Nature Neuroscience,
16*(9), 1348-1355. doi:10.1038/nn.3470
Corbetta, M., & Shulman, G. L. (2011). Spatial neglect and attention networks. *Annual Review of
Neuroscience, 34*, 569-599.

Corbetta, M., & Shulman, G. L. (2011). Spatial Neglect and Attention Networks. *Annual Review of Neuroscience, 34*(1), 569-599. doi:10.1146/annurev-neuro-061010-113731

Crossley, N. A., Mechelli, A., Scott, J., Carletti, F., Fox, P. T., McGuire, P., & Bullmore, E. T. (2014). The hubs of the human connectome are generally implicated in the anatomy of brain disorders. *Brain, 137*(8), 2382-2395. doi:10.1093/brain/awu132

de Haan, B., Clas, P., Juenger, H., Wilke, M., & Karnath, H.-O. (2015). Fast semi-automated lesion demarcation in stroke. *NeuroImage. Clinical, 9*, 69-74. doi:10.1016/j.nicl.2015.06.013

Delis, D. C., Kramer, J. H., Kaplan, E., & Holdnack, J. (2004). Reliability and validity of the Delis-Kaplan Executive Function System: An update. *Journal of the International Neuropsychological Society, 10*(2), 301-303. doi:10.1017/S1355617704102191

Dong, Y., Sharma, V. K., Chan, B. P.-L., Venketasubramanian, N., Teoh, H. L., Seet, R. C. S., . . . Chen, C. (2010). The Montreal Cognitive Assessment (MoCA) is superior to the Mini-Mental State Examination (MMSE) for the detection of vascular cognitive impairment after acute stroke. *Journal of the Neurological Sciences, 299*(1), 15-18. doi:https://doi.org/10.1016/j.jns.2010.08.051

Dørum, E. S., Kaufmann, T., Alnæs, D., Richard, G., Kolskår, K. K., Engvig, A., . . . Westlye, L. T. (2020). Functional brain network modeling in sub-acute stroke patients and healthy controls during rest and continuous attentive tracking. *Heliyon, 6*(9), e04854. doi:https://doi.org/10.1016/j.heliyon.2020.e04854

Fassbender, C., Simoes-Franklin, C., Murphy, K., Hester, R., Meaney, J., Robertson, I. H., & Garavan, H. (2006). The Role of a Right Fronto-Parietal Network in Cognitive Control. *Journal of Psychophysiology, 20*(4), 286-296. doi:10.1027/0269-8803.20.4.286

Foulon, C., Cerliani, L., Kinkingnéhun, S., Levy, R., Rosso, C., Urbanski, M., . . . Thiebaut de Schotten, M. (2018). Advanced lesion symptom mapping analyses and implementation as BCBtoolkit. *GigaScience, 7*(3). doi:10.1093/gigascience/giy004

Fox, M. D. (2018). Mapping Symptoms to Brain Networks with the Human Connectome. *New England Journal of Medicine, 379*(23), 2237-2245. doi:10.1056/NEJMra1706158

Geranmayeh, F., Leech, R., & Wise, R. J. S. (2016). Network dysfunction predicts speech production after left hemisphere stroke. *Neurology, 86*(14), 1296. doi:10.1212/WNL.0000000000002537

Ghaziri, J., Tucholka, A., Girard, G., Houde, J.-C., Boucher, O., Gilbert, G., . . . Nguyen, D. K. (2017). The Corticocortical Structural Connectivity of the Human Insula. *Cerebral cortex, 27*(2), 1216-1228. doi:10.1093/cercor/bhv308

Goulden, N., Khusnulina, A., Davis, N. J., Bracewell, R. M., Bokde, A. L., McNulty, J. P., & Mullins, P. G. (2014). The salience network is responsible for switching between the default mode network and the central executive network: Replication from DCM. *NeuroImage, 99*, 180-190. doi:https://doi.org/10.1016/j.neuroimage.2014.05.052

Griffis, J. C., Metcalf, N. V., Corbetta, M., & Shulman, G. L. (2019). Structural Disconnections Explain Brain Network Dysfunction after Stroke. *Cell Reports, 28*(10), 2527-2540.e2529. doi:https://doi.org/10.1016/j.celrep.2019.07.100

Griffis, J. C., Metcalf, N. V., Corbetta, M., & Shulman, G. L. (2020). Damage to the shortest structural paths between brain regions is associated with disruptions of resting-state functional connectivity after stroke. *NeuroImage, 210*, 116589. doi:https://doi.org/10.1016/j.neuroimage.2020.116589

Guo, J., Chen, N., Li, R., Wu, Q., Chen, H., Gong, Q., & He, L. (2014). Regional homogeneity abnormalities in patients with transient ischaemic attack: A resting-state fMRI study. *Clinical Neurophysiology, 125*(3), 520-525. doi:https://doi.org/10.1016/j.clinph.2013.08.010

Hilger, K., Ekman, M., Fiebach, C. J., & Basten, U. (2017). Efficient hubs in the intelligent brain:
Nodal efficiency of hub regions in the salience network is associated with general
intelligence. *Intelligence, 60*, 10-25. doi:https://doi.org/10.1016/j.intell.2016.11.001
Horn, D. I., Yu, C., Steiner, J., Buchmann, J., Kaufmann, J., Osoba, A., . . . He, H. (2010).
Glutamatergic and resting-state functional connectivity correlates of severity in major
depression–the role of pregenual anterior cingulate cortex and anterior insula.
Frontiers in systems neuroscience, 4, 33.
Horstmann, S., Rizos T Fau - Rauch, G., Rauch G Fau - Arden, C., Arden C Fau - Veltkamp, R., &
Veltkamp, R. (2014). Feasibility of the Montreal Cognitive Assessment in acute stroke
patients. *Eur J Neurol, 21(11)*(1468-1331 (Electronic)). doi:10.1111/ene.12505
Jenkinson, M., & Smith, S. (2001). A global optimisation method for robust affine registration
of brain images. *Medical Image Analysis, 5*(2), 143-156.
doi:https://doi.org/10.1016/S1361-8415(01)00036-6
Julayanont, P., & Nasreddine, Z. S. (2017). Montreal Cognitive Assessment (MoCA): Concept
and Clinical Review. In A. J. Larner (Ed.), *Cognitive Screening Instruments: A Practical
Approach* (pp. 139-195). Cham: Springer International Publishing.
Jung, W. H., Jang, J. H., Park, J. W., Kim, E., Goo, E.-H., Im, O.-S., & Kwon, J. S. (2014). Unravelling
the Intrinsic Functional Organization of the Human Striatum: A Parcellation and
Connectivity Study Based on Resting-State fMRI. *PloS one, 9*(9), e106768.
doi:10.1371/journal.pone.0106768
Klein, A., Andersson, J., Ardekani, B. A., Ashburner, J., Avants, B., Chiang, M.-C., . . . Parsey, R. V.
(2009). Evaluation of 14 nonlinear deformation algorithms applied to human brain
MRI registration. *NeuroImage, 46*(3), 786-802.
doi:https://doi.org/10.1016/j.neuroimage.2008.12.037
Klingbeil, J., Wawrzyniak, M., Stockert, A., & Saur, D. (2019). Resting-state functional
connectivity: An emerging method for the study of language networks in post-stroke
aphasia. *Brain and Cognition, 131*, 22-33.
doi:https://doi.org/10.1016/j.bandc.2017.08.005
Kolskår, K. K., Richard, G., Alnæs, D., Dørum, E. S., Sanders, A.-M., Ulrichsen, K. M., . . . Westlye,
L. T. (2020). Reliability, sensitivity, and predictive value of fMRI during multiple object
tracking as a marker of cognitive training gain in combination with tDCS in stroke
survivors. *Human Brain Mapping, n/a*(n/a). doi:https://doi.org/10.1002/hbm.25284
Krishnan, A., Williams Lj Fau - McIntosh, A. R., McIntosh Ar Fau - Abdi, H., & Abdi, H. (2010).
Partial Least Squares (PLS) methods for neuroimaging: a tutorial and review.
NeuroImage, 15;56(2)(1095-9572 (Electronic)), 455-475.
doi:10.1016/j.neuroimage.2010.07.034.
Krishnan, A., Williams, L. J., McIntosh, A. R., & Abdi, H. (2011). Partial Least Squares (PLS)
methods for neuroimaging: a tutorial and review. *Neuroimage, 56*(2), 455-475.
doi:10.1016/j.neuroimage.2010.07.034
Ktena Sofia, I., Schirmer Markus, D., Etherton Mark, R., Giese, A.-K., Tuozzo, C., Mills Brittany,
B., . . . Rost Natalia, S. (2019). Brain Connectivity Measures Improve Modeling of
Functional Outcome After Acute Ischemic Stroke. *Stroke, 50*(10), 2761-2767.
doi:10.1161/STROKEAHA.119.025738
Leh, S. E., Ptito, A., Chakravarty, M. M., & Strafella, A. P. (2007). Fronto-striatal connections in
the human brain: A probabilistic diffusion tractography study. *Neuroscience Letters,
419*(2), 113-118. doi:https://doi.org/10.1016/j.neulet.2007.04.049
Lim, J.-S., & Kang, D.-W. (2015). Stroke Connectome and Its Implications for Cognitive and
Behavioral Sequela of Stroke. *Journal of stroke, 17*(3), 256-267.
doi:10.5853/jos.2015.17.3.256
Lopes, R., Bournonville, C., Kuchcinski, G., Dondaine, T., Mendyk, A.-M., Viard, R., . . . Bordet, R.
(2021a). Prediction of Long-term Cognitive Function After Minor Stroke Using

Functional Connectivity. *Neurology, 96*(8), e1167.
doi:10.1212/WNL.0000000000011452

Lopes, R., Bournonville, C., Kuchcinski, G., Dondaine, T., Mendyk, A.-M., Viard, R., . . . Bordet, R.
(2021b). Prediction of Long-term Cognitive Functions after Minor Stroke, Using
Functional Connectivity. *Neurology*, 10.1212/WNL.0000000000011452.
doi:10.1212/WNL.0000000000011452

Lorenz, R., Johal, M., Dick, F., Hampshire, A., Leech, R., & Geranmayeh, F. (2021). A Bayesian
optimization approach for rapidly mapping residual network function in stroke. *Brain,
144*(7), 2120-2134. doi:10.1093/brain/awab109

Lu, T., Wang, Z., Cui, Y., Zhou, J., Wang, Y., & Ju, S. (2021). Disrupted Structural Brain
Connectome Is Related to Cognitive Impairment in Patients With Ischemic
Leukoaraiosis. *Frontiers in Human Neuroscience, 15*. doi:10.3389/fnhum.2021.654750

Macciocchi, S. N., Diamond, P. T., Alves, W. M., & Mertz, T. (1998). Ischemic stroke: Relation of
age, lesion location, and initial neurologic deficit to functional outcome. *Archives of
Physical Medicine and Rehabilitation, 79*(10), 1255-1257.
doi:https://doi.org/10.1016/S0003-9993(98)90271-4

Majerus, S., Péters, F., Bouffier, M., Cowan, N., & Phillips, C. (2018). The Dorsal Attention
Network Reflects Both Encoding Load and Top–down Control during Working
Memory. *Journal of Cognitive Neuroscience, 30*(2), 144-159. doi:10.1162/jocn_a_01195

Marek, S., Tervo-Clemmens, B., Calabro, F. J., Montez, D. F., Kay, B. P., Hatoum, A. S., . . .
Dosenbach, N. U. F. (2020). Towards Reproducible Brain-Wide Association Studies.
bioRxiv, 2020.2008.2021.257758. doi:10.1101/2020.08.21.257758

MathWorks, I. (2018). *MATLAB : the language of technical computing : computation,
visualization, programming*: Natwick : Math Works Inc., 2018.

McIntosh, A. R., & Lobaugh, N. J. (2004). Partial least squares analysis of neuroimaging data:
applications and advances. *NeuroImage, 23 Suppl 1*, S250-263.
doi:10.1016/j.neuroimage.2004.07.020

McKiernan, K. A., Kaufman, J. N., Kucera-Thompson, J., & Binder, J. R. (2003). A Parametric
Manipulation of Factors Affecting Task-induced Deactivation in Functional
Neuroimaging. *Journal of Cognitive Neuroscience, 15*(3), 394-408.
doi:10.1162/089892903321593117

Megías, A., Cándido, A., Maldonado, A., & Catena, A. (2018). Neural correlates of risk
perception as a function of risk level: An approach to the study of risk through a daily
life task. *Neuropsychologia, 119*, 464-473.
doi:https://doi.org/10.1016/j.neuropsychologia.2018.09.012

Menon, V., & Uddin, L. Q. (2010). Saliency, switching, attention and control: a network model
of insula function. *Brain Structure and Function, 214*(5), 655-667. doi:10.1007/s00429-
010-0262-0

Munsch, F., Sagnier, S., Asselineau, J., Bigourdan, A., Guttmann Charles, R., Debruxelles, S., . . .
Tourdias, T. (2016). Stroke Location Is an Independent Predictor of Cognitive
Outcome. *Stroke, 47*(1), 66-73. doi:10.1161/STROKEAHA.115.011242

Munthe-Kaas, R., Aam, S., Saltvedt, I., Wyller Torgeir, B., Pendlebury Sarah, T., Lydersen, S., &
Ihle-Hansen, H. (2021). Test Accuracy of the Montreal Cognitive Assessment in
Screening for Early Poststroke Neurocognitive Disorder. *Stroke, 52*(1), 317-320.
doi:10.1161/STROKEAHA.120.031030

Nakaoku, Y., Oishi, N., Hase, Y., Hase, M., Saito, S., Mitsueda, T., . . . Takahashi, R. (2018).
Montreal Cognitive Assessment score correlates with regional cerebral blood flow in
post-stroke patients. *Clinical Neurology and Neurosurgery, 174*, 68-74.
doi:https://doi.org/10.1016/j.clineuro.2018.09.004

Nasreddine, Z. S., Phillips, N. A., Bédirian, V., Charbonneau, S., Whitehead, V., Collin, I., . . .
Chertkow, H. (2005). The Montreal Cognitive Assessment, MoCA: A Brief Screening

Tool For Mild Cognitive Impairment. *Journal of the American Geriatrics Society, 53*(4), 695-699. doi:10.1111/j.1532-5415.2005.53221.x

Paulus, M. P., Feinstein, J. S., Leland, D., & Simmons, A. N. (2005). Superior temporal gyrus and insula provide response and outcome-dependent information during assessment and action selection in a decision-making situation. *NeuroImage, 25*(2), 607-615. doi:https://doi.org/10.1016/j.neuroimage.2004.12.055

Pendlebury, S. T. (2009). Stroke-related dementia: Rates, risk factors and implications for future research. *Maturitas, 64*(3), 165-171. doi:https://doi.org/10.1016/j.maturitas.2009.09.010

Ptak, R., Bourgeois, A., Cavelti, S., Doganci, N., Schnider, A., & Iannotti, G. R. (2020). Discrete Patterns of Cross-Hemispheric Functional Connectivity Underlie Impairments of Spatial Cognition after Stroke. *The Journal of Neuroscience, 40*(34), 6638. doi:10.1523/JNEUROSCI.0625-20.2020

Rehme, A. K., & Grefkes, C. (2013). Cerebral network disorders after stroke: evidence from imaging-based connectivity analyses of active and resting brain states in humans. *The Journal of Physiology, 591*(1), 17-31. doi:10.1113/jphysiol.2012.243469

Richard, G., Kolskår, K. K., Ulrichsen, K. M., Kaufmann, T., Alnæs, D., Sanders, A.-M., . . . Westlye, L. T. (2020). Brain age prediction in stroke patients: Highly reliable but limited sensitivity to cognitive performance and response to cognitive training. *NeuroImage: Clinical, 25*, 102159. doi:https://doi.org/10.1016/j.nicl.2019.102159

Rise, H. H., Brune, S., Chien, C., Berge, T., Bos, S. D., Andorra, M., . . . Høgestøl, E. A. (2021). Brain disconnectome mapping and serum neurofilament light levels in multiple sclerosis. *medRxiv*, 2021.2004.2021.21255887. doi:10.1101/2021.04.21.21255887

Rottschy, C., Langner, R., Dogan, I., Reetz, K., Laird, A. R., Schulz, J. B., . . . Eickhoff, S. B. (2012). Modelling neural correlates of working memory: A coordinate-based meta-analysis. *NeuroImage, 60*(1), 830-846. doi:https://doi.org/10.1016/j.neuroimage.2011.11.050

Sagnier, S., & Sibon, I. (2019). The new insights into human brain imaging after stroke. *Journal of Neuroscience Research, n/a*(n/a). doi:https://doi.org/10.1002/jnr.24525

Salvalaggio, A., De Filippo De Grazia, M., Zorzi, M., Thiebaut de Schotten, M., & Corbetta, M. (2020). Post-stroke deficit prediction from lesion and indirect structural and functional disconnection. *Brain, 143*(7), 2173-2188. doi:10.1093/brain/awaa156

Shi, L., Zhao, L., Yeung, F. K., Wong, S. Y., Chan, R. K. T., Tse, M. F., . . . Mok, V. C. T. (2018). Mapping the contribution and strategic distribution patterns of neuroimaging features of small vessel disease in poststroke cognitive impairment. *Journal of Neurology, Neurosurgery & Psychiatry, 89*(9), 918. doi:10.1136/jnnp-2017-317817

Siegel, J. S., Ramsey, L. E., Snyder, A. Z., Metcalf, N. V., Chacko, R. V., Weinberger, K., . . . Corbetta, M. (2016). Disruptions of network connectivity predict impairment in multiple behavioral domains after stroke. *Proceedings of the National Academy of Sciences, 113*(30), E4367. doi:10.1073/pnas.1521083113

Stam, C. J. (2014). Modern network science of neurological disorders. *Nature Reviews Neuroscience, 15*, 683. doi:10.1038/nrn3801

Stolwyk Renerus, J., O'Neill Megan, H., McKay Adam, J. D., & Wong Dana, K. (2014). Are Cognitive Screening Tools Sensitive and Specific Enough for Use After Stroke? *Stroke, 45*(10), 3129-3134. doi:10.1161/STROKEAHA.114.004232

Thiebaut de Schotten, M., Foulon, C., & Nachev, P. (2020). Brain disconnections link structural connectivity with function and behaviour. *Nature Communications, 11*(1), 5094. doi:10.1038/s41467-020-18920-9

Uddin, L. Q., Yeo, B. T. T., & Spreng, R. N. (2019). Towards a Universal Taxonomy of Macro-scale Functional Human Brain Networks. *Brain topography, 32*(6), 926-942. doi:10.1007/s10548-019-00744-6

Ulrichsen, K. M., Kolskår, K. K., Richard, G., Alnæs, D., Dørum, E. S., Sanders, A.-M., . . . Westlye,
L. T. (2020). Brain disconnectivity mapping of post-stroke fatigue. *bioRxiv*,
2020.2011.2013.380972. doi:10.1101/2020.11.13.380972
van den Heuvel, M. P., & Sporns, O. (2019). A cross-disorder connectome landscape of brain
dysconnectivity. *Nature Reviews Neuroscience, 20*(7), 435-446. doi:10.1038/s41583-
019-0177-6
Van Roon, P., Zakizadeh, J., & Chartier, S. (2014). Partial least squares tutorial for analyzing
neuroimaging data. *Tutorials in Quantitative Methods for Psychology, 10*, 200-215.
Vatansever, D., Manktelow, A., Sahakian, B. J., Menon, D. K., & Stamatakis, E. A. (2018). Default
Mode Network Engagement Beyond Self-Referential Internal Mentation. *Brain*
Connectivity, 8(4), 245-253. doi:10.1089/brain.2017.0489
Vicentini, J. E., Weiler, M., Casseb, R. F., Almeida, S. R., Valler, L., de Campos, B. M., & Li, L. M.
(2021). Subacute functional connectivity correlates with cognitive recovery six months
after stroke. *NeuroImage. Clinical, 29*, 102538-102538. doi:10.1016/j.nicl.2020.102538
Vossel, S., Geng, J. J., & Fink, G. R. (2014). Dorsal and ventral attention systems: distinct neural
circuits but collaborative roles. *The Neuroscientist : a review journal bringing*
neurobiology, neurology and psychiatry, 20(2), 150-159.
doi:10.1177/1073858413494269
Wang, R., Benner, T., Sorensen, A. G., & Wedeen, V. J. (2007). Diffusion Toolkit: A Software
Package for Diffusion Imaging Data Processing and Tractography *Proceedings of the*
International Society for Magnetic Resonance in Medicine, 15.
Warren, D. E., Power, J. D., Bruss, J., Denburg, N. L., Waldron, E. J., Sun, H., . . . Tranel, D. (2014).
Network measures predict neuropsychological outcome after brain injury. *Proceedings*
of the National Academy of Sciences, 111(39), 14247. doi:10.1073/pnas.1322173111
Wilk, H. A., Ezekiel, F., & Morton, J. B. (2012). Brain regions associated with moment-to-
moment adjustments in control and stable task-set maintenance. *NeuroImage, 59*(2),
1960-1967. doi:https://doi.org/10.1016/j.neuroimage.2011.09.011
Willer, L., Pedersen, P. M., Forchhammer, H. B., & Christensen, H. (2016). Cognitive assessment
at bedside for iPad: A preliminary validation of a novel cognitive test for stroke
patients. *European Stroke Journal, 1*(4), 294-301. doi:10.1177/2396987316665233
Ystad, M., Hodneland, E., Adolfsdottir, S., Haász, J., Lundervold, A. J., Eichele, T., & Lundervold,
A. (2011). Cortico-striatal connectivity and cognition in normal aging: A combined DTI
and resting state fMRI study. *NeuroImage, 55*(1), 24-31.
doi:https://doi.org/10.1016/j.neuroimage.2010.11.016
Zhao, L., Biesbroek, J. M., Shi, L., Liu, W., Kuijf, H. J., Chu, W. W. C., . . . Wong, A. (2017). Strategic
infarct location for post-stroke cognitive impairment: A multivariate lesion-symptom
mapping study. *Journal of Cerebral Blood Flow & Metabolism, 38*(8), 1299-1311.
doi:10.1177/0271678X17728162
635